

A Service of



Leibniz-Informationszentrum Wirtschaft Leibniz Information Centre

Biewen, Martin; Jenkins, Stephen P.

Working Paper

Estimation of Generalized Entropy and Atkinson Inequality Indices from Complex Survey Data

IZA Discussion Papers, No. 763

Provided in Cooperation with:

IZA - Institute of Labor Economics

Suggested Citation: Biewen, Martin; Jenkins, Stephen P. (2003): Estimation of Generalized Entropy and Atkinson Inequality Indices from Complex Survey Data, IZA Discussion Papers, No. 763, Institute for the Study of Labor (IZA), Bonn

This Version is available at: https://hdl.handle.net/10419/21414

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.





IZA DP No. 763

Estimation of Generalized Entropy and Atkinson Inequality Indices from Complex Survey Data

Martin Biewen Stephen P. Jenkins

April 2003

Forschungsinstitut zur Zukunft der Arbeit Institute for the Study of Labor

Estimation of Generalized Entropy and Atkinson Inequality Indices from Complex Survey Data

Martin Biewen

University of Mannheim, DIW Berlin and IZA Bonn

Stephen P. Jenkins

ISER, University of Essex, DIW Berlin and IZA Bonn

Discussion Paper No. 763 April 2003

IZA

P.O. Box 7240 D-53072 Bonn Germany

Tel.: +49-228-3894-0 Fax: +49-228-3894-210 Email: iza@iza.org

This Discussion Paper is issued within the framework of IZA's research area *The Future of Labor*. Any opinions expressed here are those of the author(s) and not those of the institute. Research disseminated by IZA may include views on policy, but the institute itself takes no institutional policy positions.

The Institute for the Study of Labor (IZA) in Bonn is a local and virtual international research center and a place of communication between science, politics and business. IZA is an independent, nonprofit limited liability company (Gesellschaft mit beschränkter Haftung) supported by the Deutsche Post AG. The center is associated with the University of Bonn and offers a stimulating research environment through its research networks, research support, and visitors and doctoral programs. IZA engages in (i) original and internationally competitive research in all fields of labor economics, (ii) development of policy concepts, and (iii) dissemination of research results and concepts to the interested public. The current research program deals with (1) mobility and flexibility of labor, (2) internationalization of labor markets, (3) welfare state and labor market, (4) labor markets in transition countries, (5) the future of labor, (6) evaluation of labor market policies and projects and (7) general labor economics.

IZA Discussion Papers often represent preliminary work and are circulated to encourage discussion. Citation of such a paper should account for its provisional character. A revised version may be available on the IZA website (www.iza.org) or directly from the author.

ABSTRACT

Estimation of Generalized Entropy and Atkinson Inequality Indices from Complex Survey Data*

Applying a method suggested by Woodruff (1971), we derive the sampling variances of Generalized Entropy and Atkinson inequality indices when estimated from complex survey data. It turns out that this method also greatly simplifies the calculations for the i.i.d. case when compared to previous derivations in the literature. Both cases are illustrated with examples from the German Socio-Economic Panel Study and the British Household Panel Survey.

JEL Classification: C14, D31

Keywords: inequality, statistical inference, complex surveys

Corresponding author:

Martin Biewen Department of Economics University of Mannheim Verfügungsgebäude L7, 3-5 68131 Mannheim Germany

Fax: +49 621 1811 841

Email: biewen@rhein.vwl.uni-mannheim.de

* Financial support from the Deutsche Forschungsgesellschaft (DFG) and core funding to ISER from the ESRC and the University of Essex is gratefully acknowledged.

1 Introduction

Probability weighting, clustering, and stratification, are survey design features underlying much of the survey data that economists and others use. It is well known that these features have a potentially large impact on the sampling variability of statistics computed from such surveys. Nevertheless, they are rarely taken into account in practical work, the measurement of inequality being no exception. We derive estimates for the sampling variance of two commonly-used classes of inequality indices, the Generalized Entropy and the Atkinson family of indices, using the approach of Woodruff (1971).² It turns out that Woodruff's method also greatly simplifies the computation of variance estimates in an i.i.d. framework when compared to previous derivations in the literature. In order to assess the error made by not taking into account clustering and stratification we apply our results to a sample extracted from the German Socio-Economic Panel (GSOEP) and the British Household Panel Survey (BHPS).

2 Estimation from complex surveys

Generalized Entropy and Atkinson inequality measures can be written, as we show below, as functions I = f(T) of population totals $T = (T_1, \ldots, T_K)$. Population total T_k , $k = 1, \ldots, K$ is given by the summation of an observational variable t_{hijk} over the different stages of the sampling design, i.e. $T_k = \sum_{n=1}^L \sum_{i=1}^{N_h} \sum_{j=1}^{M_i} t_{hijk}$ where L denotes the number of strata, N_h the number of clusters in stratum h and M_i the number of individuals in cluster i. If the sampling design involves more than one stage of clustering it suffices to consider the first stage only (see e.g. Cochran, 1977). Replacing totals T by their estimates \hat{T} , the index is then estimated as $\hat{I} = f(\hat{T})$ with $\hat{T}_k = \sum_{n=1}^L \sum_{i=1}^{n_h} \sum_{j=1}^{m_i} w_{hij} t_{hijk}$ where n_h is the number of actually sampled first stage clusters and m_i the number of actually sampled individuals in cluster i. The sampling weight of individual hij is given by w_{hij} . Assuming that the sample is large enough that a Taylor approximation of $f(\cdot)$ holds, the variance of \hat{I} can be approximated by the variance of the first order residual $\sum_{k=1}^K (\partial f(T)/\partial T_k) \hat{T}_k$. Woodruff (1971) observed that this variance can be easily determined by reversing the order of summation in the residual, i.e. $\text{var}(\hat{I}) \approx \text{var}(\sum_{n=1}^L \sum_{i=1}^{n_h} \sum_{j=1}^{m_i} w_{hij} \left[\sum_{k=1}^K (\partial f(T)/\partial T_k) t_{hijk}\right]) = \text{var}(\hat{S})$. Note that \hat{S} is of the

²An alternative but conceptually less straightforward approach to variance estimation in complex surveys is the estimating equations approach described in Binder (1983) and Binder and Patak (1994). It turns out that this approach leads to the same estimators derived here. Calculations are available from the authors on request.

same form as the \hat{T}_k s so that the problem is reduced to the estimation of the sampling variance of a total estimator for which well-known formulas exist (see Cochran, 1977, or Deaton, 1997). Using these formulas (and replacing T by \hat{T} in the derivative), the variance estimate for \hat{I} is

$$\widehat{\text{var}}(\widehat{I}) = \sum_{n=1}^{L} \frac{n_h}{n_h - 1} \sum_{i=1}^{n_h} \left(\sum_{j=1}^{m_i} w_{hij} \widetilde{s}_{hij} - \frac{\sum_{i=1}^{n_h} \sum_{j=1}^{m_i} w_{hij} \widetilde{s}_{hij}}{n_h} \right)^2$$
(1)

with $\tilde{s}_{hij} = \sum_{k=1}^{K} (\partial f(\hat{T})/\partial \hat{T}_k) t_{hijk}$.

If y_{hij} represents the income of individual hij, then population Generalized Entropy and Atkinson indices are given by

$$I_{GE}^{\alpha} = \left(\alpha^2 - \alpha\right)^{-1} \left[U_0^{\alpha - 1} U_1^{-\alpha} U_{\alpha} - 1 \right], \quad \alpha \in \mathbb{R} \setminus \{0, 1\}$$
 (2)

$$I_{Theil} = T_{1,1}U_1^{-1} - \log(U_1U_0^{-1}), \quad \alpha \to 1$$
 (3)

$$I_{MLD} = -T_{0,1}U_0^{-1} + \log(U_1U_0^{-1}), \quad \alpha \to 0$$
 (4)

$$I_A^{\varepsilon} = 1 - U_0^{-\varepsilon/(1-\varepsilon)} U_1^{-1} U_{1-\varepsilon}^{1/(1-\varepsilon)}, \quad \varepsilon \ge 0, \quad \varepsilon \ne 1,$$
 (5)

$$I_A^1 = 1 - U_0 U_1^{-1} \exp(T_{0,1} U_0^{-1}), \quad \varepsilon \to 1$$
 (6)

for totals $U_{\alpha} = \sum_{n=1}^{L} \sum_{i=1}^{N_h} \sum_{j=1}^{M_i} (y_{hij})^{\alpha}$ and $T_{\alpha} = \sum_{n=1}^{L} \sum_{i=1}^{N_h} \sum_{j=1}^{M_i} (y_{hij})^{\alpha} (\log y_{hij})$. Note that U_0 is the population size. Estimates of these indices are obtained by replacing U_{α} , T_{α} with $\hat{U}_{\alpha} = \sum_{n=1}^{L} \sum_{i=1}^{n_h} \sum_{j=1}^{m_i} w_{hij} (y_{hij})^{\alpha}$ and $\hat{T}_{\alpha} = \sum_{n=1}^{L} \sum_{i=1}^{n_h} \sum_{j=1}^{m_i} w_{hij} (y_{hij})^{\alpha} (\log y_{hij})$.

The corresponding variances are then given by substituting \tilde{s}_{hij} in (1) with

$$\tilde{s}_{hij}^{GE} = \frac{1}{\alpha} \hat{U}_{\alpha} \hat{U}_{1}^{-\alpha} \hat{U}_{0}^{\alpha-2} - \frac{1}{\alpha - 1} \hat{U}_{\alpha} \hat{U}_{1}^{-\alpha - 1} \hat{U}_{0}^{\alpha - 1} \cdot y_{hij} + \frac{1}{\alpha^{2} - \alpha} \hat{U}_{0}^{\alpha - 1} \hat{U}_{1}^{-\alpha} \cdot (y_{hij})^{\alpha}$$
(7)

$$\tilde{s}_{hij}^{Theil} = \hat{U}_{1}^{-1} \cdot y_{hij} \log y_{hij} - \hat{U}_{1}^{-1} \left(\hat{T}_{1,1} \hat{U}_{1}^{-1} + 1 \right) \cdot y_{hij} + \hat{U}_{0}^{-1}$$
(8)

$$\tilde{s}_{hij}^{MLD} = -\hat{U}_0^{-1} \cdot \log y_{hij} + \hat{U}_1^{-1} \cdot y_{hij} + U_0^{-1} \left(\hat{T}_{0,1} \hat{U}_0^{-1} - 1 \right)$$

$$\tag{9}$$

$$\tilde{s}_{hij}^{A} = \frac{\varepsilon}{1-\varepsilon} \hat{U}_{1}^{-1} \hat{U}_{1-\varepsilon}^{\frac{1}{1-\varepsilon}} \hat{U}_{0}^{-\frac{1}{1-\varepsilon}} + \hat{U}_{0}^{\frac{-\varepsilon}{1-\varepsilon}} \hat{U}_{1-\varepsilon}^{\frac{1}{1-\varepsilon}} \hat{U}_{1}^{-2} \cdot y_{hij} - \frac{1}{1-\varepsilon} \hat{U}_{0}^{\frac{-\varepsilon}{1-\varepsilon}} \hat{U}_{1}^{-1} \hat{U}_{1-\varepsilon}^{\frac{\varepsilon}{1-\varepsilon}} \cdot (y_{hij})^{1-\varepsilon} (10)$$

$$\tilde{s}_{hij}^{1} = \left(\hat{I}_{A}^{1} - 1\right)\hat{U}_{0}^{-1}\left(1 - \hat{U}_{0}^{-1}\hat{T}_{0,1}\right) + \left(1 - \hat{I}_{A}^{1}\right)\hat{U}_{1}^{-1} \cdot y_{hij} + \left(\hat{I}_{A}^{1} - 1\right)\hat{U}_{0}^{-1} \cdot \log y_{hij}. \tag{11}$$

These variance estimators allow arbitrary correlations between observations below the first-stage clusters. They are therefore an alternative to the estimators developed by Schluter and Trede (2002).

3 Application to the i.i.d. case

In an i.i.d. framework the above indices are usually treated as follows. Income x_i and weight w_i of observation $i=1,\ldots,n$ are regarded as i.i.d. draws from a population (x,w). The index in question can then be represented as a function $I=g(\mu)$ of population moments $\mu=E(X_i)$, where X_i is a vector-valued function of (x_i,w_i) . It is estimated as $\hat{I}=g(\bar{X})$ and its sampling variance as $n^{-1}\nabla g(\bar{X})'\widehat{\text{var}}(X_i)\nabla g(\bar{X})$ (Cowell, 1989, or for the case without weights, Thistle, 1990). By contrast, Woodruff's method would yield a variance estimate $n^{-1}\widehat{\text{var}}(\nabla g(\bar{X})'X_i)$. It is easy to see that both estimates are identical. However, Woodruff's method leads to much simpler expressions as the problem is reduced to estimating the sampling variance of a scalar. In particular, no covariances need to be computed. Defining $\mu_{\alpha}=E(w_ix_i^{\alpha})$, $\tau_{\alpha}=E(w_ix_i^{\alpha}(\log x_i))$, $\hat{\mu}_{\alpha}=n^{-1}\sum_{i=1}^n w_ix_i^{\alpha}$ and $\hat{\tau}_{\alpha,\gamma}=n^{-1}\sum_{i=1}^n w_ix_i^{\alpha}(\log x_i)$, the sampling variances of the indices

$$I_{GE}^{\alpha} = \left(\alpha^2 - \alpha\right)^{-1} \left[\mu_0^{\alpha - 1} \mu_1^{-\alpha} \mu_\alpha - 1\right], \quad \alpha \in \mathbb{R} \setminus \{0, 1\}$$

$$\tag{12}$$

$$I_{Theil} = \tau_{1,1}\mu_1^{-1} - \log(\mu_1\mu_0^{-1}), \quad \alpha \to 1$$
 (13)

$$I_{MLD} = -\tau_{0,1}\mu_0^{-1} + \log(\mu_1\mu_0^{-1}), \quad \alpha \to 0$$
 (14)

$$I_A^{\varepsilon} = 1 - \mu_0^{-\varepsilon/(1-\varepsilon)} \mu_1^{-1} \mu_{1-\varepsilon}^{1/(1-\varepsilon)}, \quad \varepsilon \ge 0, \quad \varepsilon \ne 1, \tag{15}$$

$$I_A^1 = 1 - \mu_0 \mu_1^{-1} \exp(\tau_{0,1} \mu_0^{-1}), \quad \varepsilon \to 1$$
 (16)

can therefore simply be estimated by substituting

$$\tilde{z}_{i}^{GE} = \frac{1}{\alpha} \hat{\mu}_{\alpha} \hat{\mu}_{1}^{-\alpha} \hat{\mu}_{0}^{\alpha-2} - \frac{1}{\alpha - 1} \hat{\mu}_{\alpha} \hat{\mu}_{1}^{-\alpha - 1} \hat{\mu}_{0}^{\alpha - 1} \cdot x_{i} + \frac{1}{\alpha^{2} - \alpha} \hat{\mu}_{0}^{\alpha - 1} \hat{\mu}_{1}^{-\alpha} \cdot x_{i}^{\alpha}$$
(17)

$$\tilde{z}_{i}^{Theil} = \hat{\mu}_{1}^{-1} \cdot x_{i} \log x_{i} - \hat{\mu}_{1}^{-1} \left(\hat{\tau}_{1,1} \hat{\mu}_{1}^{-1} + 1 \right) \cdot x_{i} + \hat{\mu}_{0}^{-1}$$
(18)

$$\tilde{z}_i^{MLD} = -\hat{\mu}_0^{-1} \cdot \log x_i + \hat{\mu}_1^{-1} \cdot x_i + \mu_0^{-1} \left(\hat{\tau}_{0,1} \hat{\mu}_0^{-1} - 1\right)$$
(19)

$$\tilde{z}_{i}^{A} = \frac{\varepsilon}{1-\varepsilon}\hat{\mu}_{1}^{-1}\hat{\mu}_{1-\varepsilon}^{\frac{1}{1-\varepsilon}}\hat{\mu}_{0}^{-\frac{1}{1-\varepsilon}} + \hat{\mu}_{0}^{\frac{-\varepsilon}{1-\varepsilon}}\hat{\mu}_{1-\varepsilon}^{\frac{1}{1-\varepsilon}}\hat{\mu}_{1}^{-2} \cdot x_{i} - \frac{1}{1-\varepsilon}\hat{\mu}_{0}^{\frac{-\varepsilon}{1-\varepsilon}}\hat{\mu}_{1}^{-1}\hat{\mu}_{1-\varepsilon}^{\frac{\varepsilon}{1-\varepsilon}} \cdot x_{i}^{1-\varepsilon}$$

$$(20)$$

$$\tilde{z}_{i}^{1} = (\hat{I}_{A}^{1} - 1) \hat{\mu}_{0}^{-1} (1 - \hat{\mu}_{0}^{-1} \hat{\tau}_{0,1}) + (1 - \hat{I}_{A}^{1}) \hat{\mu}_{1}^{-1} \cdot x_{i} + (\hat{I}_{A}^{1} - 1) \hat{\mu}_{0}^{-1} \cdot \log x_{i}$$
 (21)

for \tilde{z}_i in

$$\widehat{\operatorname{var}}(\widehat{I}) = \frac{1}{n(n-1)} \sum_{i=1}^{n} \left(w_i \widetilde{z}_i - \frac{\sum_{i=1}^{n} w_i \widetilde{z}_i}{n} \right)^2.$$
 (22)

Equation (22) has a similar structure to equation (1), but note that the weights are treated differently in the complex survey and i.i.d. cases.⁴

³If the distribution of household income among individuals is analyzed, then observational units i are households and observations are replicated at the household level. In this case weights are $w_i = w_i'w_i''$ with household size w_i' and sample weight w_i'' . See Biewen (2002).

⁴For more discussion of different weighting concepts, see Cowell and Jenkins (2003).

4 Empirical illustration

We contrast the variance estimators using data from the first waves of the German Socio-Economic Panel (SOEP Group, 2001), and the British Household Panel Survey (Taylor et al., 2002).⁵ The GSOEP and the BHPS are widely used to analyze the income distribution in these two countries. Moreover, they provide information on primary sampling units (clusters) and strata identification variables. The first wave of each survey was chosen to avoid complications of the panel design. We considered the distribution of income among individuals. Following convention, each person was assumed to receive the equivalent household income of the household to which she belonged. (The equivalence scale was the square root of household size.)

The survey estimates shown in column (1) of Table 1 take into account sampling weights, stratification and clustering. Replication of observations at the household level is automatically accounted for, as this represents a form of clustering below the first stage clusters. The i.i.d. estimates shown in column (2) only take into account sampling weights and replication of observations at the household level, but not stratification or clustering. A comparison of columns (1) and (2) indicates that ignoring clustering and stratification makes surprisingly little difference for these data sets. By contrast, the estimates shown in columns (3) and (4) suggest that taking into account the replication of observations at the household level is much more important. The results in column (3) ignore the replication of observations at the household level, whereas those in column (4) take it into account. (Both (3)and (4) ignore first-stage clustering and stratification.) For the German data, this leads to standard errors that are about twice as large. This shows that survey design *can* matter. However, the precise effect appears to depend on the survey analysed: corresponding estimates in columns (3) and (4) differ little when BHPS data are used, by contrast with the GSOEP case.

— Table 1 near here —

The fact that ignoring stratification and first-stage clustering has only a small impact might be interpreted as good news for practitioners using these data, or for those using surveys in which primary sampling unit and strata identification variables are not made available, but it is not clear that this empirical finding can be generalized. Whatever the case, our variance estimators provide a straightforward means by which researchers can accommodate a range of design effects in their analysis.

⁵Stata programs to compute the estimators are available from the authors on request.

5 References

Biewen, M., 2002, Bootstrap Inference for Inequality, Mobility and Poverty Measurement. *Journal of Econometrics* 108, 317 – 342.

Binder, D.A., 1983, On the Variances of Asymptotically Normal Estimators from Complex Surveys, *International Statistical Review* 51, 293 – 300.

Binder, D.A. and Z. Patak, 1994, Use of Estimating Functions for Estimation From Complex Surveys, *Journal of the American Statistical Association* 89, 1035 – 1043.

Cochran, W.G., 1977, Sampling Techniques, 3d ed. (Wiley New York).

Cowell, F.A., 1989, Sampling Variance and Decomposable Inequality Measures, *Journal of Econometrics* 42, 27-41.

Cowell, F.A. and Jenkins, S.P., 2003, Estimating Welfare Indices: Household Weights and Sample Design, in: Y. Amiel and J.A. Bishop, eds., *Inequality, Welfare and Poverty: Theory and Measurement*, Volume 9 (Elsevier Science, Amsterdam) 147 – 172, .

Deaton, A., 1997, The Analysis of Household Surveys: a Microeconometric Approach to Development Policy (Johns Hopkins University Press, Baltimore).

Schluter, C. and M. Trede, 2002, Statistical Inference for Inequality and Poverty Measures with Dependent Data, *International Economic Review* 43, 185 – 200.

SOEP-Group, 2001, The German Socio-Economic Panel (GSOEP) after more than 15 years - Overview, in: Holst, E., D.R. Lillard and T.A. DiPrete, eds., Proceedings of the 2000 Fourth International Conference of German Socio-Economic Panel Study Users (GSOEP2000), Vierteljahreshefte zur Wirtschaftsforschung (Quarterly Journal of Economic Research) Vol. 70, 7 – 14.

Taylor, M., ed., 2002, *British Household Panel Survey User Manual*, Institute for Social and Economic Research, Colchester.

Thistle, P.D., 1990, Large Sample Properties of Two Inequality Indices, *Econometrica 58*, 725 – 728.

Woodruff, R.S., 1971, A Simple Method for Approximating the Variance of a Complicated Estimate, *Journal of the American Statistical Association* 66, 411 – 414.

6 Tables

Table 1. Income inequality¹ in West Germany (1984) and Britain (1991)

Index	(1) Survey ²		(2) I.i.d. ³		(3) Survey, i.i.d. ⁴		(4) Survey ⁵			
	estimate	std. err.	estimate	std. err.	estimate	std. err.	estimate	std. err.		
German Socio-Economic Panel ⁶ (1984)										
GE(-1)	0.4397	0.2971	0.3992	0.2573	0.4397	0.1333	0.4397	0.2978		
MLD	0.1339	0.0153	0.1338	0.0136	0.1339	0.0087	0.1339	0.0152		
Theil	0.1540	0.0276	0.1540	0.0246	0.1540	0.0187	0.1540	0.0276		
GE(2)	0.3673	0.1586	0.3502	0.1400	0.3673	0.1102	0.3673	0.1584		
Atkinson(0.5)	0.0658	0.0079	0.0661	0.0071	0.0658	0.0052	0.0658	0.0079		
Atkinson(1)	0.1253	0.0134	0.1253	0.0119	0.1253	0.0076	0.1253	0.0133		
Atkinson(1.5)	0.2153	0.0477	0.2101	0.0419	0.2153	0.0219	0.2153	0.0478		
Atkinson(2)	0.4679	0.1682	0.4439	0.1591	0.4679	0.0754	0.4679	0.1686		
British Household Panel Survey ⁷ (1991)										
GE(-1)	0.3656	0.0605	0.3627	0.0541	0.3656	0.0571	0.3655	0.0601		
MLD	0.1907	0.0051	0.1914	0.0050	0.1907	0.0030	0.1906	0.0050		
Theil	0.1779	0.0050	0.1784	0.0050	0.1779	0.0030	0.1779	0.0049		
GE(2)	0.2064	0.0085	0.2071	0.0086	0.2064	0.0054	0.2064	0.0085		
Atkinson(0.5)	0.0873	0.0022	0.0875	0.0022	0.0873	0.0013	0.0873	0.0021		
Atkinson(1)	0.1736	0.0042	0.1741	0.0041	0.1736	0.0025	0.1736	0.0041		
Atkinson(1.5)	0.2685	0.0084	0.2693	0.0082	0.2685	0.0058	0.2684	0.0082		
Atkinson(2)	0.4223	0.0404	0.4204	0.0363	0.4223	0.0381	0.4223	0.0401		

¹ Income refers to monthly equivalent household net income distributed among individuals (equivalence scale = square root of household size).

² Survey estimator, individual data, weight = individual sample weight, accounting for clustering and stratification; replication of observations at household level automatically accounted for.

³ I.i.d. estimator, household data, weight = household size * household sample weight (thus accounting for replication of observations at household level), ignoring clustering and stratification.

⁴ Survey estimator, individual data, weight = individual sample weight, ignoring clustering, stratification and replication of observations at household level (identical to i.i.d. estimator, individual data, weight = individual sample weight).

⁵ Survey estimator, individual data, weight = individual sample weight, ignoring clustering and stratification but accounting for replication of observations at household level (households are interpreted as clusters).

⁶ 110 strata, 516 clusters, 4232 households, 9441 individuals.

⁷ 75 strata, 250 clusters, 4814 households, 11616 individuals.

IZA Discussion Papers

No.	Author(s)	Title	Area	Date
747	J. Messina	Sectoral Structure and Entry Regulations	3	03/03
748	M. Pflüger	Economic Integration, Wage Policies and Social Policies	2	04/03
749	V. Jakobsen N. Smith	The Educational Attainment of the Children of the Danish 'Guest Worker' Immigrants	1	04/03
750	R. Hujer M. Caliendo S. Thomsen	New Evidence on the Effects of Job Creation Schemes in Germany - A Matching Approach with Threefold Heterogeneity	6	04/03
751	L. Borghans B. ter Weel	Are Computer Skills the New Basic Skills? The Returns to Computer, Writing and Math Skills in Britain	5	04/03
752	N. Malchow-Møller J. R. Skaksen	Skill-Biased Technological Change in Denmark: A Disaggregate Perspective	5	04/03
753	S. Burgess H. Turon	Unemployment Equilibrium and On-the-Job Search	1	04/03
754	H. Turon	Separability of Duration Dependence and Unobserved Heterogeneity	1	04/03
755	T. Dohmen	In Support of the Supporters? Do Social Forces Shape Decisions of the Impartial?	7	04/03
756	N. Datta Gupta R. Oaxaca N. Smith	Swimming Upstream, Floating Downstream: Comparing Women's Relative Wage Positions in the U.S. and Denmark	5	04/03
757	J. Hunt	Teen Births Keep American Crime High	5	04/03
758	J. Lise S. Seitz J. Smith	Equilibrium Policy Experiments and the Evaluation of Social Programs	6	04/03
759	E. P. Lazear	The Peter Principle: A Theory of Decline	1	04/03
760	E. P. Lazear	Entrepreneurship	5	04/03
761	E. P. Lazear	Output-Based Pay: Incentives, Retention or Sorting?	5	04/03
762	M. Gerfin R. E. Leu	The Impact of In-Work Benefits on Poverty and Household Labour Supply - A Simulation Study for Switzerland	3	04/03
763	M. Biewen S. P. Jenkins	Estimation of Generalized Entropy and Atkinson Inequality Indices from Complex Survey Data	5	04/03

An updated list of IZA Discussion Papers is available on the center's homepage www.iza.org.