

Candeal, Juan Carlos; Induráin, Esteban; Molina, José Alberto

Working Paper

Interpersonal comparisons of utility: an algebraic characterization of projective preorders and some welfare consequences

IZA Discussion Papers, No. 2594

Provided in Cooperation with:

IZA – Institute of Labor Economics

Suggested Citation: Candeal, Juan Carlos; Induráin, Esteban; Molina, José Alberto (2007) : Interpersonal comparisons of utility: an algebraic characterization of projective preorders and some welfare consequences, IZA Discussion Papers, No. 2594, Institute for the Study of Labor (IZA), Bonn, <https://nbn-resolving.de/urn:nbn:de:101:1-20080422267>

This Version is available at:

<https://hdl.handle.net/10419/34120>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.

IZA DP No. 2594

**Interpersonal Comparisons of Utility:
An Algebraic Characterization of Projective
Preorders and Some Welfare Consequences**

Juan Carlos Candeal
Esterban Induráin
José Alberto Molina

January 2007

Interpersonal Comparisons of Utility: An Algebraic Characterization of Projective Preorders and Some Welfare Consequences

Juan Carlos Candeal
University of Zaragoza

Esteban Induráin
Universidad Pública de Navarra

José Alberto Molina
University of Zaragoza and IZA

Discussion Paper No. 2594
January 2007

IZA

P.O. Box 7240
53072 Bonn
Germany

Phone: +49-228-3894-0
Fax: +49-228-3894-180
E-mail: iza@iza.org

Any opinions expressed here are those of the author(s) and not those of the institute. Research disseminated by IZA may include views on policy, but the institute itself takes no institutional policy positions.

The Institute for the Study of Labor (IZA) in Bonn is a local and virtual international research center and a place of communication between science, politics and business. IZA is an independent nonprofit company supported by Deutsche Post World Net. The center is associated with the University of Bonn and offers a stimulating research environment through its research networks, research support, and visitors and doctoral programs. IZA engages in (i) original and internationally competitive research in all fields of labor economics, (ii) development of policy concepts, and (iii) dissemination of research results and concepts to the interested public.

IZA Discussion Papers often represent preliminary work and are circulated to encourage discussion. Citation of such a paper should account for its provisional character. A revised version may be available directly from the author.

ABSTRACT

Interpersonal Comparisons of Utility: An Algebraic Characterization of Projective Preorders and Some Welfare Consequences

It is shown that any completely preordered topological real algebra admits a continuous utility representation which is an algebra-homomorphism (i.e., it is linear and multiplicative). As an application of this result, we provide an algebraic characterization of the *projective (dictatorial) preorders* defined on \mathbb{R}^n . We then establish some welfare implications derived from our main result. In particular, the connection with the normative property called *independence of the relative utility pace* is discussed.

JEL Classification: C60, C65, D60, D63

Keywords: algebraic utility, projective preorders

Corresponding author:

José Alberto Molina
Department of Economic Analysis
Faculty of Economics
University of Zaragoza
Gran Vía 2
50005 Zaragoza
Spain
E-mail: jamolina@unizar.es

1 Introduction

The main purpose of this paper is to provide some new insights that explain the algebraic nature of some normative properties that naturally appear in utility theory, social choice theory and welfare economics. To that end, we present a utility theorem, in addition to some of its applications in economics, for completely preordered algebraic structures which involve more algebraic machinery than that often encountered in the (economics) literature. Namely, we pay attention to completely preordered topological real algebras, i.e., algebraic structures equipped with three binary operations and endowed with a complete and continuous preorder which is compatible with these operations. As far as we can judge, no results of representability in this context have been considered in the economics literature before. Some related material to that presented here can be found in the classical works by Birkhoff (1948) and Fuchs (1963).

We show that any completely preordered topological real algebra admits a continuous utility representation which is an algebra-homomorphism, i.e., it preserves the algebraic structure. In addition to the own algebraic interest of this result, as a consequence we obtain an algebraic characterization of the *projective preorders* defined on \mathbb{R}^n . If we interpret each vector $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ as the utility-vector of a society which consists of n agents, this result can be described as follows. Let \preceq be a complete preorder defined on \mathbb{R}^n , representing the utility levels of n agents. Then \mathbb{R}^n , endowed with the usual operations defined *coordinatewise*, is a completely preordered topological real algebra under \preceq , if there is a kind of *dictator* for the society which means that the social welfare is measured just in terms of the solely welfare of that individual; that is, there is $j \in \{1, \dots, n\}$ such that $x \preceq y \iff x_j \leq y_j$ where $x = (x_1, \dots, x_n)$, $y = (y_1, \dots, y_n) \in \mathbb{R}^n$. In other words, the existence of a dictator can be stated in terms of algebraic conditions. In the next section we also offer an interpretation of the binary operations involved in terms of welfare normative properties.

The relationship between projective preorders and what we call the *independence of the relative utility pace* (I.R.U.P.) property is also discussed. This condition can be described as follows. Let $x, y \in \mathbb{R}^n$ be two utility-vectors of an n -agents society. Then, $x = (x_1, \dots, x_j, \dots, x_n) \preceq y = (y_1, \dots, y_j, \dots, y_n)$ iff $(x_1, \dots, f(x_j), \dots, x_n) \preceq (y_1, \dots, f(y_j), \dots, y_n)$ for every $j \in \{1, \dots, n\}$, and every increasing map $f : \mathbb{R} \rightarrow \mathbb{R}$. In words, I.R.U.P. means that individual increase of utility levels does not necessarily increase the welfare of society. Intuitively, this loss of welfare can be imputed to the existence of dominating coalitions. In fact, we prove that if \preceq is non-trivial and increasing then the independence of the relative utility pace assumption implies the existence of a dictator. We conclude with a remark explaining the role that algebraic issues of the kind considered in this note play in *social choice theory*.

2 Previous definitions and notations

Throughout the paper, \preceq will denote a complete preorder defined on a (non-empty) set X , i.e., a binary relation on X which is *reflexive*, ($x \preceq x$ for all $x \in X$), *transitive* ($x \preceq y, y \preceq z$ implies $x \preceq z$); and *decisive* or *complete* (either $x \preceq y$ or $y \preceq x$ for every $x, y \in X$). We also say that (X, \preceq) is a *completely preordered set*. Two elements $x, y \in X$ are indifferent if $x \preceq y$ and $y \preceq x$ (briefly, $x \sim y$). If, in addition, \preceq is anti-symmetric ($x \preceq y$ and $y \preceq x$ implies $x = y$) then \preceq is said to be a *total order*.

Given \preceq , we consider as usual the *strict binary relation* \prec on X defined as: $x \prec y$ iff no $(y \preceq x)$. The preorder \preceq is said to be *non-trivial* if there are $x, y \in X$ such that $x \prec y$.

Let (X, \preceq) be a completely preordered set, a function $u : X \longrightarrow \mathbb{R}$ is said to be a *utility function* for \preceq if, for every $x, y \in X$, it holds that $x \preceq y \iff u(x) \leq u(y)$.

We will focus our attention on spaces X having a rich algebraic structure, namely, rings and real algebras. For the sake of completeness, we only recall the definition of a real algebra.¹

Definition 2.1. A *real algebra* $(X, +, \cdot_{\mathbb{R}}, *)$ is a space X endowed with three binary operations such that:

- (i) $(X, +, \cdot_{\mathbb{R}})$ is a real vector space.
- (ii) $(X, +, *)$ is a ring.
- (iii) $\lambda \cdot (x * y) = (\lambda \cdot x) * y = x * (\lambda \cdot y), \forall x, y \in X, \forall \lambda \in \mathbb{R}$.

Let $(X, +, \cdot_{\mathbb{R}}, *)$ be a real algebra. A function $v : X \longrightarrow \mathbb{R}$ is said to be an *algebra-homomorphism* if it is linear and multiplicative, i.e.:

- (a) $v(\lambda \cdot x + \beta \cdot y) = \lambda v(x) + \beta v(y), \forall x, y \in X, \forall \lambda, \beta \in \mathbb{R}$.
- (b) $v(x * y) = v(x)v(y), \forall x, y \in X$.

Definition 2.2. A *completely preordered real algebra* $(X, \preceq, +, \cdot_{\mathbb{R}}, *)$ is a real algebra equipped with a complete preorder \preceq , which is compatible with the operations $+, \cdot_{\mathbb{R}}$ and $*$, i.e.:

- (1) $x \preceq y$ implies $x + z \preceq y + z, \forall z \in X$.
- (2) $x \preceq y, 0 \leq \lambda$ implies $\lambda \cdot x \preceq \lambda \cdot y$.
- (3) $x \preceq y, \mathbf{0} \preceq z$ imply $z * x \preceq z * y$ and $x * z \preceq y * z$ ($\mathbf{0}$ denotes the null element with respect to $+$).

¹The reader will find the textbook of Hungerford (2003) a useful reference for algebraic issues.

Following Moulin(1988) let us provide some welfare interpretations of axioms (1), (2) and (3) above. Condition (1) says that \succsim is *zero independent* which means that comparisons between utility streams depend upon absolute variations of utilities or, equivalently, \succsim is invariant to independent changes of origins. Condition (2) states that \succsim is *homothetic* (see, e.g., Candeal and Induráin (1995 a)) which reflects the independence of a common utility scale (or common relative variations of utility). Finally, condition (3) means that \succsim is *independent of arbitrarily relative variations of utilities* or, equivalently, it is invariant to independent changes of units (in which, say, utility is measured). Of course, all of these welfare interpretations for \succsim are easily established provided that $X = \mathbb{R}^n$ and the operations $+$, $\cdot_{\mathbb{R}}$ and $*$ are the usual ones defined coordinatewise. However, they are still meaningful in broader contexts like l_{∞} , the space of bounded real sequences, which naturally arises in intertemporal decision problems.

As we are interested in obtaining strong properties of representability for \succsim , we allow X to be a topological space endowed with a topology τ . In order for this to be made in a natural way, we require $(X, \tau, +, \cdot_{\mathbb{R}})$ to be a *topological vector space* (note that no assumptions involving topological considerations are made on $*$). We are now ready to introduce a key definition.

Definition 2.3. We say that $(X, \succsim, \tau, +, \cdot_{\mathbb{R}}, *)$ is a *completely preordered topological real algebra* if

- (i) $(X, \succsim, +, \cdot_{\mathbb{R}}, *)$ is a completely preordered real algebra.
- (ii) $(X, \tau, +, \cdot_{\mathbb{R}})$ is a topological real vector space.
- (iii) The preorder \succsim is *continuous*, i.e., for every $x \in X$, the lower and upper contour sets $L(x) = \{y \in X; y \succsim x\}$ and $G(x) = \{y \in X; x \succsim y\}$, respectively, are closed in X (i.e., the complements of $L(x)$ and $G(x)$ belong to τ for every $x \in X$).

A complete preorder \succsim defined on a subset $X \subseteq \mathbb{R}^n$ is said to be *increasing* if for every $x, y \in X$, such that that $x \leq y$ it holds that $x \succsim y$, where by $x \leq y$ we mean $x_i \leq y_i, \forall i \in \{1, \dots, n\}$, $x = (x_1, \dots, x_i, \dots, x_n)$, $y = (y_1, \dots, y_i, \dots, y_n)$.

Let us state now the so-called in the mathematical utility literature Eilenberg-Debreu theorem which will be used later on.

Theorem 2.4. (Eilenberg(1941), Debreu(1964)) *Let (X, τ) be a second countable topological space. Then every τ -continuous complete preorder \succsim defined on X can be represented by a continuous utility function.*

3 The main result

In this section we obtain a theorem of representability of a completely preordered topological real algebra by means of a continuous utility function which is an

algebra-homomorphism, i.e., it is linear and multiplicative. It is important to say that the result is also valid in an infinite-dimensional context and nothing about the commutativity of the binary operation $*$ is assumed.

Before stating the theorem, we need a preliminary lemma.

Lemma 3.1. *Suppose that \mathbb{R} is endowed with both a binary operation $*$ and a non-trivial continuous total order \preceq , in such a way that $(\mathbb{R}, \preceq, \tau_{Euclidean}, +, *)$ becomes a completely preordered topological ring². Then \preceq is representable by a continuous utility function which is a ring-homomorphism. Moreover, there exists $a \in \mathbb{R} \setminus \{0\}$; $x * y = axy$, $\forall x, y \in \mathbb{R}$.*

Proof. Since \preceq is continuous on \mathbb{R} , it follows from Theorem 2.4 that there exists an (injective) continuous utility function $u : \mathbb{R} \rightarrow \mathbb{R}$ for \preceq . Actually, u is a homeomorphism. Thus, without loss of generality, we can assume that either $u(x) = x$ or $u(x) = -x$ ($x \in \mathbb{R}$). Suppose that the first holds, the other case being analyzed in a similar way.

Now by a result by Pickert and Hion (see Birkhoff (1948), p. 398-399), \preceq is representable by a utility function $v : \mathbb{R} \rightarrow \mathbb{R}$ which is a ring-homomorphism. So, because u and v are utility functions for \preceq , there exists an increasing function $g : v(\mathbb{R}) \rightarrow \mathbb{R}$, such that $gv(x) = x$, $\forall x \in \mathbb{R}$. Let us see that v is also increasing. For that, let $x, y \in \mathbb{R}$; $x < y$. Then, because v is injective, either $v(x) < v(y)$ or $v(y) < v(x)$. If $v(y) < v(x)$ then, since g is increasing, $gv(y) < gv(x)$, i.e., $y < x$, which is a contradiction. So, $v(x) < v(y)$, and v is an increasing function. Remember that v is an additive function and it is well known that if v is discontinuous at some point then it is discontinuous at every point of \mathbb{R} . This last fact, being v increasing, leads to $v(x) = ax$ (with $a > 0$). Because $a(x*y) = v(x*y) = v(x)v(y) = a^2xy$, this type of representation gives $x*y = axy$ $\forall x, y \in \mathbb{R}$. \square

Remark 3.2. As the lemma shows, the operation $*$ is essentially the usual multiplicative one in \mathbb{R} . Note that $(\mathbb{R}, +, *)$ is a *field* and the map $(x, y) \in \mathbb{R} \times \mathbb{R} \rightarrow x * y \in \mathbb{R}$ is continuous.

Let us now prove the main result of the paper, which involves a very particular kind of utility representation for completely preordered topological real algebras.

Theorem 3.3. *Let $(X, \preceq, \tau, +, \cdot_{\mathbb{R}}, *)$ be a completely preordered topological real algebra. Then \preceq admits a continuous utility function $\Gamma : X \rightarrow \mathbb{R}$ which is, in addition, an algebra-homomorphism.*

Proof. If \preceq is trivial, then $\Gamma \equiv 0$ works. Otherwise, consider the set $I(\mathbf{0}) = \{x \in X; x \sim \mathbf{0}\}$. First, let us see that $I(\mathbf{0})$ is an ideal of X , for which we need to prove the following two properties:

- (i) $I(\mathbf{0})$ is a real vector subspace of X .

² $(X, \preceq, +, *)$ is a completely preordered ring if $(X, +, *)$ is a ring and \preceq is a complete preorder on X which satisfies condition (1) and (3) of Definition 2.2

(ii) For every $x \in I(\mathbf{0})$, $y \in X$, it holds that $y * x \in I(\mathbf{0})$ and $x * y \in I(\mathbf{0})$.

Let $x, y \in I(\mathbf{0})$. Because \preceq is compatible with $+$, it follows that $x + y \sim x + \mathbf{0} = x \sim \mathbf{0}$. So, in order to prove (i), it is enough to see that, given $x \in I(\mathbf{0})$ and $\lambda \in \mathbb{R}$, then $\lambda \cdot x \sim \mathbf{0}$. To that end, denote by $S(x)$ the linear subspace generated by x , i.e, $S(x) = \{\lambda \cdot x; \lambda \in \mathbb{R}\}$, and consider the preorder \preceq restricted to $S(x)$. Since $(X, \tau, +, \cdot_{\mathbb{R}})$ is a topological vector space, endowed with the relative topology of X , $S(x)$ is homomorphic to \mathbb{R} and therefore it is a connected and separable topological space. Because \preceq is continuous it follows, by Theorem 2.4, that there exists a continuous utility function $u : S(x) \rightarrow \mathbb{R}$ which represents \preceq . By identifying $S(x)$ with \mathbb{R} , the function u clearly satisfies the following functional equation $u(\lambda_1) = u(\lambda_2)$ iff $u(\lambda_1 + \alpha) = u(\lambda_2 + \alpha)$, $\forall \alpha \in \mathbb{R}$ or, equivalently, the function $s \in \mathbb{R}_{++} \rightarrow v(s) = u(\log s)$ satisfies the functional equation: $v(s_1) = v(s_2)$ iff $v(\lambda s_1) = v(\lambda s_2)$, $\forall \lambda \in \mathbb{R}_{++}$. It was shown in Candeal and Induráin (1993) that the only continuous solutions of the latter functional equation are the constant functions. Thus, every element in $S(x)$ is indifferent to $\mathbf{0}$.

To prove (ii), let $y \in X$ and $x \in I(\mathbf{0})$. If $\mathbf{0} \preceq y$ then, since $(X, +, \cdot_{\mathbb{R}}, *)$ is a completely preordered real algebra, $y * x \preceq y * \mathbf{0} = \mathbf{0}$ and $\mathbf{0} = y * \mathbf{0} \preceq y * x$. Therefore, $y * x \sim \mathbf{0}$. If $y \preceq \mathbf{0}$, then $\mathbf{0} \preceq -y$ and so $-y * x \sim \mathbf{0}$. Because $y * x = -(-y) * x$, and $I(\mathbf{0})$ is a vector subspace of X , it holds that $y * x \sim \mathbf{0}$. Similarly, we can prove that $x * y \sim \mathbf{0}$. Thus (ii) holds and therefore $I(\mathbf{0})$ is an ideal of X .

Now consider the quotient space $X/I(\mathbf{0})$. Because $X/I(\mathbf{0})$ coincides with the quotient space X/\sim , $X/I(\mathbf{0})$ is a totally ordered set. We denote the natural order on $X/I(\mathbf{0})$ by \preceq' . Since $I(\mathbf{0})$ is an ideal of X , the operations $+$, $\cdot_{\mathbb{R}}$ and $*$ pass to the quotient in such a way that $(X/I(\mathbf{0}), \preceq', +, \cdot_{\mathbb{R}}, *)$ becomes a totally ordered real algebra. Moreover, endowed with the quotient topology ς , $X/I(\mathbf{0})$ is a topological vector space because $I(\mathbf{0})$ is a closed subspace of X which follows from the fact that \preceq is continuous. Therefore, $(X/I(\mathbf{0}), \preceq', \varsigma, +, \cdot_{\mathbb{R}}, *)$ is a totally ordered topological real algebra. By Theorem 2.4 again, a standard connectedness argument shows that the co-dimension of $I(\mathbf{0})$ is one or, equivalently, $X/I(\mathbf{0})$ is homeomorphic and isomorphic as a vector space to \mathbb{R} . Now, by the previous lemma, \preceq' is representable by a continuous utility function $u : X/I(\mathbf{0}) \rightarrow \mathbb{R}$, which is linear and multiplicative. Let us denote by $p : X \rightarrow X/I(\mathbf{0})$ the projection map. It is easy to see that p is linear, multiplicative and continuous. Then it is enough to consider the composition $u \circ p : X \rightarrow \mathbb{R}$ to obtain the desired result. \square

4 Applications

In this section we provide some applications of our theorem. The first one is concerned with the characterization of all (continuous) complete preorders \preceq defined on \mathbb{R}^n for which $(\mathbb{R}^n, \preceq, \tau_{Euclidean})$, under the usual operations defined coordinatewise, is a completely preordered topological real algebra. This will lead us to the notion of *projective* preorder. The second one links the algebraic

condition of being a completely preordered topological real algebra with a normative property; namely, the *independence of the relative utility pace*, which turns out to be a generalization of the property known as the *independence of the common utility pace* introduced by D'Aspremont and Gevers (1977). For a survey of this, and other normative properties and their welfare significance see Moulin (1988) and, more recently, Bossert and Weymark (2004) and Fleurbaey and Hammond (2004).

Definition 4.1. Let \succsim be a complete preorder defined on \mathbb{R}^n and let $j \in \{1, \dots, n\}$. We say that \succsim is *j-projective* if for every $x = (x_1, \dots, x_j, \dots, x_n)$, $y = (y_1, \dots, y_j, \dots, y_n)$, it holds that $x \succsim y$ if and only if $x_j \leq y_j$. \succsim is said to be *projective* if it is j-projective for some $j \in \{1, \dots, n\}$.

Let \mathbb{R}^n be endowed with the usual operations $+$, $\cdot_{\mathbb{R}}$ and $*$ defined coordinatewise. It is then clear that $(\mathbb{R}^n, +, \cdot_{\mathbb{R}}, *)$ is a real algebra. Now, we are interested in finding all complete preorders \succsim defined on \mathbb{R}^n for which $(\mathbb{R}^n, \succsim, \tau_{Euclidean}, +, \cdot_{\mathbb{R}}, *)$ becomes a completely preordered topological real algebra.

Proposition 4.2. *The only complete preorders \succsim defined on \mathbb{R}^n for which $(\mathbb{R}^n, \succsim, \tau_{Euclidean}, +, \cdot_{\mathbb{R}}, *)$ is a completely preordered topological real algebra are the projective ones.*

Proof. Let \succsim be a preorder defined on \mathbb{R}^n such that $(\mathbb{R}^n, \succsim, \tau_{Euclidean}, +, \cdot_{\mathbb{R}}, *)$ is a completely preordered topological real algebra. By using the representation result (Theorem 3.3), it follows that there is a continuous, linear and multiplicative function $\psi : \mathbb{R}^n \rightarrow \mathbb{R}$ which represents \succsim . Because ψ is linear and continuous, ψ is of the form $\psi(x_1, \dots, x_n) = \sum_{i=1}^n a_i x_i$, for some $a_i \in \mathbb{R}$, $i = 1, \dots, n$. Let us prove that all, but at most one, coefficients a_i are zero. Suppose by way of contradiction that there exist $i \neq j$ such that $a_i \cdot a_j \neq 0$. Then, because ψ is also multiplicative, $\psi(0, \dots, 0, \dots, 0) = \psi((0, \dots, 1_i, \dots, 0) * (0, \dots, 1_j, \dots, 0)) = \psi(0, \dots, 1_i, \dots, 0) \psi(0, \dots, 1_j, \dots, 0)$. That is, $0 = a_i \cdot a_j$. This contradiction proves that there is $j \in \{1, \dots, n\}$ such that $\psi(x_1, \dots, x_n) = a_j x_j$. It should be noted in addition that $\psi((1, \dots, 1) * (1, \dots, 1)) = \psi(1, \dots, 1) = \psi(1, \dots, 1) \psi(1, \dots, 1)$. In other words, $a_j = a_j^2$, hence $a_j = 0$ or 1 . Thus, if $a_j = 0$, then \succsim is trivial (all the utility vector are indifferent) and \succsim is j-projective, for all $j \in \{1, \dots, n\}$. If $a_j = 1$, then $\psi(x_1, \dots, x_n) = x_j$ and \succsim is a j-projective complete preorder defined on \mathbb{R}^n . \square

Remarks 4.3. (i) Proposition 4.2 can be re-stated in the following equivalent terms: *The only complete and continuous preorders defined on \mathbb{R}^n which satisfy zero-independence, homotheticity and independence of arbitrarily relative variations of utilities are the projective ones.*

(ii) It should be noted that the conclusion of the previous proposition need not be true in the infinite-dimensional context. For example, if the “utility vectors” represent streams of utility with (a countable number of) infinite coordinates like in the intertemporal decision problems, there could exist non-projective continuous complete preorders defined on the space of utility

streams in such a way that this space can be given a structure of a completely preordered topological real algebra. This is the case for the space l_∞ , which consists of all real bounded sequences, and that is often encountered in the economics literature related to decision problems over time involving an infinite horizon. It is easy to check that with the usual operations defined coordinatewise, $(l_\infty, \|\cdot\|_\infty, +, \cdot_{\mathbb{R}}, *)$ is a topological real algebra (in fact, it is a Banach real algebra); the topology given by the supremum norm $\|x\|_\infty = \|(x_1, \dots, x_n, \dots)\|_\infty = \sup\{|x_n|; n \in \mathbb{N}\}$. Let us denote by $\beta(\mathbb{N})$ the Stone-Ćech compactification of the set of the natural numbers \mathbb{N} (see Dugundji (1966)). Let $p \in \beta(\mathbb{N}) \setminus \mathbb{N}$. Then the evaluation map at p defines a continuous, linear and multiplicative map $x \in l_\infty \rightsquigarrow e_p(x) \in \mathbb{R}$. In other words, e_p belongs to the spectrum of the Banach algebra $(l_\infty, \|\cdot\|_\infty, +, \cdot_{\mathbb{R}}, *)$ (see, Rickart, (1960)). Thus, the complete preorder \succsim_p defined on l_∞ as $x \succsim_p y \Leftrightarrow e_p(x) \leq e_p(y)$ is continuous and allows the structure $(l_\infty, \succsim_p, \|\cdot\|_\infty, +, \cdot_{\mathbb{R}}, *)$ to become a completely preordered topological real algebra. It should be noted that, from the point of view of an economic interpretation, \succsim_p can be considered as the existence of an *invisible dictator or social planner*.

(iii) Although the conclusion of Proposition 4.2 does not hold, in general, in the infinite-dimensional setting (see Remarks (ii) above), it can still be generalized. Indeed, the statement remains true for the space of real sequences which vanish at infinity, usually denoted by c_0 . Equipped with the usual operations defined coordinatewise and the topology given by the supremum norm, c_0 is a topological real algebra (in fact, a Banach subalgebra of l_∞). The reason for the statement of Proposition 4.2 to be true in this case is that the spectrum of c_0 , the set of all linear and multiplicative functionals defined on it, can be easily seen to coincide with the set of natural numbers \mathbb{N} .

Definition 4.4. Let \succsim be a complete preorder defined on \mathbb{R}^n . We say that \succsim is *independent of the relative utility pace* (briefly, I.R.U.P.) if $(x_1, \dots, x_j, \dots, x_n) \succsim (y_1, \dots, y_j, \dots, y_n)$ implies $(x_1, \dots, f(x_j), \dots, x_n) \succsim (y_1, \dots, f(y_j), \dots, y_n)$ for every $j \in \{1, \dots, n\}$ and every increasing map $f : \mathbb{R} \rightarrow \mathbb{R}$.

On the basis of this definition we can state the following result.

Proposition 4.5. *Let \succsim be a non-trivial increasing ($x \leq y$ implies $x \succsim y$) and continuous complete preorder defined on \mathbb{R}^n . Then the following statements are equivalent:*

(i) \succsim is projective

(ii) \succsim is I.R.U.P.

Proof. (i) \Rightarrow (ii) is obvious. (ii) \Rightarrow (i). By induction on n . For $n = 1$ it is obvious since \succsim is the usual ordering on \mathbb{R} . Let us prove it for $n = 2$. Let $z = (z_1, z_2)$, $x = (x_1, x_2)$ and $y = (y_1, y_2) \in \mathbb{R}^2$. Because \succsim is I.R.U.P., we have that $(x_1, x_2) \succsim (y_1, y_2)$ if and only if $(x_1 + z_1, x_2) \succsim (y_1 + z_1, y_2)$ by taking the increasing function $f(r) = r + z_1$ ($r \in \mathbb{R}$). By repeating this process over the second coordinate, we obtain $x \succsim y \Leftrightarrow x + z \succsim y + z$, i.e., \succsim is compatible

with respect to $+$ (property (1) of Definition 2.2). This fact, together with \succsim being continuous, implies that \succsim can be represented by a linear utility function (see Trockel, (1992) or Candeal and Induráin (1995 b)). Since \succsim is non-trivial and increasing, there is a non-null $\mathbf{0} \leq p = (p_1, p_2) \in \mathbb{R}^2$ such that $x \succsim y \Leftrightarrow (p, x) \leq (p, y)$, where (\cdot) denotes the usual inner product in \mathbb{R}^n . Let $\mathbf{0} \neq z = (z_1, z_2) \sim (0, 0)$. Because the set $I(\mathbf{0})$ is a linear subspace and \succsim is increasing, we can have neither $z_1, z_2 > 0$ nor $z_1, z_2 < 0$. Suppose now, without loss of generality, that $z_1 \geq 0, z_2 \leq 0$. If $z_1 > 0$, then by taking an increasing function $f: \mathbb{R} \rightarrow \mathbb{R}$ such that $f(z_1) = k$ ($k > 0$) and $f(0) = 0$, by I.R.U.P., we have that $(f(z_1), z_2) \sim (f(0), 0)$; i.e., $(k, 0) \sim (0, 0)$. So $p_1 = 0$ (and therefore $z_2 = 0$), hence \succsim is 2-projective. A similar argument holds for $z_2 < 0$. In any case, \succsim is projective.

To show the general case, note first that, by the same arguments as for $n = 2$, there exists a non-null $\mathbf{0} \leq p = (p_1, \dots, p_n) \in \mathbb{R}^n$ such that $x \succsim y \Leftrightarrow (p, x) \leq (p, y)$ ($x, y \in \mathbb{R}^n$). Consider $\mathbb{R}^{n-1} = \{(x_1, \dots, x_n) \in \mathbb{R}^n; x_n = 0\}$. If \succsim is trivial on \mathbb{R}^{n-1} , then $p_1 = \dots = p_{n-1} = 0$ and $p_n \neq 0$. Then, \succsim is n-projective. Suppose, otherwise, that \succsim is non-trivial on \mathbb{R}^{n-1} . Then, by induction hypothesis (note that the continuity, increasingness and I.R.U.P. are induced over linear subspaces), there is $j \in \{1, \dots, n-1\}$ such that $p_j \neq 0$ and $p_i = 0 \forall i \in \{1, \dots, n-1\} \setminus \{j\}$. Now, it is enough to consider $\mathbb{R}^{j,n} = \{(0, \dots, x_j, 0, \dots, x_n); x_j, x_n \in \mathbb{R}\}$ and apply again the induction hypothesis for $n = 2$ to obtain that, in this case, necessarily $p_n = 0$. Thus, \succsim is j-projective. \square

The next corollary follows immediately and shows an algebraic characterization of those complete preorders defined on \mathbb{R}^n which are independent of the relative utility pace. Assume that \mathbb{R}^n is endowed with the usual operations defined coordinatewise.

Corollary 4.6. *Let \succsim be a non-trivial increasing preorder defined on \mathbb{R}^n . Then $(\mathbb{R}^n, \succsim, \tau_{\text{Euclidean}}, +, \cdot_{\mathbb{R}}, *)$ is a completely preordered topological real algebra if and only if \succsim is I.R.U.P.*

Remark 4.7. In this paper we have presented a mathematical result concerning the existence of continuous linear and multiplicative utility for the class of ordered structures called *completely preordered topological real algebras*. In particular, we have offered an algebraic characterization of the class of social welfare orderings called *projective preorders* defined on \mathbb{R}^n . Since this latter class plays an important role in social choice theory, it seems natural to ask whether or not the approach followed can result into a different methodology to obtain, say, Arrow-like theorems. Indeed, this is not new in the literature. It is well known (see, e.g., D'Aspremont and Gevers (1977)) that a social welfare functional, defined on profiles of individual preference orderings, induces a social welfare functional defined on profiles of individual utilities. Moreover, independence properties of the former rules pass to invariance properties of the latter. This process yields social welfare functionals that can be represented by social welfare orderings. Now, it would be expected that our algebraic characterization of the dictatorial preorders (defined even in more general settings

than \mathbb{R}^n , see Remarks 4.3, (iii)) can result into a characterization of dictatorial social welfare functionals, say, à la Arrow. The nub point here is the continuity of the social welfare ordering induced by the social welfare functional.

References

- [1] Birkhoff, G., 1948, *Lattice Theory*, American M. Society, Providence RI
- [2] Bossert, W. and J.A. Weymark, 2004, *Utility in social choice*, Chapter **20** of the Handbook of Utility Theory (Vol.2), Edited by S. Barberà, P.J. Hammond and Ch. Seidl. Kluwer Academic Publishers. Amsterdam.
- [3] Candeal, J.C and E. Induráin, 1993, *On the structure of homothetic functions*, Aequationes Mathematicae **45**, 207-218.
- [4] Candeal, J.C. and E. Induráin, 1995 a, *Homothetic and weakly homothetic preferences*, Journal of Mathematical Economics **24**, 147-158.
- [5] Candeal, J.C. and E. Induráin, 1995 b, *A note on linear utility*, Econ. Theory **6**, 519-522.
- [6] D'Aspremont, C. and L. Gevers, 1977, *Equity and the informational basis of collective choice*, Rev. Econ. Stud. **44**, 199-209.
- [7] Debreu, G., 1954, *Representation of a preference ordering by a numerical function*, in: R. Thrall, C. Coombs and R. Davis, eds. Decision processes (Wiley, New York) 159-166.
- [8] Dugundji, J., 1966, *Topology*, Allyn and Bacon, Boston.
- [9] Eilenberg, S., 1941, *Ordered topological spaces*, American Journal of Mathematics **63**, 39-45.
- [10] Fleurbaey, M. and P.J. Hammond, 2004, *Interpersonally comparable utility*, Chapter **21** of the Handbook of Utility Theory (Vol.2), Edited by S. Barberà, P.J. Hammond and Ch. Seidl. Kluwer Academic Publishers. Amsterdam.
- [11] Fuchs, L., 1963, *Partially ordered algebraic systems*, Pergamon Press
- [12] Hungerford, T.W., 2003, *Algebra*, Graduate Texts in Mathematics, Vol. **73** (5th ed.), Springer.
- [13] Moulin, H., 1988, *Axioms for cooperative decision making*, Econometric Society Monographs.
- [14] Rickart, C.E., 1960, *General theory of Banach algebras*, D. Van Nostrand Company, Inc., Princeton, N.J.
- [15] Trockel, W., 1992, *An alternative proof for the linear utility representation theorem*, Econ. Theory **2**, 298-302.