

Bozgå, Ion; Pandia, Olimpia; Sărăcin, Ion

**Conference Paper**

## Physiological proprieties from corn influenced by the differentiate application of nitrogen and phosphorus

**Provided in Cooperation with:**

The Research Institute for Agriculture Economy and Rural Development (ICEADR), Bucharest

*Suggested Citation:* Bozgå, Ion; Pandia, Olimpia; Sărăcin, Ion (2014) : Physiological proprieties from corn influenced by the differentiate application of nitrogen and phosphorus, In: Agrarian Economy and Rural Development - Realities and Perspectives for Romania. 5th Edition of the International Symposium, November 2014, Bucharest, The Research Institute for Agricultural Economy and Rural Development (ICEADR), Bucharest, pp. 141-145

This Version is available at:

<https://hdl.handle.net/10419/111623>

**Standard-Nutzungsbedingungen:**

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

**Terms of use:**

*Documents in EconStor may be saved and copied for your personal and scholarly purposes.*

*You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.*

*If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.*

# PHYSIOLOGICAL PROPRIETIES FROM CORN INFLUENCED BY THE DIFFERENTIATE APPLICATION OF NITROGEN AND PHOSPHORUS

ION BOZGĂ<sup>1</sup>, OLIMPIA PANDIA<sup>1</sup>, ION SĂRĂCIN<sup>2</sup>

**Abstract:** *The paper enumerates several physiological processes that take place at the Champion corn hybrid and the interaction between the hybrid in the irrigated and un-irrigated culture, as well as the appliance of NP fertilizers. To obtain performances at this type of corn culture, soil analyses were made before the establishment of experimental parcels and it was aimed the moment when the corn plants had already formed eight leaves to emphasize the results better. The determinations and the laboratory analyses emphasizes the most important physiological processes from plant life, where the water and fertilizer dosages had a crucial role in obtain of positive results.*

**Key words:** *physiological processes, respiration, transpiration, carotene, hybrid*

## INTRODUCTION

In our country, the maize crop (culture) occupies an important area of the arable land and it represents one of the main cereal crops because of its importance in nourishment, feeding the animals and in industry.

The demographic growth of the population as well as the animal effectives imposed an extension of crop areas and a growth of production/ area; these two aspects were possible by an intensification of maize crop by using chemical fertilizers and irrigation systems.

Because maize culture is the most extended culture in the world, not only in our country, research into discovering new productive and qualitative hybrids have been performed, for human beings, animals and also industry, and different properties and processes were had in view.

Because corn can adapt at almost all types of soil in Romania, numerous varieties and hybrids were created, thus it occupies the largest culture surface from Romania. The corn culture has a vast area of spreading and in the southern and western parts can be made big productions of corn, even for satisfying the needs necessary of national economy.

Because in the human nutrition, corn grains are consumed in different forms, it is important to have healthier corn cultures, so that the products that derives from them to be beneficial for our organisms.

In the human nutrition only corn grains are considered and they are an important aliment consumed in various ways, specific to each nation. The grains are used as they are, as boiled or baked corn (very indicated for this type is the sweet corn). Sweet corn grains detached from the cob in milk maturation are used boiled or canned like grained peas.

## MATERIAL AND METHOD

The study was performed using an Champion maize hybrid, when the 8 full-grown leaves became visible and this hybrid was cultivated in the north-western part of Olt District, at Grădinari, on chernozem, and both irrigated and non-irrigated systems were a used, after a precursory wheat culture in 2012, and physiological analysis were performed in order to point out the changes of the chemical composition of plants when applying different dosage of NP.

Analysis were performed by using variants disposed in four repetitions having as an example one row graduated multi-staged plots of land method. The sowed area of the variant is of 22.4 m<sup>2</sup>, and its density is of 50,000 plants/hectare. Analyses of soil were effectuated at 0-25 cm and 23-34 cm. depth.

---

<sup>1</sup>Conf.univ.dr. Ion Bozgă: University of Agricultural Sciences and Veterinary Medicine Bucharest, Romania, e-mail: veterinar\_serv@yahoo.com, <sup>1</sup>Sl.univ.dr.ing. Olimpia Pandia: e-mail: olimpia\_pandia@yahoo.com

<sup>2</sup>Prof.univ.dr.ing. Ion Sărăcin: University of Craiova, Faculty of Agriculture, Romania, e-mail: ion\_saracin@yahoo.com

Table 1

Chemical properties of the argic chernosiomus from the Grădinari-Olt

Genetic level	Depth(cm)	pH value (H <sub>2</sub> O)	S.B. m.e/100g	S.H. m.e	Humus %	N total %	P p.p.m.	K p.p.m.
ApI	0-25	7,02	24,37	3,43	2,78	0,141	22,7	166,8
Ap1	0-25	7,03	24,39	3,47	2,77	0,143	22,9	166,5
Ap2	0-25	7,01	24,36	3,45	2,77	0,142	22,7	166,7
Ap3	0-25	7,02	24,38	3,43	2,76	0,141	22,8	166,5
ApII	25-34	7,19	26,85	2,32	29,18	0,133	28,2	172,9
Ap1	25-34	7,16	26,84	2,34	29,19	0,135	28,2	172,7
Ap2	25-34	7,18	26,85	2,31	29,18	0,133	28,4	172,8
Ap3	25-34	7,19	26,86	2,33	29,17	0,134	28,3	172,8

### Field research method

Establishing an optimum fertilization system which positive influences the quantity and the quality of the culture and which implies an improving of the soil fertility potential represents an important way of increasing the economic efficiency. The influence of chemical fertilizers upon the maize culture is related to physiological processes which take place at a plant level in certain moments of its growth.

The experiment has in view two important factors:

Factor **A**: irrigating system

B1 – irrigated

B2 – non-irrigated

Factor **B**: applying the dosage of fertilizer

Within the experimental filed an important moment was followed, respective the 8 full-grown leaves phase, and lab tests were kept in order to perform physiological determinations.

Table 2

Influence of the nitrogen and phosphorus on some physiological processes of the crop hybrid Campion - 20 june non irrigated system 2012

Variant	Photosynthesis mgCO/cm <sup>2</sup>	Respiration mgCO <sup>2</sup> /100g m.v.	Absorption capacity g/apă	Suction force atm	Chlorophyll pigments			
					Chlorophyll A mg/dm <sup>2</sup>	Chlorophyll B mg/dm <sup>2</sup>	Caroten mg/dm <sup>2</sup>	Total mg/dm <sup>2</sup>
N <sub>0</sub> P <sub>0</sub>	228,6	183,6	5,89	4,1	0,363	0,168	0,199	0,728
N <sub>60</sub> P <sub>40</sub>	253,5	189,8	5,94	4,2	0,369	0,173	0,21	0,75
N <sub>80</sub> P <sub>60</sub>	256,6	201,4	5,87	4,13	0,368	0,175	0,205	0,746
N <sub>100</sub> P <sub>80</sub>	260,9	210,5	6,03	4,13	0,372	0,174	0,202	0,746
N <sub>120</sub> P <sub>100</sub>	253,5	206,6	5,95	4,17	0,377	0,172	0,207	0,753


Table 3

Influence of the nitrogen and phosphorus on some physiological processes of the crop hybrid Campion - 20 June irrigated system 2012

Variant	Photosynthesis mgCO/cm <sup>2</sup>	Respiration mgCO <sup>2</sup> /100g m.v.	Absorption capacity g/apă	Suction force atm	Chlorophyll pigments			
					Chlorophyll A mg/dm <sup>2</sup>	Chlorophyll B mg/dm <sup>2</sup>	Caroten mg/dm <sup>2</sup>	Total mg/dm <sup>2</sup>
N <sub>0</sub> P <sub>0</sub>	239,8	186,6	6,02	4,1	0,376	0,177	0,212	0,763
N <sub>60</sub> P <sub>40</sub>	262,5	192,4	6,11	4,19	0,39	0,179	0,219	0,777
N <sub>80</sub> P <sub>60</sub>	268,4	205,5	6,22	4,3	0,387	0,182	0,23	0,788
N <sub>100</sub> P <sub>80</sub>	270,6	215,8	6,16	4,3	0,389	0,188	0,228	0,803
N <sub>120</sub> P <sub>100</sub>	276,9	220,5	6,24	4,27	0,378	0,186	0,225	0,789

After lab determinations of physiological processes which took place in the case of this hybrid in two different systems and after applying different doses of fertilizers, the results were also graphically represented.

Influence of the nitrogen and phosphorus doses on some physiological processes of the crop hybrid Campion- 20 june year 2012



The chlorophyll pigments represented by 'a' and 'b' chlorophyll and carotene are very influenced by irrigated variants compared to non-irrigated ones (figure 1.)


An increase of the content of chlorophyll 'a' in the case of the irrigated system is noticeable, no matter what dose of fertilizer was applied, significant values are observed when applying N<sub>100</sub>P<sub>80</sub> and N<sub>120</sub>P<sub>100</sub>.

Chlorophyll 'b' has values that have a weak fluctuation no matter what dose of fertilizers was applied, an import factor in this case is the irrigating the area which led to a significant increase of this assimilative pigment.

The content of carotene has significant values towards the control, compared to variants where fertilizers were applied. Irrigating, when applying N<sub>100</sub>P<sub>80</sub> and N<sub>120</sub>P<sub>100</sub>, led to an important increase of this pigment.

Combined irrigation and applying moderate dose of fertilizers (N<sub>100</sub>P<sub>80</sub>) led to a maximum carotene quantity.

Influence of the nitrogen and phosphorus doses on some physiological processes of the crop hybrid Campion- 20 june year 2012



The evolution of photosynthesis (picture 2 and 3) expressed by mg CO<sub>2</sub>/dm<sup>2</sup> that was determined in the first moment, that is the 8 full-grown leaves, has an important increase when comparing the non-irrigated and the irrigated variants, and also within the same variant.

Thus, a 20% increase of the photosynthesis intensity towards the control in the case of irrigated variant towards the non-irrigated control is noticeable.

The optimum dose, as it is noticed in figure 2, was recorded in the case of the irrigated variant, when using N<sub>120</sub>P<sub>100</sub>, compared to the non-irrigated variant where better results were obtained when applying a dose of N<sub>100</sub>P<sub>80</sub>.

Regarding the respiration process, expressed as mg CO<sub>2</sub>/100 g.m.v., significant values on June, 20 are ascertain within the irrigated system as well as the non-irrigated system as well, the quantity of eliminated CO<sub>2</sub> being significant in the case of the studied variant.

The capacity of absorption, g/H<sub>2</sub>O, has minimum values in the case of non-irrigated variants, excepting the N<sub>100</sub>P<sub>80</sub> variant which has the highest value, that is 6.02 g/H<sub>2</sub>O.

Within the irrigated variants, the difference is important and it is static assured once the content of fertilizer in the case of N<sub>120</sub>P<sub>100</sub> variant increases.

The suction force (figure 2d), expressed as atmospheres, has a tendency of slow increase in the case of the first non-irrigated variants, after which it starts to increase and it reaches from 4 to 4.16 atmospheres. In the case of variants where irrigation was applied, the most evident suction force was registered when applying a N<sub>120</sub>P<sub>100</sub> dose.

## CONCLUSIONS

The combined influence between hybrid and culture technology (irrigation and applying the right dose of fertilizers) determines modifications of the main physiological processes which lead to obtaining maximum production when rationally applying dose of fertilizers.

The studied hybrid acted differently from the culture technology point of view, the registered values from the physiological point of view were different, each of them using the right quantity of fertilizer.

The effects of simultaneously applying variable doses of fertilizers and irrigation were studied and its effect upon the main physiological processes which depend on them.

All analyzed physiological processes lead us to the conclusion that the Champion hybrid has a high physiologic and biochemical potential, and the acquired information situates it among the most productive hybrids that have been studied.

The combined influence of culture system (irrigated non-irrigated) as well as the applied doses of fertilizers determined, from the quantitative point of view, remarkable differences in the case of all qualitative features of the Champion hybrid.

We highly recommend Champion hybrid for production, when using an irrigated system and applying a maximum dose of N<sub>120</sub>P<sub>100</sub> and N<sub>100</sub>P<sub>80</sub> for an economic efficiency.

## BIBLIOGRAPHY

1. Abu-Awwad A. M. (1999). *Irrigation water management for efficient water use in mulched onion*. Journal of Agronomy and Crop Science vol. 183.
2. Goian M. F. Sala, Adina Berbecea, Isidora Radulov, C. Gherban . (2000). *The effect of mineral fertilizers upon the bean production of maize, on cambic chernozem within the Timisoara Didactic Research Station, between 1997 and 1999*. Scientific research U.S.A.M.V. Timisoara, Agriculture Category. volume XXXII, part I. Agroprint Publishing House. Timisoara.
3. Hera C. (1972). *The influence of fertilizers upon certain maize hybrids sowed on different types of soils in Romania*. An. ICCPT. Fundulea XXXVII.
4. Marghidanu, N. *et al.* (1985). *Economia mecanizării agriculturii*. Editura Ceres. București.
5. Mărghitaș Marilena, M. Rusu. (2003). *Utilizarea îngrășămintelor în agricultură*. Ed. AcademicPres. Cluj-Napoca.
6. Oancea, I. (1998). *Tratat de tehnologii agricole*. Editura Tehnică. București.
7. Olimpia Pandia. (2006) *Fertilizer doses, different methods applied for establishing the quality of two hybrids Danubiu and Minerva*. Scientific research I.N.M.A.T.E.H.I. Bucharest.
8. Sărăcin, I. (2005). *Energetic Base for Agriculture. Transmissions*. Unversitaria Publishing House. Craiova.
9. Sima E. (2002). *Ecological Agriculture and Changing Perception in the Case of Quantity and Quality*. The Information and Economic Reference Material Center. Bucharest.