

López, Fernando A.; Angulo, Ana M.; Mur, Jesús; Artal Tur, Andrés

Conference Paper

The distribution of the economic activity in European region: identification of cluster

51st Congress of the European Regional Science Association: "New Challenges for European Regions and Urban Areas in a Globalised World", 30 August - 3 September 2011, Barcelona, Spain

Provided in Cooperation with:

European Regional Science Association (ERSA)

Suggested Citation: López, Fernando A.; Angulo, Ana M.; Mur, Jesús; Artal Tur, Andrés (2011) : The distribution of the economic activity in European region: identification of cluster, 51st Congress of the European Regional Science Association: "New Challenges for European Regions and Urban Areas in a Globalised World", 30 August - 3 September 2011, Barcelona, Spain, European Regional Science Association (ERSA), Louvain-la-Neuve

This Version is available at:

<https://hdl.handle.net/10419/120071>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.

The distribution of the economic activity in European regions: Identification of cluster

*Fernando López (**)*

Ana Angulo ()*

Jesús Mur ()*

*Andrés Artal(**)*

(*) Department of Economic Analysis.
University of Zaragoza
Gran Vía, 2-4. (50005). Zaragoza. Spain.
e-mails: aangulo@unizar.es, jmur@unizar.es

(**) Department of Quantitative Methods and Computing
Technical University of Cartagena.
Paseo Alfonso XIII, 50 - 30203 Cartagena. Spain.
e-mail: fernando.lopez@upct.es

Abstract

Mur et al (2008a) show that it is relatively simple to obtain symptoms of instability from a model with problems of heterogeneity in cross-sectional spatial model. In this sense, Lopez et al. (2009) propose the use of the local estimation for detecting such situation of instabilities. In the line of previous works, in this paper we try to analyse the capacity of spatial panel data models to deal with the problem of heterogeneity in spatial data. Furthermore, we try to asset whether or not the local estimation technique can be of help for the case of panel data models. We pay special attention to the forecast performance of several alternatives models. The empirical application refers to the explanation of employment in European Regional at NUTS II administrative level in terms of Eurostat. Panel data models are estimated on the basis of annual data (1980-2004) and data for 2005-2008 are gathered for evaluating the forecasting performance of the alternative models. The obtained results show that although panel data models are indeed designed for capturing the unobservable heterogeneity of data, the local estimation technique can also be of great help in the context of panel data models. From a forecasting point of view, the best model is the dynamic fixed effect with a spatial lag structure in the equation estimated through local estimation techniques.

Keywords: Dynamic spatial panel data models, Local estimation, European employment, forecast.

JEL Classification: C21; C22; C23; C53; R15

1. Introduction

The field of panel data models has received considerable attention during the last decade. Static panel data literature offers the opportunity of allowing for unobservable cross-sectional and time-period specific effects. In addition, dynamic panel data models also offer the possibility of considering the serial dependence between observations on each cross-sectional unit over time. Other advantages of panel data are that they are generally more informative and contain more variation and less collinearity between variables. The use of panel data leads to a greater availability of degrees of freedom and, hence, increases the efficiency of the estimation. Panel data also allow for the specification of more complicated behavioural hypotheses, including effects that cannot be addressed using pure cross-sectional or time-series data (Wooldridge, 2002; Arellano, 2003; Hsiao, 2003; Baltagi, 2005).

When cross-sectional data refers to spatial units (municipalities, provinces, regions or countries) the spatial dependence between cross-sectional units at each point in time is also important. Spatial dependence implies that, due to spillover effects (e.g., commuter labour and trade flows), neighbouring regions may have similar economic performance. Hence, we expect to improve traditional panel data models by paying attention to the location of the spatial units. There has been growing interest in the estimation of static panel data models with spatial dependence: see Kelejian and Prucha (2002), Elhorst (2003), Yang et al. (2006), Baltagi et al. (2006), Kapoor et al. (2006), Kelejian et al. (2006) or Pesaran (2006). Prediction with these types of models is analysed in Baltagi and Li (2004, 2006) for predicting per-capita cigarette and liquor consumption in the United States, respectively, and in Longhi and Nijkamp (2007) for forecasting the regional labour market in West German regions. The extension of the traditional dynamic panel data model to include spatial effects has been worked on by Elhorst (2005) and Su and Yang (2007), who have derived the ML estimator of a dynamic panel data model extended to include cross-sectionally correlated error terms; Elhorst (2008), who derived the ML estimator for a spatially lagged dependent variable model (endogenous interaction effects); and Korniotis (2005) and Yu et al. (2007), who considered a dynamic panel data model extended to include both endogenous and lagged endogenous interaction effects.

The main objective of this paper is to evaluate the performance of the dynamic spatial panel data models in the forecasting of regional series. In our application, we use

data on employment for 267 European regions (NUTS II administrative spatial unit in terms of Eurostat) from 1980 to 2008. The period 1980-2004 will allow us to estimate and check the models which, in a second step, will be used to forecast the series of employment by regions for the years 2005-2008. In this sense, our objective is similar to that of Kholodilin et al. (2008) when forecasting the GDP of German Länder although, in our paper, we use more efficient estimation techniques. Furthermore, a novelty in our paper is that we evaluate the capacity of local estimation techniques for capturing any type of heterogeneity still present after estimating any type of panel data models.

The structure of the present paper is as follows. In Section 2, we provide a description of the panel data models we consider in our application. Section 3 is devoted to the presentation of the data and the main estimation results. In Section 4, we present the forecast performance of the dynamic panel data models. Finally, the paper finishes with a section of concluding remarks.

2. Spatial data models

In this section, we describe a battery of models for panel data. We denote by R the number of spatial units (in our case, provinces) we observe as cross-sectional data ($i=1,2, \dots, R$) and T denotes the total number of observations in the time dimension ($t=1,2,\dots,T$). Let's start with the simple model, a *pooled* panel model:

$$\left. \begin{aligned} y_t &= x_t \beta + \eta_t \\ \eta_t &\sim N \left[0, \sigma_\eta^2 I_R \right] \end{aligned} \right\} \quad (1)$$

$$\text{with } y_t = \begin{bmatrix} y_{1t} \\ y_{2t} \\ y_{3t} \\ \vdots \\ y_{Rt} \end{bmatrix}; x_t = \begin{bmatrix} 1 & x_{21t} & \cdots & x_{k1t} \\ 1 & x_{22t} & \cdots & x_{k2t} \\ 1 & x_{23t} & \cdots & x_{k3t} \\ \vdots & \vdots & \cdots & \vdots \\ 1 & x_{2Rt} & \cdots & x_{kRt} \end{bmatrix}; \beta = \begin{bmatrix} \beta_1 \\ \beta_2 \\ \beta_3 \\ \vdots \\ \beta_k \end{bmatrix}$$

which imposes the homogeneity restriction on both the intercept and slope coefficients across all the regions.

However, model (1) does not consider the probably presence of cross-sectional dependence among the observations at each point in time. To this respect, recently Anselin et al. (2006) developed, in the context of panel data, the Lagrange Multiplier tests for a spatially lagged dependent variable, for spatial error correlation and their

counterparts robustified versions. When considering the spatial interaction among observations we obtain the so-called *spatial panel data* models, which mainly adopt two forms: i) if we introduce the spatially lagged dependent variable as an explicative variable, we obtain the Spatial Lag Model (SLM) version defined as:

$$\left. \begin{aligned} y_t &= \rho W y_t + x_t \beta + \eta_t \\ \eta_t &\sim N[0, \sigma_\eta^2 I_R] \end{aligned} \right\} \quad (2)$$

And, ii) if a spatially autorregressive process is incorporated into the error term, we obtain the Spatial Error Model (SEM), defined as:

$$\left. \begin{aligned} y_t &= x_t \beta + \varepsilon_t \\ \varepsilon_t &= \rho W \varepsilon_t + \eta_t \\ \eta_t &\sim N[0, \sigma_\eta^2 I_R] \end{aligned} \right\} \quad (3)$$

where W in equations (2) and (3) is the spatial weight matrix. As is well known, this matrix is pre-specified, nonnegative and of order $R \times R$ and describes the arrangement of the cross-sectional units in the sample (Anselin, 1988, 2007).

Models (2) and (3) consider the spatial interactions effects. However, they could also be improved when also considering the spatial specific effects, μ_i ($i=1,2,\dots,R$), in order to account for the heterogeneity among spatial units. In fact, these terms represent the effect of omitted variables that are space-specific time-invariant variables that affect the dependent variable but are difficult to measure or hard to obtain, obtaining the following specifications:

SLM + Spatial specific effects	SEM + Spatial specific effects
$\left. \begin{aligned} y_t &= \rho W y_t + x_t \beta + \mu + \eta_t \\ \eta_t &\sim N[0, \sigma_\eta^2 I_R] \\ \mu &= [\mu_1, \mu_2, \dots, \mu_R]' \end{aligned} \right\}$	$\left. \begin{aligned} y_t &= x_t \beta + \mu + \varepsilon_t \\ \varepsilon_t &= \rho W \varepsilon_t + \eta_t \\ \eta_t &\sim N[0, \sigma_\eta^2 I_R] \\ \mu &= [\mu_1, \mu_2, \dots, \mu_R]' \end{aligned} \right\}$

(4)

The spatial specific effects may be treated as fixed effects or as random effects. In the fixed effects model, a dummy variable is introduced for each spatial unit, while in the random effect model, μ_i is treated as a random variable that is independently and identically distributed with zero mean and variance σ_μ^2 . Furthermore, it is assumed that the random variable μ and ε_t are independent of each other. The random effects model

can be tested against the fixed effects model using Hausman's specification test (Baltagi, 2005). If the hypothesis is rejected, the random effects models must be rejected in favour of the fixed effects model. However, discussion about random or fixed effects goes further than the only use of the Hausman's specification test. In the context of spatial data, the situation may be summarized according to two different positions.

On the one hand, models including a spatial structure need a very large sample (a large R , number of regions, in our case), because the convergence results are obtained with R tending to infinite. But, on the other hand, if the omitted effects are non-random, a problem of incidental parameters appears (the number of parameters grows at the same rate as the number of observations); in that case, a large T and small R are preferable. The last observation leads Anselin et al. (2006) to discard the use of fixed effects in mechanisms of spatial dependence: *'Since spatial models rely on asymptotics in the cross-sectional dimension (...), this would preclude the fixed effects model from being extended with a spatial lag or spatial error term'*. These authors prefer the random effect framework, where the inference is conditional and we only need a very large R (the improvements with T are of minor importance).

Elhorst (2003) does not share that view when he states that: *'The spatial units of observation should be representative of a larger population, and the number of units should potentially be able to go to infinity in a regular fashion. Moreover, the assumption of zero correlation between μ_r and the explanatory variables is particularly restrictive. Hence, the fixed effects model is compelling, even when R is large and T is small'*.

Further improvement of the models could be obtained by introducing the serially lagged dependent variable for capturing the inertia present in the temporal data or, in other words, for taking into account the serial dependence between observations on each cross-sectional unit over time. By doing this we obtain the following dynamic specifications:

Dynamic SLM + Spatial specific effects	Dynamic SEM + Spatial specific effects
---	---

$$\left. \begin{array}{l} y_t = \tau y_{t-1} + \rho W y_t + x_t \beta + \mu + \eta_t \\ \eta_t \sim N[0, \sigma_\eta^2 I_R] \\ \mu = [\mu_1, \mu_2, \dots, \mu_R]' \end{array} \right\} \quad \left. \begin{array}{l} y_t = \tau y_{t-1} + x_t \beta + \mu + \varepsilon_t \\ \varepsilon_t = \rho W \varepsilon_t + \eta_t \\ \eta_t \sim N[0, \sigma_\eta^2 I_R] \\ \mu = [\mu_1, \mu_2, \dots, \mu_R]' \end{array} \right\} \quad (5)$$

Estimation of static panel data models are explained by Elhorst (2003) and dynamic spatial panel data models can be estimated as explained by Elhorst (2005, 2008).

3. Data and estimation

For our estimation and forecasting we use data on employment for each of the 217 European regions (NUTS II administrative spatial unit in terms of Eurostat). Employment will be explained by investment and remuneration. The three variables are gathered for the period 1980 to 2008 from the Cambridge Database. The spatial distribution of the employment in the total economy in four different years appears in Figure 1.

(Insert Figure 1)

In order to define the proper specification for the models described in Section 2, we start by analyzing the statistical properties of data for each region, in logarithms (to reduce heterogeneity). The evolution of the log of employment in level and in its first differences, for each region, is displayed in Figures 2 and 3, respectively.

(Insert Figures 2 and 3)

The non-stationarity of the variables is confirmed by the applications of the common tests for unit roots in panel data. From the results obtained, shown in Table 1, we can conclude that the series are integrated of order one. As a consequence, the dependent variable for all panel data models will be the first differences of the logarithm of employment; that is, we will explain the growth rate of employment. The similar approach applied to the explicative variables reaches us to consider as explicative variables the growth rates of both variables, investment and remuneration. Furthermore, a temporal lag of such explicative variables will be included in order to avoid simultaneity problems.

(Insert Table 1)

In order to estimate spatial models, we must specify one or several weight matrices to reflect the network of cross-sectional relationships in the system of regions. To this respect, we have decided to develop a mixed neighbourhood criterion, which consists of the following. In the first place, we have used a criterion of neighbourhood based on the distance between the centroids of the regions. Furthermore, to avoid situations of excessive imbalance, we have opted to qualify the distance criterion by also incorporating the r nearest neighbours. Thus, we define the binary matrix W^b as:

$$w_{ij}^b(k, r) = \begin{cases} \text{if } \min_{i \neq s} \{d_{is}\} > k \Rightarrow \begin{cases} w_{ij}^b(k, r) = 1 & \text{if } j \in N_r(i) \\ w_{ij}^b(k, r) = 0 & \text{if } j \notin N_r(i) \end{cases} \\ \text{if } \min_{i \neq s} \{d_{is}\} \leq k \Rightarrow \begin{cases} w_{ij}^b(k, r) = 1 & \text{if } d_{ij} \leq k \\ w_{ij}^b(k, r) = 0 & \text{if } d_{ij} > k \end{cases} \end{cases} \quad (6)$$

where d_{ij} is the distance in kilometres between the centroids of regions i and j , k the interaction radii and $N_r(i)$ the set of the r regions closest to region i . As usual, $w_{ii}=0$ for all i . In the paper we offer the results obtained for $k=600$ and $r=2$ but we have checked that results are very consistent with other values.

Firstly, we confirm the present of instability in the spatial cross-sectional model specified for several years. Results for the robust instability tests defined in Angulo et al (2008) derived from their counterpart non-robust versions obtained in Mur et. al (2008) are shown in Table 2. As expected, there are clear symptoms of instability in the three considered dimension: regression parameters, dispersion parameter and spatial dependence parameter. These results show that it is likely to improve specification of cross-sectional s with the use of panel data models that introduce spatial specific effect to consider the unobservable heterogeneity of the data.

(Insert Table 2)

The proper specification of spatial panel data model is derived through the estimation of alternative models. Results are gathered in Table 3.

(Insert Table 3)

Firstly, we estimate the pool model. However, according to the LM test for testing the null of no spatial effects (Anselin et al., 2006), the model suffers from misspecification, being a Spatial Lag Model the spatial structure that underline the data. The pooled SLM model (second column of results) outperform previous model, but

either the FE-SLM or the RE-SLM are better specifications. From Hausman's test the RE-SLM model cannot be rejected. However, we support Elhorst's point of view, since we believe that the unobservable effects (or the omitted variables they are representing) are probably correlated with our explanatory variables. Moreover, FE-SLM model present a higher R^2 value and therefore a higher goodness of fit. Nevertheless, we will also compare the forecast performance of both models in next section.

Finally, we introduce temporal dynamic into the FE-SLM model. Results show that this model outperforms the previous one. However, a centre-periphery pattern can still be observed in the residual terms of the model¹. As a consequence, a more flexible model is estimated by introducing a dummy variable (that takes the value of one for the periphery regions and zero value, otherwise) interacting with the two explicative variables. As deduced from results displayed in Table 3, this last model represent, till now, the best specification for our data set.

The question that we try to answer now it is to what respect all heterogeneity present initially in the data has been gathered with our flexible dynamic FE-SLM panel data model with centre-periphery interacted dummy. In order to solve this question, we propose a simple exercise consisting of using the local estimation technique with the reference model Dynamic FE+SLM. If results from local estimation technique outperform previous ones, we can conclude that spatial panel data models cannot capture all heterogeneity inherent in spatial data and therefore, they can be benefit from the use of local estimation techniques.

Briefly stated, the local estimation technique consists of fitting individual regressions to selected points in the sample, with more weight assigned to observations that are closer to the point of interest (McMillen 1996). Repeating this exercise for every point in the sample, we can construct estimation surfaces in order to discuss the nonstationarity of each parameter in the model. The concept of '*closeness*' is flexible and must be adapted to the objectives of the study. Moreover, the distribution of the weights among the neighboring observations with respect to point r is determined by the kernel function (Cressie 1991). In the case of the GWR, this is a decreasing function of the distance between the points, and the bandwidth determines how rapidly the weights decline with distance. We decided to use a rectangular or uniform kernel with a fixed

¹ Results are available from the authors.

bandwidth of m for every point. This means that the m nearest neighbors will receive a weight of one, and the other points zero.

In our case we have to resolve the local estimation of an Dynamic FE+SLM for which it is not advisable to use the OLS algorithm. Following the example of Brunson et al. (1998) and of Pace and Lesage (2004), we will obtain the local estimators from the ML estimation of the local model:

$$\left. \begin{aligned} y_{t,r}^{(m)} &= \tau_r^{(m)} y_{t-1,r}^{(m)} + \rho_r^{(m)} W_r^{(m)} y_{t,r}^{(m)} + x_{t,r}^{(m)} \beta_r^{(m)} + u_r^{(m)} + \eta_{t,r}^{(m)} \\ \eta_{t,r}^{(m)} &\sim N[0, \sigma_{\eta,m}^2 I_m] \\ \mu_r^{(m)} &= [\mu_1, \mu_2, \dots, \mu_m]' \end{aligned} \right\} \quad (7)$$

The indexes r and m mean that the data correspond to the local system defined by m elements around point r . Therefore, $y_{t,r}^{(m)} = (y_{t,r}, y_{t,i_1}, y_{t,i_2}, \dots, y_{t,i_{m-1}})$ where $i_k \in N(r)$, being $N(r)$ the bundle of indexes of the $m-1$ neighbours nearest to the point r . The same criterion is used to define $x_{t,r}^{(m)}$. Matrix $W_r^{(m)}$ refers to the weighting matrix obtained for this local system, defined with the same connectivity criteria that are used to obtain the global W matrix, specified following standard criteria. Finally $\rho_r^{(m)}$, $\beta_r^{(m)}$ and $\sigma_{\eta,m}^2$ are the local parameters of interest. This is what we call the *Zoom* estimation (different to the SALE algorithm of Pace and Lesage, 2004, in that, in each local system, we use the matrix $W_r^{(m)}$ specific for the local network around point r). We refer to m as the *Zoom size* (equivalent to *window size* in nonparametric literature).

Results for the estimated parameters through the Local estimation techniques is shown in Figure 4. As can be observed, there exists important differences among regions not only in magnitudes but also, in the cases of parameters associated to remuneration and investment, even in sign.

(Insert Figure 4)

Finally, in the next section, we analyze the forecast performance of all the spatial panel data models.

4. Forecast performance of the different models

The purpose of this section is to obtain and evaluate the employment forecast for all the regions for the period 2005-2008 derived from the different spatial panel data models.

Godberger (1962) shows that the best linear unbiased predictor (BLUP) for the cross-sectional units in a linear regression model $Y = X\beta + \eta$ with disturbance covariance matrix Ω at a future period $T+C$ is given by:

$$\widehat{Y_{T+C}} = X_{T+C}\hat{\beta} + \Psi'\Omega^{-1}e \quad (8)$$

where $\Psi = E(\eta_{T+C}\eta)$ is the covariance between the future disturbance η_{T+C} and the sample disturbance η , X represents the independent variables of the model, $\hat{\beta}$ is the estimator of β , and e denotes the residual vector of the model. Elhorst (2009) derived the prediction formulas for the fixed effects and random effects model with a spatially lagged dependent variable.

Formulas for the fixed effect models are straightforward as $\Psi = 0$ provided that error terms are not serially correlated over time. Hence, predictions for the FE+SLM model can be derived as:

$$\widehat{Y_{T+C}} = (I_R - \hat{\rho}W)^{-1} (X_{T+C}\hat{\beta} + \hat{\mu}) \quad (9)$$

and prediction for all variants of such model can be derived by defining the X matrix accordingly.

Unlike the fixed effects model, the correction term $\Psi'\Omega^{-1}e$ in the random effect model is not zero. In the random effects spatial lag model, RE-SLM, predictions can be calculated as follows:

$$\widehat{Y_{T+C}} = (I_R - \hat{\rho}W)^{-1} \left[X_{T+C}\hat{\beta} + (1 - \hat{\theta}^2) \left(\frac{1}{T} \sum_{t=1}^T \begin{pmatrix} e_{1t} \\ \dots \\ e_{Nt} \end{pmatrix} \right) \right] \quad (10)$$

$$\text{where } (1 - \hat{\theta}^2) = \frac{T\sigma_\mu^2}{T\sigma_\mu^2 + \sigma_\eta^2}$$

That is, for the RE-SLM model to calculate the correction term $\Psi'\Omega^{-1}e$, the residual of each spatial unit are first averaged over the sample period and then multiplied with $(1 - \hat{\theta}^2)$, a factor that can take values between 0 and 1.

Forecast performance of each spatial panel data model is evaluated through the Mean Absolute Percent Error (MAPE), which for each time period t (forecast horizon equal to $t-2004$) is defined as:

$$\text{MAPE}_t = \frac{1}{N} \sum_{r=1}^N \frac{\left| \text{employment}_{r,t} - \widehat{\text{employment}}_{r,2004}(t-2004) \right|}{\text{employment}_{r,t}} * 100 \quad (11)$$

$t = 2005 - 2008$

Table 4 shows figures for the different MAPE quantitative magnitudes while Figure 5 shows the spatial distributions of the temporal mean of the absolute percent prediction errors.

(Insert Table 4 and Figure 5)

Results from Table 4 shows that predictions from the FE+SLM model clearly outperform that obtained from the RE+SLM model. Hence, these results confirm our intuitive decision on the preference of fixed effects over random effects. As regards as results of the other more ample FE+SLM models considered, we have conclude the following. Both, the dynamic FE+SLM model and its extension (when we incorporate the structural break associated to the centre-periphery situation of regions) offer better results in terms of the MAPE. However, the local estimation dynamic FE+SLM referred to spatial units outperforms previous results. Consequently, we can conclude that zoom-estimation also can be of help in the context of panel data in order to capture the remaining heterogeneity of models. Furthermore, we can observe that improvement is especially important as the prediction horizon increases.

Finally, we analyse the possibility of further improvement in predictions with the application of local estimation referred not only to the spatial unit but also to the time dimension. That is, whether or not we can forecast better a region when using not only its neighboring spatial units but only its most recent observations in time. The obtained forecast results are shown in the last column of the Table (Dynamic FE+SLM+spatial and time zoom). From the obtained magnitudes we can conclude that Mean Absolute Prediction Error are larger than for the previous model and hence, in this particular case, the use of the zoom estimation associated to the cross-sectional dimension of data is the best option.

5. Concluding remarks

Econometric literature clearly accepts the good performance of panel data models for being able to capture the unobservable heterogeneity of data. The empirical application offered in this paper has shown clear evidence on the fact that forecast results of flexible panel data models can still be improved by making use of the local estimation techniques.


Further research will be directed towards the evaluation of the effect on results of certain decision about the size of the spatial or time zoom, as well as the possible effect of the selected functional form of the base model.

References

- Anselin L (1988) Spatial econometrics: methods and models. Kluwer, Dordrecht
- Anselin L (2007) Spatial Econometrics. In: Mills TC, Patterson K (eds) Handbook of econometrics, Vol. 1. Palgrave, Basingstoke, pp 901-969
- Arellano M (2003) Panel data econometrics. Oxford University Press, Oxford
- Arellano M, Bond (1991) Some tests of specification for panel data: monte carlo evidence and application to employment equations. *Review of Economic Studies* 58: 277-297
- Aznar A, Trávez FJ (1993) Métodos de predicción en Economía. Ariel, Barcelona
- Baltagi BH (2005) *Econometric Analysis of Panel Data* (3rd Edition). Wiley, Chichester
- Baltagi BH, Bresson G, Griffin JM, Pirotte A (2003) Homogeneous, heterogeneous or shrinkage estimators? some empirical evidence from French regional gasoline consumption. *Empirical Economics* 28: 795-811
- Baltagi BH, Bresson G, Pirotte A (2002) Comparison of forecast performance for homogeneous, heterogeneous and shrinkage estimators: some empirical evidence from US electricity and natural gas. *Economics Letters* 76: 375-382
- Baltagi BH, Bresson G, Pirotte A (2004) Tobin q: forecast performance for hierarchical Bayes, shrinkage, heterogeneous and homogeneous panel data estimators. *Empirical Economics* 29: 107-113
- Baltagi BH, Griffin JM (1997) Pooled estimators vs. their heterogeneous counterparts in the context of dynamic demand for gasoline. *Journal of Econometrics* 77: 303-327
- Baltagi BH, Griffin JM, Xiong W (2000) To pool or not to pool: homogeneous versus heterogeneous estimations applied to cigarette demand. *Review of Economics and Statistics* 82: 117-126
- Baltagi BH, Li D (2004) Prediction in the panel data model with spatial correlation. In: Anselin L, Florax RJGM, Rey S (Eds) *Advanced in spatial econometrics: methodology, tools and application*. Springer-Verlag, Heidelberg (Germany), pp 283-295
- Baltagi BH, Li D (2006) Prediction in the panel data model with spatial correlation: the case of liquor. *Spatial Economic Analysis* 1(2): 175-185
- Baltagi BH, Song SH, Jung BC, Koh W (2006) Testing for serial correlation, spatial autocorrelation and random effects using panel data. *Journal of Econometrics* 140: 5-51
- Brücker H, Siliverstovs B (2006) On the estimation and forecasting of international migration: how relevant is heterogeneity across countries?. *Empirical Economics* 31: 735-754
- Burridge P (1981) Testing for a common factor in a spatial autoregression model. *Environment and Planning A* 13: 795-800
- Box G, Jenkins GM, Reinsel G (2008) *Time Series Analysis: Forecasting and Control*. 4th Edition. John Wiley and Sons, New York

- Elhorst JP (2003) Specification and Estimation of Spatial Panel Data Models. *International Regional Sciences Review* 26: 244-268
- Elhorst JP (2005) Unconditional maximum likelihood estimation of linear and log-linear dynamic models for spatial panels. *Geographical Analysis* 37: 62-83
- Elhorst JP (2008) Estimation of dynamic panels with endogenous interaction effects. II Wold Conference of the Spatial Econometrics Association. New York
- Hamilton JD (1994) *Time Series Analysis*. Princeton University Press, Princeton
- Hsiao C (2003) *Analysis of panel data* (2nd Edition). Cambridge University Press, Cambridge
- Hsiao C, Pesaran MH, Tahmiscioglu AK (2002) Maximum Likelihood Estimation of Fixed Effects Dynamic Panel Data Models Covering Short Time Periods. *Journal of Econometrics* 109: 107-150
- Instituto Nacional de Estadística, INE (several years). *Encuesta de población activa*. Madrid
- Kapoor M, Kelejian HH, Prucha IR (2006) Panel data models with spatially correlated error components. *Journal of Econometrics* 140: 97-130
- Kelejian HH, Prucha IR (2002) 2SLS and OLS in a spatial autoregressive model with equal spatial weights. *Regional Science and Urban Economics* 32: 691-707
- Kelejian HH, Prucha IR, Yuzefovich Y (2006) Estimation problems in models with spatial weighting matrices which have blocks of equal elements. *Journal of Regional Science* 46: 507-515
- Kholodilin KA, Siliverstovs B, Kooths S (2008). A dynamic panel data approach to the forecasting of the GDP of German Länder. *Spatial Economic Analysis* 3(2): 195-207
- Korniotis GM (2005) A dynamic panel estimator with both fixed and spatial effects. <http://www.uni-kiel.de/ifw/konfer/spatial/korniotis.pdf>
- Layton A, Defris L, Zehnwrith B (1986) An International Comparison of Economic Leading Indicators of Telecommunication Traffic. *International Journal of Forecasting* 2: 413-425
- Longhi S, Nijkamp P (2007) Forecasting regional labor market developments under spatial heterogeneity and spatial correlation. *International Regional Science Review* 30: 100-119
- Magnus JR, Neudecker H (1988) *Matrix differential calculus with applications in statistics and econometrics*. John Wiley, Chichester
- Nerlove M (1999) Properties of Alternative Estimators of Dynamic Panel models: An Empirical Analysis of Cross-Country Data for the Study of Economic Growth. In: Hsiao C, Lahiri K, Lee LF, Pesaran MH (eds) *Analysis of Panels and Limited Dependent Variable Models*. Cambridge University Press, Cambridge
- Nerlove M, Balestra P (1996) Formulation and estimation of econometric models for panel data. In: Mátyás L, Sevestre P *The econometrics of panel data*, 2nd revised edition. Kluwer, Dordrecht (The Netherlands)
- Nickell S (1981) Biases in dynamic models with fixed effects. *Econometrica* 49: 1417-1426
- Pesaran MH (2006) Estimation and inference in large heterogeneous panels with a multifactor error structure. *Econometrica* 74: 967-1012
- Sevestre P, Trognon A (1996) Dynamic linear models. In: Mátyás L, Sevestre P *The econometrics of panel data*, 2nd revised edition. Kluwer, Dordrecht (The Netherlands)
- Su L, Yang Z (2007) QML estimation of dynamic panel data models with spatial errors. <http://fp.paceprojects.f9.co.uk/Su.pdf>
- Wooldridge J (2002) *Econometric Analysis of Cross Section and Panel Data*. The MIT Press, Cambridge
- Yang Z, Li C, Tse YK (2006) Functional form and spatial dependence in spatial panels. *Economics Letters* 91: 138-145
- Yaffee RE, McGee M (2000) *Introduction to Time Series Analysis and Forecasting*. Academic Press, Inc. San Diego
- Yu J, de Jong R, Lee L (2007) Quasi-maximum likelihood estimators for spatial dynamic panel data with fixed effects when both n and T are large. *Journal of Econometrics* 146(1):118-134
- Zou H, Yang Y (2004) Combining Time Series Models for Forecasting. *International Journal of Forecasting* 20-69-84

Figure 1. Spatial distribution of the employment



In all cases:




Figure 2. Evolution of the log(employment) in each of the 217 cross-sectional units (1980-2008)




Figure 3. Evolution of the $\Delta\log(\text{employment})$ in each of the 217 cross-sectional units (1980-2008)




Figure 4. Spatial distribution of zoom parameters




Figure 5. Spatial distribution of the temporal mean of the absolute percent predictions errors




Table 1. Unit root test, under the assumption of specific mean and trend for each province

	Log (employment)		Log (investment)		Log (remuneration)	
	H₀: I(1) H₁: I(0)	H₀: I(2) H₁: I(1)	H₀: I(1) H₁: I(0)	H₀: I(2) H₁: I(1)	H₀: I(1) H₁: I(0)	H₀: I(2) H₁: I(1)
	Statistic (Prob.)	Statistic (Prob.)	Statistic (Prob.)	Statistic (Prob.)	Statistic (Prob.)	Statistic (Prob.)
H₀: Unit root (equal for all cross-sections)						
Levin, Lin & Chu (2002), t*	0.33 (0.63)	-23.09 (0.00)	-3.37 (0.00)	-27.67 (0.00)	-3.38 (0.00)	-33.9 (0.00)
Breitung	0.55 (0.71)	-17.39 (0.00)	-5.30 (0.00)	-28.77 (0.00)	-11.74 (0.00)	-24.36 (0.00)
H₀: Unit root (specific for each cross-sections)						
Im, Pesaran and Shin W-stat	7.83 (1.00)	-30.18 (0.00)	6.76 (1.00)	-37.42 (0.00)	2.76 (0.99)	-35.68 (0.00)
ADF - Fisher χ^2	360 (1.00)	1918 (0.00)	369 (1.00)	2403 (0.00)	492 (0.90)	2282 (0.00)
PP - Fisher χ^2	245 (1.00)	2917 (0.00)	316 (1.00)	4038 (0.00)	546 (0.35)	3528 (0.00)
Conclusion	I(1)		I(1)		I(1)	

Table 2. Cross-sectional spatial dependence and instability tests

	1982	1995	2000	2004
Moran	6.84*	4.76*	10.67*	9.75*
Lmerr	34.38*	17.12*	92.16*	76.99*
Lmlag	25.33*	20.08*	98.44*	84.39*
Robust Lmerr	10.03*	0.68	0.06	0.07
Robust Lmlag	0.98	4.64*	6.35*	7.47*
Sarma	35.36*	20.76*	98.51*	84.46*
LM ^{SLM(SEM)} _{Break-Chow-Het}	32.88*	90.69*	645.69*	628.12*
LM ^{SLM(SEM)*} _{Break}	8.48*	15.89*	27.51*	32.38*
LM ^{SLM(SEM)*} _{Het}	17.97*	67.71*	542.34*	532.04*
LM ^{SLM(SEM)*} _{Chow}	13.90*	22.76*	84.85*	82.03*
LM ^{SLM(SEM)*} _{Break-Het}	18.97*	67.93*	560.84*	546.09*
LM ^{SLM(SEM)*} _{Break-Chow}	22.39*	38.66*	112.37*	114.42*
LM ^{SLM(SEM)*} _{Chow-Het}	31.87*	90.47*	627.20*	614.07*

Table 3: Results obtained for the estimation of the different models

	OLSQ	SLM	RE+SLM	FE+SLM	Dynamic FE+SLM (a)	Dynamic FE+SLM With C-P interacting dummy (a)
CONSTANT	0.002* (4.89)	0.000 (0.33)	0.000 (0.15)	0.000 (0.00)	0.000 (0.00)	0.000 (0.00)
$\Delta \text{LOG}(\text{INVESTMENT})_{t-1}$						
All	0.037* (10.60)	0.016* (5.37)	0.015 (5.09)	0.015* (4.93)	0.018* (2.92)	
Centre						0.007 (0.95)
Periphery- Centre						0.043 (3.02)
$\Delta \text{LOG}(\text{REMUNERATION})_{t-1}$						
All	0.001 (0.17)	0.044 (1.18)	0.006 (1.78)	0.008* (2.32)	-0.004 (-0.54)	
Centre						-0.008 (-0.83)
Periphery- Centre						-0.001 (-0.06)
SPATIAL TERM: ρ		0.779* (51.79)	0.755 (46.61)	0.731* (42.22)	0.01* (11.53)	0.01* (11.53)
TIME DYNAMIC: τ					0.282* (11.03)	0.279* (10.89)
R^2	0.02	0.29	0.33	0.36	0.40	0.42
LM for no spatial error	3367.7*					
LM for no spatial lag	3500.4*					
Robust LM for no spatial error	21.89*					
Robust LM test for no spatial lag	154.64*					
LR for testing the null of no specific regional effects				633.3*		
Hausman Test			2.588			

^(a) Dynamic panel data model must be estimated with a symmetric weight matrix, W. The range of ρ is in this case of (-0.091;0.017)

Table 4. Forecast performance of the different models, Mean of the Absolute Percent Error (MAPE), (%)

	RE+SLM	FE+SLM	Dynamic FE+SLM	Dynamic FE+SLM With regional dummies	Dynamic FE+SLM + spatial zoom	Dynamic FE+SLM + spatial and time zoom
MAPE(2005)	1.791	1.638	1.480	1.482	1.263	1.556
MAPE (2006)	2.954	2.696	2.512	2.437	1.974	2.726
MAPE (2007)	4.171	3.821	3.662	3.541	2.889	4.137
MAPE (2008)	4.937	4.323	4.155	4.006	3.356	5.126
MAPE	3.463	3.120	2.952	2.866	2.370	3.386

(a) EAPM(t) denote de mean of the absolute percent error for predicting the

employment of period t in all the regions, $EAPM(t) = \frac{\sum_{i=1}^N EAPM_i(t)}{N}$. The term EAPM denotes the mean over the four predictions.