

Andersen, Lykke E.; Reis, Eustáquio J.

Working Paper

Carbon emissions from deforestation in the Brazilian Amazon

Documento de Trabajo, No. 02/00

Provided in Cooperation with:

Instituto de Investigaciones Socio-Económicas (IISEC), Universidad Católica Boliviana

Suggested Citation: Andersen, Lykke E.; Reis, Eustáquio J. (2000) : Carbon emissions from deforestation in the Brazilian Amazon, Documento de Trabajo, No. 02/00, Universidad Católica Boliviana, Instituto de Investigaciones Socio-Económicas (IISEC), La Paz

This Version is available at:

<https://hdl.handle.net/10419/145403>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.



Instituto de Investigaciones Socio Económicas

Documento de Trabajo No. 02/00
Marzo 2000

**Carbon Emissions from Deforestation in the
Brazilian Amazon**

por
Lykke E. Andersen,
& Eustáquio J. Reis

Carbon Emissions from Deforestation in the Brazilian Amazon

Eustáquio J. Reis
Institute for Applied Economics Research
Rio de Janeiro, Brazil

Lykke E. Andersen^{*}
Institute for Socio-Economic Research
Universidad Católica Boliviana
La Paz, Bolivia
E-mail: landersen@ucb.edu.bo

March 14, 2000

Abstract

This paper provides an estimate of the amount of carbon emissions that resulted from large scale deforestation in the Brazilian Amazon during the aggressive development period 1970 - 1985.

The estimate is derived by combining a dynamic carbon model with a municipality level data set describing natural vegetation cover and land uses in 316 municipalities in Legal Amazonia.

The results show average carbon emissions from Legal Amazonia of 168 million tons per year during the period 1970-1985, or about 95 tC per hectare cleared. This is lower than other estimates found in the literature because we take into account the uneven spatial distribution of clearing.

The aggressive development policies implied rapid economic growth in the region. On average, there was generated GDP of approximately \$44 per ton carbon emitted.

JEL classification: Q23, Q28, R12, R14.

Keywords: Amazon deforestation, carbon emissions.

^{*} This research was **carried out while the second author** was visiting the Institute for Applied Economics Research in Rio de Janeiro. The hospitality of the institute is gratefully acknowledged as is the financial support received from the NEMESIS/PRONEX project.

1. Introduction

Deforestation has costs at both local and international levels, and one of the biggest international concerns seems to be its contribution to global warming through the release of the carbon that used to be stored in the biomass of the forests. Since Brazil contains vast amounts of forest and has been deforesting rapidly in the name of development, it is perceived as one of the main contributors to global warming. Deforestation in the Brazilian Amazon alone has been estimated to account for 1 to 9% of total CO₂ emissions (Schroeder & Winjum 1994; World Bank 1991; Brown & Lugo 1992).

It has been argued that a reduction in tropical deforestation rates would be a relatively cheap way of curbing global CO₂ emissions compared to the cost of reducing fossil fuel consumption in the industrialized world (Nordhaus 1991; Schneider 1993; Kolk 1996; FACE 1993). To assess whether this is a reasonable argument one would need better estimates of carbon emissions from deforestation as well as better estimates of the benefits of deforestation. This paper attempts to provide such estimates for the Brazilian Amazon by the use of detailed municipality level data for the period 1970—1985. This period was characterized by the implementation of unusually aggressive development policies in the region, resulting in rapid deforestation as well as rapid economic growth.

The remainder of the paper is organized as follows. Section 2 describes the carbon emissions model and the parameter values used for simulations. Section 3 describes the data on original vegetation cover and changes in land use and summarizes the estimated age structure of deforested land. Section 4 summarizes deforestation measures and the calculated carbon emissions and compares these to the economic activity generated by the aggressive policies. Section 5 discusses factors contributing to uncertainty in the carbon estimates and makes several sensitivity analyses. Section 6 concludes.

2. The carbon emission model

A typical dense forest of the Brazilian Amazon contains a biomass of about 350 tons per hectare with a dry weight carbon content of approximately 50%, i.e. about 175 tC/ha. Typical agricultural land, on the other hand, contains less than 10 tC/ha. Thus, when forests are converted to agricultural land uses, massive amounts of carbon is released to the atmosphere, thus contributing to the increasing concentration of CO₂ in our atmosphere.

Houghton *et al* (1983) has developed a carbon emissions model that keeps track of the carbon that was originally stored in the biomass of the forest when the forest is converted to agricultural land use through slash-and-burn methods.

Various versions of this model has been used by Schroeder & Winjum (1994a, 1994b) and Reis (1996) to calculate carbon emissions from changes in land use in Brazil.

The carbon model has two important parts. First, a set of equilibrium values for the carbon content in different types of forests and in different types of land uses. Second, a set of response functions that indicate how carbon is decomposed and recomposed after changes in land use. The model is a bookkeeping type of model, and is thus purely mechanical. The parameters used to run the model are compiled from the literature.

2.1. Carbon contents

Table 1 presents central¹ estimates of carbon contents in different vegetation types and for different land uses in the Amazon². The dry weight carbon content is everywhere assumed to be 50% of biomass (Fearnside *et al* 1993; Brown & Lugo 1984).

Aboveground estimates of carbon content is taken from Reis (1996) which is based on Bohrer's (1993) survey. Belowground carbon content is assumed to be 17% of above ground carbon content for dense forest, seasonal forest, and campinarana (Reis 1996; Brown & Lugo 1992; Schroeder & Winjum 1994; Fearnside 1992b). For savanna and ecological transition (cerrado woodland) belowground carbon content is assumed to be 160% of aboveground carbon content (Schroeder & Winjum 1994a; Singh & Joshi 1979). For wetlands this ratio is assumed to be 75% (Schroeder & Winjum 1994a; Cannell 1982).

Reis (1996) assumes that all secondary vegetation is identical and reaches a modest climax carbon content of 43.9 tC/ha after 40 years. This is a very conservative estimate which results in a worst case scenario for carbon emissions. In this paper we allow for different kinds of secondary vegetation and more realistic climax values. Following Houghton *et al* (1983), we assume that secondary vegetation in dense forest, seasonal forest, and campinarana areas reaches 75% of the original carbon content after 50 years. Secondary vegetation in savanna, wetlands, and ecological transition areas is assumed to reach 100% of original carbon content after 50 years (Houghton *et al* 1983).

For agropastoral land use (crops and planted pasture) it is assumed, according to Sanchez *et al* (1989), that belowground carbon content is 18% of aboveground carbon content.

Table 1: Carbon contents for different vegetation types and land uses

| Vegetation type/ land use | Above-ground ^a (tC/ha) | Below-ground (tC/ha) | Total (tC/ha) |
|---------------------------------------|--------------------------------------|-------------------------|------------------|
| <i>Primary vegetation:</i> | | | |
| Dense forest (62.5%) | 150.0 | 25.5 | 175.5 |
| Seasonal forest (6.1%) | 93.0 | 15.8 | 108.8 |
| Savannas (15.4%) | 37.5 | 60.0 | 97.5 |
| Ecological transition (7.3%) | 65.0 | 104.0 | 169.0 |
| Wetlands (2.5%) | 57.5 | 43.1 | 100.6 |
| Campinarana (5.4%) | 60.0 | 10.2 | 70.2 |
| <i>Secondary vegetation: (climax)</i> | | | |
| Dense forest | 112.5 | 19.1 | 131.6 |
| Seasonal forest | 69.7 | 11.9 | 81.6 |
| Savannas | 37.5 | 60.0 | 97.5 |
| Ecological transition | 65.0 | 104.0 | 169.0 |
| Wetlands | 57.5 | 43.1 | 100.6 |
| Campinarana | 45.0 | 7.7 | 52.7 |
| Agropastoral land use | 5.0 | 0.9 | 5.9 |

Source: Reis (1996) + assumptions in text.

a: Including litter.

¹ Reis (1996, Table 3), in contrast, presents worst case numbers in the sense that carbon contents for natural vegetation are from the upper end of the range found in the literature, while carbon contents for agropastoral land and secondary forests are in the low end.

² Carbon contents vary greatly from plot to plot and from region to region. This has resulted in a wide variation in estimate of average carbon contents. See Brown & Lugo (1984), Fearnside (1986, 1987), Brown *et al* (1989), Brown & Lugo (1992), and Fearnside (1992) for a debate about biomass content in the Brazilian Amazon.

2.2 Carbon Dynamics

Typical slash-and-burn agriculture in the Amazon involves three phases: i) land clearing and forest burning, ii) agropastoral use of land, iii) soil exhaustion, land abandonment, and the recomposition of secondary vegetation. Figure 1 illustrates the carbon cycle of vegetation in this pattern of agricultural settlement.

Carbon emissions from deforestation can usefully be divided into the three phases identified for slash-and-burn agriculture. Evidence from Seiler and Crutzen (1980), Fearnside (1992), Fearnside *et al* (1993), Carvalho Jr. *et al* (1994), and Houghton *et al* (1991) suggests that about 30% of aboveground carbon is released in the first phase of forest burning, while belowground carbon is released in this phase. In the second phase, the remaining aboveground carbon plus belowground carbon is slowly decomposed. It is assumed that the decomposition follows an exponential decay function:

$$cav_{v,i} = 0.7 \cdot cav_{v,0} \cdot e^{-r_v \cdot i}$$


where $cav_{v,i}$ is aboveground carbon content in the original vegetation of type v , i periods after burning, $cav_{v,0}$ is aboveground carbon content in original vegetation type y before burning, r_v is the rate of carbon decomposition after burning un vegetation type y , and i is time elapsed since forest burning.

For belowground carbon we do not observe any immediate decline during forest burning, so the decay function for belowground carbon in natural vegetation becomes:

$$cbv_{v,i} = cbv_{v,0} \cdot e^{-r_v \cdot i}$$

where $cbv_{v,i}$ is belowground carbon content in the original vegetation of type v , i periods after burning,

Figure 1: Carbon contents during a slash-and-burn cycle



Reis (1996) uses the same decay rates for above and belowground carbon in which case the total carbon content in original vegetation becomes:

$$cv_{v,i} = cav_{v,i} + cbv_{v,i} = (0.7 \cdot cav_{v,0} + cbv_{v,0}) \cdot e^{-r_v \cdot i} \quad (1)$$

Houghton *et al* (1991) estimate rates of carbon decomposition of 0.5 for closed forests and 0.3 for savannas. Following Reis (1996) we assume a rate of 0.4 for other vegetation types. These rates imply that 94% of aboveground carbon in dense forests is released within 5 years, with the corresponding numbers for savanna and other vegetation types being 84%

and 91%, respectively. Within 15 years more than 99% of aboveground carbon will be emitted for all vegetation types.

In the third phase there will be a recomposition of biomass and carbon stock as secondary species invade the abandoned land. Recomposition will be relatively rapid in the beginning due to the fast growth of pioneer species (Uhl *et al* 1988, Uhl 1987, Houghton *et al* 1983) but will then slow down and converge to a new climax. Odum (1988) suggest the following specification for the recomposition of biomass and carbon content in abandoned areas:

$$cf_{v,i} = \frac{cm_v}{1 + e^{(a_v - s_v \cdot i)}}, \quad (2)$$

where $cf_{v,i}$ is carbon content (both above- and belowground) in fallow land of age i on vegetation type v , cm_v is the climax carbon content of vegetation type v , and a_v and s_v are parameters to be estimated.

To estimate the parameters a_v and s_v , we make the following assumptions: i) The carbon content at the time of abandonment is equal to that of agropastoral land use, i.e. $cf_{v,0} = 5.9$ tC/ha for ah vegetation types. ii) Fallow lands take 50 years to reach 99% of their climax carbon content.

The climax carbon contents of different vegetation types are given in Table 1. When combined with the assumptions above they yield the following values for a_v and s_v :

Table 2: Decay parameters

| Vegetation type | a_v | s_v |
|-----------------------|--------|--------|
| Dense forest | 3.0589 | 0.1530 |
| Seasonal forest | 2.5518 | 0.1429 |
| Savanna | 2.7425 | 0.1468 |
| Ecological transition | 3.3194 | 0.1583 |
| Wetlands | 2.7758 | 0.1474 |
| Campinarana | 2.0701 | 0.1333 |

2.3 Carbon Stocks

The carbon stock in a given municipality at time t is the sum of the carbon stock in original vegetation, CV_t , the carbon stock in crops, CK_t and the carbon stock in fallow lands, CF_t :

$$C_t = CV_t + CK_t + CF_t$$

The carbon stock in original vegetation consists of two parts. First, the carbon stock in undisturbed areas, and second the decaying carbon stock in deforested areas:

$$CV_t = \sum v_{v,0} \cdot (A_{v,0} - D_{v,t}) + \sum_v \sum_i cv_{v,i} \cdot A_{v,i,t}^c,$$

where $A_{v,0}$ is area of vegetation type v at time 0, $D_{v,t}$ is deforested area on vegetation type v at time t , $A_{v,i,t}^c$ is crop area of age i on vegetation type v at time t , and $CV_{v,i}$, the decreasing carbon contents of deforested natural vegetation given by Equation 1.

The carbon content in crops is assumed to be identical for ah crops (including planted pasture), so the carbon stock in crops becomes:

$$CK_t = ck \cdot \sum_v \sum_i A_{c,i,t}^c,$$

where ck is the carbon content of all types and ages of crops.

The carbon stock in fallow areas is the sum of the carbon stock in fallow on different vegetation types and of different ages:

$$CF_t = \sum_v \sum_i cf_{v,i} \cdot F_{v,i,t},$$

where $F_{v,i,t}$ is fallow land of age i at time t on vegetation type v , and $cf_{v,i}$ is the increasing carbon content of fallow areas given by Equation 2.

The sum of crop areas and fallow areas add up to total deforested area. Thus:

$$\sum_v D_{v,t} = \sum_v \sum_i A_{v,i,t}^c + \sum_v \sum_i F_{v,i,t},$$

2.4 The Age Structure of Fallow Lands

Information about the age structure of fallow lands is not available in the Amazon data set (see section 3 below), but it can be estimated, if we are willing to make some assumptions. We need assumptions both about the initial age structure of fallow lands and about the dynamic process that guides the aging and use of fallow lands.

To estimate the initial age structure, we make the assumption that the average age of fallow area in period 0 in municipality j , I_{0j} , is determined by the share of fallow in total agropastoral areas, D_{0j} , in such a way that a higher share of fallow implies a higher average age of fallow:

$$I_{0j} = K \cdot \frac{F_{0j}}{D_{0j}},$$

where K is a parameter determined by the condition that the average age of fallow over all municipalities should equal the assumed average for Legal Amazonia, I_{0A} :

$$I_{0A} = \sum_j I_{0j} \cdot \frac{F_{0j}}{F_{0A}},$$

and F_{0A} is total fallow area in Legal Amazonia at time 0, i.e. $F_{0A} = \sum_j F_{0j}$.

The dynamic changes in the age structure is believed to be well characterized by the following two assumptions. First, when fallow areas are expanding, the increase must be fallow of age 1, while the rest of the fallow area just gets one period older. The extent of fallow of age i at time t , in a given municipality, is thus determined by:

$$\left. \begin{aligned} F_{t,i} &= \Delta F_t && \text{for } i = 1 \\ F_{t,i} &= F_{t-1,i-1} && \text{for } i > 1 \end{aligned} \right\} \text{when } \Delta F_t \geq 0$$

Second, when fallow areas are contracting, it is because the oldest fallow is being converted to crop land again. This assumption can be following equations:

$$\left. \begin{aligned} F_{t,i} &= F_{t-1,i-1} && \text{for } i > 1 \quad \text{if } \sum_{j=2}^i F_{t-1,j-1} \leq F_t \\ F_{t,i} &= F_t - \sum_{j=2}^{i^*} F_{t-1,j-1} && \text{for } i = 1 \end{aligned} \right\} \text{when } \Delta F_t < 0$$

where i^* is the maximum age of fallow that is not converted to crop land (as determined by the condition in the first equation).

3. The Data

The data for this project comes from a big data base compiled and maintained at the Institute for Applied Economic Research in Rio de Janeiro, Brazil. It covers all the municipalities in Legal Amazonia, spans the period 1970-1985, and measures hundreds of variables that might be relevant for modeling deforestation and economic development in the region. For this paper, we will use only a small subset of all the variables available. These are the variables specifically describing land use and vegetation cover.

The data on original vegetation cover is provided by IBGE, who operate with 6 main categories: I) dense forest, ii) seasonal forest, iii) savanna, iv) ecological transition, v) wetlands, and vi) Campinarana. There are big regional variations in the density of the natural vegetation cover.

The agricultural censuses conducted every five years by IBGE provide the data about changes in land use. Private land used for annual crops, planted forest, planted pasture, and fallow land is considered cleared areas, while all public land plus private land kept as natural forest and natural pasture is considered virgin areas.

3.1 Summary of the Age Structure of Deforested Areas

3.1.1 The age structure of fallow lands

1970 is the initial years of our analysis and the remaining observations come at five year intervals. We assume that the average age of fallow lands in 1970 was 3.5 years and that all fallow land was 10 years old.³ Using that assumption and the equations of section 2.4, we get an age structure as described in Tables 3 and 4.

Table 3: Estimated age structure of fallow areas in Legal Amazonia

| Age group (years) | Area (million hectares) | | | |
|-------------------|-------------------------|------|------|------|
| | 1970 | 1975 | 1980 | 1985 |
| 0 - 5 | 9.2 | 2.4 | 5.0 | 7.4 |
| 5 - 10 | 3.8 | 7.1 | 1.7 | 4.6 |
| 10 - 15 | 0 | 1.0 | 5.9 | 1.5 |
| 15 - 20 | 0 | 0 | 0.5 | 5.4 |
| 20 - 25 | 0 | 0 | 0 | 0.4 |
| Total | 13.0 | 10.4 | 13.1 | 19.4 |

Table 4: Summary statistics for the average age of fallow areas across municipalities.

| | 1970 | 1975 | 1980 | 1985 |
|-----------|------|------|------|------|
| Mean | 3.5 | 6.6 | 8.6 | 9.6 |
| Std. dev. | 0.8 | 1.7 | 3.2 | 4.0 |
| Minimum | 2.5 | 2.5 | 2.5 | 2.7 |
| Maximum | 5.0 | 9.8 | 14.0 | 18.8 |
| Num. obs. | 316 | 315 | 316 | 316 |

³ Reis (1996) chose an initial average age 2.5 in 1975, but noted that this seemed to be too low. It also seemed to be too low for 1970, so we chose 3.5 instead. The carbon emissions results are not sensitive to changes in this parameter.

3.1.2 The age structure of crop Lands

The productivity of annual crops on tropical forest soils is very high the first couple of years while there is an abundance of nutrients from the ashes and while the land is relatively pest free after burning. Thereafter productivity drops sharply, and a long fallow period is needed before the process can be repeated. The alternative is to plant perennial crops and supply the necessary fertilizers and pesticides.

We assume that areas with annual crops and planted pasture have been cleared for 2.5 years, on average. Areas with perennial crops and planted forest is assumed to have been cleared for 7.5 years, on average.

4. Deforestation, Carbon Emissions and Economic Growth 1970—1985

4.1 Deforestation

Table 5 shows that deforestation has been very unevenly distributed across Legal Amazonia. The largest state, with the most dense forest, had lost less than 1 percent of its natural vegetation by 1985 while the more accessible states in the eastern and southern part of Legal Amazonia had lost between 15 and 30%. This pattern also shows in Table 6 which gives the percentage of each vegetation type cleared by 1985. More than 20% of ah savanna has been converted for agropastoral purposes and much of the remaining savanna is used as natural pastures. Less than 6% of the dense forest had been cleared by 1985.

Table 5: Accumulated deforestation in Legal Amazonia by 1985

| State | State area (million ha) | Share deforested % |
|-------------|----------------------------|-----------------------|
| Acre | 15.3 | 3.7 |
| Amapá | 14.3 | 2.5 |
| Amazonas | 157.8 | 0.9 |
| Goiás | 29.2 | 27.1 |
| Maranhao | 28.4 | 30.0 |
| Mato Grosso | 90.7 | 15.5 |
| Pará | 125.3 | 7.8 |
| Rondônia | 23.9 | 7.3 |
| Roraima | 22.5 | 1.6 |
| Total | 507.4 | 8.8 |

Source: The DESMAT data set. Variables: GEAREA, PASPLA85, FLOPLA85, LATEM85, LAPER85, FAL85.

Table 6: Accumulated deforestation in Legal Amazonia by 1985

| Natural vegetation | Area (million ha) | Share deforested % |
|-----------------------|----------------------|-----------------------|
| Dense forest | 317.0 | 5.6 |
| Seasonal forest | 30.7 | 13.4 |
| Savanna | 78.3 | 20.4 |
| Ecological transition | 36.9 | 13.5 |
| Wetlands | 13.1 | 12.8 |
| Campinarana | 31.4 | 0.4 |
| Total | 507.4 | 8.8 |

Source: The DESMAT data set. Variables: PASPLA85, FLOPLA85, LATEM85, LAPER85, FAL85, FODAR, FESAR, SAVAR, TECAR, FOPAR, CAMAR.

It is important to take into account the uneven distribution of clearing when calculating carbon emissions. In this paper, we assume that clearing is randomly distributed across vegetation types *within each municipality*, but because of the large variations in vegetation types and clearing rates across municipalities, we get non-randomly distributed clearing across vegetation types in the whole region (see Table 6).

4.2 Carbon emissions

According to our carbon emissions model, the total carbon stock in Legal Amazonia declined from 74.7 billion tons in 1970 to 72.1 billion tons in 1985 implying average annual emissions of 168 million tC. This is a net effect arising from carbon release from the destroyed original vegetation (198 million tC/year) and carbon sequestration in secondary vegetation (-30 million tC/year). Globally, annual carbon emissions amount to approximately 7.0 billion tC implying a contribution from Amazon deforestation of about 2.4%. The largest emissions came from the southern and eastern states of Pará, Mato Grosso, Maranhão, and Goiás. See Table 7. The table also shows that the latter three states had the lowest emissions per hectare of cleared vegetation. This is due to the generally thinner natural vegetation in these states.

Table 7: Carbon emissions in Legal Amazonia 1970—1985

| State | Accumulated carbon emissions 1970-1985 (million tC) | Carbon emissions per hectare cleared 1970-1985 (tC/ha) |
|-------------|---|--|
| Acre | 54.5 | 135.5 |
| Amapá | 31.1 | 119.0 |
| Amazonas | 67.1 | 128.4 |
| Goiás | 335.5 | 76.0 |
| Maranhão | 388.8 | 112.2 |
| Mato Grosso | 689.8 | 68.5 |
| Pará | 776.4 | 130.0 |
| Rondonia | 150.2 | 119.0 |
| Roraima | 22.8 | 101.0 |
| Total | 2,516.0 | 94.6 |

Source: The DESMAT data set. Variables: PASPLAyy, FLOPLAyy, LATEMyy, LAPERyy, FALyy, FODAR, FESAR, SAVAR, TECAR, FOPAR, CAMAR + assumptions in text

4.3 Economic growth

Total agricultural output in Legal Amazonia increased from US\$ 654 million in 1970 to 2,274 million in 1985 (measured in fixed 1985-US\$). Urban GDP grew even faster from US\$ 1,522 million in 1970 to 11,206 million. This implies an impressive average annual GDP growth rate of 12.9%. Table 9 shows the cumulated GDP over the 1970-1985 period for the fine states in Legal Amazonia and compares it with the carbon emitted.

On average, the states obtained an accumulated GDP of \$44. 1 per ton of carbon they omitted. There are big variations, however. The densely forested and largely inaccessible state of Amazonas obtained a much better trade-off between economic output and carbon emissions than the more accessible and less densely forested south-eastern states like Goiás and Mato Grosso. There are several explanations for these large differences. Goiás and Mato Grosso attracted a lot of cattle ranchers who mainly speculated in rising land prices and tax

evasion while the cattle raising activity itself gave very low, if not negative, yields. The activities in Amazonas, on the other hand, were much more diversified including extractivism (mainly rubber and Brazil nuts) and a big industry related to the Manaus Free Zone.

Table 8: Carbon emissions in Legal Amazonia 1970—1985

| State | Accumulated carbon emissions 1970-1985 (million tC) | Carbon emissions per hectare cleared 1970-1985 (tC/ha) |
|-----------------------|---|--|
| Dense forest | 1,446.1 | 133.6 |
| Seasonal forest | 185.7 | 71.8 |
| Savanna | 499.5 | 52.5 |
| Ecological transition | 354.0 | 113.9 |
| Wetlands | 27.6 | 56.5 |
| Campinarana | 3.2 | 41.7 |
| Total | 2,516.0 | 94.6 |

Source: The DESMAT data set. Variables: PASPLAyy, FLOPLAyy, LATEMyy, LAPERYy, FALyy, FODAR, FESAR, SAVAR, TEOAR, FOPAR, CAMAR ± assumptions in text.

5. Discussion

5.1. Comparisons with other carbon emission studies

Fearnside (1992b) conducted an intensive study of carbon dynamics in the Brazilian Amazon. Rather than using a bookkeeping model, as is done in this paper, he uses a committed carbon approach, thus attributing a carbon that would ultimately be released to the year where deforestation takes place. He concludes that clearing of 1.38 million hectares per year of primary forest was responsible for the release of approximately 270 million tC/year in the period 1989-1990. This corresponds to 196 tC/ha, which is considerably higher than our average estimate for the 1970-1985 period (94.6 tC/ha).

Table 9: Accumulated GDP in Legal Amazonia 1970—1985

| State | Accumulated GDP 1970-1985 (billion fixed 1985-US\$) | Average GDP per ton carbon emitted 1970-1985 (\$/tC) |
|-------------|---|--|
| Acre | 2.8 | 50.9 |
| Amapá | 2.1 | 67.1 |
| Amazonas | 25.5 | 379.4 |
| Goiás | 4.5 | 13.3 |
| Maranhao | 18.9 | 48.5 |
| Mato Grosso | 13.9 | 20.1 |
| Pará | 35.8 | 46.1 |
| Rondonia | 6.6 | 44.3 |
| Roraima | 1.1 | 47.5 |
| Total | 111.0 | 44.1 |

Source: The DESMAT data set. Variables: PASPLAyy, FLOPLAyy, LATEMyy, LAPERYy, FALyy, FODAR, FESAR, SAVAR, TECAR, FOPAR, CAMAR, PIBAGyy, PIBURByy ± assumptions in text.

Schroeder & Winjum (1994b) gives a much lower estimate of net carbon emissions from land use changes in all of Brazil. Using a bookkeeping model with gradual decomposition of carbon in cleared areas and gradual recomposition in secondary forest, they get net carbon emissions for Brazil in the range of -44 to +10 million tC/year for 1990. The large discrepancy between Fearnside (1992b) and Schroeder & Winjum (1994b) is mainly attributed to the massive carbon recomposition in secondary forests in the latter study (+245 million tC/year).

5.2 Uncertainties

5.2.1 *Biomass content*

Biomass contents of Amazonian forest patches vary greatly due to differences in soil characteristics, rainfall regimes, altitude, and previous natural and antropogenic disturbances. In addition, there is a large statistical variation when sampling forest patches, because biomass content is very sensitive to the presence of large trees.

There has been a vigorous debate on the forest biomass in the Brazilian Amazon (Brown Lugo 1984, 1990, 1992; Fearnside 1985, 1986, 1987, 1992; Lugo & and Brown 1986; Brown *et al* 1989). Brown *et al* (1995) notes that the estimates during the last decade differ by more than a factor of two. They provide a new study for biomass content in Rondônia and gives an estimate of the uncertainty associated with the biomass estimate. Through Monte Carlo Simulation they find that measurement errors are approximately +20% around the mean (for a 95% confidence interval).

If we use carbon contents that are 20% lower than our mean values we get a lower estimate of annual carbon release of 128 million tC per year or 72.2 tC per hectare cleared. Similarly, we get a high estimate of 207 million tC per year or 117 tC per hectare cleared. See Table 10.

Table 10: Biomass sensitivity analysis

| | Low biomass content | Central biomass content | High biomass content |
|--|---------------------|-------------------------|----------------------|
| Average annual emissions (million tC) | 128 | 168 | 207 |
| Average emissions per hectare cleared (tC/ha) | 72.2 | 94.6 | 117.0 |
| Average GDP per tC emitted (1985-US\$/tC) | 57.8 | 44.1 | 35.7 |

5.2.2 *Steady state assumption*

The calculations done in this paper assumes that undisturbed primary forest, which still account for the major part of the Brazilian Amazon, is in a steady state equilibrium with no net carbon gain or loss. Lugo & Brown (1992) questions this assumption of a steady state equilibrium, arguing that the rate of biomass accumulation of tropical forests is changing over time as a result of natural climatic changes, catastrophes, and past human disturbances. They cite plot level field data from Venezuela and Puerto Rico that show mature forests

accumulating carbon at rates of 1 - 2 tC/ha/year (Schroeder & Winjum 1994b).

A possible explanation for the increase in steady state carbon levels is CO₂ fertilization. Studies of agricultural and wild plant species have shown typically 20-40% higher photosynthesis and growth under doubled CO₂ conditions (Rocheffort & Bazzaz 1992; Körner & Arnone 1992; Idso & Kimball 1993). This effect suggests faster forest regrowth and can also help explaining a possible increase in steady state carbon stocks. Schimel (1995) notes, however, that long term, ecosystem responses may be substantially smaller than plant-level responses measured in laboratories.

Because of the huge area of primary forest in the Amazon, even the slightest change in biomass per hectare would result in significant effects on total carbon storage. If the approximately 350 million hectares of primary dense forest in the Brazilian Amazon, for example, increased its biomass with just 0.5% per year, then about 300 million tC would be sequestered every year. This number is larger than our estimate of annual carbon emitted from deforested areas. That makes it difficult to say anything certain about the net emissions from the whole of Legal Amazonia. Our analysis only applies to the net contributions from disturbed areas.

6. Conclusions

The calculations in this paper indicate that average annual carbon emissions from deforestation in the Brazilian Amazon falls in the range 128 - 207 million tC/year during the rapid development period 1970 - 1985. This corresponds to 1.8 - 3.0% of global carbon emissions from fossil fuel combustion and deforestation. The estimated range is within the range provided by other studies, but falls in the low end. This is due to two factors. First, we take into account the uneven spatial distribution of deforestation: Lower clearing costs causes deforestation to take place in the most accessible and least dense forests as long as these are available. Second, we take into account the considerable secondary regrowth and carbon sequestration that takes place when cleared land is abandoned.

During the same period, the Amazonian economy grew rapidly resulting in average annual GDP growth rates of 12.9%. This is impressive by any measure, and several studies show that settlers in the Amazon did well, especially when you take into account their generally poor background (Mattos, Uhl & Goncalves 1992; Ozorio de Almeida 1992; FAO/UNDP/MARA 1992; Jones *et al* 1992; Schneider 1992). On average during the period, there was generated \$44 of GDP per ton carbon emitted. There were big variations in the benefit, however. In the state of Amazonas the benefit was as high as \$379/tC. For comparison, the CO₂ tax implemented in Denmark in 1992 require households to pay \$16 per ton carbon emitted and VAT-registered firms to pay half the tax (Svendsen 1996 quoted in Young 1997). It is therefore not at all clear that stopping deforestation is the cheapest (and most just) way to curb carbon emissions. Especially not when we take into account that it will also require substantial direct costs to protect the Amazon against slash-and-burn deforestation.

References

- Almeida, A. d. (1992), Deforestation and turnover in Amazon colonization, Discussion paper (draft), Washington, D.C.: The World Bank.
- Bohrer, C. B. A. (1993), Base de dados municipais sobre o volume de madeira e a biomassa forestal da Amazônia Legal, Memo, 8 pp, IPEA/Rio.
- Brown, I. F., Martinelli, L. A., Thomas, W. W., Moreira, M. Z., Ferreira, C. A. C. & Victoria, R. A. (1995), 'Uncertainty in the biomass of Amazonian forests: An example from Rondônia, Brazil', *Forest Ecology and Management* 75, 175—189.
- Brown, S., Gillespie, A. J. R. & Lugo, A. E. (1989), 'Biomass estimation methods for tropical forests with applications to forest inventory data', *Forest Science* 35, 881-902.
- Brown, S. & Lugo, A. E. (1984a), 'Biomass estimates for tropical forests: A new estimate based on forest volumes', *Science* 223, 1290-1293.
- Brown, S. & Lugo, A. E. (1984b), 'The storage and production of organic matter in tropical forests and their role in the global carbon cycle', *Biotropica* 14, 161-187.
- Brown, S. & Lugo, A. E. (1992a), 'Aboveground biomass estimates for tropical moist forest of the Brazilian Amazon', *Interciencia* 17, 8-18.
- Brown, S. & Lugo, A. E. (1992b), Biomass estimates for Brazil's Amazonian moist forests., in 'Forests '90: Annals of the First International Symposium on Environmental Studies on Tropical Rain Forests', Manaus, pp. 46-52.
- Cannell, M. G. R. (1982), *World forest biomass and primary production data*, London: Academic Press.
- Carvalho Jr., J. A., Santos, J. C., Santos, J. M. & Higuchi, N. (1994), 'Estimate of combustion efficiency in a forest clearing experiment in the Manaus region', *Revista Brasileira da Geofísica* 12, 45-48.
- FACE (1993), *Jaarverslag*, Arnhem: The FACE Foundation.
- FAO/UNDP/MARA (1992), Principais indicadores sócio-econômicos dos assentamentos de reforma agrária, Project BRA-87/022, Ministerio da Agricultura e Reforma Agrária.
- Fearnside, P. M. (1985a), 'Brazil's Amazon forest and the global carbon problem', *Interciencia* 10, 179-186.
- Fearnside, P. M. (1985b), 'Brazil's Amazon forest and the global carbon problem: Reply to Lugo and Brown', *Interciencia* 11, 58-64.
- Fearnside, P. M. (1987), Summary of progress in quantifying the potential contribution of Amazonian deforestation to the global carbon problem, in D. Athié, T. E. Lovejoy & P. de M. Oyens, eds, 'Proceedings of the Workshop on Biogeochemistry of Tropical Rain Forests: Problems for Research', Piracicaba, Sao Paulo: Universidade de Sao Paulo, pp. 75-82.
- Fearnside, P. M. (1992a), 'Forest biomass in Brazilian Amazonia: Comments on the estimate by Brown and Lugo', *Interciencia* 17, 19-27.
- Fearnside, P. M. (1992b), Greenhouse gas emissions from deforestation in the Brazilian Amazon, in W. Makundi & J. Sathaye, eds, 'Tropical forestry and global climate change: land use policy, emission and sequestration', Proceedings of an international workshop held at the Lawrence Berkeley Laboratory, Berkeley, California, 29-31 May, 1991.
- Fearnside, P. M., Leal Jr., N. & Moreira Fernandes, M. (1993), 'Rainforest burning and the global carbon budget: biomass, combustion efficiency, and charcoal formation in the Brazilian Amazon', *Journal of Geophysical Research* 98, 16733-16743.
- Houghton, R. A., Hobbie, J. E., Melillo, J. M., Moore, B., Peterson, B. J., Shaver, G. R. & Woodwell, G. M. (1983), 'Changes in the carbon content of terrestrial biota and soils between 1860 and 1980: A net release of CO₂ to the atmosphere', *Ecological Monographs* 53, 235-262.
- Houghton, R. A., Skole, D. L. & Lefkowitz, D. S. (1991), 'Changes in the landscape of Latin America between 1850 and 1985. II. net release of CO₂ to the atmosphere', *Forest Ecology and Management* 38, 173-199.
- Idso, S. B. & Kimball, B. A. (1993), 'Tree growth in carbon dioxide enriched air and its implications

- for global carbon cycling and maximum levels of atmospheric CO₂, *Global Biogeochemical Cycles* 7, 537-555.
- Jones, D. W. et al. (1992), *Farming in Rondônia*, Blacksburg, Virginia: Oak Ridge National Laboratory for the U.S. Department of Energy and Department of Urban and Regional Planning, Virginia Polytechnic Institute and State University.
- Kolk, A. (1996), *Forests in International Environmental Politics: International Organizations, NGOs and the Brazilian Amazon*, Utrecht: International Books.
- Körner, C. & Amone III, J. A. (1992), 'Responses to elevated carbon dioxide in artificial tropical ecosystems', *Science* 257, 1672-1675.
- Lugo, A. E. & Brown, S. (1992), 'Tropical forests as sinks of atmospheric carbon', *Forest Ecology and Management* 54, 239-255.
- Mattos, M.M., C. U. & de Almeida Goncalves, D. (1992), Economic and ecological perspectives on ranching in eastern Amazon in the 1990s, Paper submitted to world development, Instituto do Homem e Meio Ambiente da Amazonia, Brazil; EMBRAPA, Penn State University.
- Nordhaus, W. D. (1991), 'A survey of estimates of the costs of controlling greenhouse gases', *The Energy Journal* 12, 37-66.
- Odum, E. P. (1988), *Ecologia*, Rio de Janeiro: Guanabara S.A.
- Reis, E. J. (1996), Padrões geográficos de incidência dos efeitos econômicos e ambientais do programa Grande Carajás, in R. Varsano, ed., 'Perspectivas da Economia Brasileira - 1996', Rio de Janeiro: IPEA.
- Rocheffort, L. & Bazzaz, F. A. (1992), 'Growth response to elevated CO₂ in seedlings of four co-occurring birch species', *Canadian Journal of Forest Research* 22, 1583-1587.
- Sanchez, P. A., Palm, C. A., Szott, L. T., Cuevas, E. & Lal, R. (1989), Organic input management in tropical agroecosystems, in D. C. Coleman et al., eds, 'Dynamics of soil organic matter in tropical ecosystems', Honolulu: University of Hawaii Press.
- Schimel, D. S. (1995), 'Terrestrial ecosystems and the carbon cycle', *Global Change Biology* 1, 77-91.
- Schneider, R. R. (1993), The potential for trade with the Amazon in greenhouse gas reduction, Latent Dissemination Note II. From <http://www.worldbank.org/html/lat/english/papers/env/note.2.pdf>, The World Bank.
- Schneider, R. R. (1995), Government and the economy on the Amazon frontier, World Bank Environment Paper 11, The World Bank.
- Schroeder, P. E. & Winjum, J. K. (1994a), Assessing Brazil's carbon budget: II. biotic carbon pools, Working paper, U.S. EPA Environmental Research Laboratory, Oregon.
- Schroeder, P. E. & Winjum, J. K. (1994b), Assessing Brazil's carbon budget: II. biotic fluxes and net carbon balance, Working paper, U.S. EPA Environmental Research Laboratory, Oregon.
- Seiler, W. & Crutzen, P. J. (1980), 'Estimates of gross and net fluxes of carbon between the biosphere and the atmosphere from biomass burning', *Climatic Change* 2, 207-247.
- Singh, J. S. & Joshi, M. C. (1979), Primary production, in R. T. Coupland, ed., 'Grassland ecosystems of the world: Analysis of grasslands and their uses', Cambridge: Cambridge University Press.
- Svendsen, G. T. (1992), Public choice and environmental regulation: Tradable permit systems in the United States and CO₂ taxation in Europe, Technical report, PhD-dissertation presented at the Aarhus School of Business, Aarhus, Denmark.
- Uhl, C. (1987), 'Factors controlling succession following slash-and-burn agriculture in Amazonia', *Journal of Ecology* 75, 377-407.
- Uhl, C., Buschbacher, R. & Serrão, E. A. S. (1988), 'Abandoned pastures in eastern Amazonia. patterns of plant succession', *Journal of Ecology* 76, 663-681.
- World Bank (1991), The forest sector, A world bank policy paper, Washington: The World Bank.
- Young, C. E. F. (1997), International policy issues on carbon fluxes and forests in the South, Draft, IPEA, Rio de Janeiro, Brazil.