

Langbein, Jörg; Peters, Jörg; Vance, Colin

**Working Paper**

## Outdoor cooking prevalence in developing countries and its implication for clean cooking policies

Ruhr Economic Papers, No. 680

**Provided in Cooperation with:**

RWI – Leibniz-Institut für Wirtschaftsforschung, Essen

*Suggested Citation:* Langbein, Jörg; Peters, Jörg; Vance, Colin (2017) : Outdoor cooking prevalence in developing countries and its implication for clean cooking policies, Ruhr Economic Papers, No. 680, ISBN 978-3-86788-788-5, RWI - Leibniz-Institut für Wirtschaftsforschung, Essen, <https://doi.org/10.4419/86788788>

This Version is available at:

<https://hdl.handle.net/10419/155771>

**Standard-Nutzungsbedingungen:**

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

**Terms of use:**

*Documents in EconStor may be saved and copied for your personal and scholarly purposes.*

*You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.*

*If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.*



# RUHR

ECONOMIC PAPERS

Jörg Langbein  
Jörg Peters  
Colin Vance

## **Outdoor Cooking Prevalence in Developing Countries and its Implication for Clean Cooking Policies**

# Imprint

## Ruhr Economic Papers

Published by

Ruhr-Universität Bochum (RUB), Department of Economics  
Universitätsstr. 150, 44801 Bochum, Germany

Technische Universität Dortmund, Department of Economic and Social Sciences  
Vogelpothsweg 87, 44227 Dortmund, Germany

Universität Duisburg-Essen, Department of Economics  
Universitätsstr. 12, 45117 Essen, Germany

RWI Leibniz-Institut für Wirtschaftsforschung  
Hohenzollernstr. 1-3, 45128 Essen, Germany

## Editors

Prof. Dr. Thomas K. Bauer  
RUB, Department of Economics, Empirical Economics  
Phone: +49 (0) 234/3 22 83 41, e-mail: thomas.bauer@rub.de

Prof. Dr. Wolfgang Leininger  
Technische Universität Dortmund, Department of Economic and Social Sciences  
Economics – Microeconomics  
Phone: +49 (0) 231/7 55-3297, e-mail: W.Leininger@tu-dortmund.de

Prof. Dr. Volker Clausen  
University of Duisburg-Essen, Department of Economics  
International Economics  
Phone: +49 (0) 201/1 83-3655, e-mail: vclausen@vwl.uni-due.de

Prof. Dr. Roland Döhrn, Prof. Dr. Manuel Frondel, Prof. Dr. Jochen Kluge  
RWI, Phone: +49 (0) 201/81 49-213, e-mail: presse@rwi-essen.de

## Editorial Office

Sabine Weiler  
RWI, Phone: +49 (0) 201/81 49-213, e-mail: sabine.weiler@rwi-essen.de

## Ruhr Economic Papers #680

Responsible Editor: Manuel Frondel

All rights reserved. Bochum, Dortmund, Duisburg, Essen, Germany, 2017

ISSN 1864-4872 (online) – ISBN 978-3-86788-788-5

The working papers published in the Series constitute work in progress circulated to stimulate discussion and critical comments. Views expressed represent exclusively the authors' own opinions and do not necessarily reflect those of the editors.

---

**Ruhr Economic Papers #680**

Jörg Langbein, Jörg Peters, and Colin Vance

**Outdoor Cooking Prevalence in Developing  
Countries and its Implication for Clean  
Cooking Policies**

## **Bibliografische Informationen der Deutschen Nationalbibliothek**

---

Die Deutsche Bibliothek verzeichnet diese Publikation in der deutschen Nationalbibliografie; detaillierte bibliografische Daten sind im Internet über:  
*<http://dnb.d-nb.de>* abrufbar.

Das RWI wird vom Bund und vom Land Nordrhein-Westfalen gefördert.

<http://dx.doi.org/10.4419/86788788>

ISSN 1864-4872 (online)

ISBN 978-3-86788-788-5

---

Jörg Langbein, Jörg Peters, and Colin Vance<sup>1</sup>

# Outdoor Cooking Prevalence in Developing Countries and its Implication for Clean Cooking Policies

## Abstract

*More than 3 billion people use wood fuels for their daily cooking needs, with detrimental health implications related to smoke emissions. Global initiatives to disseminate clean cooking stoves emphasize technologies that are either expensive, such as electricity and gasifier stoves, or for which supply chains hardly reach rural areas, such as LPG. This emphasis neglects that many households in the developing world cook outdoors. Our calculations demonstrate that for such households, already the use of less expensive biomass cooking stoves can substantially reduce smoke exposure. The cost-effectiveness of clean cooking policies can thus be improved by taking cooking location and ventilation into account.*

*JEL Classification: Q53, I12, O13*

*Keywords: Air pollution; health behavior; energy access*

*March 2017*

---

<sup>1</sup> Jörg Langbein, RWI; Jörg Peters, RWI and AMERU Johannesburg, South Africa; Colin Vance, RWI and Jacobs University Bremen. – We would like to thank Gunther Bensch for valuable comments and suggestions. Peters and Langbein gratefully acknowledge the support of a special grant (Sonderetatbestand) from the German Federal Ministry for Economic Affairs and Energy and the Ministry of Innovation, Science, and Research of the State of North Rhine-Westphalia. – All correspondence to: Jörg Peters, RWI, Hohenzollernstr. 1/3, 45128 Essen, Germany; joerg.peters@rwi-essen.de

# 1 Introduction

In recent years, the promotion of clean cookstoves to reduce smoke exposure has received much attention in both academic and policy discussions. Indeed, much is at stake: More than 3 billion people in developing countries rely on firewood and charcoal for their daily cooking purposes. According to the World Health Organisation (WHO), the emitted smoke from household air pollution kills 4.3 million people every year - more deaths than are caused by malaria, tuberculosis and HIV combined - making it one of the most lethal environmental health risks (WHO, 2016; Martin, 2011).

Under the auspices of the United Nations Initiative Sustainable Energy for All (SE4All) and spearheaded by the Global Alliance for Clean Cookstoves (GACC), the international development community is currently embarking on a massive effort to spur universal adoption of clean cookstoves and fuels (GACC, 2011; SE4ALL, 2015). Achieving universal adoption is a laudable outcome, but one that faces substantial organizational and financial constraints. This raises the question of whether policies should concentrate on technologies and fuels that qualify as absolutely clean from a public health perspective, such as electricity, LPG, or advanced gasifier biomass stoves, or whether intermediate technologies such as simple improved biomass stoves should also be promoted (Simon et al. 2014). Notwithstanding their considerably higher costs and often fragmented supply chains, a recent WHO report advocates “energy solutions that are clean for health at the point-of-use” (WHO, 2016, p. 87), these being primarily LPG and electricity or advanced gasifier biomass stoves.

In the present paper, we argue for an alternative prioritization that takes into account how smoke exposure is impacted by the interaction of cookstove technologies and cooking behaviors (Jeuland et al., 2015). In this regard, where people cook - whether indoors

or outdoors - has important implications for ventilation and thus smoke exposure (see Bensch and Peters, 2015; Dasgupta et al., 2006; Yu, 2011), but has nonetheless been widely neglected in debates about clean stove distribution. Impact potentials of stoves are higher if meals are prepared indoors. Conversely, if meals are prepared outdoors, natural ventilation reduces exposure considerably, with an associated reduction in the beneficial impact of the clean cookstove.<sup>1</sup> Scarce public resources should consequently concentrate on distributing the most advanced cookstoves among households where indoor cooking prevails and hence exposure is highest. In areas where outdoor cooking dominates, much simpler - and cheaper - improved biomass stoves are potentially more cost effective in reducing the adverse effects of biomass cooking.

We develop this argument in two steps. Drawing on data from the Demographic and Health Surveys (DHS), we first document cooking behavior by country, which reveals a sizeable incidence of outdoor cooking. Next, we calculate hypothetical exposure reductions for different stove types and ventilation scenarios and then categorize the stoves into different internationally recognized emissions categories, or tiers. This exercise demonstrates that depending on the scenario, stoves that would otherwise be categorized in the lowest tier (Tier-Zero), are re-categorized in higher tiers when used outdoors. Based on the documented heterogeneity in cooking patterns, we suggest that the distribution of cheaper biomass stoves should be given serious consideration as a cost-effective instrument to bring down exposure levels among households that cook outdoors.

---

<sup>1</sup> See Grabow et al. (2013) for results in a laboratory environment and Rosa et al. (2014) for results in a field environment.



## 2 Policy and literature background

### 2.1 Health effects of household air pollution and cooking ventilation

Exposure to particulate matter induced by biomass cooking affects health in various ways and may lead to acute respiratory infections, stunted growth in children, pneumonia, chronic bronchitis in women, chronic obstructive pulmonary disease (COPD), cataracts and other visual impairments, cardiovascular diseases, lung cancer, tuberculosis and perinatal diseases (Po et al., 2011; Ezzati and Kammen, 2002; Amegah et al., 2014; Dherani et al., 2008; McCracken et al., 2012; Hosgood et al., 2010; Bruce et al., 2013; Smith et al., 2014). The WHO's Global Burden of Disease/Comparative Risk Assessment Project estimated that the exposure to household air pollution from cooking with solid fuels caused 4.3 million premature deaths in 2012 (WHO, 2016).

There are only two studies that systematically analyze the effect of outdoor cooking on health. Rehfuess et al. (2009) and Buchner and Rehfuess (2015) conduct cross country studies among 16 African countries and 9 Sub-Saharan countries, respectively, finding that the effect of cooking with solids fuels on acute lower respiratory infections varies among children with regards to ventilation practices and the cooking location. Bensch and Peters (2015) observe a surprising improvement in self-reported health indicators for an ICS whose design is not expected to generate health effects. They provide explorative evidence for the transmission channel and find that a reduction in smoke exposure due to a shorter cooking duration and increased outside cooking might explain this result.

A few studies drawn from cross-sectional field surveys examine the particulate matter (PM) concentration level once the cooking location is outdoors. The suggested range

of the effect is broad. Balakrishnan et al. (2002) find a reduction of particulate matter concentration between 40 and 44 percent in India, while Rosa et al. (2014) find a reduction of 57 percent in Rwanda. The highest estimate of which we are aware is from Albalak et al. (1999), who find a 77 percent reduction in Bolivia.

## 2.2 Policy background

Improved cooking is high on the agenda of international policy, spearheaded by WHO and the Global Alliance for Clean Cookstoves. The different levels of cleanliness of stoves are accounted for in SE4All’s Global Tracking Framework (GTF), which uses a four-tier system to categorize ICS and track the progress towards universal access to modern energy. These four tiers, defined according to measurements that are done under standardized indoor conditions, are also used as a reference by WHO, GACC and other actors in the clean cooking policy scene. The GTF evaluates cookstoves in the four categories of efficiency, safety, indoor emissions, and total emissions for a high- and low-power scenario, with the latter categories and their respective tiers shown in Table 1.

**Table 1:** Emissions and indoor emissions tiers of performance levels

	Tier 0	Tier 1	Tier 2	Tier 3	Tier 4
Indoor emission PM <sub>2.5</sub> (mg/min)	>40	≤40	≤17	≤8	≤2
Emissions in high power scenario PM <sub>2.5</sub> (mg/MJ <sub>d</sub> )	>979	≤979	≤386	≤168	≤41
Emissions in low power scenario PM <sub>2.5</sub> (mg/min/L)	>8	≤8	≤4	≤2	≤1

*Source:* ISO, 2012

While all stakeholders are dedicated to eradicate energy poverty and to provide households with improved cookstoves, the understanding of what exactly constitutes an improved cookstove differs between the different actors. Many non-governmental organizations and most African governments focus on affordable simple technologies that are

designed to save fuels in order to reduce deforestation pressures and improve livelihoods through reduced firewood collection time or charcoal expenditures. These stoves, which fall under Tiers 1, 2, and 3, are not designed to completely eliminate smoke emissions. WHO and GACC, by contrast, clearly concentrate on the adverse health effects of wood-fuel cooking and thus only consider an ICS as improved if it is classified as Tier 4. The rationale behind this is the so-called non-linear particulate exposure-response relation found in medical research, which suggests that large reductions in smoke exposure are required in order to ensure positive health effects (see, for example Ezzati and Kammen, 2002; Pope et al., 2011; Burnett et al., 2014).

The present paper argues that cooking behavior that affects ventilation, particularly outdoor cooking, can have a considerable effect on particulate matter exposure and should be taken into account when decisions are taken on whether to consider a certain stove as clean and, consequently, whether to consider it for promotion.

### 3 Data

We use data from the latest waves of the nationally representative Demographic and Health Surveys (DHS). The data have been regularly collected in around 90 low-and middle-income countries since 1984. For our purpose, we only included low and lower middle income countries in Africa, Latin America and South-East Asia as defined by the World Bank, thereby excluding Brazil and the Maldives. Due to data regulations, not all countries that fit this classification could be included in the analysis.<sup>2</sup> Information on the cooking location is only available for those countries where the latest available wave (wave 6) or the second latest available wave (wave 5) of the standard DHS questionnaire

---

<sup>2</sup> This excludes Cambodia, Eritrea, Equatorial Guinea, Samoa, Sri Lanka, Vietnam, and Yemen.

was conducted.<sup>3</sup> If information in two waves were available for one country, we used the latest wave. This leaves us with a sample of 40 countries and 650,723 household observations for the years 2006 to 2014. Most of the included countries are situated in Africa (30), followed by Asia (6) and Latin America (4).<sup>4</sup>

The DHS questionnaires contain questions regarding cooking behavior, including stove usage, cooking fuels, and cooking location. We restrict our interest to the question on the cooking place. Households that cook at home were asked whether they usually cook in the house, in a separate building, or outside. It was not possible to give multiple answers.

We divide the sample between rural and urban areas, since we expect different outdoor cooking patterns for these two groups. All results are furthermore weighted to ensure nationally representative results, with the weights provided by the DHS.

## 4 Outdoor cooking prevalence

As seen from Figures 1 and 2, outside cooking is prevalent in both the urban and rural areas of many developing countries, reaching a high of nearly 80 percent in rural Niger. Notwithstanding substantial heterogeneity, a few patterns in the data can be discerned. Out of the 20 countries with the highest outdoor cooking rates, 18 are located in Africa. Further differencing within the African continent shows that West African countries have the highest share of outdoor cooking. Among the ten countries with the highest outdoor cooking rates, seven are in West Africa. At the other end of the spectrum, the four countries with the lowest outdoor cooking rates are spread across South America, the

---


<sup>3</sup> This excludes Botswana, Cape Verde, Central African Republic, Colombia, Guatemala, Guyana, Laos, Mauritania, Paraguay, Sao Tome and Principe, South Africa, Swaziland, and Tanzania.

<sup>4</sup> See Table A.1 in the Appendix for a list of included countries and respective number of observations.

Caribbean, South East Asia and Asia, with Pakistan registering the lowest rate of about 1 percent.

Large differences between urban and rural outside cooking patterns are evident in some countries. We take a closer look at only those countries with more than 15 percentage points difference in rural and urban outdoor cooking patterns. This yields two different types of countries, all based in Africa: those in which more households cook outside in rural areas than in urban areas (Benin, Gabon, Lesotho and Namibia) and those in which more households cook less outside in rural areas than in urban areas (Burundi, Republic of the Congo, Democratic Republic of the Congo, Guinea, Liberia, Madagascar, Malawi and Uganda). For all other countries, no major difference between household cooking patterns in rural and urban areas is observed.


**Figure 1: Cooking place in rural areas in developing countries**



*Note:* DRC refers to Democratic Republic of Congo and Congo refers to Republic of Congo.

*Source:* Demographic and Health surveys (2006-2014)

**Figure 2:** Cooking place in urban areas in developing countries



Note: DRC refers to Democratic Republic of Congo and Congo refers to Republic of Congo.

Source: Demographic and Health surveys (2006-2014)

## 5 Implications for air pollution - a stylized numerical comparison

The variation in cooking location has considerable implications for the emission-exposure nexus of cooking induced smoke. In this section, we provide a back-of-the-envelope calculation of particulate matter levels for different stove types according to whether the stove is used indoors or outdoors. The aim is to show that the effective cleanliness of a stove is profoundly impacted by this distinction. We use as cleanliness categories the tiers as defined in the SE4All Global Tracking Framework (see Table 1 in Section 2.2). Our analysis includes stoves from tiers zero to three. Tier four stoves are mostly those that run on electricity and LPG, so virtually free of smoke emissions. All stoves have in common that they are non-traditional, portable, household biomass stoves without a chimney and not used for commercial purposes.<sup>5</sup>

For the cookstoves examined, we rely on emissions figures from Jetter et al. (2012), who analyze the emission of 22 cookstoves in a controlled environment in the laboratory. The selection of stoves in Jetter et al. (2012) is based on availability, which excludes a large number of other non-standard stoves and chimney stoves, but covers those most widely disseminated. The authors measure the emission (in mg/minute) from a low power and a high power scenario as defined in the Water Boiling Test Protocol. Although Water Boiling Tests undoubtedly diverge from actual field use, they have the virtue of allowing comparison of many cookstoves under identical circumstances.<sup>6</sup>

---

<sup>5</sup> See Table A.2 in the Appendix for a list of the cookstoves, their categories, fuel and retail price.

<sup>6</sup> A typical WBT consists of three phases that immediately follow each other: A cold start high power phase, in which a measured quantity of water is boiled. After the first phase, the water is replaced by new water. This is called the high power, hot start phase. After the water is again boiled, in the last phase (low power), the water simmers just below boiling point for 45 minutes. For the



We focus on the high power scenario results, since emissions tend to be higher during this phase. Results for the low power scenario are presented in Table A.3 in the Appendix. Whereas the high power scenario simulates the actual high power use of the cookstove, such as quickly boiling water, the low power scenario simulates the long simmering of legumes or pulses (GACC, 2014).

The first four columns of Table 2 show the cooking device, associated cooking fuel, indoor emissions and their tiers for the high power scenario. Among the cooking devices, values are presented for both a minimally tended and carefully tended three stone fire, as this is the most prevalent cooking technology in developing countries. Jetter et al. (2012) report a minimally tended three-stone fire to be closer to the values that are observed in the field. Indoor emission rates vary considerably for the high power scenario, as can be seen in column 3 of Table 2.

Based on the cookstove and their respective indoor emissions (measured in mg/min) depicted in Table 2, we calculate average PM<sub>2.5</sub> concentration levels (measured in  $\mu\text{g}/\text{m}^3$ ) in the kitchen during cooking time under different scenarios. To this end, we apply a variant of the single zone box model developed by Johnson et al. (2011) that was refined for easier implementation by the Aprovecho Research Center (2016) in the form of a spreadsheet tool.<sup>7</sup> The model abstracts from different concentration levels in different parts of a room or house and has been used in the analysis of biomass cooking emissions (e.g. WHO, 2014b). In line with Johnson et al. (2011), further assumptions are four hours of cooking per day (1 hour in the morning, 1 hour at lunch time, and two hours for dinner) as well a kitchen volume of 30m<sup>3</sup> and 25 air exchanges per hour. Plugging the

---

high power values the average is calculated for the emissions from the two high power phases (see GACC, 2014 for a detailed description of the procedure).

<sup>7</sup> The reliability of the tool was corroborated by comparing the results to those obtained by Johnson et al. (2011) and WHO (2014b). The results were similar.

values of the indoor emissions of the respective cookstoves into the spreadsheet yields the respective indoor PM<sub>2.5</sub> concentration levels ( $\mu\text{g}/\text{m}^3$ ), presented in column 5 of Table 2.

As discussed in section 2.1, effects from moving the location outdoors on particulate matter concentration level occur on a broad range. Accordingly, we account for this variability by showing exposure reductions for three scenarios: 40, 60, and 80 percent reductions (see Table 2, column 6, 7, 8).<sup>8</sup>

---

<sup>8</sup> Note that the outside-cooking studies we are referring to observe differences in PM concentrations under real-world conditions, while the Jetter et al. (2012) emission measurements are determined in standardized low- and high-power scenario.

**Table 2: Cookstove emissions in the high power scenario**


Cooking device	Fuel	Indoor			Outdoor		
		Indoor emissions, tiers and		Assumed outdoor reduction	level and PM <sub>2.5</sub> average		
		PM <sub>2.5</sub> average concentration	level during cooking time		concentration level during	cooking time ( $\mu\text{g}/\text{m}^3$ )	
mg/min	Tier	Concentration	percent	percent	percent	percent	
3-stone minimally tended	Wood	93.8	0	7373	4424	2949	1475
3-stone carefully tended	Wood	56.9	0	4473	2684	1789	895
Envirofit-G330	Wood	52.6	0	4135	2481	1654	827
Philips HD4008 Natural Draft	Wood	53.8	0	4229	2537	1692	846
Saampada	Wood	56.9	0	4473	2684	1789	895
StoveTec Greenfire	Wood	46.3	0	3639	2183	1456	728
Upesi Portable	Wood	69.2	0	5440	3264	2176	1088
GERES	Charcoal	44.2	0	3474	2084	1390	695
Gyapa	Charcoal	26.0	1	2044	1226	818	409
Jiko-Ceramic	Charcoal	22.6	1	1776	1066	710	355
Jiko-Metal	Charcoal	17.5	1	1376	826	550	275
KCJ Standard	Charcoal	18.3	1	1438	863	575	288
Kenya Uhai	Charcoal	20.8	1	1635	981	654	327
StoveTec Charcoal	Charcoal	26.3	1	2067	1240	827	413
StoveTec Greenfire, reduced fuel fee	Wood	25.1	1	1973	1184	789	395
Mayon Turbo	Rice Hulls	31.3	1	2460	1476	984	492
Berkeley Darfur	Wood	18.4	1	1446	868	578	289
Envirofit-G3300, reduced fuel feed	Wood	23.2	1	1824	1094	730	365
Protos	Plant oil	34.5	1	2712	1627	1085	542
Belonio	Rice hulls	8.2	2	645	387	258	129
Philips HD4012 fan	Wood	6.6	3	519	311	208	104
Oorja stove	Biomass pellets	2.9	3	228	137	91	46
StoveTec TLUD	Wood pellets	4.4	3	346	208	138	69

*Note:* High power scenario refers to a scenario of the Water Boiling Test where the indoor emission is measured in the time from the start of the cooking process until a 5 liter pot of water is boiling. This is done with a cold start, where the cookstove has not been used for some time before and a hot start where the stove was used immediately before. For the high power scenario values here, the average is taken for the values obtained in the hot start and cold start scenario as it was done by Jetter et al. (2012). Cooking time is assumed to be 4 hours and the average value during cooking time is taken for the concentration level.

Source: Jetter et al. (2012)

Since the  $PM_{2.5}$  concentration levels ( $\mu g/m^3$ ) are directly proportional to the emission (mg/min), we can convert the concentration figures from Table 2 into the emission levels, thereby yielding the yardstick used in the SE4All-Tier system. Figure 3 shows that there is a strong effect of outdoor cooking on how the stove should be categorized. Most stoves would improve by one tier in the 40 and 60 percent reduction scenarios and by two tiers for the 80 percent reduction scenario. The effect is similar in a low power scenario (see Table A.3 in the Appendix).

**Figure 3:** Indoor emissions, outdoor cooking reduction and tiers



Importantly, the difference in tier categorization applies even to very simple and inexpensive cooking devices, such as the KCJ Standard, a charcoal stove that costs six US-Dollars. This device advances up the scale when used outdoors, from tier 1 to tier 3 under the 80 percent reduction scenario. As an example for a fuelwood driven stove, the Berkeley Darfur stove is within tier 1 with indoor emissions, but tier 2 assuming an outdoor reduction of 40 percent and tier 3 in case of a reduction of 60 or 80 percent. Its cost amounts to 25 US-Dollars. These examples illustrate that when scarce resources constrain the coverage of an intervention to disseminate clean cookstoves, which can cost upward of 90 US-Dollar, consideration of cooking location should be at least one of the factors that bears on the decision of which region is targeted, prioritizing those regions where indoor cooking predominates. This prioritization applies equally to within country contexts in instances where cooking patterns differ between rural and urban areas.

## **6 Conclusion**

Although large cookstove initiatives are currently slated for implementation, reaching the target of universal adoption of clean fuels and improved cookstoves is a long-term endeavor that will require massive investments extending well beyond current commitments. Given the urgency and breadth of the challenges, including LPG-supply chain bottlenecks, it behooves development agencies to chart a course of improved cookstove distribution that accounts for the interaction of this new technology with cooking behaviors. This paper has argued that the cooking location - whether indoors or outdoors - is a key mediating factor on the effectiveness of clean cookstove adoption. We further

document that outdoor cooking rates are high but vary tremendously between countries and continents as well as between rural and urban areas.

Given this heterogeneity, we regard the fixation on the dissemination of Tier 4 stoves evident in much of the donor community as unfortunate, as it risks missing opportunities to substantially reduce exposure through the distribution of lower cost stoves among households that cook outdoors. It is important to emphasize that these simple improved biomass stoves, which are often a fraction of the cost of more advanced models, also generate additional benefits of improved cooking that are related to deforestation, climate, time and monetary savings (see for example Bensch and Peters, 2013, 2015; Beyene et al., 2015; Jagger and Perez-Heydrich, 2016; Jeuland and Pattanayak, 2012; Martin et al., 2011). Furthermore, affordability is already one of the documented barriers to adoption of lower-cost cooking technologies using market based dissemination (Bensch et al., 2015; Mobarak et al., 2012; Lewis and Pattanayak, 2012), a barrier that would even be higher for Tier 4 stoves. To increase the effectiveness of policy measures, the following lessons should be taken from these results:

In prioritizing regions and stove technologies for dissemination, the effectiveness of a program can be increased by taking the cooking locations into account. While clean cookstoves are likely to be the best option to reduce the exposure to air pollution among households that cook indoors, improved biomass stoves are potentially the more cost-efficient policy intervention in regions where outdoor cooking prevails. Our results give indications for hot spot regions where exposure is the highest owing to cooking location.

Further research is needed on smoke exposure under different ventilation conditions as well as cooking locations using rigorous evaluation methods. For example, negative health effects may also result from disseminating bricked stoves installed in kitchens because

people switch from outside to inside cooking.<sup>9</sup> Furthermore, there may also be a negative impact from ambient air pollution to those cooking outside, though this effect seems to be negligible in comparison to indoor air pollution. Behavioral change interventions, such as health education including sensitization to ventilation, and the coupling of those interventions with cookstove interventions could be one promising avenue for the future but still requires further research (e.g. Barnes, 2014; Grabow et al., 2013; Zhou et al., 2006).

---

<sup>9</sup> Note that this may be aggravated for chimney stoves if the chimneys are not well maintained (see Hanna et al. (2016) and Grimm and Peters (2012)).

## References

- Albalak, R., Frisancho, A. and Keeler, G. (1999). Domestic biomass fuel combustion and chronic bronchitis in two rural Bolivian villages, *Thorax* **54**(11): 1004–1008.
- Amegah, A. K., Quansah, R. and Jaakkola, J. J. K. (2014). Household air pollution from solid fuel use and risk of adverse pregnancy outcomes: a systematic review and meta-analysis of the empirical evidence, *PloS one* **9**(12): e113920.
- Aprovecho Research Center (2016). Stove-emission-model-realtime-indoor\_outdoor-realtime-3.8.16. URL: [http://aprovecho.org/stove-emissions-model-indoor\\_outdoor-realtime-3-8-16/](http://aprovecho.org/stove-emissions-model-indoor_outdoor-realtime-3-8-16/) [last accessed on 16/11/2016].
- Balakrishnan, K., Sankar, S., Parikh, J., Padmavathi, R., Srividya, K., Venugopal, V., Prasad, S. and Pandey, V. L. (2002). Daily average exposures to respirable particulate matter from combustion of biomass fuels in rural households of southern India, *Environmental Health Perspectives* **110**(11): 1069–1075.
- Barnes, B. R. (2014). Behavioural change, indoor air pollution and child respiratory health in developing countries: a review, *International Journal of Environmental Research and Public Health* **11**(5): 4607–4618.
- Bensch, G., Grimm, M. and Peters, J. (2015). Why do households forego high returns from technology adoption? Evidence from improved cooking stoves in Burkina Faso, *Journal of Economic Behavior & Organization* **116**: 187–205.
- Bensch, G. and Peters, J. (2013). Alleviating deforestation pressures? Impacts of improved stove dissemination on charcoal consumption in urban Senegal, *Land Economics* **89**(4): 676–698.
- Bensch, G. and Peters, J. (2015). The intensive margin of technology adoption-experimental evidence on improved cooking stoves in rural Senegal, *Journal of Health Economics* **42**: 44–63.
- Beyene, A. D., Bluffstone, R., Dissanayake, S., Gebreegziabher, Z., Martinsson, P., Mekonnen, A. and Toman, M. (2015). Can Improved Biomass Cookstoves Contribute to REDD+ in Low-Income Countries? Evidence from a Controlled Cooking Test Trial with Randomized Behavioral Treatments, *World Bank Policy Research Paper* (7394).
- Bruce, N. G., Dherani, M. K., Das, J. K., Balakrishnan, K., Adair-Rohani, H., Bhutta, Z. A. and Pope, D. (2013). Control of household air pollution for child survival: estimates for intervention impacts, *BMC Public Health* **13 Suppl 3**: S8.
- Buchner, H. and Rehfuess, E. A. (2015). Cooking and season as risk factors for acute lower respiratory infections in African children: a cross-sectional multi-country analysis, *PloS one* **10**(6): e0128933.
- Burnett, R. T., Pope, C. A. r., Ezzati, M., Olives, C., Lim, S. S., Mehta, S., Shin, H. H., Singh, G., Hubbell, B., Brauer, M. and et al. (2014). An integrated risk function for estimating the global burden of disease attributable to ambient fine particulate matter exposure, *Environmental Health Perspectives* **122**(4): 397–403.



- Dasgupta, S., Huq, M., Khaliqzaman, M., Pandey, K. and Wheeler, D. (2006). Who suffers from indoor air pollution? Evidence from Bangladesh, *Health Policy and Planning* **21**(6): 444–458.
- Dherani, M., Pope, D., Mascarenhas, M., Smith, K. R., Weber, M. and Bruce, N. (2008). Indoor air pollution from unprocessed solid fuel use and pneumonia risk in children aged under five years: a systematic review and meta-analysis, *Bulletin of the World Health Organization* **86**(5): 390–398.
- Ezzati, M. and Kammen, D. M. (2002). The health impacts of exposure to indoor air pollution from solid fuels in developing countries: knowledge, gaps, and data needs, *Environmental Health Perspectives* **110**(11): 1057–1068.
- GACC, Global Alliance on Clean Cookstoves (2011). Igniting change: a strategy for universal adoption of clean cookstoves and fuels.
- GACC, Global Alliance on Clean Cookstoves (2014). The water boiling test – version 4.2.3. URL: <http://cleancookstoves.org/technology-and-fuels/testing/protocols.html> [last accessed on 14/11/2016].
- Grabow, K., Still, D. and Bentson, S. (2013). Test Kitchen studies of indoor air pollution from biomass cookstoves, *Energy for Sustainable Development* **17**(5): 458–462.
- Grimm, M. and Peters, J. (2012). *Improved cooking stoves that end up in smoke?*, Vol. 52 of *RWI: Positionen*, RWI, Essen.
- Hanna, R., Duflo, E. and Greenstone, M. (2016). Up in smoke: the influence of household behavior on the long-run impact of improved cooking stoves, *American Economic Journal: Economic Policy* **8**(1): 80–114.
- Hosgood, H. D., Boffetta, P., Greenland, S., Lee, Y.-C. A., McLaughlin, J., Seow, A., Duell, E. J., Andrew, A. S., Zaridze, D., Szeszenia-Dabrowska, N., Rudnai, P. and et al. (2010). In-home coal and wood use and lung cancer risk: a pooled analysis of the International Lung Cancer Consortium, *Environmental Health Perspectives* **118**(12): 1743–1747.
- ISO, International Organisation for Standardisation (2012). IWA 11 — Guidelines for evaluating cookstove performance. URL: [http://www.iso.org/iso/catalogue\\_detail?csnumber=61975](http://www.iso.org/iso/catalogue_detail?csnumber=61975) [last accessed on 05/11/2016].
- Jagger, P. and Perez-Heydrich, C. (2016). Land use and household energy dynamics in Malawi, *Environmental Research Letters* **11**(12): 125004.
- Jetter, J., Zhao, Y., Smith, K. R., Khan, B., Yelverton, T., Decarlo, P. and Hays, M. D. (2012). Pollutant emissions and energy efficiency under controlled conditions for household biomass cookstoves and implications for metrics useful in setting international test standards, *Environmental Science & Technology* **46**(19): 10827–10834.
- Jeuland, M. A. and Pattanayak, S. K. (2012). Benefits and costs of improved cookstoves: assessing the implications of variability in health, forest and climate impacts, *PLoS one* **7**(2): e30338.
- Jeuland, M., Pattanayak, S. K. and Bluffstone, R. (2015). The economics of household air pollution, *Annual Review of Resource Economics* **7**(1): 81–108.

- Johnson, M., Lam, N., Brant, S., Gray, C. and Pennise, D. (2011). Modeling indoor air pollution from cookstove emissions in developing countries using a Monte Carlo single-box model, *Atmospheric Environment* **45**(19): 3237–3243.
- Lewis, J. J. and Pattanayak, S. K. (2012). Who adopts improved fuels and cookstoves? A systematic review, *Environmental Health Perspectives* **120**(5): 637–645.
- Martin, W. J. n., Glass, R. I., Balbus, J. M. and Collins, F. S. (2011). Public health. A major environmental cause of death, *Science* **334**(6053): 180–181.
- McCracken, J. P., Wellenius, G. A., Bloomfield, G. S., Brook, R. D., Tolunay, H. E., Dockery, D. W., Rabadan-Diehl, C., Checkley, W. and Rajagopalan, S. (2012). Household air pollution from solid fuel use: evidence for links to CVD, *Global Heart* **7**(3): 223–234.
- Mobarak, A. M., Dwivedi, P., Bailis, R., Hildemann, L. and Miller, G. (2012). Low demand for nontraditional cookstove technologies, *Proceedings of the National Academy of Sciences of the United States of America* **109**(27): 10815–10820.
- Po, J. Y. T., FitzGerald, J. M. and Carlsen, C. (2011). Respiratory disease associated with solid biomass fuel exposure in rural women and children: systematic review and meta-analysis, *Thorax* **66**(3): 232–239.
- Pope, C. A. r., Burnett, R. T., Turner, M. C., Cohen, A., Krewski, D., Jerrett, M., Gapstur, S. M. and Thun, M. J. (2011). Lung cancer and cardiovascular disease mortality associated with ambient air pollution and cigarette smoke: shape of the exposure-response relationships, *Environmental Health Perspectives* **119**(11): 1616–1621.
- Rehfuess, E. A., Tzala, L., Best, N., Briggs, D. J. and Joffe, M. (2009). Solid fuel use and cooking practices as a major risk factor for ALRI mortality among African children, *Journal of Epidemiology and Community Health* **63**(11): 887–892.
- Rosa, G., Majorin, F., Boisson, S., Barstow, C., Johnson, M., Kirby, M., Ngabo, F., Thomas, E. and Clasen, T. (2014). Assessing the impact of water filters and improved cook stoves on drinking water quality and household air pollution: a randomised controlled trial in Rwanda, *PloS one* **9**(3): e91011.
- SE4ALL, Sustainable Energy for All (2015). *Progress toward sustainable energy 2015 – Global tracking framework report*, The World Bank, Washington, D.C.
- Smith, K. R., Bruce, N., Balakrishnan, K., Adair-Rohani, H., Balmes, J., Chafe, Z., Dherani, M., Hosgood, H. D., Mehta, S., Pope, D. and Rehfuess, E. (2014). Millions dead: how do we know and what does it mean? Methods used in the comparative risk assessment of household air pollution, *Annual Review of Public Health* **35**: 185–206.
- WHO, World Health Organization (ed.) (2014). *Indoor air quality guidelines: household fuel combustion*, Geneva, Switzerland.
- WHO, World Health Organization (ed.) (2016). *Burning opportunity: clean household energy for health, sustainable development and wellbeing of women and children*, Geneva, Switzerland.
- Yu, F. (2011). Indoor air pollution and children’s health: net benefits from stove and behavioral interventions in rural China, *Environmental and Resource Economics* **50**(4): 495–514.

Zhou, Z., Jin, Y., Liu, F., Cheng, Y., Liu, J., Kang, J., He, G., Tang, N., Chen, X., Banis, E. and Ezzati, M. (2006). Community effectiveness of stove and health education interventions for reducing exposure to indoor air pollution from solid fuels in four Chinese provinces, *Environmental Research Letters* **1**(1): 014010.

# Appendix

**Table A.1:** Sample description

Country	Continent(Region)	Survey year(s)	Number of observations	
			Rural areas	Urban areas
Bangladesh	Asia	2011	12,823	4,291
Benin	Africa(West)	2012	9,631	7,599
Burkina Faso	Africa (West)	2010	10,590	3,444
Burundi	Africa (East)	2010	7,711	718
Cameroon	Africa (Central/South)	2011	6,820	6,951
Comoros	Africa (East)	2012	2,936	1,467
Cote d'Ivoire	Africa (West)	2012	4,921	4,064
Dominican Republic	Latin America	2013	2,909	7,987
Democratic Republic of the Congo	Africa (Central/South)	2014	12,344	5,695
Ethiopia	Africa (East)	2011	12,809	3,569
Gabon	Africa (Central/South)	2012	1,591	7,656
Gambia	Africa (West)	2013	2,480	3,330
Ghana	Africa (West)	2008	5,997	5,385
Guinea	Africa (West)	2012	4,715	2,205
Haiti	Latin America	2012	4,715	2,205
Honduras	Latin America	2012	10,021	10,785
India	Asia	2006	73,293	35,309
Indonesia	Asia	2012	22,156	20,688
Kenya	Africa (East)	2009	6,662	2,315
Lesotho	Africa (Central/South)	2009	6,595	2,771
Liberia	Africa (West)	2013	4,015	5,145
Madagascar	Africa (East)	2009	15,091	2,719
Malawi	Africa (East)	2010	20,676	4,104
Mali	Africa (West)	2013	7,825	2,105
Mozambique	Africa (East)	2011	9,697	4,141
Namibia	Africa (Central/South)	2013	4,718	5,092
Nepal	Asia	2011	9,212	1,513
Niger	Africa (West)	2012	8,815	1,707
Nigeria	Africa (West)	2013	21,344	16,099
Pakistan	Asia	2013	8,529	4,370
Peru	Latin America	2011	7,965	17,366
Philippines	Asia	2013	7,671	7,049
Republic of the Congo	Africa (Central/South)	2012	4,238	7,190
Rwanda	Africa (East)	2010	10,675	1,701
Senegal	Africa (West)	2011	4,016	3,770
Sierra Leone	Africa (West)	2013	8,531	3,845
Togo	Africa (West)	2014	5,285	4,096
Uganda	Africa (East)	2011	7,222	1,578
Zambia	Africa (East)	2014	9,259	6,631
Zimbabwe	Africa (East)	2011	6,463	3,287
<b>Total</b>			405,806	244,917

Source: DHS all country dataset from 2006–2014.

**Table A.2:** Characteristics of the included cookstoves

Cooking device	Category	Fuel	Retail price <i>in US-Dollar</i>
3-stone minimally tended	No stove	Wood	0
3-stone carefully tended	No stove	Wood	0
Envirofit-G330	Natural draft stove	Wood	31
Philips HD4008 Natural Draft	Natural draft stove	Wood	31
Sampada	Natural draft stove	Wood	38
StoveTec Greenfire	Natural draft stove	Wood	9
Upesi Portable	Natural draft stove	Wood	9.5
GERES	Charcoal stove	Charcoal	3.5
Gyapa	Charcoal stove	Charcoal	N/A
Jiko-Ceramic	Charcoal stove	Charcoal	N/A
Jiko-Metal	Charcoal stove	Charcoal	N/A
KCJ Standard	Charcoal stove	Charcoal	6
Kenya Uhai	Charcoal stove	Charcoal	11
StoveTec Charcoal	Charcoal stove	Charcoal	N/A
StoveTec Greenfire, reduced fuel fee	Natural draft stove	Wood	9
Mayon Turbo	Natural draft stove	Rice Hulls	15
Berkeley Darfur	Natural draft stove	Wood	25
Envirofit-G3300, reduced fuel feed	Natural draft stove	Wood	31
Protos	Liquid fuel stove	Plant oil	50
Belonio	Forced draft stove	Rice hulls	40
Philips HD4012 fan	Forced draft stove	Wood	89
Oorja stove	Forced draft stove	Biomass pellets	N/A
StoveTec TLUD	Natural draft stove	Wood pellets	N/A

Source: Jetter et al. (2012)

**Table A.3:** Cookstove emission for the low power scenario

Cooking device	Fuel		Indoor				Outdoor			
			Indoor emissions, tiers and PM <sub>2.5</sub> average concentration level during cooking time		Assumed outdoor reduction level and PM <sub>2.5</sub> average concentration level during cooking time ( $\mu\text{g}/\text{m}^3$ )					
	mg/min	Tier	Concentration level $\mu\text{g}/\text{m}^3$	Concentration level $\mu\text{g}/\text{m}^3$	percent	percent	percent	percent		
3-stone minimally tended	Wood	0	5518	3311	2207	1104				
3-stone carefully tended	Wood	0	3364	2018	1346	673				
Envirofit-G330	Wood	2	888	533	355	178				
Philips HD4008 Natural Draft	Wood	1	2280	1368	912	456				
Sampada	Wood	1	2594	1556	1038	519				
StoveTec Greenfire	Wood	2	1053	632	421	211				
Upesi Portable	Wood	1	2460	1476	984	492				
GERES	Charcoal	3	385	231	154	77				
Gyapa	Charcoal	3	503	302	201	101				
Jiko-Ceramic	Charcoal	3	369	221	148	74				
Jiko-Metal	Charcoal	4	102	61	41	20				
KCJ Standard	Charcoal	4	149	89	60	30				
Kenya Uhai	Charcoal	4	94	56	38	19				
StoveTec Charcoal	Charcoal	3	519	311	208	104				
StoveTec Greenfire, reduced fuel fee	Wood	2	1156	694	462	231				
Mayon Turbo	Rice Hulls	1	2956	1774	1182	591				
Berkeley Darfur	Wood	2	1321	793	528	264				
Envirofit-G3300, reduced fuel feed	Wood	2	731	439	292	146				
Protos	Plant oil	1	2735	1641	1094	547				
Belonio	Rice hulls	2	1234	740	494	247				
Philips HD4012 fan	Wood	2	220	132	88	44				
Oorja stove	Biomass pellets	3	479	287	192	96				
StoveTec TLUD	Wood pellets	3	181	109	72	36				

*Note:* Compared to the high power scenario is the water in the low power scenario not boiled but simmers for 45 minutes.

*Source:* Jetter et al. (2012)