

A Service of



Leibniz-Informationszentrum Wirtschaft Leibniz Information Centre

Chang, Chia-Lin; McAleer, Michael

Working Paper
The Fiction of Full BEKK

Tinbergen Institute Discussion Paper, No. 17-015/III

Provided in Cooperation with:

Tinbergen Institute, Amsterdam and Rotterdam

Suggested Citation: Chang, Chia-Lin; McAleer, Michael (2017): The Fiction of Full BEKK, Tinbergen Institute Discussion Paper, No. 17-015/III, Tinbergen Institute, Amsterdam and Rotterdam

This Version is available at: https://hdl.handle.net/10419/162281

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.



TI 2017-015/III
Tinbergen Institute Discussion Paper



The Fiction of Full BEKK*

Chia-Lin Chang¹ Michael McAleer²

¹Department of Applied Economics, Department of Finance, National Chung Hsing University, Taiwan

²Department of Quantitative Finance, National Tsing Hua University, Taiwan Econometric Institute; Erasmus School of Economics, Erasmus University Rotterdam, The Netherlands; Department of Quantitative Economics, Complutense University of Madrid Spain; Institute of Advanced Sciences Yokohama National University Japan

Tinbergen Institute is the graduate school and research institute in economics of Erasmus University Rotterdam, the University of Amsterdam and VU University Amsterdam.

Contact: <u>discussionpapers@tinbergen.nl</u>

More TI discussion papers can be downloaded at http://www.tinbergen.nl

Tinbergen Institute has two locations:

Tinbergen Institute Amsterdam Gustav Mahlerplein 117 1082 MS Amsterdam The Netherlands

Tel.: +31(0)20 598 4580

Tinbergen Institute Rotterdam Burg. Oudlaan 50 3062 PA Rotterdam The Netherlands

Tel.: +31(0)10 408 8900

The Fiction of Full BEKK*

Chia-Lin Chang

Department of Applied Economics
Department of Finance
National Chung Hsing University
Taiwan

Michael McAleer

Department of Quantitative Finance National Tsing Hua University Taiwan and **Econometric Institute** Erasmus School of Economics Erasmus University Rotterdam The Netherlands and Department of Quantitative Economics Complutense University of Madrid Spain and Institute of Advanced Sciences Yokohama National University Japan

January 2017

^{*} For financial support, the first author wishes to thank the National Science Council, Ministry of Science and Technology (MOST), Taiwan, and the second author acknowledges the Australian Research Council and the National Science Council, Ministry of Science and Technology (MOST), Taiwan.

Abstract

The purpose of the paper is to show that univariate GARCH is not a special case of multivariate

GARCH, specifically the Full BEKK model, except under parametric restrictions on the off-

diagonal elements of the random coefficient autoregressive coefficient matrix, provides the

regularity conditions that arise from the underlying random coefficient autoregressive process,

and for which the (quasi-) maximum likelihood estimates have valid asymptotic properties under

the appropriate parametric restrictions. The paper provides a discussion of the stochastic

processes, regularity conditions, and asymptotic properties of univariate and multivariate

GARCH models. It is shown that the Full BEKK model, which in practice is estimated almost

exclusively, has no underlying stochastic process, regularity conditions, or asymptotic properties.

Keywords: Random coefficient stochastic process, Off-diagonal parametric restrictions,

Diagonal and Full BEKK, Regularity conditions, Asymptotic properties, Conditional volatility, Univariate and multivariate models.

JEL: C22, C32, C52, C58

2

1. Introduction

The most widely estimated univariate and multivariate models of time-varying volatility for financial data, as well as any high frequency data that are measured in days, hours and minutes, is the conditional volatility model. The stochastic processes, regularity conditions and asymptotic properties of the most popular univariate conditional volatility models, such as GARCH (see Engle (1982) and Bollerslev (1986)) and GJR (see Glosten et al. (1993)) are well established in the literature, though McAleer and Hafner (2014) have raised caveats regarding the existence of the stochastic process underlying exponential GARCH (EGARCH) (see Nelson (1990, 1991)).

However, the same cannot be said about multivariate conditional volatility models, specifically Full BEKK (see Baba et al. (1985) and Engle and Kroner (1995)), for which the underlying stochastic processes, regularity conditions and asymptotic properties have either not been established, or are simply assumed rather than derived. These conditions are essential for valid statistical analysis of empirical estimates.

The purpose of the paper is to show that the stochastic process underlying univariate GARCH is not a special case of that underlying multivariate GARCH, except under parametric restrictions on the off-diagonal elements of the random coefficient autoregressive coefficient matrix. The paper provides the regularity conditions that arise from the underlying random coefficient autoregressive process, and for which the (quasi-) maximum likelihood estimates (QMLE) have valid asymptotic properties under the appropriate parametric restrictions.

The Full BEKK model is estimated almost exclusively in practice, despite the fact that it has no underlying stochastic process, regularity conditions, or asymptotic properties, as shown in the proposition and three corollaries.

The plan of the paper is as follows. Section 2 provides a discussion of the stochastic processes, regularity conditions, and asymptotic properties of univariate and multivariate GARCH models. Section 3 shows that the Full BEKK model has no underlying stochastic process, regularity conditions, or asymptotic properties. Section 3 gives some concluding comments.

2. Univariate and Multivariate GARCH Models

2.1 Univariate Conditional Volatility Models

Consider the conditional mean of financial returns for commodity i, in a financial portfolio of m assets, as follows:

$$y_{it} = E(y_{it}|I_{t-1}) + \varepsilon_{it}, \qquad i = 1, 2, ..., m,$$
 (1)

where the returns, $y_{it} = \Delta log P_{it}$, represent the log-difference in financial commodity prices, P_t , I_{t-1} is the information set for all financial assets at time t-I, $E(y_{it}|I_{t-1})$ is the conditional expectation of returns, and ε_{it} is a conditionally heteroskedastic error term.

In order to derive conditional volatility specifications, it is necessary to specify the stochastic processes underlying the returns shocks, ε_{it} . The most popular univariate conditional volatility model, GARCH model, is discussed below.

For a portfolio of m financial assets, consider the random coefficient autoregressive process underlying the returns shocks, ε_{it} , as follows:

$$\varepsilon_{it} = \phi_{it}\varepsilon_{it-1} + \eta_{it}, \qquad i = 1, 2, ..., m, m > 1, \qquad (2)$$

where

$$\phi_{it} \sim iid(0, \alpha_i), \alpha_i \geq 0,$$

$$\eta_{it} \sim iid(0, \omega_i), \, \omega_i \geq 0,$$

 $\eta_{it} = \varepsilon_{it}/\sqrt{h_{it}}$ is the standardized residual,

 h_{it} is the conditional volatility of financial asset i.

Tsay (1987) derived the following conditional volatility of financial asset *i* as an ARCH process (see Engle, 1982):

$$E(\varepsilon_{it}^2|I_{t-1}) \equiv h_{it} = \omega_i + \alpha_i \varepsilon_{it-1}^2, \quad i = 1, 2, ..., m, m > 1,$$
 (3)

where h_t represents conditional volatility, and I_{t-1} is the information set available at time t-1. A lagged dependent variable, h_{t-1} , is typically added to equation (3) to improve the sample fit:

$$h_{it} \equiv E(\varepsilon_{it}^2 | I_{t-1}) = \omega_i + \alpha_i \varepsilon_{it-1}^2 + \beta_i h_{t-1}, \quad \beta_i \in (-1, 1). \tag{4}$$

From the specification of equation (2), it is clear that both ω_i and α_i should be positive as they are the unconditional variances of two different stochastic processes. In equation (4), which is a GARCH(1,1) model for commodity i (see Bollerslev, 1986), the stability condition requires that $\beta_i \in (-1,1)$.

The stochastic process can be extended to asymmetric conditional volatility models (see, for example, McAleer (2014)), and to give higher-order lags and a larger number of alternative commodities, namely up to *m*-1. However, the symmetric process considered here is sufficient to focus the key ideas associated with the purpose of the paper.

As the stochastic process in equation (2) follows a random coefficient autoregressive process, under normality (non-normality) of the random errors, the maximum likelihood estimators (quasi- maximum likelihood estimators, QMLE) of the parameters will be consistent and asymptotically normal. It is worth emphasizing that the regularity conditions include invertibility, which is obvious from equation (2), as:

$$\varepsilon_{it} - \phi_{it}\varepsilon_{it-1} = \eta_{it}$$

so that the standardized shocks can be expressed in terms of the empirical data through equations (1) and (2).

Ling and McAleer (2003) and McAleer et al. (2008) provide general proofs of the asymptotic properties of univariate and multivariate conditional volatility models based on satisfying the regularity conditions in Jeantheau (1998) for consistency, and in Theorem 4.1.3 in Amemiya (1985) for asymptotic normality.

2.2 Multivariate Conditional Volatility Models

The multivariate extension of the univariate ARCH and GARCH models is given in Baba et al. (1985) and Engle and Kroner (1995). It is useful to define the multivariate extension of the relationship between the returns shocks and the standardized residuals, that is, $\eta_{it} = \varepsilon_{it}/\sqrt{h_{it}}$. The multivariate extension of equation (1), namely $y_t = E(y_t|I_{t-1}) + \varepsilon_t$, can remain unchanged by assuming that the three components are now $m \times 1$ vectors, where m is the number of financial assets.

Consider the vector random coefficient autoregressive process of order one, which is the multivariate extension of the univariate process given in equation (2):

$$\varepsilon_t = \Phi_t \varepsilon_{t-1} + \eta_t, \tag{5}$$

where

 ε_t and η_t are $m \times 1$ vectors,

 Φ_t is an $m \times m$ matrix of random coefficients,

 $\Phi_t \sim iid(0, A)$, A is positive definite,

 $\eta_t \sim iid(0, C)$, C is an $m \times m$ matrix.

Vectorization of a full matrix A to $vec\ A$ can have dimension as high as $m^2 \times m^2$, whereas vectorization of a symmetric matrix A to $vech\ A$ can have a smaller dimension of $m(m+1)/2 \times m(m+1)/2$.

In the case where A is a diagonal matrix, with $a_{ii} > 0$ for all i = 1,...,m and $|b_{jj}| < 1$ for all j = 1,...,m, so that A has dimension $m \times m$, McAleer et al. (2008) showed that the multivariate extension of GARCH(1,1) from equation (5) is given as the Diagonal BEKK model, namely:

$$Q_t = CC' + A\varepsilon_{t-1}\varepsilon'_{t-1}A' + BQ_{t-1}B', \tag{6}$$

where A and B are both diagonal matrices. The diagonality of the positive definite matrix A is essential for matrix multiplication as $\varepsilon_{t-1}\varepsilon'_{t-1}$ is an $m \times m$ matrix; otherwise equation (6) could not be derived from the vector random coefficient autoregressive process in equation (5).

McAleer et al. (2008) showed that the QMLE of the parameters of the Diagonal BEKK model were consistent and asymptotically normal, so that standard statistical inference on testing hypotheses is valid.

3. The Fiction of Full BEKK

Consider element *i* of equation (5), which is given as:

$$\varepsilon_{it} = \sum_{j=1}^{m} \phi_{ijt} \varepsilon_{ijt-1} + \eta_{it}, \quad i = 1, 2, \dots, m, m > 1$$
 (7)

which is not equivalent to equation (2) unless $\phi_{ijt} = 0 \ \forall j \neq i$. The stochastic equation (7) is not a random coefficient autoregressive process because of the presence of an additional m-1 random coefficients. Importantly, equation (7) is **not** invertible as the random processes cannot be connected to the data, which requires m equations, such as in equation (5). Consequently, the stochastic process underlying univariate ARCH is not a special case of that underlying multivariate ARCH unless $\phi_{ijt} = 0 \ \forall j \neq i$.

The same condition holds for all i = 1, ..., m, which leads to the following:

Proposition: For the stochastic process underlying univariate ARCH to be a special case of the

stochastic process underlying multivariate ARCH requires the restrictions:

$$\phi_{ijt} = 0 \ \forall j \neq i.$$

A similar condition holds for univariate GARCH and multivariate GARCH.

The Proposition leads to the following corollaries:

Corollary 1: The $m \times m$ matrix of random coefficients, Φ_t , is a diagonal matrix.

Corollary 2: From Corollary 1, it follows that the $m \times m$ weight matrix of (co)variances, A, is a diagonal matrix.

Corollary 3: Corollaries 1 and 2 show that a Full BEKK model, namely where there are no restrictions on the off-diagonal elements in Φ_t , and hence no restrictions in the off-diagonal elements in A, is not possible if univariate ARCH is to be a special case of its multivariate counterpart, BEKK.

Corollary 4: As there are no underlying regularity conditions for Full BEKK, including invertibility, the model cannot be estimated. Therefore, there is no likelihood function, and hence there are also no asymptotic properties of the QMLE of the unknown parameters in Full BEKK.

Corollary 4 is consistent with the proof in McAleer et al. (2008) that the QMLE of Full BEKK has no asymptotic properties, whereas the QMLE of Diagonal BEKK is consistent and asymptotically normal.

For all intents and purposes, Full BEKK does not exist, except by assumption.

4. Conclusion

The Full BEKK model in Baba et al. (1985) and Engle and Kroner (1995), who do not derive the model from an underlying stochastic process, was presented as equation (6), with A and B given as full matrices, with no restrictions on the off-diagonal elements. The Full BEKK model is estimated almost exclusively in practice, despite the fact that it has no underlying stochastic process, regularity conditions, or asymptotic properties, as shown in the proposition and three corollaries.

The full BEKK model can be replaced by the triangular or Hadamard (element-by-element multiplication) BEKK models, with similar problems of identification and (lack of) existence. The full, triangular and Hadamard BEKK models cannot be derived from any known underlying stochastic processes, which means there are no regularity conditions (except by assumption) for checking the internal consistency of the alternative models, and consequently no valid asymptotic properties of the QMLE of the associated parameters (except by assumption).

Moreover, as the number of parameters in a full BEKK model can be as much as 3m(m+1)/2, the "curse of dimensionality" will be likely to arise, which means that convergence of the estimation algorithm can become problematic and less reliable when there is a large number of parameters to be estimated. As a matter of fact, estimation of the full BEKK can be problematic even when m is as low as 5 financial assets. Such computational difficulties do not arise for the diagonal BEKK model. Convergence of the estimation algorithm is more likely when the number of commodities is less than 4, though this is nevertheless problematic in terms of interpretation.

The purpose of the paper was to show that univariate GARCH is not a special case of multivariate GARCH, specifically the Full BEKK model, except under parametric restrictions on a random coefficient autoregressive coefficient matrix. The paper provided the regularity conditions that arise from the underlying random coefficient autoregressive process, and for which the (quasi-) maximum likelihood estimates have valid asymptotic properties under the appropriate parametric restrictions, for the univariate and multivariate GARCH models. It was shown that the Full BEKK model has no underlying stochastic process, regularity conditions, or asymptotic properties.

References

Amemiya, T. (1985), Advanced Econometrics, Harvard University Press, Cambridge, MA, USA.

Baba, Y., R.F. Engle, D. Kraft, and K.F. Kroner (1985), Multivariate simultaneous generalized ARCH, Unpublished manuscript, Department of Economics, University of California, San Diego, CA, USA.

Bollerslev, T. (1986), Generalised autoregressive conditional heteroscedasticity, Journal of Econometrics, 31, 307-327.

Engle, R.F. (1982), Autoregressive conditional heteroscedasticity with estimates of the variance of United Kingdom inflation, Econometrica, 50, 987-1007.

Engle, R.F. and K.F. Kroner (1995), Multivariate simultaneous generalized ARCH, Econometric Theory, 11(1), 122-150.

Glosten, L.R., R. Jagannathan, and D.E. Runkle (1993), On the relation between the expected value and volatility of nominal excess return on stocks, Journal of Finance, 48(5), 1779-1801.

Jeantheau, T. (1998), Strong consistency of estimators for multivariate ARCH models, Econometric Theory, 14, 70-86.

Ling, S. and M. McAleer (2003), Asymptotic theory for a vector ARMA-GARCH model, Econometric Theory, 19, 278-308.

McAleer, M. (2014), Asymmetry and leverage in conditional volatility models, Econometrics, 2(3), 145-150.

McAleer, M., F. Chan, S. Hoti and O. Lieberman (2008), Generalized autoregressive conditional correlation, Econometric Theory, 24(6), 1554-1583.

McAleer, M. and C. Hafner (2014), A one line derivation of EGARCH, Econometrics, 2(2), 92-97.

Nelson, D.B. (1990), ARCH models as diffusion approximations, Journal of Econometrics, 45(1-2), 7-38.

Nelson, D.B. (1991), Conditional heteroskedasticity in asset returns: A new approach, Econometrica, 59(2), 347-370.

Tsay, R. S. (1987), Conditional heteroscedastic time series models, Journal of the American Statistical Association, 82(398), 590-604.