

Obedy, Eric; Ayuya, Oscar Ingasia; Owuor, George; Bokelmann, Wolfgang

Article

Consumption intensity of leafy African indigenous vegetables: towards enhancing nutritional security in rural and urban dwellers in Kenya

Agricultural and Food Economics

Provided in Cooperation with:

Italian Society of Agricultural Economics (SIDEA)

Suggested Citation: Obedy, Eric; Ayuya, Oscar Ingasia; Owuor, George; Bokelmann, Wolfgang (2017) : Consumption intensity of leafy African indigenous vegetables: towards enhancing nutritional security in rural and urban dwellers in Kenya, Agricultural and Food Economics, ISSN 2193-7532, Springer, Heidelberg, Vol. 5, Iss. 14, pp. 1-16, <https://doi.org/10.1186/s40100-017-0082-0>

This Version is available at:

<https://hdl.handle.net/10419/179098>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.



<https://creativecommons.org/licenses/by/4.0/>

RESEARCH

Open Access



Consumption intensity of leafy African indigenous vegetables: towards enhancing nutritional security in rural and urban dwellers in Kenya

Eric Obedy Gido^{1,2*}, Oscar Ingasia Ayuya², George Owuor² and Wolfgang Bokelmann¹

* Correspondence:
gidoeric@gmail.com

¹Department of Agricultural Economics and Social Sciences, Humboldt University of Berlin, Invalidenstrasse 42, 10115 Berlin, Germany

²Department of Agricultural Economics and Agribusiness Management, Egerton University, P.O. Box 536-20115, Egerton, Kenya

Abstract

Estimation of consumption intensity of African indigenous vegetables (AIVs) is important to understand how their utilization can be enhanced at the household level. The study evaluated consumption intensity of leafy AIVs using the zero-inflated negative binomial regression model. A multistage sampling technique was used to select a random sample of 450 rural and urban respondents, and data were collected using a pre-tested semi-structured questionnaire. The findings revealed that consumption intensity of leafy AIVs were higher in rural than in urban dwellers with a mean of four and two times a week, respectively. Age, occupation, household size, diversity of AIV leaves, market distance, awareness of AIV's medicinal benefits and proportion of income allocated to food purchases significantly influenced consumption intensity of leafy AIVs. Strategies that could promote the transfer of AIVs' traditional knowledge to uninformed consumer segments such as male and younger decision-makers could increase the consumption intensity of leafy AIVs in rural dwellers. Similarly, the consumption intensity of leafy AIVs in urban dwellers could increase through the promotion of the value addition activities of sorting and plucking vegetable leaves from their stalks before marketing. Finally, in both rural and urban dwellers, promotion of AIV diversity in food systems through diversified production and well-coordinated market supply chains could increase consumption intensity of leafy AIVs.

Keywords: African indigenous vegetables, Consumption intensity, Micronutrients, Negative binomial regression, Rural, Urban, Vegetable leaves

Background

The nutritional condition of any population depends on the consumption of fruits and vegetables (van der Lans et al. 2012). Health complications of non-communicable diseases are clear indications of inadequate intake of fruits and vegetables (WHO 2002, 2003, 2015; Smith and Eyzaguirre 2007). About 400 g of fruits and vegetables are recommended for consumption per person per day by the World Health Organization (Ruel et al. 2005). However, households in Kenya consume inadequate fruits and vegetables, with the lowest levels reported in the poorest people (Ruel et al. 2005; Ayieko et al. 2008; van der Lans et al. 2012). Urban and rural dwellers in Kenya consume

about 147 and 73 kg of vegetables per capita per year, respectively (Onim and Mwaniki 2008). Additionally, about 26% of the household food budget is expended on fruits and vegetables by urban dwellers in Kenya (Ayieko et al. 2008).

Vegetable consumption in Kenya is constrained by factors related to demographics, socio-psychological, and food characteristics in addition to government and industrial actions (van der Lans et al. 2012). For instance, more women than men prefer vegetables for consumption (Kimiye et al. 2007; Vorster et al. 2007). A rise in income increases the quantity of fruits and vegetables consumed per adult equivalent, with a prominent increase in fruits and sweet and Irish potatoes than for vegetables (Ayieko et al. 2008). Van der Lans et al. (2012) confirmed that lower disposable income hinders vegetable intake. Moreover, diversity in vegetable consumption is a necessary condition for a healthy diet, which requires a proper integration of dark, deep yellow, and starchy vegetables as well as tomatoes and beans in household meals (Oniang'o et al. 2003).

African indigenous vegetables (AIVs) are sources of high-quality nutrition (Abukutsa 2007). They are easily accessible, inexpensive for poor people, and contain minerals and vitamins in levels exceeding those found in most exotic vegetables (Abukutsa 2007; Odhav et al. 2007; Uusiku et al. 2010; Singh et al. 2013). In this regard, AIVs are anticipated to contribute significantly to the global initiative of WHO in increasing the consumption of fruits and vegetables in African countries (Smith and Eyzaguirre 2007). For several decades, AIVs have been consumed in African diets (Smith and Eyzaguirre 2007). Their expansion in production, marketing, and consumption could be attributed to increasing consumer awareness about their health and nutritional benefits (Schippers 2000). High marketing returns have motivated commercialization of AIVs by small-scale farmers, who produce and supply them either individually or collectively in groups (Ngugi et al. 2007; Muhanji et al. 2011; Weinberger et al. 2011). Currently, most food retail outlets sell AIV leaves, and their availability and diversity in high-valued retail outlets such as supermarkets have further induced their consumption in urban areas (Ngugi et al. 2007; Irungu et al. 2008). Changes in lifestyle and availability of cooling-storage facilities have also boosted their consumption levels in urban dwellers (Ruel et al. 2005). Moreover, ethnic background, cultural preferences, inter-marriages, and urbanization are important cultural interactions for enhancing AIV consumption among people from different ethnic origins (Maundu 1997; Kimiye et al. 2007).

A greater proportion of fresh AIV leaves sold in urban markets is supplied from various rural areas at different periods of the year due to variation in production seasons (Ngugi et al. 2007). Thus, availability of fresh leafy AIVs in urban areas is fairly constant across the year (Yang and Keding 2009). Their regular supply in urban markets is also enhanced through irrigation systems, commonly practiced by urban and peri-urban farmers (Yang and Keding 2009). Besides agro-ecological variations, oscillations in AIV production in rural farmers are prompted by a high dependency on rain-fed agriculture (Nekesa and Meso 1997). Significantly, this contributes to supply fluctuations, resulting in a cyclical reduction in the consumption of fresh leafy AIVs, especially in rural dwellers (Smith and Eyzaguirre 2007; Amaza 2009). According to Banwat et al. (2012), seasonal availability influence the

consumption of fresh vegetables. Fresh leafy AIVs are usually consumed in large quantities during rainy seasons due to increased diversity and affordable prices (Kimiye et al. 2007). Even though the supply of fresh vegetables is limited during dry spells, preserved and dried AIV leaves are major sources of nutrition in rural dwellers (Vorster et al. 2007; Yang and Keding 2009). However, nutritional benefits are limited from the consumption of dried rather than fresh leafy AIVs if they are poorly preserved during the drying process (Mibei and Ojijo 2011).

More often, AIV leaves accompany starchy food stuff as side dishes (Maundu 1997; Kimiye et al. 2007; Vorster et al. 2007). Thus, consumption frequency of starch meals could be important in explaining their regularity in household diets. Despite this, AIV consumption depends on household socio-economic variables, vegetable seasonality, retail price, market accessibility, availability of exotic vegetables in food systems, and degree of urbanization (Nekesa and Meso 1997; Yang and Keding 2009). Weinberger and Msuya (2004) revealed that AIV consumption frequency ranges from daily to several times a week during peak seasons and once a week to several times a week during off-peak seasons. Similarly, Amaza (2009) indicated an average AIV consumption frequency of two to three times a week.

Strategies for enhancing food security have emphasized on the production of cereal-based crops (Dimara and Skuras 2003; Cavatassi et al. 2011). Even though cereals significantly contribute to dietary energy requirements (Oniang'o et al. 2003; Koenders 2010; Schipmann and Qaim 2010), they contain inadequate levels of micronutrients (Uusiku et al. 2010; Aaron et al. 2012). Consumption of cereals alone could accelerate occurrence of micronutrient-related diseases. Increasing vegetable consumption intensity is acknowledged as a strategic intervention for addressing micronutrient deficiency and reduces risks associated with degenerative diseases (Hughes and Keatinge 2013; Kamga et al. 2013; Birol et al. 2015). Leafy AIVs are superior vegetables with adequate dietary micronutrients (Singh et al. 2012, 2013). However, their consumption levels and frequency are still low to guarantee such benefits (Ruel et al. 2005; Kimiye et al. 2007; Ngugi et al. 2007).

It is from the foregoing that the study evaluated determinants of consumption intensity of leafy AIVs in Kenya. The study provides empirical evidence comparison on how socio-economic variables determine consumption intensity of leafy AIVs in rural and urban dwellers to cater for spatial heterogeneity in the population. Findings from this study could be important in informing policy formulation and implementation as well as vegetable marketers as they strive to increase AIV consumption to fight micronutrient deficiency and degenerative diseases.

Methods

Study area and sampling design

A stratified multistage sampling approach was used in selecting respondents. In the first stage, a purposive sample of Nairobi, Nakuru, Kisii, and Kakamega counties in Kenya was obtained. Due to heterogeneity in consumer characteristics, Nairobi and Nakuru counties were chosen to represent urban dwellers and Kisii and Kakamega counties were chosen to represent rural dwellers. Kisii and Kakamega are among the counties with large AIV production levels in Kenya; Nakuru and Nairobi

counties are some of the final major markets where AIVs from different production zones are sold. The second and third stages were stratified based on information obtained from sub-county agricultural offices. In the second stage, one sub-county from each county identified as major areas where large volumes of AIVs are produced and/or consumed was chosen.

The third stage involved stratification of market outlets. In urban areas, markets were stratified into supermarkets, green groceries, and local open-air retail outlets. In rural areas, farm gate, green groceries, and local open-air retail outlets were identified. In the fourth stage, simple random sampling method was used to select an equal number of respondents from each retail outlet. Determination of sample size was based on a proportionate to size sampling formulae (Groebner and Shannon 2005). Eventually, 450 respondents were selected, distributed proportionately to the population size at the county level resulting to 168 and 282 respondents in rural and urban areas, respectively.

Data were obtained through a consumer survey in July 2015 using a semi-structured questionnaire. Respondents were interviewed at retail outlets after purchasing leafy vegetables. Respondents were interviewed to obtain information on age, gender, education level, occupation, household size, household income, market distance, duration of AIV consumption, diversity of leafy AIVs at retail outlets, and perceptions regarding AIV retail prices as well as medicinal and nutritional awareness. Nutritional and medicinal awareness of leafy AIVs were assessed using the questions “Is consumption of a diet with leafy AIVs considered; (a) highly nutritious? (b) of great health benefits?” The answers were classified as “Yes” (indicating awareness) and “No” (indicating unawareness). This was followed by further interrogations on positive response, where respondents were asked to name key nutrients found in various leafy AIVs and identify medicinal uses associated with commonly consumed leafy AIVs. The term “nutritional awareness” as used in this study is meant to capture the “participant self-perception of the importance of nutritious leafy AIVs in household diet,” which was adapted from Alkerwi et al. (2015). Similarly, “medicinal awareness” is meant to capture the respondent’s self-perception on therapeutic importance associated with leafy AIVs, which was adapted from previous studies (Olembo et al. 1995; Vaishali and Varsha 2013; Muhammad and Shinkafi 2014).

The coded data were analyzed using the Stata 12 (StataCorp, 2011) computer program. Leafy AIVs identified as commonly consumed vegetables were cowpea (*Vigna unguiculata* L. Walp.), amaranth (*Amaranthus cruentus* L.), spider plant (*Cleome gynandra* L.), African night shade (*Solanum scabrum* Mill.), jute mallow (*Corchorus olitorius* L.), and slender leaf (*Crotalaria brevidens* Benth).

Econometric estimation of consumption intensity of leafy AIVs

Consumption intensity in this study was measured as a number of times a household had consumed leafy AIV per week regardless of the type of AIV crop due to variation in consumer taste and preferences. Daily vegetable consumption is a health recommendation. Adequate benefits from the consumption of leafy AIVs can be realized if a more frequent consumption interval is maintained, and this can increase the vegetable consumption level, which is found to be low in

developing countries. A duration of 1 week was presumed an adequate period for analyzing consumption intensity, and the results generated could enhance the formulation of reliable policies for improving vegetable consumption in developing countries. In this regard, count data models were more suitable for evaluating consumption intensity in this study (Gujarati 2004). From count data model analysis, the probability distribution function of standard Poisson regression follows Greene (2002) and Gujarati (2004) as illustrated in Eq. (1).

$$f(Y_i = y_i | X_i) = \frac{\mu_i^{Y_i} e^{-\mu_i}}{Y_i!} \quad Y_i = 0, 1, 2, \dots; \quad \lambda > 0 \quad (1)$$

where $f(\cdot)$ is the probability that the Y_i value takes a non-negative integer, y_i is the consumption outcome made by household i , X_i is a vector of explanatory variables, and μ_i is a parameter of Poisson distribution associated with X_i . The Y factorial means:

$$Y_i! = Y * (Y-1) * (Y-2) * \dots * 1 \quad (2)$$

The main limitation of standard Poisson regression is the assumed equality between the conditional mean of the data and the variance function (Greene 2002; Gujarati 2004). However, this problem is overcome by estimating the negative binomial regression (NBR) model, in which a cross-section heterogeneity is naturally formulated by introducing an unobserved effect into the conditional mean (Greene 2002). The NBR model is equally inadequate in circumstances where zero outcomes are qualitatively different from positive ones (Greene 2002). Hence, the zero-inflated Poisson (ZIP) model becomes superior to NBR due to its ability to correct the latter problem. The ZIP model has two processes. A binary regression, which characterizes zero outcomes in stage one and a truncated Poisson regression that describes positive outcomes in stage two (Lambert 1992; Greene 2002). Lambert (1992) specified the probability function of the ZIP model as follows:

$$f(y_i) = P\delta_0(y_i) + (1-P)q(y_i) \quad (3)$$

where P is the probability of $\delta_0(y_i)$, $1 - P$ is the probability of $q(y_i)$, $q(y_i)$ is the probability function on non-negative integers, and $\delta_0(y_i)$ is the probability function of delta distribution on zero (distribution that takes only zero value) such that:

$$\delta_0(y_i) = \begin{cases} 1 & \text{if } y_i = 0 \\ 0 & \text{if } y_i = 1, 2, 3, \dots \end{cases} \quad (4)$$

The characterization process distinguishes zero outcomes into two regimes. The first regime comprises of true zeros (perfect state), which indicates a household is a real non-consumer of leafy AIVs. The second regime (imperfect state) indicates a household often consumes leafy AIVs. However, he/she did not consume these vegetables during the study period, and such an outcome is considered a count (Lambert 1992).

Even though NBR and ZIP models correct for equality of the conditional mean and variance of the distribution, they also induce an over-dispersion problem (Mullahy 1986; Greene 2002). Over-dispersion is a condition in which observed variance of a response is greater than the conditional mean while excess zeros arise

from the non-consumption of leafy AIVs (Gurmu and Trivedi 1996). Therefore, over-dispersion and excess zero limitations can simultaneously be resolved by zero-inflated negative binomial (ZINB) regression, which is a more flexible extension of the ZIP model (Minami et al. 2007). Just like the ZIP model, the ZINB model has two combined hurdles that generate the expected counts. These include a binary logit regression to identify zero outcomes associated with count data and NBR to model the count process. The probability distribution function for the ZINB model follows Minami et al. (2007) as shown in Eq. (5).

$$f(y_i|B_i, G_i, \beta, \gamma, \theta) = \begin{cases} P_i + (1-P_i) q(0|\mu_i, \theta) & \text{for } y_i = 0 \\ (1-P_i) q(y_i|\mu_i, \theta) & \text{for } y_i = 1, 2, 3, \dots \end{cases} \quad (5)$$

where B_i and G_i are row vectors of covariate values for the i th observation in imperfect and perfect states, respectively. β and γ are parameter estimates for imperfect and perfect states respectively, θ is the precision or size parameter, and μ_i is the conditional mean for the count data, which is defined as $\mu_i = e^{X_i'\beta}$. From the ZINB regression, the first hurdle (binary logit model) is given as:

$$\text{Logit}(P_i) = \ln\left(\frac{P_i}{1-P_i}\right) = G_i'\gamma \quad (6)$$

while a second hurdle is a distribution for the imperfect state, which assumes the density for a truncated NBR as follows:

$$q(y_i|\mu_i, \theta) = \frac{\Gamma(\theta + y_i)}{\Gamma(\theta)\Gamma(y_i + 1)} \left(\frac{\theta}{\theta + \mu_i}\right)^\theta \left(\frac{\mu_i}{\theta + \mu_i}\right)^{y_i} \quad \text{for } y_i = 0, 1, 2, 3, \dots \quad (7)$$

in which $\Gamma(\cdot)$ is a gamma distribution function and the log-likelihood functions of μ_i are given as $\ln\mu_i = B_i'\beta$. To generate the β , γ , and θ estimates, a log-likelihood function for the ZINB regression is optimized using the maximum likelihood method as in Eq. (8).

$$L(\beta, \gamma, \theta|Y, B, G) = \sum_{i=1}^n \ln f(y_i|B_i, G_i, \beta, \gamma, \theta) \quad (8)$$

Therefore, to identify the determinants influencing the consumption intensity of leafy AIVs, the ZINB model was used in this study. The ZINB model has previously been adopted in studies that involved count data analysis (Yau et al. 2003; Sheu et al. 2004; Minami et al. 2007; Williams 2012; Gido et al. 2015). Explanatory variables used in analyzing the determinants of the consumption intensity of leafy AIVs in rural and urban dwellers (Table 1) were derived from previous studies (Modi et al. 2006; Dovie et al. 2007; Vorster et al. 2007; Amaza 2009; Faber et al. 2010; Weinberger et al. 2011; Matenge et al. 2012; Ayanwale et al. 2016; Gido et al. 2017). These variables were found to significantly influence acceptance, choice, demand, consumption, and other utilization of indigenous vegetables in Sub-Saharan Africa.

Table 1 Definition of variables used in econometric analysis and descriptive statistics

Variable	Definition of variables and its measurement	Rural dwellers	Urban dwellers	Significance
Continuous variables		Mean		t value ^a
Age	Age of the decision-maker ^b in years	43.22	40.62	2.03**
Educ	Years of schooling of the decision-maker	9.45	10.37	-1.88*
H_size	Number of members in the household	5.56	4.38	5.65***
Yr_cons	Years of AIV consumption by the household	23.45	18.78	3.62***
P_income	Proportion of income allocated to food items (in Kenyan shillings)	0.45	0.35	2.29**
V_diversity	Number of different AIV leaves stocked at preferred retail outlets	5.67	6.00	-1.69*
Mrkt_dist	Distance to the nearest preferred retail outlets (in walking minutes)	35.39	18.00	7.41***
Categorical variables		Percentage		χ^2 ratio ^c
Gender	% of male decision-makers	34.32	30.08	0.72
Occup	% of respondents with household cook ^d formally employed	19.26	21.77	0.43
Nutrit	% of respondents informed about nutritional benefits of leafy AIVs	45.49	43.94	0.36
Medic	% of respondents informed about medicinal benefits allied to leafy AIVs	53.42	46.27	4.83*
Price_Per	% of respondents who perceive prices of leafy AIVs are affordable	74.63	85.22	5.37***

*, **, ***Significant at 10, 5, and 1%, respectively

^at test was used to determine significant differences in continuous variables, between rural and urban dwellers

^bDecision-maker is a household member responsible for making key decisions on matters concerning food consumption

^c χ^2 ratio was used to determine relationships in categorical variables, between rural and urban dwellers

^dHousehold cook is the person responsible for preparing household meals

Results and discussion

Descriptive results

Descriptive and summary statistics between the rural and urban dwellers varied (Table 1). On average, rural decision-makers were significantly elderly and less educated. Rural dwellers had more members with wider experience in AIV consumption and allocated larger proportions of income for food use. While access to a greater diversity of leafy AIVs was higher in urban dwellers, distance to preferred market outlets was significantly longer in rural areas. Moreover, rural dwellers were more informed about the medicinal benefits associated with indigenous vegetables, and the perceived retail prices for leafy AIVs were more affordable.

The average consumption intensity of leafy AIVs in both rural and urban dwellers was thrice a week (Table 2). Rural dwellers had a significantly higher

Table 2 Weekly consumption intensity of leafy AIVs among rural and urban dwellers

Sample	Mean	Minimum	Maximum	Standard deviation
Rural dwellers	3.92	0	7	1.89
Urban dwellers	2.32	0	4	1.34
Rural and urban dwellers	2.89	0	7	1.74
t value	8.80*			

*Significant at 1%

consumption intensity of about four times a week compared to urban dwellers with a mean of twice a week. Interestingly, the highest consumption intensity in urban dwellers was about four times a week while some rural dwellers were found to consume leafy AIVs on daily basis during the study period. To the lower end, zero consumption level was revealed in both rural and urban dwellers.

Determinants of consumption intensity of leafy AIVs

Four count data models were consecutively estimated to determine a regression that best fits the data in explaining the consumption intensity of leafy AIVs. Firstly, the standard Poisson regression was estimated (Appendix 1). To determine the appropriateness of the standard Poisson regression, there was a need to estimate the NBR model (Appendix 1). Results from the NBR revealed evidence of over-dispersion, since alpha was greater than zero with a significant likelihood ratio test, implying that NBR was more appropriate for the data than the standard Poisson model. Thirdly, ZIP regression was estimated (Appendix 2). Results showed that z -test for Vuong was significant indicating the ZIP model was more appropriate than the standard Poisson regression. Lastly, ZINB regression was estimated (Table 2), where the statistical suitability of the model was tested using the zip option test for ZINB over ZIP and the Vuong test for ZINB over standard NBR. The zip option test had a significant likelihood ratio test for an alpha of zero, suggesting that ZINB regression was more appropriate than ZIP regression. The z -test for Vuong was also significant, implying the ZINB model was better than standard NBR. A comparison of model statistics from the four regressions indicated that the ZINB model was more suitable for the data collected in this study.

The bottom half of Table 3 shows the logistic model results on evaluation of the perfect state. Gender and education level of the decision-maker significantly predicted true zeros in consumption intensity of leafy AIVs. In rural dwellers, male decision-makers increased the likelihood of a count being zero for the consumption intensity of leafy AIVs. This finding was not surprising since earlier studies (Amaza 2009; Weinberger et al. 2011) revealed that traditional knowledge about AIVs was more likely in women than men. Women are more informed about healthier diets and are found to consume more vegetables than men (Baker and Wardle 2003). Moreover, vegetable preparation activities of sorting and plucking leaves from their stalks are presumed more tedious and time involving (Abukutsa 2010; Matenge et al. 2012). Thus, men are unlikely to devote sufficient time for AIV preparation compared to women.

Likewise, a count for consumption intensity of leafy AIVs was less likely a true zero in urban dwellers with more educated decision-makers. This result was interesting, indicating that earlier campaign efforts of promoting consumption of leafy AIVs in urban areas were not in vain. Perhaps this is because better-educated households are more likely to attain dietary information on nutritious and healthier food items like indigenous vegetables (Sanlier and Karakus 2010; Gido et al. 2017). Contrary to findings in this study, Gido et al. (2017) revealed that African night shade was less likely to be accepted for consumption by more-educated urban decision-makers. However, spider plant, African night shade, and slender leaf were

Table 3 ZINB model results on determinants of consumption intensity of leafy AIVs

Variables	Rural dwellers		Urban dwellers	
	Coef.	Std. err.	Coef.	Std. err.
Negative binomial regression				
Gender	-0.0871	0.1124	-0.0436	0.0944
H_size	-0.0438*	0.0240	-0.0142	0.0258
Age	0.0173***	0.0058	0.0066	0.0046
Educ	-0.0068	0.0130	-0.0030	0.0114
Occup	-0.2771	0.1418	-0.3346**	0.1125
P_income	-0.1736***	0.0599	0.0857	0.0591
Yr_cons	0.0001	0.0045	-0.0008	0.0046
V_diversity	0.0407*	0.0241	0.0504**	0.0219
Mrkt_dist	0.0128	0.0534	0.0998*	0.0586
Nutrit	-0.1557	0.2685	0.0705	0.2199
Medic	0.4704*	0.2622	0.0869	0.1450
Price_Per	-0.1817	0.1311	-0.0737	0.1118
Constant	0.9820**	0.4204	1.3299***	0.4012
Logistic regression for zero-inflation				
Gender	1.4748*	0.7858	-0.2638	0.4344
H_size	0.1195	0.1334	-0.0651	0.1138
Educ	-0.0640	0.0810	-0.1119***	0.0432
P_income	-0.1796	0.3252	-0.2035	0.2594
Price_Per	-1.4133	0.8750	0.6698	0.6915
Constant	-1.3626	1.8718	-0.0859	1.5666
/lnalpha	-2.2581***	0.3486	-2.2572***	0.2687
Alpha	0.1046	0.0364	0.1046	0.0281
	$\chi^2(01) = 19.79$		$\chi^2(01) = 32.33$	
LR test of alpha = 0	$\Pr \geq \chi^2 = 0.0000***$		$\Pr \geq \chi^2 = 0.0000***$	
Vuong test	$z = 2.70$		$z = 5.58$	
	$\Pr > z = 0.0034***$		$\Pr > z = 0.0000***$	

*, **, ***Significant at 10, 5, and 1%, respectively

more likely to be accepted by their rural counterparts for consumption. There is a need for further investigation on this aspect.

The top portion of Table 3 contains NBR results on counties that were an imperfect state. Several variables significantly influenced the consumption intensity of leafy AIVs. Household size significantly and negatively influenced the consumption intensity of leafy AIVs in rural dwellers. This indicates that the consumption intensity of leafy AIVs was less likely in large rural households and this outcome was as predicted due to the anticipated wider variation in taste and preferences for different leafy AIVs. Large household size indicates that more quantities of AIV leaves are required, and this implies that more income would be necessary for such voluminous purchases (Dovie et al. 2007; Ayanwale et al. 2016). Consequently, more time is required to pluck sufficient quantities of leaves from AIV stalks in readiness for cooking, thereby reducing consumption intensity in large households due to the tedious process of vegetable preparation. Findings by Ayanwale et al. (2016) indicated that households with large membership were more likely to

increase the consumption of fluted pumpkin and reduce the consumption of English spinach because it was less prevalent in the market compared to other underutilized indigenous vegetables. Similarly, the bitter taste in slender leaf constrained its acceptance for consumption in urban dwellers with a large household size (Gido et al. 2017).

In rural dwellers, the intensity of AIV consumption significantly increased with advancement in age of the decision-maker. Elderly rural dwellers are more likely to possess adequate cultural knowledge on AIVs as opposed to the youth and urbanized dwellers (Jansen van Rensburg et al. 2007; Faber et al. 2010; Matenge et al. 2012). Moreover, traditional knowledge of AIV preparation and cooking alongside their medicinal and nutritional benefits are likely higher in elderly rural people. Perhaps adequacy in indigenous knowledge enhances consumption intensity of leafy AIVs in elderly rural dwellers (Okeno et al. 2003; Smith and Eyzaguirre 2007; Waudo et al. 2007). Ayanwale et al. (2016) found similar results, where the demand for fluted pumpkin and English spinach were high in elderly consumers. However, demand for garden egg was lower in elderly consumers because the process of preparing it for consumption was tedious.

Formal employment of the household cook significantly and negatively influenced the consumption intensity of leafy AIVs in urban dwellers. Even though household meals have traditionally been prepared by women in most African countries, such cultural obligations are slowly changing due to changes in gender roles (WHO 2000; Fontana and Natali 2008). Most urban dwellers with formal employment attain such opportunities far away from their residential homes and more often return home late from work. This implies that they are left with little time for food preparation (Gido et al. 2017). Given that leafy AIV crops require more time for preparation and cooking (Ruel et al. 2005; Matenge et al. 2012), their consumption intensity is less likely in households where the family cook has employment opportunity. According to Kimiywe et al. (2007), business and full-time employed people consume less indigenous vegetables compared to non-employed people and casual laborers. Gido et al. (2017) found similar results, where amaranth and spider plant vegetables were less accepted for consumption by rural dwellers with the household cook formally employed.

In rural dwellers, consumption intensity was negatively influenced by the proportion of income allocated for food use. This indicates that households with larger budgetary allocation for food use reduce consumption intensity of leafy AIVs. However, this does not imply AIV leaves are inferior goods since the quantity purchased was not emphasized in this study. Past studies found that AIVs are perceived by wealthier people and urbanized dwellers as food meant for “poor-rural man” (Modi et al. 2006; Jansen van Rensburg et al. 2007; Faber et al. 2010). Similar findings were revealed by Frazao et al. (2007), where an increase in household income does not necessarily increase vegetable consumption. This suggests that an improvement in household welfare, through a general increase in income, might not favor consumption of leafy AIVs as opposed to other food commodities like meat products (Bett et al. 2012).

The diversity of leafy AIV crops at retail outlets significantly increased their consumption intensity in rural and urban dwellers. These findings demonstrate that

availability of numerous AIV crops in markets stimulates their regularity in consumption. Probably this is because access to more AIV leaves presents a wider range of alternative crops for households to choose vegetables preferable by all household members (Gido et al. 2017). A higher diversity of leafy AIVs encourages consumption of crops which substitute or complement each other in a vegetable recipe. Moreover, it enhances vegetable rotation in household diets, thereby reducing the monotony of consuming a few vegetable crops. Increasing diversity of AIV leaves in consumption has advantages of ensuring that key micronutrients are attained given that each crop has unique micronutrient composition (Singh et al. 2012, 2013).

The time taken to reach preferred vegetable retail outlets was positive and significantly influenced consumption intensity of leafy AIVs in urban dwellers. The findings were surprising since distant market outlets are expected to constrain consumer access to household goods (Vorster et al. 2007). Markets are major sources of food for most urban dwellers, and consumers prefer higher-quality vegetables, which are perceived to be sold in high-valued retail outlets such as supermarkets and groceries (Ngugi et al. 2007; Irungu et al. 2008). Such retail outlets are sparsely distributed in urban areas, subjecting consumers to walk longer distances to access quality vegetables. Results by Gido et al. (2017) complement these findings where consumers were more likely to obtain complementary leafy AIVs from distant retail outlets. Contrary to these findings, consumers are less willing to shop from far distant markets that involve more time for traveling (Maruyama and Wu 2014; Gido et al. 2016).

Awareness of medicinal benefits associated with indigenous vegetables was positive and significantly influenced the consumption intensity of leafy AIVs in rural dwellers. These findings were in agreement with the study expectations and imply that informed rural dwellers about curative and therapeutic components found in AIVs were likely to consume them more regularly. Such traditional knowledge regarding the selection and utilization of leafy AIVs is highly expected in rural areas (Gido et al. 2017). In particular, rural dwellers are more equipped with skills relating to identifying vegetables that contain oxidant compounds for protecting and healing malnutrition-related diseases (Yang and Keding 2009; Singh et al. 2013).

Conclusions

The study evaluated determinants of consumption intensity of leafy AIVs in rural and urban dwellers using the zero-inflated negative binomial regression model. Findings indicated a higher consumption intensity of leafy AIVs in rural dwellers compared to urban dwellers with a mean of four and two times a week, respectively. Higher diversity of leafy AIVs at retail outlets increased their consumption intensity in both rural and urban dwellers. In urban dwellers, formal employment of the household cook reduced the consumption intensity of leafy AIVs while increased distance to retail outlets had a contrary effect. In rural dwellers, elderly decision-makers with more information on AIVs' medicinal benefits increased vegetable consumption intensity. However, large households with a greater proportion of income allocated for food use reduced vegetable consumption intensity.

The findings from this study have relevant policy implications regarding value addition strategies, sensitization of consumers on traditional knowledge about AIV utilization and increasing diversity of indigenous vegetables in food systems.

Interventions that could promote awareness programs where traditional knowledge regarding indigenous vegetables is transferred to uninformed consumer segments such as male and younger decision-makers could increase the consumption intensity of leafy AIVs in rural dwellers. This can be achieved through the circulation of brochures written in commonly used dialects and an establishment of AIV “food clinics,” where consumers can seek information about AIVs and other indigenous food items in general. In addition, integrating awareness programs on local or ethnic radio and television stations, where consumers are informed about traditional knowledge regarding AIVs in languages they understand well, could be important.

Strategies that could promote vegetable value addition activities can increase consumption intensity of leafy AIVs in urban dwellers. This could involve retailers to stock AIV leaves already sorted and plucked from their stalks, thereby reducing the time required for the preliminary stages of vegetable preparation before cooking. Finally, policies that could increase the diversity of AIV leaves at retail outlets through diverse production and well-coordinated market supply chains could increase vegetable consumption intensity in both rural and urban dwellers. This study recommends the need for a similar future research that could consider seasonal variation in vegetable availability, which might affect consumer behavior regarding the consumption intensity of leafy AIVs.

Appendix 1

Table 4 Standard Poisson regression and NBR results on determinants of consumption intensity of leafy AIVs

Variable	Standard Poisson regression				Negative binomial regression (NBR)			
	Rural dwellers		Urban dwellers		Rural dwellers		Urban dwellers	
	Coef.	Std. err.	Coef.	Std. err.	Coef.	Std. err.	Coef.	Std. err.
Gender	-0.1840**	0.0836	-0.1918	0.1428	0.0347	0.1495	0.0144	0.0695
H_size	-0.0502***	0.0174	-0.0488	0.0308	0.0050	0.0408	0.0113	0.0179
Age	0.0141***	0.0041	0.0140*	0.0075	0.0029	0.0073	0.0039	0.0034
Educ	0.0015	0.0094	0.0002	0.0165	0.0121	0.0184	0.0110	0.0084
Occup	-0.2884***	0.1077	-0.2440	0.1845	-0.1626	0.1797	-0.1824**	0.0869
P_income	-0.1313***	0.0408	-0.1274*	0.0755	0.1396	0.0942	0.1455***	0.0448
Yr_cons	0.0019	0.0031	0.0016	0.0057	-0.0007	0.0073	-0.0024	0.0034
V_diversity	0.0458***	0.0173	0.0472	0.0320	0.0472	0.0365	0.0495	0.0166
Mrkt_dist	0.0177	0.0369	-0.0029	0.0656	0.0667	0.0913	0.0652	0.0440
Nutrit	-0.1143	0.1919	-0.1642	0.3221	0.1191	0.3486	0.1390	0.1625
Medic	0.7309***	0.1874	0.7140**	0.2981	0.0775	0.2408	0.0756	0.1135
Price_Per	-0.0459	0.0944	-0.0774	0.1687	-0.1673	0.1833	-0.1630*	0.0842
Constant	0.5966*	0.3117	0.7503	0.5266	1.1863*	0.6343	1.1532***	0.2990
/lnalpha					-1.1629	0.2321	-0.5039	0.1512
Alpha					0.3126	0.0725	0.6042	0.0914
					$\chi^2(01) = 73.83$		$\chi^2(01) = 247.79$	
LR test of alpha = 0					$Pr \geq \chi^2 = 0.0000***$		$Pr \geq \chi^2 = 0.0000***$	

*, **, *** indicates significance level at 10, 5, and 1%, respectively

Appendix 2

Table 5 ZIP model results on determinants of consumption intensity of leafy AIVs

Variables	Rural dwellers		Urban dwellers	
	Coef.	Std. err.	Coef.	Std. err.
Negative binomial regression				
Gender	-0.0818	0.0842	-0.0522	0.0702
H_size	-0.0431**	0.0175	-0.0097	0.0191
Age	0.0169***	0.0042	0.0066*	0.0034
Educ	-0.0055	0.0095	-0.0032	0.0083
Occup	-0.2795***	0.1073	-0.3406***	0.0869
P_income	-0.1734***	0.0445	0.0833*	0.0441
Yr_cons	0.0001	0.0032	-0.0008	0.0034
V_diversity	0.0372**	0.0175	0.0488***	0.0158
Mrkt_dist	0.0200	0.0399	0.1115**	0.0445
Price_Per	-0.1594*	0.0959	-0.0687	0.0838
Medic	0.4328**	0.1952	0.0757	0.1081
Nutrit	-0.1583	0.1961	0.0824	0.1607
Constant	1.0022***	0.3151	1.2892***	0.3014
Logistic regression for zero-inflation				
Gender	1.3484*	0.7102	-0.2624	0.4251
H_size	0.1120	0.1260	-0.0632	0.1113
Educ	-0.0586**	0.0744	-0.1094	0.0425
P_income	-0.1313	0.3038	-0.2074	0.2547
Price_Per	-1.3257	0.8059	0.6444	0.6600
Constant	-1.2665	1.7958	-0.0565	1.5180
Vuong test	z = 3.03; Pr ≥ 0.0012***		z = 3.03; Pr ≥ 0.0012***	

*, **, *** indicates significance level at 10, 5, and 1%, respectively

The z-test is significant across, indicating that the ZIP model is more appropriate for the data than the standard Poisson regression model

Acknowledgements

The authors are grateful for research grants from the HORTINLEA (Horticultural Innovations and Learning for Improved Nutrition and Livelihood in East Africa) project funded by the Federal Ministry of Education and Research and the Federal Ministry for Economic Cooperation and Development of Germany. We recognize the cooperation received from respondents during the consumer surveys. The study was undertaken through a collaboration between Humboldt University of Berlin, Germany, and Egerton University, Kenya. The views expressed herein are solely those of the authors and not of the affiliated institutions.

Authors' contributions

Four authors contributed to the success of this work. EOG and WB conceptualized the paper. EOG and OIA managed the literature searches and designed the methodology. EOG, OIA and WB designed the questionnaire. EOG coordinated the field survey, data analysis and write up of the first draft. EOG, OIA, GO and WB managed interpretation of the analysis. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 30 March 2016 Accepted: 24 June 2017

Published online: 14 July 2017

References

- Aaron GJ, Laillou A, Wolfson J, Moench-Pfanner R (2012) Fortification of staple cereal flours with iron and other micronutrients: cost implications of following World Health Organization-endorsed recommendations. *Food Nutr Bull* 33(4):336–343
- Abukutsa OMO (2007) The diversity of cultivated African leafy vegetables in three communities in western Kenya. *Afr J Food Agric Nutr Dev* 7(3):1–15

- Abukutsa OMO (2010) African indigenous vegetables in Kenya: Strategic repositioning in the horticultural sector. Inaugural Lecture, Jomo Kenyatta University of Agriculture and Technology, Nairobi, Kenya. 30th April.
- Alkerwi A, Sauvageot N, Malan L, Shivappa N, Hébert JR (2015) Association between nutritional awareness and diet quality: evidence from the observation of cardiovascular risk factors in Luxembourg (ORISCAV-LUX) study. *Nutrients* 7:2823–2838
- Amaza SP (2009) An analysis of traditional African vegetables and sweet potato consumer demand in Kenya and Tanzania. Farm Concern, The World Vegetable Centre, International Potato Centre and Urban Harvest, Arusha, Tanzania, Project report
- Ayanwale AB, Amusan CA, Adeyemo VA, Oyedele DJ (2016) Analysis of household demand for underutilized indigenous vegetables. *Int J Vegetable Sci* 22(6):570–577
- Ayieko MW, Tschirley DL, Mathenge MW (2008) Fresh fruit and vegetable consumption patterns and supply chain systems in urban Kenya. Implications for policy and investment priorities. Tegemeo Institute of Agricultural Policy and Development, Egerton University, Working Paper 16.
- Baker AH, Wardle J (2003) Sex differences in fruit and vegetable intake in older adults. *Appetite* 40:269–275
- Banwati ME, Lar LA, Daber J, Audo S, Lassa S (2012) Knowledge and intake of fruit and vegetable consumption among adult in urban community in north central, Nigeria. *Nigeria Health JI* 12(1):12–15
- Bett HK, Musyoka MP, Peters KJ, Bokelmann W (2012) Demand for meat in the rural and urban areas of Kenya: a focus on the indigenous chicken. *Economics Research International*. doi.org/10.1155/2012/401472.
- Birof E, Meenakshi JV, Oparinde A, Perez S, Tomlins K (2015) Developing country consumers' acceptance of biofortified foods: a synthesis. *Food Sec* 7:555–568
- Cavatassi R, Lipper L, Narloch U (2011) Modern variety adoption and risk management in drought prone areas: insights from sorghum farmers of eastern Ethiopia. *Agric Econ* 42(3):279–292
- Dimara E, Skuras D (2003) Adoption of agricultural innovations as a two-stage partial observability process. *Agric Econ* 28(3):187–196
- Dovie DBK, Shackleton CM, Witkowski ETF (2007) Conceptualizing the human use of wild edible herbs for conservation in South African communal lands. *J Environ Manag* 84:146–156
- Faber M, Oelofse A, van Jaarsveld PJ, Wenhold FAM, van Rensburg WJ (2010) African leafy vegetables consumed by households in the Limpopo and KwaZulu-Natal provinces in South Africa. *S Afri J Clin Nutri* 23(1):30–38
- Fontana M, Natali L (2008) Gendered patterns of time use in Tanzania: public investment in infrastructure can help? Paper prepared for the IFPRI Project on evaluating the long-term impact of gender-focused policy interventions. University of Sussex, Brighton
- Frazao E, Andrews M, Smallwood D, Prell M (2007) Food spending patterns of low-income households: will increasing purchasing power results in healthier food choices? Economic Information Bulletin number 29-4. Economic Research Service and the United States Department of Agriculture, Washington, DC
- Gido OE, Ayuya IO, Owuor G, Bokelmann W (2016) Consumer's choice of retail outlets for African indigenous vegetables: empirical evidence among rural and urban households in Kenya. *Cogent Food Agric* 2(1):1–14
- Gido OE, Ayuya IO, Owuor G, Bokelmann W (2017) Consumer acceptance of leafy African indigenous vegetables: comparison between rural and urban dwellers. *International Journal of Vegetable Science*. doi.org/10.1080/19315260.2017.1293758.
- Gido OE, Sibiko WK, Ayuya IO, Mwangia KJ (2015) Demand for agricultural extension services among small-scale maize farmers: micro-level evidence from Kenya. *J Agric Educ Ext* 21(2):177–192
- Greene WH (2002) *Econometric analysis*, fifth edition. Prentice-Hall, Upper Saddle River, New York
- Groebner DF, Shannon PW (2005) *Business statistics, A decision making approach*. Pentice Hall, Upper Saddle River, New Jersey
- Gujarati DN (2004) *Basic econometrics*, 4th edn. McGraw-Hill/Irwin, New York
- Gurmu S, Trivedi PK (1996) Excess zeros in count models for recreational trips. *J Bus Econ Stat* 14(4):469–477
- Hughes JA, Keatinge JDH (2013) The nourished millennium: how vegetables put global goals for healthy, balanced diets within reach. In: Holmer R, Linwattana G, Nath P, Keatinge JDH (eds) *Proceedings of regional symposium on high value vegetables in Southeast Asia: production, supply and demand (SEAVEG 2012)*. AVRDC, Chiang Mai, Thailand
- Irungu C, Mburu J, Maundu P, Grum M, Hoeschle ZI (2008) The effect of market development on-farm conservation of diversity of African leafy vegetables around Nairobi. 12th EAAE Congress, People, food and environments: Global trends and European strategies, 26-29 August 2008, Gent, Belgium.
- Jansen van Rensburg WS, Van Averbeke W, Slabbert R, Faber M, Van Jaarsveld P, Van Heerden I, Wenhold F, Oelofse A (2007) African leafy vegetables in South Africa. *Water South Africa* 33:317–326
- Kamga RT, Kouame C, Atangana AR, Chagomoka T, Ndango R (2013) Nutritional evaluation of five African indigenous vegetables. *J Horticultural Res* 21(1):99–106
- Kimiywe J, Waudo J, Mbithe D, Maundu P (2007) Utilization and medicinal value of indigenous leafy vegetables consumed in urban and peri-urban Nairobi. *Afr J Food Agric Nutr Dev* 7(4):1–15
- Koenders D (2010) Feasibility study to include sorghum and meat into the WFP basket in Kenya. Report for the WFP, Kenya
- Lambert D (1992) Zero-inflated Poisson regression, with an application to defects in manufacturing. *Technometrics* 34(1):1–14
- Maruyama M, Wu L (2014) Quantifying barriers impeding the diffusion of supermarkets in China: the role of shopping habits. *J Retail Consum Serv* 21:383–393
- Matenge STP, van der Merwe D, De Beer D, Bosman MJC, Kruger A (2012) Consumers' beliefs on indigenous and traditional foods and acceptance of products made with cow pea leaves. *Afr J Agric Res* 7(14):2243–2254
- Maundu PM (1997) The status of traditional vegetable utilization in Kenya. In: Guarino L (ed) *Proceedings of the IPGRI International workshop on genetic resources of traditional vegetables in Africa: conservation and use*. ICRAF-HQ, Nairobi, Kenya

- Mibei EK, Ojijo NKO (2011) Effects of processing on chemical composition of four African leafy vegetables. *Elec J Env Agricult Food Chem Title* 10(11):3121–3131
- Minami M, Lennert-Cody CE, Gao W, Roman-Verdesoto M (2007) Modelling shark by catch: the zero-inflated negative binomial regression model with smoothing. *Fish Res* 84(2):210–221
- Modi M, Modi AT, Hendriks S (2006) Potential role for wild vegetables in household food security: a preliminary case study in Kwazulu-Natal, South Africa. *Afr J Food Agric Nutr Dev* 6:1–13
- Muhammad S, Shinkafi MA (2014) Ethnobotanical survey of some medicinal important leafy vegetables in North Western Nigeria. *J Med Plant Res* 8(1):6–8
- Muhanji G, Roothaert RL, Webo C, Mwangi S (2011) African indigenous vegetable enterprises and market access for small-scale farmers in East Africa. *Int J Agric Sustain* 9(1):194–202
- Mullahy J (1986) Specification and testing of some modified count data models. *J Econ* 33:341–365
- Nekesa P, Meso B (1997) Traditional African vegetables in Kenya: Production, marketing and utilization. Promoting the conservation and use of underutilized and neglected crops. 16. In: Guarino L (ed) *Traditional African vegetables. Proceedings of the IPGRI International Workshop on genetic resources of traditional vegetables in Africa: Conservation and use 29-31 August 1995, ICRAF-HQ, Nairobi, Kenya. IPGRI, Rome, Italy*
- Ngugi IK, Gitau R, Nyoro J (2007) Access to high value markets by smallholder farmers of African indigenous vegetables in Kenya: re-governing markets innovative practice series. International Institute for Environment and Development, London
- Odhav B, Beekrum S, Akula U, Baijnath H (2007) Preliminary assessment of nutrition value of traditional vegetables in Kwa Zulu-Natal, South Africa. *J Food Consumption Analy* 20:430–435
- Okeno JA, Chebet DK, Mathenge PW (2003) Status of indigenous vegetables in Kenya. *Acta Hort* 621:95–100
- Olembi NK, Fedha SS, Ngaira ES (1995) Medicinal and agricultural plants of Ikolomani, Kakamega. District. Development Partners
- Oniang'o RK, Mutuku JM, Malaba SJ (2003) Contemporary African food habits and their nutritional and health implications. *Asia Pac J Clin Nutr* 12:231–236
- Onim M, Mwaniki P (2008) Cataloguing and evaluation of available community/farmers-based seed enterprises on African indigenous vegetables (AIVs) four ECA countries, Entebe, Uganda
- Ruel MT, Minot N, Smith L (2005) Patterns and determinants of fruit and vegetable consumption in sub-Saharan Africa: a multicounty comparison. *FAO/WHO workshop on fruit and vegetable for health, 1 -3 September 2004. Kobe, Japan.*
- Sanlier N, Karakus SS (2010) Evaluation of food purchasing behaviour of consumers from supermarkets. *Br Food J* 112(2):140–150
- Schipmann C, Qaim M (2010) Spillovers from modern supply chains to traditional markets: product innovation and adoption by smallholders. *Agric Econ* 41(3-4):361–371
- Schippers RR (2000) African indigenous vegetables. An overview of the cultivated species. Natural Resources Institute/ACP-EU Technical Centre for Agricultural and Rural Cooperation, Chatham, UK
- Sheu M, Hu T, Keeler TE, Ong M, Sung H (2004) The effect of a major cigarette price change on smoking behaviour in California: a zero-inflated negative binomial model. *Health Econ* 13(8):781–791
- Singh S, Singh DR, Salim KM, Srivastava A, Singh LB, Srivastava RC (2012) Estimation of proximate composition, micronutrients and phytochemical compounds in traditional vegetables from Andaman and Nicobar Islands. *Int J Food Sci Nutr* 62(7):765–773
- Singh S, Singh DR, Singh LB, Chand S, Dam RS (2013) Indigenous vegetables for food and nutritional security in Andaman and Nicobar Islands, India. *Int J Agric Food Sci Technol* 4(5):503–512
- Smith IF, Eyzaguirre P (2007) African leafy vegetables: their role in World Health Organization's global fruit and vegetable initiative. *Afr J Food Agric Nutr Dev* 7(3):1–17
- StataCorp (2011) *Stata statistical software: release 12. StataCorp. LP, College station, TX*
- Uusiku PN, Oelofse A, Duodu KG, Bester MJ, Faber M (2010) Nutritional value of leafy vegetables of sub-Saharan Africa and their potential contribution to human health: a review. *J Food Compos Anal* 23(6):499–509
- Vaishali SK, Varsha DJ (2013) Traditional leafy vegetables: a future herbal medicine. *Int J Agric Food Sci* 3(2):56–58
- Van der Lans C, Snoek H, de Boer F, Elings A (2012) Vegetable chains in Kenya: production and consumption of vegetables in the Nairobi metropolis. Wageningen UR Centre for Development Innovation Rapport GTB-1130, Wageningen, The Netherlands. (<http://edepot.wur.nl/216710>). Accessed 29 Nov 2014
- Vorster IHJ, van Rensburg WJ, Van Zijl J, Venter SL (2007) The importance of traditional leafy vegetables in South Africa. *Afr J Food Agric Nutr Dev* 7(4):1–13
- Waudu J, Kimeywe J, Mbithe D, Maundu P (2007) Utilization and medical value of indigenous leafy vegetables consumed in urban and peri-urban Nairobi. *Afr J Food Agric Nutr Dev* 7(4):27–32
- Weinberger K, Msuya J (2004) Indigenous vegetables in Tanzania—significance and prospects. Asian Vegetable Research and Development Center, Technical Bulletin No 31, Publication 04-600, Shanhua, Taiwan.
- Weinberger K, Pasquini M, Kasambula P, Abukutsa OMO (2011) Supply chains for indigenous vegetables in urban and peri-urban areas of Uganda and Kenya: a gendered perspective. In: Mithoefer D, Waibel H (eds) *Vegetable production and marketing: socio-economic research. CAB International, Wallingford, UK*, pp 169–181
- Williams RK (2012) The external costs of wind farm development on the high plains: are developers making an effort to minimize these costs? Selected paper prepared for presentation at the Southern Agricultural Economics Association annual meeting, Birmingham, AL, February 4-7.
- World Health Organization (WHO) (2000) Healthy nutrition: the role of women. Report on a WHO meeting, Murmask, Russian Federation, 14th- 15th June, 2000.
- World Health Organization (WHO) (2002) The world health report 2002—reducing risks, promoting healthy life. World Health Organisation, Geneva, Switzerland
- World Health Organization (WHO) (2003) Fruit and vegetable promotion initiative—report of the meeting, 25-27/08/03. World Health Organisation, Geneva, Switzerland

- World Health Organization (WHO) (2015) Healthy diet. Fact sheet number 394. World Health Organization, Geneva, Switzerland. Accessed 10 Mar 2017.
- Yang RY, Keding GB (2009) Nutritional contributions of important African indigenous vegetables, pp. 105-143. In: Shackleton CM, Pasquini MW, Drescher AW (eds) African indigenous vegetables in urban agriculture. Earthscan, London
- Yau KW, Wang K, Lee AH (2003) Zero-inflated negative binomial mixed regression modelling of over-dispersed count data with extra zeros. *Biometrical J* 45(4):437-452

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ springeropen.com
