

A Service of

ZBW

Leibniz-Informationszentrum Wirtschaft Leibniz Information Centre for Economics

Zahedi, Mortaza; Mondani, Farzad; Eshghizadeh, Hamid Reza

#### Article

# Analyzing the energy balances of double-cropped cereals in an arid region

**Energy Reports** 

#### **Provided in Cooperation with:** Elsevier

*Suggested Citation:* Zahedi, Mortaza; Mondani, Farzad; Eshghizadeh, Hamid Reza (2015) : Analyzing the energy balances of double-cropped cereals in an arid region, Energy Reports, ISSN 2352-4847, Elsevier, Amsterdam, Vol. 1, pp. 43-49, https://doi.org/10.1016/j.egyr.2014.11.001

This Version is available at: https://hdl.handle.net/10419/187803

#### Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

#### Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.



NC ND https://creativecommons.org/licenses/by-nc-nd/4.0/



### WWW.ECONSTOR.EU

#### Energy Reports 1 (2015) 43-49

Contents lists available at ScienceDirect

**Energy Reports** 

journal homepage: www.elsevier.com/locate/egyr

## Analyzing the energy balances of double-cropped cereals in an arid region



<sup>a</sup> Department of Agronomy and Plant Breeding, Faculty of Agriculture, Isfahan University of Technology, Isfahan, Iran

<sup>b</sup> Department of Agronomy and Plant Breeding, Campus of Agriculture and Natural Resources, Razi University, Kermanshah, Iran

#### ARTICLE INFO

Article history: Received 28 September 2014 Received in revised form 19 November 2014 Accepted 22 November 2014 Available online 21 January 2015

Keywords: Cereal Double cropping Environmental health Energy efficiency Productivity Specific energy

#### ABSTRACT

Efficient use of energy in agroecosystems will reduce environmental problems, prevent destruction of natural resources and serve to promote sustainable agriculture as an economical production system. The aim of this study was to investigate the energy use efficiency in four double cropping systems including: wheat-silage corn (W–SC), barely–silage corn (B–SC), barely–grain corn (B–GC) and barely–rice (B–R) in the arid regions of Isfahan province, Iran. Data used in this study were collected from 73, 45, 38, 18, 18 wheat, barley, silage corn, grain corn and rice farms, respectively, personal interview using semi-structured questionnaire during 2010. The results indicated that the total energy consumed were 140,422, 128,979, 121,360 and 172,962 MJ ha<sup>-1</sup> for the W–SC, the B–GC, the B–SC and the B–R cropping systems, respectively. The share of diesel fuel by 43.36% (W–SC), 43.93% (B–GC), 42.82% (B–SC) and 49.40 % (B–R) was the highest input. This was followed by fertilizer (W–SC: 24.70%, B–GC: 25.12%, B–SC: 27.05 and B–R: 16.11) and water (W–SC: 10.54%, B–GC: 11.76%, B–SC: 10.73 and B–R: 13.85), respectively. The energy use efficiency was found as 1.70 for W–SC, 1.65 for B–GC, 1.64 for B–SC and 1.03 for B–R double cropping systems, respectively. According to the research results the W–SC, B–SC, B–GC and B–R double cropping systems were more efficient in terms of energy, respectively.

© 2014 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

#### 1. Introduction

Energy is a critical input in agricultural production systems. The energy used in agriculture was directly related to environmental factors such as soil and climatic conditions, amount of inputs and techniques employed in production (Esengun et al., 2007). The link between agriculture and energy is very close. Agriculture itself uses energy and is also a supplier of energy in the form of bio-energy (Alam et al., 2005). Energy used in agriculture has developed in response to increasing populations, the limited supply of arable land and a desire for increasing standards of living (Shahan et al., 2008). All inputs and outputs of a cropping system can be expressed in terms of energy. Hence, energy input and output are essential factors for determining energy efficiency and the environmental impact of crop production. However, energy utilization and output differs widely among crops, production systems and management intensity (Rathke et al., 2007).

\* Corresponding author. E-mail address: f.mondani@razi.ac.ir (F. Mondani).

Changes in farm technology over time have increased the amount of energy used in crop production (Rathke and Diepenbrock, 2006). The predominant feature for increasing crop production is the use of a large amount of energy either directly or indirectly in the form of fuel, electricity and fertilizers (Haj-SeyedHadi et al., 2009). Environmental problems such as those associated with soil, water pollution and CO<sub>2</sub> and N<sub>2</sub>O emissions that contribute to global warming are related to intensive use of energy. Energy analysis of agricultural ecosystems seems to be a promising approach to investigate and assess efficiency, environmental problems and their relations to sustainability (Khan et al., 2007). It is also used to compare different production systems (Ghasemi-Mobtaker et al., 2010). Efficient use of energy in agriculture will minimize environmental problems, prevent destruction of natural resources and serve to promote sustainable agriculture as an economical production system (Esengun et al., 2007; Erdal et al., 2007). The relation of energy input and energy output in the agroecosystems have been investigated by many researchers for many crops such as sugar beet (Asgharipour et al., 2012; Yousefi et al., 2014), tomato (Rezvani Moghaddam et al., 2011), pulses (Koocheki et al., 2011) and cotton (Zahedi et al., 2014).

In the Mediterranean regions such as Isfahan province when irrigation water is available, the double cropping systems can be

http://dx.doi.org/10.1016/j.egyr.2014.11.001







<sup>2352-4847/© 2014</sup> The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4. 0/).

improved income of farmers and might be helped to sustainability of agricultural activities. Double cereal systems differences in management practices such as farm technology, tillage and intensity, have considerable effects on energy input and efficiency of crop production systems. Browning (Browning, 2011) indicated that soybean double-cropped after barley has the potential to yield equal to or greater than full-season soybean or doublecropped soybean following wheat, but its relative yield is very dependent on growing conditions in Virginia and the Mid-Atlantic, USA. Therefore, aims of this study were (i) to determine the total amount of input–output energy used in four double cropping systems (wheat–silage corn, barely–silage corn, barely–grain corn and barely–rice), (ii) to determine energy use efficiency, (iii) to determine the best double-cropped cereals based on energy efficiency, in Isfahan province of Iran as a Mediterranean region.

#### 2. Material and methods

The present study was conducted in Isfahan province located in central Iran (geographical coordinates 30°43' and 34°27'N and  $49^{\circ}36'$  and  $55^{\circ}31'E$ ). The total area of the province is 105,937 km<sup>2</sup> and the total farming area is 360,181 km<sup>2</sup>, of that the share of cereal (wheat (Triticum aestivum), barley (Hordeum vulgare), rice (Oryza sativa) and corn (Zea mays)) is about 57% (206,172 ha). Four double cropping systems consist: wheat-silage corn (W-SC), barely-silage corn (B-SC), barely-grain corn (B-GC) and barely-rice (B-R) were determining energy use, to investigate the energy use efficiency, and to make an economical analysis. Information was collected from cereal farmers using a face-to-face questionnaire during 2010. In addition to the data obtained by surveys, previous studies of related organizations such as the Ministry of Agriculture of Iran (MAI) (Browning, 2011) were also used for this research. The number of operations involved in cereal rotation production systems and their energy requirements influence the final energy balance. The sample size was calculated using the Neyman method (Newbold, 1994):

$$n = \frac{N \times S^2}{(N-1)S_x^2 + S^2}$$
(1)

where: *n*, is the required sample size; *N*, is number of farmers in the target population; *S* is standard deviation,  $S_X$ , is standard deviation of sample mean ( $S_X = d/z$ ), *d*, is the permissible error in the sample size, and was determined as 10% of the mean for a 95% confidence interval and *z* is the reliability coefficient (1.96, which represents 95% reliability). Based on this calculation the size of 73 for wheat, 45 for barley, 38 for silage corn, 18 for grain corn, and 18 for rice farms were considered as sampling sizes.

Energy efficiency of the agricultural system has been evaluated by the energy ratio between output and input. Human labor, machinery, diesel oil, fertilizer, pesticides and seed amounts and output yield values of cereal production systems have been used to estimate energy ratios (Alam et al., 2005). Energy equivalents shown in Table 1 were used for estimations (Haj-SeyedHadi et al., 2009; Khan et al., 2007; Erdal et al., 2007). The sources of mechanical energy used on the selected farms included tractors and diesel fuel. Mechanical energy was computed on the basis of total fuel consumption (l ha<sup>-1</sup>) in different operations. Therefore, the energy consumed was calculated using conversion factors and expressed in MJ ha<sup>1</sup> (Tsatsarelis, 1991). The energy of a tractor and its equipment reveals the amount of energy needed for unit weights and calculates repair and care energy, transport energy, total machine weight and average economic life. Based on energy equivalents of inputs and outputs (Table 1), energy use efficiency, energy productivity, specific energy, energy intensiveness and net energy were

#### Table 1

Energy equivalent of inputs and outputs in agricultural production.

Particulars	Unit	Energy equivalent (MJ unit <sup>-1</sup> )
A. Inputs		
1. Human labor	h	1.95
2. Machinery	h	62.7
3. Diesel fuel	1	50.23
4. Chemical fertilizers		
(a) Nitrogen (N)	kg	75.46
(b) Phosphate (P <sub>2</sub> O <sub>5</sub> )	kg	13.07
(c) Potassium (K <sub>2</sub> O)	kg	11.15
(d) Micro	kg	120.00
5. Manure	kg	0.30
6. Chemicals		
(a) Herbicides	kg or l	238.3
(b) Pesticide	1	101.2
(c) Fungicide	kg	181.9
7. Electricity	kWh	3.6
8. Water for irrigation	m <sup>3</sup>	1.02
9. Seeds (wheat)	kg	20.10
10. Seeds (barely)	kg	14.7
11. Seeds (corn)	kg	14.7
12. Seeds (rice)	kg	14.7
B. Outputs		
1. Wheat grain yield	kg	14.7
2. Wheat straw yield	kg	2.25
3. Barely grain yield	kg	14.7
4. Barely straw yield	kg	2.25
5. Corn grain yield	kg	14.7
6. Corn straw yield	kg	2.25
7. Rice grain yield	kg	14.7
8. Rice straw yield	kg	2.25

calculated by the following equations (Demircan et al., 2006):

Energy use efficiency = 
$$\frac{Energy \ output \ (MJ \ ha^{-1})}{Energy \ input \ (MJ \ ha^{-1})}$$
 (2)

Energy productivity 
$$= \frac{crops \ output \ (Kg \ ha^{-1})}{Energy \ input \ (MJ \ ha^{-1})}$$
(3)

Specific energy = 
$$\frac{Energy input (MJ ha^{-1})}{crops output (Kg ha^{-1})}$$
 (4)

Energy intensiveness = 
$$\frac{Energy input (MJ ha^{-1})}{cost of cultivation (\$ ha^{-1})}$$
 (5)

Indirect energy included energy embodied in seeds, chemical fertilizers, herbicide, pesticide, fungicide, farmyard manure and machinery; while direct energy was evaluated in terms of human labor, diesel, electricity and water for irrigation used in the cereal rotation production systems. Non-renewable energy included diesel, electricity, chemical fertilizers, herbicides; pesticides, fungicides and machinery; and renewable energy consisted of human labor, farmyard, seeds and water for irrigation, farmyard manure.

#### 3. Results and discussion

#### 3.1. Structures of farms

The average field size was about 20.2 ha for wheat, 14.4 ha for barley, 5.5 ha for silage corn, 2.6 ha grain corn, and 0.5 ha for rice in according to information provided by the survey. Planting areas for wheat, barley, rice and grain corn were 139,426, 47,288, 17,452 and 2006 ha, and the production of these crops was 561,652, 177,893, 99,407 and 13,838 tons, respectively. All necessary cultural practices such as soil tillage, seedbed preparation, planting

Management practices of wheat, barley, rice and corn for grain and silage.

Operations	Wheat	Barely	Seed corn	Seed silage	Rice	Millet
Names of varieties	Rosahan, Kavir, Pishtaz, Native cultivars	Valphajr, Reyhan, Native cultivars	SC704, SC700	SC704, SC700	Native cultivars (Sorkhe, Gerdesefid)	Jam, Kermanshahi, Karaj 12-60-31
Land preparation tractor used: 285 MF	Moldboard plow, Disc harrows,	Moldboard plow, Disc harrows,	Moldboard plow, Disc harrows,	Chisel, Disc harrows	Moldboard plow, Disc harrows	Chisel-Disc harrows
Land preparation period	Late October–Mid September	Early October–Early September	Mid May	Late June	Early June	June
Average tilling number	2.2	2.2	2.2	1.2	2.2	1.0
Planting period	Early December–Late September	Mid October–Mid September	Mid June	Early July	Late June	June
Fertilization period (Before planting)	Late October–Mid September	Early October–Early September	Mid May	Late June	Early June	June
Fertilization period (Top dressing)	Mid March–Late March	Mid March-Late March	Late July–Early August	Early August	-	-
Average number of fertilization	2.2	1.2	1.5	1.5	1.2	1.0
Irrigation period	September–Early June	September–Early June	June–Late September	June–Late September	June-Late September	June–September
Average number of irrigation	12.5	11.5	7.2	6	-	3
Spraying period	May-June	May-June	Late July–Early August	-	August	July
Average number of	1	0.6	1.8	0.8	1	1
Harvesting period	Late June	Late June	Late September	Late September	Late September	November

#### Table 3

Energy consumption and energy input-output relationship of wheat.

Energy	Quantity per unit area (ha)	Energy equivalent (MJ unit <sup>-1</sup> )	Total energy equivalent (MJ)	Percentage of total energy input (%)
Input				
Human labor (h)	495.4	1.95	966.0	1.4
Machinery (h)	52.0	62.70	3260.5	4.8
Diesel fuel (1)	579.3	50.23	29100.8	42.7
Nitrogen (kg)	161.9	75.46	12221.9	17.9
Phosphate (P <sub>2</sub> O <sub>5</sub> ) (kg)	137.4	13.07	1796.4	2.6
Potassium (K <sub>2</sub> O) (kg)	106.2	11.15	1184.7	1.7
Manure (kg)	6400	0.30	1920.0	2.8
Micro (kg or l)	4.6	120.0	558.3	0.8
Treflan (l)	3.9	238.32	928.2	1.4
Pesticide (Diazinon) (1)	1.2	101.20	121.4	0.2
Fungicide (Carboxin) (kg)	2.0	216.90	432.0	0.63
Electricity (kWh)	800.0	3.60	2880.0	4.2
Water for irrigation (m <sup>-3</sup> )	6700.0	1.02	6834.0	10.03
Seed (kg)	293.5	20.1	5899.4	8.7
Total energy input (MJ) Outputs			68104.1	100.00
Wheat grain vield (kg)	6700.50	14.90	98497.3	90.5
Bean straw yield (kg)	4600.30	2.25	10350.7	9.5
Total energy output (MI)			108848.0	
Energy efficiency			1.6	

methods, planting and harvest period were determined and presented in Table 2. Our results also indicated that about 84% and 16% of the total planting area in cereal production system was irrigated farms and dryland farms, respectively. All farms were in personal possession. The method and timing of management practices for different crops during the growing season are shown in Table 2.

#### 3.2. Analysis of input-output energy

Total input energy consumed for cereal crops with respect to different input and agronomical practices are shown in Tables 3–7. Most of the input energy for wheat production was related to diesel

fuel, nitrogen and water for irrigation by 42.7, 17.9 and 10.03%, respectively. The same trend was observed for barely, grain and silage corn but it was different in rice as inputs of water for irrigation (15.7%) and electricity (10.4%) were higher than nitrogen. So, the share of diesel fuel was the highest energy input for all crops. Similar result have been observed for wheat (Canakci et al., 2005; Ghorbani et al., 2011; Khan et al., 2009), barely (Canakci et al., 2005; Khan et al., 2009), corn (Canakci et al., 2005), rice (Khan et al., 2009) and other irrigated crops such canola and sunflower (Sheikh-Davoodi and Houshyar, 2009) despite the differences in the arrangement in the first second or third category. Given the predominant role of diesel fuel, irrigation and fertilization in accounting for sequestered energy in agricultural systems in this

Energy	Quantity per unit area (ha)	Energy equivalent (MJ unit <sup>-1</sup> )	Total energy equivalent (MJ)	Percentage of total energy input
Input				
Human labor (h)	350.3	1.95	683.1	1.2
Machinery (h)	47.3	62.70	2965.9	5.1
Diesel fuel (1)	486.9	50.23	24457.7	41.4
Nitrogen (kg)	183.4	75.46	13841.8	23.4
Phosphate $(P_2O_5)(kg)$	121.3	13.07	1584.8	2.7
Potassium (K <sub>2</sub> O) (kg)	91.7	11.15	1022.1	1.7
Manure (kg)	2600.0	0.30	780.0	1.3
Micro (kg or l)	1.4	120.0	162.6	0.3
Treflan (l)	3.1	238.3	737.8	1.3
Pesticide (Diazinon) (1)	1.7	101.20	167.9	01.3
Fungicide (Carboxin) (kg)	2.7	216.00	578.1	0.9
Electricity (kWh)	700	3.60	2520.0	4.3
Water for irrigation $(m^{-3})$	6000.0	1.02	6120.0	10.4
Seed (kg)	232.7	14.70	3420.7	5.8
Total energy input (MJ) Outputs			59042.5	100.00
Barely grain yield (kg)	5081.6	14.07	74699.5	88.5
Barely straw yield (kg)	4300.5	2.25	9676.1	11.5

Total energy output (MJ)

Energy efficiency

Energy consumption and energy input-output relationship of grain corn.

Energy	Quantity per unit area (ha)	Energy equivalent (MJ unit <sup>-1</sup> )	Total energy equivalent (MJ)	Percentage of total energy input (%)
Input				
Human labor (h)	378.9	1.95	756.6	1.1
Machinery (h)	33.4	62.70	2069.1	3.0
Diesel fuel (l)	641.2	50.23	32205.7	46.1
Nitrogen (kg)	173.1	75.46	13057.7	18.7
Phosphate (P <sub>2</sub> O <sub>5</sub> ) (kg)	90.5	13.07	1182.4	1.7
Potassium (K <sub>2</sub> O) (kg)	57.5	11.15	641.4	0.9
Manure (kg)	8125.0	0.30	2437.5	3.5
Micro (kg or l)	7.6	120.0	918.6	1.3
Treflan (l)	5.1	238.32	1213.8	1.7
Pesticide (Diazinon) (l)	1.6	101.20	161.9	0.2
Fungicide (Carboxin) (kg)	3.1	216.90	669.6	1.0
Electricity (kWh)	1330.0	3.60	4788.0	6.8
Water for irrigation (m <sup>-3</sup> )	8870.0	1.02	9047.4	12.9
Seed (kg)	51.7	14.7	759.9	1.1
Total energy input (MJ)			69936.7	100.00
Outputs				
Corn grain yield (kg)	8880.50	14.7	129360.0	100.0
Total energy output (MI)			129360.0	
Energy efficiency			1.85	

843756

1.43

region, it is evident that any attempt to reduce energy input should begin by finding to reduce these inputs. The machinery management and using efficient equipment to reduce direct use of diesel fuel energy, increasing nitrogen use efficiency (due to mainly embodied energy of nitrogen, 75.46 MJ kg<sup>-1</sup>) with different approaches such as application of nitrogen only base on the soil analysis and using nitrogen in several time as topdressing.

In this region, average of irrigation water that used for successful wheat, barley, rice, grain and silage corn production are around 6700, 6000, 17,500, 8870 and 7821 m<sup>3</sup> per each hectare, respectively. So, reduction of water irrigation may be achieved by reducing the amount of water supplied as effective use of water, deficit irrigation strategy, using, where possible, alternative irrigation systems or improving irrigation and pumping efficiency (Tsatsarelis, 1991).

The total energy input of 140,422 MJ ha<sup>-1</sup> (W: 68,104 MJ ha<sup>-1</sup> + SC: 72,318 MJ ha<sup>-1</sup>), 128,979 MJ ha<sup>-1</sup> (B: 59,043 MJ ha<sup>-1</sup> + GC: 69,937 MJ ha<sup>-1</sup> BSC), 121,360 MJ ha<sup>-1</sup> (B: 59,043 MJ ha<sup>-1</sup> + SC: 72,318 MJ ha<sup>-1</sup>) and 172,962 MJ ha<sup>-1</sup> (B: 59,043 MJ ha<sup>-1</sup> + R: 113,920 MJ ha<sup>-1</sup>) were required for different double cropping

systems, respectively (Table 8). In literature, the results showed that total energy input were 51,040 MJ ha<sup>-1</sup> for full-season wheat and 44,866 for full-season barley (Sahabi et al., 2012), and 72,743 for full-season corn (Safa et al., 2010), respectively.

(%)

Grain and straw yield of crops are shown in Tables 3–7. Total energy output per hectare was 108,848, 84,375.6, 129,360, 130,981.5, and 93,690.3 MJ ha<sup>-1</sup> in wheat, barley, grain corn, silage corn and rice production systems, respectively. The highest output energy (239,829 MJ ha<sup>-1</sup>) was obtained in W–SC and the lowest (178,066 MJ ha<sup>-1</sup>) was in the B–R double-cropped system. Energy consumption and energy input–output in different double-cropped cereals are shown in Table 9.

Energy use efficiency in wheat, barley, grain corn, silage corn, and rice was 1.6, 1.43, 1.85, 1.81 and 0.82, respectively. The doublecropped cereals, W–SC had the highest energy efficiency (1.70) and barely–rice had the lowest (1.03). The research results were consistent with finding reported by other authors, such as: 1.70 for wheat and 1.83 for barley in irrigated farming in northeast of Iran (Sahabi et al., 2012) and 2.8 for wheat and 3.8 for maize in Antalya region,

Energy consumption and energy i	nput-output relationshi	ip of silage corn.			
Energy	Quantity per unit area (ha)	Energy equivalent (MJ unit <sup>-1</sup> )	Total energy equivalent (MJ)	Percentage of total energy input (%)	
Input					
Human labor (h)	489.5	1.95	954.5	1.3	
Machinery (h)	43.1	62.70	2698.1	3.7	
Diesel fuel (1)	633.1	50.23	31798.8	44.0	
Nitrogen (kg)	214.4	75.46	16181.7	22.4	
Phosphate $(P_2O_5)(kg)$	90.5	13.07	1182.4	1.6	
Potassium (K <sub>2</sub> O) (kg)	57.5	11.15	641.4	0.9	
Manure (kg)	8125.0	0.30	2473.5	3.4	
Micro (kg or l)	7.6	120.0	918.6	1.3	
Treflan (l)	5.1	238.32	1213.8	1.7	
Pesticide (Diazinon) (l)	1.6	101.20	161.9	0.2	
Fungicide (Carboxin) (kg)	3.1	216.90	669.6	0.9	
Electricity (kWh)	1230.0	3.60	4428.0	6.1	
Water for irrigation (m <sup>-3</sup> )	7821.0	1.02	7977.4	11.1	
Seed (kg)	71.7	14.7	1054.0	1.5	
Total energy input (MJ) Outputs			72317.7	100.00	
Corn straw yield (kg) Total energy output (MJ) Energy efficiency	58214.30	2.25	130981.5 130981.5 1.81	100.0	

Table 6

Energy consumption and energy input-output relationship of rice.

Energy	Quantity per unit area (ha)	Energy equivalent (MJ unit <sup>-1</sup> )	Total energy equivalent (MJ)	Percentage of total energy input (%)	
Input					
Human labor (h)	881.3	1.95	1718.5	1.5	
Machinery (h)	58.1	62.70	3642.9	3.2	
Diesel fuel (1)	1214.9	50.23	61024.9	53.6	
Nitrogen (kg)	116.0	75.46	8750.8	7.7	
Phosphate $(P_2O_5)(kg)$	91.4	13.07	1195.3	1.1	
Potassium $(K_2O)(kg)$	56.2	11.15	627.2	0.5	
Manure (kg)	4123.0	0.30	1236.9	1.1	
Micro (kg or l)	5.6	120.0	678.6	0.6	
Treflan (l)	4.9	238.32	1166.2	1.1	
Pesticide (Diazinon) (1)	1.3	101.20	131.6	0.1	
Fungicide (Carboxin) (kg)	1.1	216.90	237.6	0.2	
Electricity (kWh)	3300.0	3.60	11880.0	10.4	
Water for irrigation $(m^{-3})$	17500.0	1.02	17850.0	15.7	
Seed (kg)	257.1	14.7	3779.4	3.3	
Total energy input (MJ) Outputs			113919.7	100.00	
Rice grain vield (kg)	5920.40	14.7	87029.9	92.9	
Rice straw vield (kg)	2960.20	2.25	6660.5	7.1	
Total energy output (MI)			93690.3		
Energy efficiency			0.82		

#### Table 8

Total energy input in the form of direct, indirect, renewable energy for different double cropping systems.

Source	Wheat-Silage corn			Barely-Se	Barely-Seed corn			Barely-Silage corn			Barely-Rice		
	W	SC	Total	В	GC	Total	В	SC	Total	В	R	Total	
Direct energy <sup>a</sup>	39,780.8	45,158.8	84,939.6	33,780.7	46,797.7	80,578.4	33,780.7	45,158.8	78,939.5	33,780.7	92,473.5	126,254.2	
Indirect energy <sup>b</sup>	28,323.2	27,159.0	55,482.2	25,261.7	23,139.0	48,400.7	25,261.7	27,159.0	52,420.7	25,261.7	21,446.3	46,708	
Renewable energy <sup>c</sup>	15,619.3	12,423.4	28,042.7	11,003.7	13,001.5	24,005.2	11,003.7	12,423.4	23,427.1	11,003.7	24,584.8	35,588.5	
Non-renewable	39,780.8	59,894.3	99,675.1	48,038.7	56,935.2	104,973.9	48,038.7	59,894.3	107,933	48,038.7	89,334.9	137,373.6	
energy <sup>d</sup>													
Total energy input	68,104	72,318	140,422	59,043	69,937	128,979	59,043	72,318	131,360	59,043	113,920	172,962	
<sup>a</sup> Indicates human labor,	diesel, elect	ricity and w	ater.										

<sup>b</sup> Indicates seeds, chemical fertilizers (NPK), herbicide (Treflan and Basagran), pesticide (Diazinon), fungicide (Carboxin) and machinery.

<sup>c</sup> Indicates human labor, seeds and water.

<sup>d</sup> Indicates diesel, electricity, chemical fertilizers (NPK), herbicide (Treflan and Basagran), pesticide (Diazinon), fungicide (Carboxin) and machinery.

Turkey (Canakci et al., 2005) under the conventional farming system, respectively. It seems that suitable condition as climate and soil properties is one of the important reasons for more efficient agriculture systems in Turkey in comparison with Iran in these reports. In wheat, barley and rice farms overall energy input–output ratio is very low compared to farms at Australia (Khan et al., 2009), where it was 9.21, 8.21 and 6.70, respectively. In case of wheat, energy use efficiency in other parts of the world as well as New Zealand (Barber, 2004), Turkey (Canakci et al., 2005), India

Energy input-output ratio in four double cropping systems consist: wheat-silage corn (W-SC), barely-silage corn (B-SC), barely-grain corn (B-GC) and barely-rice (B-R).

Source	Unit	Wheat-S	ilage corn		Barely–Grain corn			Barely–Silage corn			Barely-Rice		
		W	SC	Total	В	GC	Total	В	SC	Total	В	R	Total
Total energy input	MJ ha <sup>-1</sup>	68,104	72,318	140,422	59,043	69,937	128,979	59,043	72,318	131,360	59,043	113,920	172,962
Total energy output	MJ ha <sup>-1</sup>	108,848	130,981	239,829	84,376	129,360	213,736	84,376	130,982	215,357	84,376	93,690	178,066
Energy efficiency	-	1.6	1.81	1.70	1.43	1.85	1.66	1.43	1.81	1.64	1.43	0.82	1.03
Energy intensiveness	MJ \$ <sup>-1</sup>	36.5	39.5	76	39.3	38.7	78	39.3	39.5	78.8	39.3	35.7	75
Specific energy	MJ kg <sup>-1</sup>	0.1	1.2	1.3	11.6	7.9	19.5	11.6	1.2	12.8	11.6	0.1	11.7
Energy productivity Net energy	kg MJ <sup>-1</sup> MJ ha <sup>-1</sup>	0.098 40,744.0	0.804 58,663.8	0.46 99,407.8	0.086 25,333.2	0.125 59,423.3	0.107 84,756.5	0.086 25,333.2	0.804 58,663.8	0.481 83,997	0.086 25,333.2	0.051 -20,229.4	0.063 5103.8

#### Table 10

Energy consumption (total energy equivalent (MJ ha<sup>-1</sup>)) and energy input–output relationship in different double cropping systems.

Source	Wheat-Silage corn			Barely-S	eed corn		Barely-S	ilage corn		Barely-Rice		
	W	SC	Total	В	GC	Total	В	SC	Total	В	R	Total
Human labor	966	954	1,920	683	757	1,440	683	955	1,638	683	1,719	2,402
Machinery	3,260	2,698	5,959	2,966	2,069	5,035	2,966	2,698	5,664	2,966	3,643	6,609
Diesel fuel	29,101	31,799	60,900	24,458	32,206	56,663	24,458	31,799	56,257	24,458	61,025	85,483
Fertilizers	15,761	18,924	34,685	16,611	15,800	32,411	16,611	18,924	35,535	16,611	11,252	27,863
Chemicals	1,482	2,045	3,527	1,484	2,045	3,529	1,484	2,045	3,529	1,484	1,535	3,019
Manure	1,920	2,473	4,393	780	2,438	3,218	780	2,474	3,254	780	1,237	2,017
Electricity	2,880	4,428	7,308	2,520	4,788	7,308	2,520	4,428	6,948	2,520	11,880	14,400
Water	6,834	7,977	14,811	6,120	9,047	15,167	6,120	7,977	14,097	6,120	17,850	23,970
Seed	5,899	1,054	6,953	3,421	760	4,181	3,421	1,054	4,475	3,421	3,779	7,200
Total energy input	68,104	72,318	140,422	59,043	69,937	128,979	59,043	72,318	131,360	59,043	113,920	172,962
Total energy output	108,848	130,981	239,829	84,376	129,360	213,736	84,376	130,982	215,357	84,376	93,690	178,066
Energy efficiency	1.6	1.81	1.70	1.43	1.85	1.65	1.43	1.81	1.64	1.43	0.82	1.03

(Singh et al., 2002) and Pakistan (Khan et al., 2007), was 2.9, 2.8, 3.2, 2.5 and 3.46, respectively, which is greater than the value obtained in this study. The main reason for this can be the higher consumption of fossil fuels and fertilizer inputs at low cost and low efficiency of the equipment used in this region.

#### 3.3. Energy intensiveness, productivity, specific and net energy

The results showed that acquired amounts of energy intensity had little difference in the various double-cropped systems as is summarized in Table 8. Average of energy productivity of wheat, barley, grain corn, silage corn, and rice was 0.098 kg  $MJ^{-1}$ , 0.086 kg  $MJ^{-1}$ , 0.125 kg  $MJ^{-1}$ , 0.804 kg  $MJ^{-1}$  and 0.052 kg  $MJ^{-1}$ , respectively. However, the highest energy productivity was observed in the B–SC (0.481 kg  $MJ^{-1}$ ) and W–SC (0.462 kg  $MJ^{-1}$ ) double cropping systems and the lowest was achieved in B–R (0.036 kg  $MJ^{-1}$ ) (Table 8).

The barely–grain corn double-cropped system had the highest specific energy followed by barely–silage corn, barely–rice and winter wheat–silage corn. Net energy was 99,407 MJ ha<sup>-1</sup>, 84,756 MJ ha<sup>-1</sup>, 83,997 MJ ha<sup>-1</sup> and 5103 MJ ha<sup>-1</sup>in wheat–silage corn, barely–grain corn, barely–silage corn and barely–rice, respectively. The highest net energy was obtained in grain corn by 59,423 MJ ha<sup>-1</sup>, whereas the lowest (-20,229 MJ ha<sup>-1</sup>) was related to rice (Table 8). It seems that high level of electricity used was due to electric pumps are old and also high consumption of chemicals and fertilizers could be due to pest invasion and lake of soil analysis which leading to unconscious usage of chemicals. On the other hand, machinery is extensively used for soil preparation, spraying activities and transportation in production process leading to high level of require diesel fuel energy (Rafiee et al., 2010; Singh et al., 2004).

#### 3.4. Energetic of producing cereals and double-cropped systems

The total energy input consumed in wheat, barley, grain corn, silage corn and rice could be classified as direct, indirect, renewable and non-renewable energy. The barely-rice double-cropped systems had the highest direct energy (126.254 MI) followed by wheat-silage corn (84,939 MJ), barely-grain corn (80,578 MJ) and barely-silage corn (78,939 MJ). The highest (55,482 MJ) indirect energy was related to wheat-silage corn and the lowest (46,708 MJ) was obtained in barely-rice double-cropped systems. Amounts of renewable and non-renewable energy in the double-cropped systems are illustrated in Table 10. The highest records for renewable and non-renewable energy were in the barely-rice. The share of indirect and non-renewable energy input was higher than direct and renewable energy in all crops and double-cropped of cereals. Therefore, it is necessary to increase the share of renewable energy for achieving high energy efficiency and energy productivity in agroecosystems. Saving in diesel fuel by changing tillage method can enhance energy use efficiency. Due to the highly mechanized agricultural system in Iran, fuel consumption has risen by 10% in recent years (BeheshtiTabar et al., 2010). Ghorbani et al. (2011) reported that the share of non-renewable energy (76%) compared to renewable energy (24%) was higher in irrigated and dry-land wheat production systems in Iran.

Change of agricultural systems towards using low inputs of fossil energy would contribute to reduce  $CO_2$  and  $N_2O$  emissions (Shahan et al., 2008; Zahid et al., 2010). Our findings indicated that fertilizers especially nitrogen were one of the main input energy that caused to emission of  $N_2O$ , so, should be applied by alternative resource such as residual crops, legumes, and manure (McLaughlin et al., 2000). Traditionally, legumes have been viewed as excellent sources of nitrogen in agriculture (Kinzig and Socolow, 1994). Crop rotations with legumes, capable for fixing atmospheric nitrogen, can maintain production levels with reduced reliance on energy intensive mineral fertilizers (Shahan et al., 2008).

#### 4. Conclusions

Based on the present study the following conclusions are drawn:

1. The total energy input of 140,422 MJ  $ha^{-1}$  (W: 68,104 MJ  $ha^{-1}$  + SC: 72,318 MJ  $ha^{-1}$ ), 128,979 MJ  $ha^{-1}$  (B: 59,043 MJ  $ha^{-1}$  + GC: 69,937 MJ  $ha^{-1}$  BSC), 121,360 MJ  $ha^{-1}$  (B: 59,043 MJ

 $ha^{-1}$  + SC: 72,318 MJ  $ha^{-1}$ ) and 172,962 MJ  $ha^{-1}$  (B: 59,043 MJ  $ha^{-1}$  + R: 113,920 MJ  $ha^{-1}$ ) were required for different double cropping systems, respectively.

- The share of diesel fuel by 43.36% (W–SC), 43.93% (B–GC), 42.82% (B–SC) and 49.40% (B–R) was the highest input. This was followed by fertilizer (W–SC: 24.70%, B–GC: 25.12%, B–SC: 27.05 and B–R: 16.11) and water (W–SC: 10.54%, B–GC: 11.76%, B–SC: 10.73 and B–R: 13.85), respectively.
- 3. The energy use efficiency was found as 1.70 for W–SC, 1.65 for B–GC, 1.64 for B–SC and 1.03 for B–R double cropping systems, respectively.
- 4. The energy productivity were found as 0.46 for W–SC, 0.11 for B–GC, 0.48 for B–SC and 0.063 for B–R double cropping systems, respectively.

#### Acknowledgment

I would like to thank Iran National Science Foundation (INSF), whose funding has made this research possible.

#### References

- Alam, M.S., Alam, M.R., Islam, K.K., 2005. Energy flow in agriculture: Bangladesh. Am. J. Environ. Sci. 1, 213–220.
- Asgharipour, M.R., Mondani, F., Riahinia, S., 2012. Energy use efficiency and economic analysis of sugar beet production system in Iran: A case study in Khorasan Razavi province. Energy 44, 1078–1084.
- Barber, A., 2004. Seven case study farms: Total energy and carbon indicators for New Zealand arable and outdoor vegetable production. In: Arable and Outdoor Vegetable Energy and Carbon Indicators. AgriLINK New Zealand Ltd.
- BeheshtiTabar, I., Keyhani, A., Rafiee, S., 2010. Energy balance in Iran's agronomy (1990–2006). Renew. Sustain. Energy Rev. 14, 849–855.
- Browning, P.W., 2011. Agronomic and economic comparison of full-season and double-cropped small grain and soybean systems in the mid-Atlantic USA (M.S. thesis), In: Crop and Soil Environmental Sciences. Faculty of the Virginia Polytechnic Institute and State University.
- Canakci, M., Topakci, M., Akini, I., Ozmerzi, A., 2005. Energy use pattern of some field crops and vegetable production: Case study for Antalya region, Turkey. Energy. Con. Manage. 46, 656–666.
- Demircan, V., Ekinci, K., Keener, H.M., Akbolat, D., Ekinci, C., 2006. Energy and economic analysis of sweet cherry production in Turkey: a case study from Isparta province. Energy Convers. Manage. 47, 1761–1769.
- Erdal, G., Esengun, K., Erdal, H., Gunduz, O., 2007. Energy use and economical analysis of sugar beet production in Tokat province of Turkey. Energy 32, 35–41.
- Esengun, K., Erdal, G., Gunduz, O., Erdal, H., 2007. An economic analysis and energy use in stake-tomato production in Tokat province of Turkey. Renew. Energy 32, 1873–1881.
- Ghasemi-Mobtaker, H., Keyhani, A., Mohammadi, A., Rafiee, S., Akram, A., 2010. Sensitivity analysis of energy inputs for barley production in Hamedan Province of Iran. Agric. Eco. Environ. 137, 367–372.
- Ghorbani, R., Mondani, F., Amirmoradi, S., Feizi, H., Khorramdel, S., Teimouri, M., Sanjani, S., Anvarkhah, S., Aghel, H., 2011. A case study of energy use and economical analysis of irrigated and dry land wheat production systems. Appl. Energy 88, 283–288.

- Haj-SeyedHadi, M., Darzi, M., Sharifi-Ashoorabadi, E., 2009. Study the effects of conventional and low input production system on energy efficiency of *Silvbumm arianum* L. World. Academy. Sci. Engin. Technol. 54, 364–366.
- Khan, S., Khan, M.A., Hanjra, M.A., Mu, J., 2009. Pathways to reduce the environmental footprints of water and energy inputs in food production. Food Policy. 34, 141–149.
- Khan, M.A., Khan, S., Mushtaq, S., 2007. Energy and economic efficiency of wheat production using different irrigation supply methods. Soil Environ. 26, 121–129.
- Kinzig, A.P., Socolow, R.H., 1994. Human impacts on the nitrogen cycle. Phys. Today. 47, 24–31.
- Koocheki, Alireza, Ghorbani, Reza, Mondani, Farzad, Moradi, Rooholla, Alizade, Yaser, 2011. Pulses production systems in term of energy use efficiency and economical analysis in Iran. Inter. J. Energy Eco. and Poli. 1, 95–106.
- McLaughlin, N.B., Hiba, A., Wall, G.J., King, D.J., 2000. Comparison of energy inputs for inorganic fertilizer and manure based corn production. Can. J. Agric. Eng. 42, 2.1–2.14.
- Ministry of Agriculture of Iran (MAJ), 2011 Portal of Iranian Agriculture, http://www.maj.ir/english/Main/Default.asp.
- Newbold, P., 1994. Statistics for Business And Economics. Englewood Cliffs, Prentice-Hall.
- Rafiee, S., Mousavi-avval, S., Mohammadi, A., 2010. Modeling and sensitivity analysis of energy inputs for apple production in Iran. Energy 35, 3301–3306.
- Rathke, G.W., Diepenbrock, W., 2006. Energy balance of winter oilseed rape (*Brassica napus* L.) cropping as related to nitrogen supply and preceding crop. Europ. J. Agron. 24, 35–44.
- Rathke, G.W., Wienhold, B.J., Wilhelm, W.W., Diepenbrock, W., 2007. Tillage and rotation effect on corn-soybean energy balances in eastern Nebraska. Soil. Till. Res. 97, 60–70.
- Rezvani Moghaddam, P., Feizi, H., Mondani, F., 2011. Evaluation of tomato production systems in terms of energy use efficiency and economical analysis in Iran. Not. Sci. Biol. 3, 58–65.
- Safa, M., Mohtasebi, S.S., Behroozi Lar, M., Ghasemi-Varnamkhasti, M., 2010. Energy consumption in production of grains prevalent in Saveh. Iran. African. J. Agric. Res. 5, 2637–2646.
- Sahabi, H., Feizi, H., Amirmoradi, S., 2012. Which crop production system is more efficient in energy use: wheat or barley? Environ. Dev. Sustain. 12, 9402–9404.
- Shahan, S., Jafari, A., Mobli, H., Rafiee, S., Karimi, M., 2008. Energy use and economical analysis of wheat production in Iran: A case study from Ardabil province. J. Agric Technol. 4, 77–88.
- Sheikh-Davoodi, M.J., Houshyar, E., 2009. Energy consumption of canola and sunflower production in Iran. American-Eurasian. J. Agric. Environ. Sci. 6, 381–384.
- Singh, H., Mishra, D., Nahar, N.M., 2002. Energy use pattern in production agriculture of a typical village in arid zone. India-part I. Energy. Convers. Manage. 43, 2275–2286.
- Singh, G., Singh, S., Singh, J., 2004. Optimization of energy inputs for wheat crop in Punjab. Energy. Convers. Manag. 45, 453–465.
- Tsatsarelis, C.A., 1991. Energy requirements for cotton production in central Greece. J. Agric. Eng. Res. 50, 239–246.
- Yousefi, M., Khoramivafa, M., Mondani, F., 2014. Integrated evaluation of energy use, greenhouse gas emissions and global warming potential for sugar beet (*Beta vulgaris*) agroecosystems in Iran. Atmos. Environ. 92, 501–505.
- Zahedi, M., Eshghizadeh, H.R., Mondani, F., 2014. Energy use efficiency and economical analysis in cotton production system in an Arid region: A case study for Isfahan Province. Iran. Inter. J. Energy Eco. and Poli. 4, 43–52.
- Zahid, H., Azam Khan, M., Irfan, M., 2010. Water energy and economic analysis of wheat production under raised bed and conventional irrigation systems: A case study from a semi-arid area of Pakistan. Soil. Tillage. Res. 109, 61–67.