

Iqbal, Zafar; Aleem, M.

## Article

# Estimation of parameters of the beta-extreme value distribution

Pakistan Journal of Commerce and Social Sciences (PJCSS)

## Provided in Cooperation with:

Johar Education Society, Pakistan (JESPK)

*Suggested Citation:* Iqbal, Zafar; Aleem, M. (2008) : Estimation of parameters of the beta-extreme value distribution, Pakistan Journal of Commerce and Social Sciences (PJCSS), ISSN 2309-8619, Johar Education Society, Pakistan (JESPK), Lahore, Vol. 1, pp. 105-117

This Version is available at:

<https://hdl.handle.net/10419/187984>

## Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

## Terms of use:

*Documents in EconStor may be saved and copied for your personal and scholarly purposes.*

*You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.*

*If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.*



<https://creativecommons.org/licenses/by-nc/4.0/>

# Estimation of Parameters of the Beta-Extreme Value Distribution

Zafar Iqbal

Lecturer, Department of statistics The Islamia University of Bahawalpur. Pakistan

E-mail: zafariqbal101@hotmail.com

Dr. M Aleem (Corresponding author)

Chairman, Department of statistics The Islamia University of Bahawalpur .Pakistan

E-mail: draleemiub@hotmail.com

## Abstract

In this research paper The Beta Extreme Value Type (III) distribution which is developed by Zafar and Aleem (2007) is considered and parameters are estimated by using moments of the Beta-Extreme Value (Type III) Distribution when the parameters 'm' & 'n' are real and moments of the Beta-Extreme Value (Type III) Distribution when the parameters 'm' & 'n' are integers and then a Comparison between  $r^{\text{th}}$  moments about origin when parameters are 'm' & 'n' are real and when parameters are 'm' & 'n' are integers. At the end second method, method of Maximum Likelihood is used to estimate the unknown parameters of the Beta Extreme Value Type (III) distribution.

## 1. Introduction

### 1.1 Beta-Extreme Value (Type III) Distribution:

The Beta-Extreme Value (Type III) Distribution Zafar and Aleem (2007), its generalized class is by defined as

$$f(x; m, n, a, b, c) = \frac{1}{\beta(m, n)} \frac{a}{b-c} \left( \frac{x-c}{b-c} \right)^{a-1} \left[ 1 - e^{-\left( \frac{x-c}{b-c} \right)^a} \right]^{m-1} \left[ e^{-\left( \frac{x-c}{b-c} \right)^a} \right]^n \quad (1.1)$$

Where  $c \leq x \leq \infty, a, b, c, m, n \geq 0$

The Beta-Extreme value type (III) distribution is one of the reliability models. This distribution is useful for analyzing reliability and survival patterns. This model is also useful in extreme situation such as hot weather, earthquakes and flooding.

### 2. Moments of the Beta-Extreme Value (Type III) Distribution when 'm' & 'n' real:

By definition the  $r^{\text{th}}$  moment about origin is given by

$$\mu_r = \int_c^{\infty} x^r \cdot f(x) dx \quad (2.1)$$

From (1.1) when 'm' and 'n' have real values.

$$f(x; m, n, a, b, c) = \frac{\Gamma m}{\beta(m, n)} \frac{a}{(b-c)^a} \sum_{i=0}^{m-1} (x-c)^{a-1} \frac{(-1)^i e^{-i\left(\frac{x-c}{b-c}\right)^a}}{i! \Gamma(m-i)} \left[ e^{-\left(\frac{x-c}{b-c}\right)^a} \right]^n \quad (2.2)$$

Using (2.2) in (2.1)

$$\mu_r = \frac{a}{\beta(m, n)} \frac{1}{(b-c)^a} \sum_{i=0}^{m-1} (-1)^i \frac{\Gamma m}{i! \Gamma(m-i)} \int_c^{\infty} x^r (x-c)^{a-1} e^{-(n+i)\left(\frac{x-c}{b-c}\right)^a} dx$$

$$y = \left( \frac{x-c}{b-c} \right)^a$$

When  $x=c$ , then  $y=0$ , When  $x=\infty$ , then  $y=\infty$

$$y.(b-c)^a = (x-c)^a \quad (x-c) = (b-c)y^{\frac{1}{a}} \quad dx = (b-c)\frac{1}{a}y^{\frac{1}{a}-1}dy$$

$$\mu_r' = \frac{a}{\beta(m,n)} \frac{1}{(b-c)^a} \sum_{i=0}^{m-1} (-1)^i \frac{\Gamma m}{\Gamma(m-i)} \frac{1}{i!} \int_0^{\infty} \left( c + (b-c)y^{\frac{1}{a}} \right)^r \left( (b-c)y^{\frac{1}{a}} \right)^{a-1} \left( e^{-(n+i)y} \right) (b-c) \frac{1}{a} y^{\frac{1}{a}-1} dy$$

$$\mu_r' = \frac{1}{\beta(m,n)} \frac{1}{(b-c)^a} \sum_{i=0}^{m-1} (-1)^i \frac{\Gamma m}{\Gamma(m-i)} \frac{1}{i!} \int_0^{\infty} \left( \sum_{j=0}^r \right)^r \left( (b-c)y^{\frac{1}{a}} \right)^{a-1} \left( e^{-(n+i)y} \right) (b-c) y^{\frac{1}{a}-1} dy$$

$$\mu_r' = \frac{1}{\beta(m,n)} \sum_{i=0}^{m-1} (-1)^i \frac{\Gamma m}{\Gamma(m-i)} \frac{1}{i!} \int_0^{\infty} \left( \sum_{j=0}^r \binom{r}{j} (C)^j (b-c)^{r-j+a-1+1-a} (y)^{\frac{r-j}{a}+1-\frac{1}{a}+\frac{1}{a}-1} \left( e^{-(n+i)y} \right) dy \quad (2.3)$$

We know that

$$(y)^{\frac{r-j}{a}} \left( e^{-(n+i)y} \right) dy = \Gamma \left( \frac{1-j}{a} + 1 \right)$$

So (2.3) becomes

$$\mu_r' = D \sum_{j=0}^r \binom{r}{j} (C)^j (b-c)^{r-j} \frac{\Gamma \left( \frac{1-j}{a} + 1 \right)}{(n+i)^{\frac{r-j}{a}+1}} \quad (2.4)$$

$$\text{Where } D = \frac{1}{\beta(m,n)} \sum_{i=0}^{m-1} (-1)^i \frac{\Gamma m}{i! \Gamma(m-i)}$$

(2.4) gives the  $r_{th}$  moment of the Beta-Extreme Value (Type III) Distribution.

From (2.4) we obtained the first four moments about origin of Beta-Extreme value (type III) distribution.

$$\text{When } r=1 \quad \mu_1' = D \sum_{j=0}^1 \binom{1}{j} (C)^j (b-c)^{1-j} \frac{\Gamma \left( \frac{1-j}{a} + 1 \right)}{(n+i)^{\frac{1-j}{a}+1}}$$

$$\text{When } r=2 \quad \mu_2' = D \sum_{j=0}^2 \binom{2}{j} (C)^j (b-c)^{2-j} \frac{\Gamma \left( \frac{2-j}{a} + 1 \right)}{(n+i)^{\frac{2-j}{a}+1}}$$

$$\text{When } r=3 \quad \mu_3' = D \sum_{j=0}^3 \binom{3}{j} (C)^j (b-c)^{3-j} \frac{\Gamma \left( \frac{3-j}{a} + 1 \right)}{(n+i)^{\frac{3-j}{a}+1}}$$

$$\text{When } r=4 \quad \mu_4' = D \sum_{j=0}^4 \binom{4}{j} (C)^j (b-c)^{4-j} \frac{\Gamma\left(\frac{4-j}{a} + 1\right)}{(n+i)^{\frac{4-j}{a}+1}}$$

$$\begin{aligned} \mu_2 &= \mu_2' - (\mu_1')^2 & \mu_3 &= \mu_3' - 3\mu_2' \mu_1' + 2(\mu_2')^3 \\ \mu_4 &= \mu_4' - 3\mu_3' \mu_1' + 6\mu_2' (\mu_1')^2 - 3(\mu_1')^4 \end{aligned}$$

### 3. Moments of the Beta-Extreme Value (Type III) Distribution when 'm' & 'n' integers:

By definition the rth moment about origin is given by

$$\mu_r' = \int_0^{\infty} x^r \cdot f(x) dx \quad (3.1)$$

By using (1.1) in (3.1), we get

$$\mu_r' = \frac{a}{\beta(m,n)} \frac{1}{(b-c)^a} \sum_{j=0}^r (-1)^j C_{(m,i)} \frac{1}{j!} \int_c^{\infty} x^r (x-c)^{a-1} e^{-\left(\frac{x-c}{b-c}\right)^a} dx$$

By simplifying we get

$$\mu_r' = D \sum_{j=0}^r \binom{r}{j} (C)^j (b-c)^{r-j} \frac{\frac{r-j}{a} + 1}{(n+i)^{\frac{r-j}{a}+1}} \quad (3.2)$$

$$\text{Where } D = \frac{1}{\beta(m,n)} \sum (-1)^i C_{(m,i)}$$

(3.2) gives the  $r_{th}$  moment of the Beta-Extreme Value (Type III) Distribution, when 'm' and 'n' are integers.

From (3.2) we obtained the first four moments about origin of Beta-Extreme value (type III) distribution.

$$\text{When } r=1 \quad \mu_1' = D \sum_{j=0}^1 \binom{1}{j} (C)^j (b-c)^{1-j} \frac{\frac{1-j}{a} + 1}{(n+i)^{\frac{1-j}{a}+1}}$$

$$\text{When } r=2 \quad \mu_2' = D \sum_{j=0}^2 \binom{2}{j} (C)^j (b-c)^{2-j} \frac{\frac{2-j}{a} + 1}{(n+i)^{\frac{2-j}{a}+1}}$$

$$\text{When } r=3 \quad \mu_3' = D \sum_{j=0}^3 \binom{3}{j} (C)^j (b-c)^{3-j} \frac{\frac{3-j}{a} + 1}{(n+i)^{\frac{3-j}{a}+1}}$$

$$\text{When } r=4 \quad \mu_4' = D \sum_{j=0}^4 \binom{4}{j} (C)^j (b-c)^{4-j} \frac{\frac{4-j}{a} + 1}{(n+i)^{\frac{4-j}{a}+1}}$$

$$\mu_2 = \mu_2' - (\mu_1')^2$$

$$\mu_3 = \mu_3' - 3\mu_2' \mu_1' + 2(\mu_2')^3$$

$$\mu_4 = \mu_4' - 3\mu_3' \mu_1' + 6\mu_2' (\mu_1')^2 - 3(\mu_1')^4$$

**4. Comparison between  $r_{th}$  moments about origin when parameters 'm' & 'n' are real and parameters 'm' & 'n' are integers.**

Case1: When 'c'=0

| $R_{th}$ moment for real values                                                                                                                     | $R_{th}$ moment for integers values                                                                                                   |
|-----------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------|
| $\mu_1' = \frac{(b)(\Gamma m)\Gamma\left(\frac{1+a}{a}\right)}{\beta(m,n)} \sum_{i=0}^{m-1} (-1)^i \frac{1}{i!(n+i)^{\frac{1+a}{a}} \Gamma(m-i)}$   | $\mu_1' = \frac{(b)\Gamma\left(\frac{1+a}{a}\right)}{\beta(m,n)} \sum_{i=0}^{m-1} (-1)^i (C_{m,i}) \frac{1}{(n+i)^{\frac{1+a}{a}}}$   |
| $\mu_2' = \frac{(b^2)(\Gamma m)\Gamma\left(\frac{2+a}{a}\right)}{\beta(m,n)} \sum_{i=0}^{m-1} (-1)^i \frac{1}{i!(n+i)^{\frac{2+a}{a}} \Gamma(m-i)}$ | $\mu_2' = \frac{(b^2)\Gamma\left(\frac{2+a}{a}\right)}{\beta(m,n)} \sum_{i=0}^{m-1} (-1)^i (C_{m,i}) \frac{1}{(n+i)^{\frac{2+a}{a}}}$ |
| $\mu_3' = \frac{(b^3)(\Gamma m)\Gamma\left(\frac{3+a}{a}\right)}{\beta(m,n)} \sum_{i=0}^{m-1} (-1)^i \frac{1}{i!(n+i)^{\frac{3+a}{a}} \Gamma(m-i)}$ | $\mu_3' = \frac{(b^3)\Gamma\left(\frac{3+a}{a}\right)}{\beta(m,n)} \sum_{i=0}^{m-1} (-1)^i (C_{m,i}) \frac{1}{(n+i)^{\frac{3+a}{a}}}$ |
| $\mu_4' = \frac{(b^4)(\Gamma m)\Gamma\left(\frac{4+a}{a}\right)}{\beta(m,n)} \sum_{i=0}^{m-1} (-1)^i \frac{1}{i!(n+i)^{\frac{4+a}{a}} \Gamma(m-i)}$ | $\mu_4' = \frac{(b^4)\Gamma\left(\frac{4+a}{a}\right)}{\beta(m,n)} \sum_{i=0}^{m-1} (-1)^i (C_{m,i}) \frac{1}{(n+i)^{\frac{4+a}{a}}}$ |

Case 2: When 'm'=1 and 'c'=0 then for different values of parameters 'a', 'b' and 'n' the moments about origin becomes

| <b>R<sub>th</sub> moment for real values</b>                                  | <b>R<sub>th</sub> moment for integers values</b>                              |
|-------------------------------------------------------------------------------|-------------------------------------------------------------------------------|
| $\mu_1' = \frac{nb\Gamma\left(\frac{1+a}{a}\right)}{(n)^{\frac{1+a}{a}}}$     | $\mu_1' = \frac{nb\Gamma\left(\frac{1+a}{a}\right)}{(n)^{\frac{1+a}{a}}}$     |
| $\mu_2' = \frac{n(b^2)\Gamma\left(\frac{2+a}{a}\right)}{(n)^{\frac{1+a}{a}}}$ | $\mu_2' = \frac{n(b^2)\Gamma\left(\frac{2+a}{a}\right)}{(n)^{\frac{1+a}{a}}}$ |
| $\mu_3' = \frac{n(b^3)\Gamma\left(\frac{3+a}{a}\right)}{(n)^{\frac{1+a}{a}}}$ | $\mu_3' = \frac{n(b^3)\Gamma\left(\frac{3+a}{a}\right)}{(n)^{\frac{1+a}{a}}}$ |
| $\mu_4' = \frac{n(b^4)\Gamma\left(\frac{4+a}{a}\right)}{(n)^{\frac{1+a}{a}}}$ | $\mu_4' = \frac{n(b^4)\Gamma\left(\frac{4+a}{a}\right)}{(n)^{\frac{1+a}{a}}}$ |

Case 3: When 'n'=1 & 'c'=0 then for different values of parameters 'a', 'b' and 'm' the moments about origin becomes

| <b>R<sub>th</sub> moment for real values</b>                                                                                      | <b>R<sub>th</sub> moment for integers values</b>                                                                                      |
|-----------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------|
| $\mu_1' = (b)\Gamma(m+1)\Gamma\left(\frac{1+a}{a}\right)\sum_{i=0}^{m-1} (-1)^i \frac{1}{i!(1+i)^{\frac{1+a}{a}} \Gamma(m-i)}$    | $\mu_1' = \frac{(b)\Gamma\left(\frac{1+a}{a}\right)}{\beta(m,1)} \sum_{i=0}^{m-1} (-1)^i (C_{m,i}) \frac{1}{(n+i)^{\frac{1+a}{a}}}$   |
| $\mu_2' = (b^2)(\Gamma(m+1)\Gamma\left(\frac{2+a}{a}\right)\sum_{i=0}^{m-1} (-1)^i \frac{1}{i!(1+i)^{\frac{2+a}{a}} \Gamma(m-i)}$ | $\mu_2' = \frac{(b^2)\Gamma\left(\frac{2+a}{a}\right)}{\beta(m,1)} \sum_{i=0}^{m-1} (-1)^i (C_{m,i}) \frac{1}{(n+i)^{\frac{2+a}{a}}}$ |
| $\mu_3' = (b^3)\Gamma(m+1)\Gamma\left(\frac{3+a}{a}\right)\sum_{i=0}^{m-1} (-1)^i \frac{1}{i!(1+i)^{\frac{3+a}{a}} \Gamma(m-i)}$  | $\mu_3' = \frac{(b^3)\Gamma\left(\frac{3+a}{a}\right)}{\beta(m,1)} \sum_{i=0}^{m-1} (-1)^i (C_{m,i}) \frac{1}{(n+i)^{\frac{3+a}{a}}}$ |

|                                                                                                                                 |                                                                                                                                      |
|---------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------|
| $\mu_4' = (b^4)\Gamma(m+1)\Gamma\left(\frac{4+a}{a}\right)\sum_{i=0}^{m-1} (-1)^i \frac{1}{i!(1+i)^{\frac{4+a}{a}}\Gamma(m-i)}$ | $\mu_4' = \frac{(b^4)\Gamma\left(\frac{4+a}{a}\right)}{\beta(m,1)}\sum_{i=0}^{m-1} (-1)^i (C_{m,i}) \frac{1}{(n+i)^{\frac{4+a}{a}}}$ |
|---------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------|

Case 4: When  $a=1$  &  $c=0$  then for different values of parameters 'b', 'n' and 'm' the moments about origin becomes

| <b>R<sub>th</sub> moment for real values</b>                                                          | <b>R<sub>th</sub> moment for integers values</b>                                         |
|-------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------|
| $\mu_1' = \frac{(b)(\Gamma m)}{\beta(m,n)}\sum_{i=0}^{m-1} (-1)^i \frac{1}{i!(n+i)^2\Gamma(m-i)}$     | $\mu_1' = \frac{(b)}{\beta(m,n)}\sum_{i=0}^{m-1} (-1)^i (C_{m,i}) \frac{1}{(n+i)^2}$     |
| $\mu_2' = \frac{2(b^2)(\Gamma m)}{\beta(m,n)}\sum_{i=0}^{m-1} (-1)^i \frac{1}{i!(n+i)^3\Gamma(m-i)}$  | $\mu_2' = \frac{2(b^2)}{\beta(m,n)}\sum_{i=0}^{m-1} (-1)^i (C_{m,i}) \frac{1}{(n+i)^3}$  |
| $\mu_3' = \frac{6(b^3)(\Gamma m)}{\beta(m,n)}\sum_{i=0}^{m-1} (-1)^i \frac{1}{i!(n+i)^4\Gamma(m-i)}$  | $\mu_3' = \frac{6(b^3)}{\beta(m,n)}\sum_{i=0}^{m-1} (-1)^i (C_{m,i}) \frac{1}{(n+i)^4}$  |
| $\mu_4' = \frac{24(b^4)(\Gamma m)}{\beta(m,n)}\sum_{i=0}^{m-1} (-1)^i \frac{1}{i!(n+i)^5\Gamma(m-i)}$ | $\mu_4' = \frac{24(b^4)}{\beta(m,n)}\sum_{i=0}^{m-1} (-1)^i (C_{m,i}) \frac{1}{(n+i)^5}$ |

Case 5: When  $b=1$  &  $c=0$  then for different values of parameters 'a', 'n' and 'm' the moments about origin becomes

| <b>R<sub>th</sub> moment for real values</b>                                                                                                 | <b>R<sub>th</sub> moment for integers values</b>                                                                                |
|----------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------|
| $\mu_1' = \frac{(\Gamma m)\Gamma\left(\frac{1+a}{a}\right)}{\beta(m,n)}\sum_{i=0}^{m-1} (-1)^i \frac{1}{i!(n+i)^{\frac{1+a}{a}}\Gamma(m-i)}$ | $\mu_1' = \frac{\Gamma\left(\frac{1+a}{a}\right)}{\beta(m,n)}\sum_{i=0}^{m-1} (-1)^i (C_{m,i}) \frac{1}{(n+i)^{\frac{1+a}{a}}}$ |
| $\mu_2' = \frac{(\Gamma m)\Gamma\left(\frac{2+a}{a}\right)}{\beta(m,n)}\sum_{i=0}^{m-1} (-1)^i \frac{1}{i!(n+i)^{\frac{2+a}{a}}\Gamma(m-i)}$ | $\mu_2' = \frac{\Gamma\left(\frac{2+a}{a}\right)}{\beta(m,n)}\sum_{i=0}^{m-1} (-1)^i (C_{m,i}) \frac{1}{(n+i)^{\frac{2+a}{a}}}$ |
| $\mu_3' = \frac{(\Gamma m)\Gamma\left(\frac{3+a}{a}\right)}{\beta(m,n)}\sum_{i=0}^{m-1} (-1)^i \frac{1}{i!(n+i)^{\frac{3+a}{a}}\Gamma(m-i)}$ | $\mu_3' = \frac{\Gamma\left(\frac{3+a}{a}\right)}{\beta(m,n)}\sum_{i=0}^{m-1} (-1)^i (C_{m,i}) \frac{1}{(n+i)^{\frac{3+a}{a}}}$ |

|                                                                                                                                                  |                                                                                                                                   |
|--------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------|
| $\mu_4' = \frac{(\Gamma m) \Gamma\left(\frac{4+a}{a}\right)}{\beta(m, n)} \sum_{i=0}^{m-1} (-1)^i \frac{1}{i!(n+i)^{\frac{4+a}{a}} \Gamma(m-i)}$ | $\mu_4' = \frac{\Gamma\left(\frac{4+a}{a}\right)}{\beta(m, n)} \sum_{i=0}^{m-1} (-1)^i (C_{m,i}) \frac{1}{(n+i)^{\frac{4+a}{a}}}$ |
|--------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------|

### 5. Method of Maximum Likelihood:

We used method of Maximum Likelihood for the estimation of parameters of Beta-Extreme Value (Type III) Distribution.

We know that

$$f(x) = \frac{1}{\beta(m, n)} \frac{a}{b-c} \left[ \frac{x-c}{b-c} \right]^{a-1} \left[ 1 - e^{-\left(\frac{x-c}{b-c}\right)^a} \right]^{m-1} \left[ e^{-\left(\frac{x-c}{b-c}\right)^a} \right]^n$$

The maximum likelihood function of the above equation is.

$$L(x; m, n, a, b, c) = \prod_{i=1}^n \frac{1}{\beta(m, n)} \frac{a}{b-c} \left[ \frac{x-c}{b-c} \right]^{a-1} \left[ 1 - e^{-\left(\frac{x-c}{b-c}\right)^a} \right]^{m-1} \left[ e^{-\left(\frac{x-c}{b-c}\right)^a} \right]^n$$

$$L(x; m, n, a, b, c) = \left[ \frac{1}{\beta(m, n)} \right]^n \left( \frac{a}{b-c} \right)^n \prod_{i=1}^n \left[ \frac{x-c}{b-c} \right]^{a-1} \prod_{i=1}^n \left[ 1 - e^{-\left(\frac{x-c}{b-c}\right)^a} \right]^{m-1} \prod_{i=1}^n \left[ e^{-\left(\frac{x-c}{b-c}\right)^a} \right]^n$$

By taking log on both sides

$$\log L(x; m, n, a, b, c) = n \left[ \log(1) - \log \beta(m, n) \right] + n \left[ \log a - \log(b-c) \right]$$

$$+ (a-1) \sum_{i=1}^n \log \left( \frac{x-c}{b-c} \right) + (m-1) \sum_{i=1}^n \log \left[ 1 - e^{-\left(\frac{x-c}{b-c}\right)^a} \right] - n \sum_{i=1}^n \left( \frac{x-c}{b-c} \right)^a$$

$$\log L(x; m, n, a, b, c) = -n \log \beta(m, n) + n \log a - n \log(b-c) + (a-1) \sum_{i=1}^n \log \left( \frac{x-c}{b-c} \right)$$

$$+ (m-1) \sum_{i=1}^n \log \left( 1 - e^{-\left(\frac{x-c}{b-c}\right)^a} \right) - n \sum_{i=1}^n \left( \frac{x-c}{b-c} \right)^a$$

$$\frac{\partial \log L}{\partial m} = -n \frac{\partial \log \beta(m, n)}{\partial m} + \sum_{i=1}^n \log \left[ 1 - e^{-\left(\frac{x-c}{b-c}\right)^a} \right]$$

$$= -n \frac{\partial \log \beta(m, n)}{\partial m} + \sum_{i=1}^n \log \left[ 1 - e^{-\left(\frac{x-c}{b-c}\right)^a} \right]$$

$$\frac{\partial \log L}{\partial m} = -n \psi(m, n) + \sum_{i=1}^n \left[ 1 - e^{-\left(\frac{x-c}{b-c}\right)^a} \right] \quad (5.1)$$

$$\frac{\partial \log L}{\partial n} = -n \frac{\partial \beta(m, n)}{\partial n} - \log \beta(m, n) + \log a - \log(b-c) - \sum_{i=1}^n \left( \frac{x-c}{b-c} \right)^a$$

$$\frac{\partial \log L}{\partial n} = -n \Psi(m, n) - \log \beta(m, n) + \log a - \log(b-c) - \sum_{i=1}^n \left( \frac{x-c}{b-c} \right)^a \quad (5.2)$$

$$\frac{\partial \log L}{\partial a} = \frac{n}{b} + \sum_{i=1}^n \log \left( \frac{x-c}{b-c} \right) + (m-1) \sum_{i=1}^n \frac{1}{\left[ 1 - e^{-\left(\frac{x-c}{b-c}\right)^a} \right]} \frac{\partial}{\partial a} \left[ 1 - e^{-\left(\frac{x-c}{b-c}\right)^a} \right]$$

$$- n \sum_{i=1}^n \left( \frac{x-c}{b-c} \right)^a \log \left( \frac{x-c}{b-c} \right) \quad (5.3)$$

$$\frac{\partial \log L}{\partial b} = -\frac{n}{b-c} - (a-1) \sum_{i=1}^n \frac{1}{b-c} + (m-1) \sum_{i=1}^n \frac{1}{1 - e^{-\left(\frac{x-c}{b-c}\right)^a}} \frac{\partial}{\partial b} \left[ 1 - e^{-\left(\frac{x-c}{b-c}\right)^a} \right]$$

$$- na \sum_{i=1}^n \left( \frac{x-c}{b-c} \right)^{a-1} \frac{\partial}{\partial b} \left( \frac{x-c}{b-c} \right)$$

$$= \frac{-n}{b-c} - (a-1) \sum_{i=1}^n \frac{1}{b-c} + (m-1) \sum_{i=1}^n \left[ \frac{-e^{-\left(\frac{x-c}{b-c}\right)^a}}{1 - e^{-\left(\frac{x-c}{b-c}\right)^a}} - a \left[ \frac{x-c}{b-c} \right]^{a-1} \frac{\partial}{\partial b} \left( \frac{x-c}{b-c} \right) \right]$$

$$- na \sum_{i=1}^n \left( \frac{x-c}{b-c} \right)^{a-1} \left[ \frac{-(x-c)}{(b-c)^2} \right]$$

$$\begin{aligned}
\frac{\partial \log L}{\partial b} &= \frac{-n}{b-c} - \frac{n(a-1)}{b-c} + a(m-1) \sum_{i=1}^n \frac{e^{-\left(\frac{x-c}{b-c}\right)^a}}{1 - e^{-\left(\frac{x-c}{b-c}\right)^a}} \left[ \frac{x-c}{b-c} \right]^{a-1} \left[ \frac{-(x-c)}{(b-c)^2} \right] \\
&+ na \sum_{i=1}^n \left( \frac{x-c}{b-c} \right)^{a-1} \frac{x-c}{(b-c)^2} \\
&= \frac{-n-na+n}{(b-c)} - a(m-1) \sum_{i=1}^n \frac{e^{-\left(\frac{x-c}{b-c}\right)^a}}{1 - e^{-\left(\frac{x-c}{b-c}\right)^a}} \left[ \frac{x-c}{b-c} \right]^{a-1} \frac{x-c}{(b-c)^2} \\
&+ na \sum_{i=1}^n \left( \frac{x-c}{b-c} \right)^{a-1} \left( \frac{x-c}{(b-c)^2} \right) \\
\frac{\partial \log L}{\partial b} &= \frac{-na}{(b-c)} + a(m-1) \sum_{i=1}^n \frac{e^{-\left(\frac{x-c}{b-c}\right)^a}}{1 - e^{-\left(\frac{x-c}{b-c}\right)^a}} \left[ \frac{x-c}{b-c} \right]^{a-1} \frac{x-c}{(b-c)^2} \\
&+ na \sum_{i=1}^n \left( \frac{x-c}{b-c} \right)^{a-1} \left( \frac{x-c}{(b-c)^2} \right) \tag{5.4}
\end{aligned}$$

$$\begin{aligned}
\frac{\partial \log L}{\partial c} &= + \frac{n}{b-c} + (a-1) \sum_{i=1}^n \frac{1}{\left( \frac{x-c}{b-c} \right)} \frac{\partial}{\partial c} \left( \frac{x-c}{b-c} \right) + (m-1) \sum_{i=1}^n \frac{1}{1 - e^{-\left(\frac{x-c}{b-c}\right)^a}} \frac{\partial}{\partial c} \left[ 1 - e^{-\left(\frac{x-c}{b-c}\right)^a} \right] \\
&= \frac{-n}{b-c} + (a-1)(b-c) \sum_{i=1}^n (x-c)^{-1} \left[ \frac{(b-c)(a-1) + (x-c)}{(b-c)^2} \right] \\
&+ (m-1) \sum_{i=1}^n \frac{1}{1 - e^{-\left(\frac{x-c}{b-c}\right)^a}} \left[ -e^{-\left(\frac{x-c}{b-c}\right)^a} - a \left( \frac{x-c}{b-c} \right)^{a-1} \right] \left[ \frac{-(b-c) + (x-c)}{(b-c)^2} \right] \\
&= \frac{-n}{b-c} + (a-1) \sum_{i=1}^n \frac{-(b-c) + (x-c)}{(b-c)} (x-c)^{-1}
\end{aligned}$$

$$\begin{aligned}
& + (m-1) \sum_{i=1}^n \frac{1}{1 - e^{-\left(\frac{x-c}{b-c}\right)^a}} \left[ -a e^{-\left(\frac{x-c}{b-c}\right)^a} \left(\frac{x-c}{b-c}\right)^{a-1} - \frac{(b-c) + (x-c)}{(b-c)^2} \right] \\
\frac{\partial \log L}{\partial c} &= \frac{-n}{b-c} + \frac{a-1}{b-c} \sum_{i=1}^n \left(\frac{x-b}{x-c}\right) \frac{-a(m-1)}{(b-c)^{a+1}} \sum_{i=1}^n \left[ 1 - e^{-\left(\frac{x-c}{b-c}\right)^a} \right]^{-1} e^{-\left(\frac{x-c}{b-c}\right)^a} \cdot (x-c)^{a-1} (x-b)
\end{aligned}
\tag{5.5}$$

Where  $\Psi(x) = \partial \ln \Gamma x / \partial x$  is the derivative of the gamma function and equating these expressions (5.1), (5.2), (5.3), (5.4), (5.5) equal to zero then solving them simultaneously yields the maximum likelihood estimates of the five parameters Beta-Extreme Value (Type III) Distribution.

Equations (5.1), (5.2), (5.3), (5.4), (5.5) are solved iteratively to obtain  $\hat{m}, \hat{n}, \hat{a}, \hat{b}$  and  $\hat{c}$  the maximum likelihood estimates (MLE) for parameters  $m, n, a, b$  and  $c$  respectively. By taking the second partial derivatives of (5.1), (5.2), (5.3), (5.4), (5.5) and the fisher information matrix can be obtained by taking the expectations of second partial derivatives with minus sign. The Inverse of the fisher information matrix provides the variances and the covariances for the maximum likelihood estimators.

### Bibliography

- Memon, A. G. and Shaikh, N.M. (2003). Fitting Gumbel and Generalized Extreme Value Probability Distributions on Floods data of Indus River at Sukkur Barrage, [unpublished report].
- Costantino, R.F., and Deshamais, R.A. (1981). Gamma Distributions of Adult Numbers for Tribolium Populations in the Region of Their Steady States. *Journal of Animal Ecology*, 50, 667-681.
- Cox, D. R. and Oakes, D. (1984). *Analysis of Survival Data*, Chapman & Hall. London, UK.
- Eugene, L. and Famoye, (2002). Beta-Normal Distribution And Its Application, *Communication in Statistics, -Theory Meth.*, 31(4), 497-512.
- Gupta, A.K. and Nadarajah, S. (2004). On the Moments of the Beta-Normal Distribution. *Communication I Statistics: Theory and Methods*, 33(1), 1-13.
- Gupta, A.K., Nadarajah, S. (2004). *Handbook of Beta Distribution and its Applications*. Marcel Dekker Inc, New York.
- Good, I.J. (1953). The Population Frequencies of the Species and the Estimation of Population Parameters. *Biometrika*. 40, 237-260.
- Iqbal, Z. (2007). Beta-Extreme Value Distribution. M.Phil Thesis [Unpublished], The Islamia University Of Bahawalpur, Pakistan.
- Holmes, J.D. and Moriarty, W.W. (1999). Application of the generalized Pareto distribution to extreme value analysis in wind engineering. *Journal of Wind Engineering and Industrial Aerodynamics*, 83(1-3), 1-10.
- Lee, C., Famoye, P. and Ohmolade, O. (2006). The Beta-Weibull Distribution: Some Properties and Applications to censorsord data.
- Lock, M. O. (1973). *Reliability, Maintainability, and Availability Assessment*. Hayden Book Co. New Jersey, [ISBN: 0-87671-173-5]
- Nadarajah, S., Kotz, S. (2004). *The Beta Gunmbel Distribution: Mathematical Problems in Engineering*, 323-332.
- Nadarajah, S., Kotz, S. (2005). *The Beta Gunmbel Distribution: Reliability Engineering and System Safety*. 689-697.
- Nelson, W. (1982). *Applied Life Data Analysis*. J. Wiley, New York. USA, [ISBN: 0-471-09458-7].
- Eugene, N., Lee, C. and Famoye, F. (2002). Beta-Normal distribution and its applications. *Comm. Statist. Theory Methods*, 4, 497-512.

Sara, L, Nadrajah and Samuel, K. (2004). *The Beta Gumbel Distribution*, *Journal of mathematical problems in Engineering*. 4, 323-332.

Hanook, S. (2006). Three Parameter Beta-Inverse Weibull Distribution. M. Phil Thesis [Unpublished] Department of Statistics, G.C. University Lahore, Pakistan.

Kotz, S. and Nadarajah, S. (2000). *Extreme Value Distributions, Theory and Applications*. Imperial College Press, London.

#### **Web Resources**

<http://www.itl.nist.gov/div898/handbook/apr/section1/apr163.htm>

[http://en.wikipedia.org/wiki/Generalized\\_extreme\\_value\\_distribution](http://en.wikipedia.org/wiki/Generalized_extreme_value_distribution)

[http://en.wikipedia.org/wiki/Beta\\_distribution](http://en.wikipedia.org/wiki/Beta_distribution)