

A Service of

ZBW

Leibniz-Informationszentrum Wirtschaft Leibniz Information Centre for Economics

Pacheco Barzallo, Diana

Working Paper Health Conditions and Social Interactions

IRENE Working Paper, No. 11-03

Provided in Cooperation with: Institute of Economic Research (IRENE), University of Neuchâtel

Suggested Citation: Pacheco Barzallo, Diana (2011) : Health Conditions and Social Interactions, IRENE Working Paper, No. 11-03, University of Neuchâtel, Institute of Economic Research (IRENE), Neuchâtel

This Version is available at: https://hdl.handle.net/10419/191463

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.



WWW.ECONSTOR.EU

University of Neuchatel Institute of Economic Research IRENE, Working paper 11-03

Health Conditions and Social Interaction

Diana Pacheco Barzallo*



Institut de recherches économiques

Health Conditions & Social Interactions

Diana Pacheco Barzallo

Department of Economics: University of Neuchatel

August 4, 2011

Abstract

This paper evaluates the impact of an antipoverty program on the health condition of individuals. The program combines a cash transfer with financial incentives for positive behavior to poor families. Its main purposes are to improve the living conditions of eligible households and to promote their investment in their children through school attendance and the provision of basic health services. Since the design of the program includes information on eligible and ineligible families, it is possible to evaluate its direct effect as well as its indirect effect.

While the direct impact is measured by the effect of cash grants on eligible individuals, its indirect impact is estimated by the effect generated for the treated neighbors on their non-treated peers. The results show that eligible and ineligible individuals significantly improved their health status due to the program's interventions: the sickness incidences decreased, the sickness spell was reduced, and people seemed to be able to manage normal activities with less difficulty.

Thus anti-poverty programs, despite the fact that constitute a big weight in the public finance of a country, have an important multiplicative effect on the population. It not only generates, in the short run, an increase in the demand for health services but also changes in a positive way the behaviour of people by educating them on the importance of health and nutrition. This result will impact treated and non-treated families in the long run due to the high interaction that characterizes poor societies: families are learning from others.

JEL-Classification: C93, I12, I28, I38

Keywords: randomization, experimental design, social interactions.

1 Introduction

Despite improvements over the last few decades, health problems and child nutrition deficiency persist in developing countries. For this reason, many governments have implemented social programs that try to intervene in poor populations through cash grants. These programs can be classified into two groups. In the first group, we have conditional programs that make cash transfers contingent on some behavior, usually investments in human capital such as sending children to school or bringing them to health centers.¹ In the second group, we find unconditional programs that mostly focus on a specific age group; these programs make transfers without any limitations or conditions imposed on eligible individuals.

This paper analyses the impact of PROGRESA on health and nutrition. Serving approximately 25 million people, PROGRESA is Mexico's principal antipoverty initiative. Launched by the government in 1997, the program awards cash grants to families living in poverty. The grants are conditioned upon criteria such as preventive health check-ups and regular school attendance for children. Its main goal is to break the intergenerational transmission of poverty by increasing investment by treated families in the human capital of their children.

Related literature has studied the effect of PROGRESA on school attendance, but few documents have evaluated the impact on health and nutrition. The main findings suggest that antipoverty programs that combine education, health, and nutrition interventions can be quite successful at improving the capacity of families to pull themselves out of poverty (Behrman & Hoddinott, 2000). Health and nutrition are very important factors, not only because they directly increase an individual's welfare but because they improve a child's physical and cognitive development (Haas et al, 1996). In particular, healthier children more regularly attend school, while healthier adults with better cognitive ability are more productive and substantially increase their wages (Gertler, 2000).

Unconditional programs, which have been implemented in countries such as Ecuador and South Africa, have had positive results. In the case of Ecuador, authors have concluded that by the age of five, children who were inadequately nourished have already

¹For more information about countries that have applied this type of program, see: Rawlings L., & Rubio G., Evaluating the Impact of Conditional Cash Transfer Programs, (2005). The World Bank Research Observer, vol. 20, no.1

fallen well behind the cognitive development of their better-nourished peers. Assuming that the disparities persist, the malnourished children will have poorer performance in school, accumulate less human capital, and make less than their peers (Paxson & Schady, 2005). In the case of South Africa, several working papers have shown that unconditional payments have bolstered early childhood nutrition. What is a general consensus among authors is that antipoverty programs that condition payments upon a specific behavior would further increase the program's effects (Kakwani & Veras, 2005).

The paper that is closest to this one is Gertler and Boyce (2001). The authors found that PROGRESA significantly increased the utilization of public health clinics for preventive care. In addition, the program lowered the number of inpatient hospitalizations and visits to private providers. Specifically, there was an important reduction in the incidence of illness, an increase in children's height, and a reduction in anemia. In the case of adults, the program significantly increased the number of kilometers they were able to walk without becoming fatigued and decreased the number of days of difficulty experienced with normal activities due to illness.

In the estimation framework, program evaluation literature use mainly a differencein- difference estimation (Gertler, 2000; Behrman, & Sengupta & Todd, 2000); however, this method is quite complicated because the survey evaluations have changed every year: the variables before and after the program's implementation are not always the same. Other authors who use the data at the individual level apply propensity score estimation methods (Dian & Handa, 2004; Diaz & Sudhanshu, 2006). Finally, some documents work with panel data-fixed effects; however, this method tends to eliminate a large number of observations because the treatment indicator is invariant over time (Gertler, 2000; Behrman & Hoddinott, 2001).

This paper estimates PROGRESA's impact using different health outcomes. Among these, we have binary and count variables. For the first case, we estimate a logit and a probit specification. In the second case, we estimate either a Poisson or a zero-inflated negative binomial model. It is important to stress the fact that the estimation is done at the locality level and in poor and non-poor subgroups. This division, besides making the program a randomly assigned experiment, allows us to estimate the indirect effect of the program on the non-targeted population.

Regarding the data, the sample is composed of household socioeconomic surveys from rural areas in Mexico. An important aspect of this paper, in contrast to related documents, is that it defines the global impact of PROGRESA on health and nutrition by different age groups. By desegregating the impact on groups, it is possible to eliminate the problem of misrepresentation: the results do not correspond to the average individual (Bolduc et al., 1996).

The results suggest that the direct effect of PROGRESA, measured by the impact of the treatment on the targeted population, significantly improved the health status of the population. Specifically, children are less likely to get sick, and if sick, they face shorter periods in this state. Children also increased the frequency of visits to the doctor to control height and weight. Adult individuals show fewer problems dealing with daily/moderate activities, and can walk more kilometers before becoming tired. Finally, older treated individuals exhibit greater probabilities of staying healthy.

In the same way, results of the effect of PROGRESA on the non-targeted population reveal a positive impact as well. Ineligible children show a significant increase in their frequency of visits to medical checkups. Not only children, but also young and old individuals exhibit shorter spells of sickness. Adult individuals improved their capabilities to deal with normal and difficult activities. In spite of this, the results also suggest that ineligible adults have more trouble when bathing and dressing. Finally, the spillover effect, measured by the effect of treated neighbors on the non-treated, show that ineligible children aged between 0 and 12 years, are less likely to get sick and have shorter spells of sickness. They also increased their visits to medical checkups.

The document is organized as follows. The next section gives a general view of the health situation in Mexico. It also describes background information on the PRO-GRESA program: the main goals, structure, and design. In the third part, the document presents data information and some descriptive statistics that explain how the sample is organized. The fourth section defines the econometric analysis. In the final section, we show the estimation results and offer a brief summary and a conclusion.

2 Background

This section describes health in Mexico. In addition, it provides background information about the PROGRESA: the main goals, design and coverage.

2.1 Health Access in Mexico

The health care system places Mexico above Latin American standards but far below OECD standards. In Figure 1, we can see that, for the year 2005, the infant mortality rate in Mexico was 22 per 1000 live births, where as, in the OECD, the rate is only 5 per 1000 live births. The average life expectancy in Mexico was 75 in 2005, while it was 80 in OECD countries. (World Bank, 2005).



Figure 1: Health Outcomes Comparison

Access to health services varies widely in the country. Some workers have access to superior health care services with significant pensions, while most people who work in the informal sector access either public providers or affordable private health care. In general, people employed in the public sector are covered by the national ISSSTE social security institute, which, on average, brings better-quality health care and more generous pension benefits. Finally, people who belong to unions, because of their bargaining power, can generally access better benefit packages and services (World Bank Report: Mexico 2006-2012, 2007, p.9).

2.2 The Program

PROGRESA is an antipoverty program run by the Mexican government. The program relies on conditional cash grants to poor households. It was introduced in 1997 in small rural communities and has gradually expanded to urban areas. Its main purposes are

Source: World Development Indicators, 2005

to improve the living conditions of eligible households, and promote investment by these households in the human capital of their children through school attendance and in the demand for basic health services.

The program beneficiaries receive bi-monthly cash transfers that represent about 20% of the household consumption. Due to important evidence that suggesting if the cash resources go to the mothers, larger shares of these resources are directed toward child health and nutrition, the money is granted to mothers, with some exceptions (Parker & Teruel, 2005 p. 206). In the case where the mother does not live with the family, the money is given to the person who is in charge of purchasing and preparing food and who takes care of the children's health.

The program's health component relies on four specific strategies:

1. Cash transfers:

Money transfers require the entire family to register at the nearest health care provider to schedule regular visits to control health and nutrition. Furthermore, the person who is receiving the money has to attend monthly lectures on health and hygiene: platicas. If a family does not fulfill one of these conditions, the cash grants are suspended. The increase in the family's income aims to result in a more frequent use of health services and an increase in food consumption. Estimates indicate that a 10% increase in income translates into a 3% to 4.5% increase in caloric availability (Technical Document Number 9 on the Evaluation of Oportunidades 2004, 2004, p.5).

2. Participation in the *platicas*:

Conducted by trained physicians and nurses, the meetings teach about nutrition and health topics. This training is delivered in the form of lectures to mothers. The emphasis is placed on preventive health care, including how to prevent disease through water treatment, safe food handling, and immunizations.

3. Nutritional supplements:

The Program not only provides daily nutritional supplements not only to mothers with infants and small children but also to children between 4 months and 2 years of age and pregnant and lactating mothers. Once a month, mothers receive six packets of supplements per eligible child. The packets provide calories, protein, vitamins, and other micro-nutrients.

4. Growth monitoring:

To receive nutritional supplements, growth monitoring of preschool children is required.

This control enables to parents to be aware of nutritional problems.

Table 1 summarizes the components, services and conditions of the program:

Components		Conditions	Services/ monthly transference		
Health	Medical attention	Registration at the nearest health unit	Basic package		
	Undernourishment prevention	Nutritional supple- ments			
	Education		Informative sessions		
Feeding		Attendance at health services and informative sessions	115^1 (household)		

Table 1: Components of the Program

 ^{1.} Values in Mexican pesos from January-June 1999. In June 1999, the peso/dolar exchange rate = 9.4409. About 12.18 USD. (Banco de México)
 Source: Analisis del Programa de Educacion, Salud y Alimentacion PROGRESA, RIMISP-FAO, p. 21, July 1999.

Family as well as community participation is one of the most important elements of PROGRESA. Therefore, people included in the program are asked to fulfill several requirements to continue receiving the money.

- Enroll people younger than 18 years old who have not finished basic education (primary or high school) to give them support to increase attendance rate and academic achievement.
- Enroll people up to 20 years old who have finished basic education to give them support to improve academic achievement.
- Enroll the family members in the designated medical unit.
- Attend scheduled appointments (for all family members) to receive medical services in the Health Package.
- Attend the monthly lectures on health and nutrition.

• Spend the money on goods and services that improve the health status of the family.

2.2.1 The design

Randomization was done at the locality level. In the first stage, a group of localities was selected that met the inclusion criteria for the program. The identification of the localities was conducted, in the first stage, by geographic focalization such that the program verified communities' deprivation and access to health services and basic school systems. The selection used a deprivation index based on social indicators.² The index classified the localities into five deprivation categories: very high, high, medium, low, and very low. Localities categorized to have a high or very high deprivation level had priority for inclusion in the program.³ Once selected, each locality was randomly allocated to the intervention or control group.





Source: General Rural Methodology Note, SEDESOL, November 2005

Stratified sampling was used for the original evaluation sample. It was conducted by locality size proportional to the number of localities. This process was implemented to generate a selection process with equal probabilities of selection for each locality

²The Deprivation Index was constructed from the sociodemographic data of the XI General Population and Household Census 1990 and 2000 and the 1996 Population and Household Count.

 $^{^3\}mathrm{For}$ further details about the construction of the index, see The Rural Methodology Note, reference 2

size. The sample was estimated over a universe of 4546 communities to select 330 for the treated group and a universe of 1850 communities to select 191 for the control group. After some adjustments, the final sample consisted of 320 treated localities, where PROGRESA first operated, and 186 control localities, where the program began to operate after some time.

During each intervention, where the locality was identified as a potential beneficiary, the program applied a poverty criterion (ENCASEH 97) to determine which households would be eligible to become beneficiaries of the program. Once the beneficiaries were identified, the head of the family received an identification card, the unique document that gives access to the program. In the final phase, the results were presented in a public assembly to the community to seek comments and make corrections and additions.

The program has survey evaluations every six months since the baseline survey in March 1998. A number of core questions about the demographic composition of house-holds and their socio-economic status were applied in each round of the survey. The surveys also asked about family background, schooling, health and nutritional status, health care utilization, consumption of food and non-food items, income, allocation of time of household members, and productive activities. The baseline includes information about 112319 individuals in 506 experimental communities, and about 60% corresponds to treated areas.

Table 2 describes the organization of each locality by beneficiary household and by their socio-economic condition. As we can see, the localities that receive PROGRESA include poor and non-poor families. In this group, only the poor families are eligible for the program and, therefore receive the bimonthly cash grants. The important thing to note is that the the poor/non-poor groups of families in the PROGRESA and in control localities have exactly the same background characteristics. Thus, before, during, and after the implementation of the program it is possible to compare the situation among the groups.

The experimental communities are located in the seven states that were among the first states to receive PROGRESA, including Veracruz, Guererro, San Luis Potosi, Hidalgo, Queretaro, Michoacan, and Puebla. Figure 1 shows the geographical localization of the beneficiary states as well as the disposable health infrastructure around the country. By January 2008, the program had around 35848 medical unities and 69 public hospitals (Lara, 2008).

PROGRESA Localites	Control Localities
poor families: receive money	poor families
non-poor families	non-poor familiess

 Table 2: PROGRESA Structure

Figure 3: Mexico Map: Location of the Experimental Communities (PROGRESA)



Source: Unidad IMSS-Oportunidades, SISPA, January-December 2007

3 Descriptive Statistics

The data corresponds to rural areas in Mexico. It contains information for about 125028 individuals interviewed in 1997 (baseline), 1998, 1999, 2000, 2003, and 2007. In total, it contains 451494 observations. The data is divided between control and treatment groups; within each group, people are categorized as poor (eligible) and non-poor (ineligible).

Table 3 describes PROGRESA's coverage. The participation of poor people in the program increased each year. In 1997 (baseline), almost 37% of individuals categorized as poor received the treatment; by 2000, this share had become 49.4%. Since 2003 all eligible individuals have boceme a part of the program, thus there is no information regarding the poor and non-poor share.

	Baseline		1998		1999		2000	
Receive treatment ¹	Yes	No	Yes	No	Yes	No	Yes	No
Non-poor Poor	$24.68 \\ 36.75$	$16.47 \\ 22.10$	$24.21 \\ 37.24$	$16.14 \\ 22.41$	$\begin{array}{c} 11.52\\ 46.61 \end{array}$	$7.980 \\ 31.90$	$11.17 \\ 49.38$	$7.521 \\ 31.93$
N	125449		130279		115889		130476	

Table 3: PROGRESA Composition

[1.] Values are percentages with respect to the total number of individuals (N). Source: PROGRESA Evaluation Data

Table 4 details the main variables of the baseline survey (ENCASEH 97). The summary statistics describe the socio-economic characteristics of the sample. Each variable is divided into the treated and control groups, as well as eligibility condition. Additionally, the table reports a t-test on the mean differences between groups (treated-control). the results suggest that the randomization worked almost perfectly in the sample.

In looking closely at the data, it is possible to get an idea of the sample's composition and social background. In the case of age, the sample is divided into five groups: baby (0-5 years), child (5-12 years), young (12-18 years), adult (18-65 years), and old individuals (older than 65 years). This division conforms to the survey composition: different questions were asked of different people depending on their age. Additionally, when analyzing health outcomes, it is important to make distinctions because a per-

Variable		Eligible =	= poor	In	eligible =	non-poor
Receive treatment	Yes	No	Diff. ¹ (abs.z-Value)	Yes	No	Diff. ¹ (abs.z-Value)
Baby	0.206	0.206	0.000 (0.935)	0.103	0.103	0.000 (0.956)
Child	0.245	0.246	(0.000) -0.001 (0.732)	0.142	0.139	0.004 (0.243)
Young	0.143	0.142	(0.001) (0.863)	0.162	0.167	(0.210) -0.003 (0.319)
Adult	0.378	0.376	(0.002) (0.638)	0.518	0.519	(0.002) (0.717)
Old	0.028	0.030	-0.001 (0.297)	0.075	0.073	(0.001) (0.540)
Male	0.500	0.500	$0.000 \\ (0.523)$	0.512	0.508	$0.004 \\ (0.431)$
Parents Ed.: Primary school ²	0.246	0.229	0.017 (0.077)	0.189	0.177	0.011 (0.199)
Actual schooling	0.378	0.375	0.003 (0.369)	0.246	0.240	0.007 (0.100)
Completed grade 5^3	0.094	0.093	(0.001) (0.844)	0.066	0.066	0.000 (0.864)
Completed grade 6^3	0.254	0.248	0.006 (0.909)	0.262	0.264	0.002 (0.748)
Secondary school ³	0.030	0.027	0.002 (0.192)	0.031	0.029	0.001 (0.525)
Health disabilities ⁴	0.062	0.063	-0.001 (0.652)	0.052	0.056	-0.004 (0.123)
$Employment^5$	0.229	0.232	-0.002 (0.450)	0.365	0.359	0.006 (0.169)
$Income^{6}$	36.01	34.44	1.572 (0.107)	49.28	51.06	-1.78 (0.286)
Mothers' age at first job	16.50	15.58	(0.919) (0.126)	17.332	17.679	-0.347 (0.585)
running water	0.175	0.176	-0.001 (0.734)	0.377	0.366	0.011 (0.525)
Wall: asbestos	0.001	0.001	`0.000 [´] (0.096)	0.002	0.001	(0.339)
Wall: metal	0.001	0.001	0.000' (0.690)	0.001	0.001	0.000' (0.777)

Table 4: Descriptive Statistics: Baseline (ENCASEH 1997)

^{1.} Difference between the treated and control groups.

² Refers to parents that completed primary school.

^{3.} Refers to individuals that have finished that grade.

⁴. Blind, mute, deaf, mentally and physically disabled people.
 ⁵. Individuals that engaged in a productive paid activity the week before the survey.

^{6.} Reported income on a daily basis.

son's health status depends on various characteristics conditioned on age.

In checking the data, it is possible to see that the eligible group contains an important percentage of young people (0-18 years), almost 60%, while the ineligible group contains a lot of people of middle age (18-65 years). In contrast, both groups contain a similar ratio of men and women.

In the case of education, parents in the eligible group mainly attended primary school. On the other hand, the attendance level at secondary school is higher for the non-poor group. The variable "Actual schooling" shows important difference between groups: poor people attend school more regularly than their peers in the non-poor group do.

A potential indicator of health status is defined by the variable "health disabilities", which indicates whether an individual has physical problems such as blindness, deafness, or the loss of a limb. The statistics show that the control and treated groups present very similar values. On average, 6.2% of the poor sample suffer one of the mentioned disabilities. This indicator is a bit lower for the second group: about 5%.

The labor conditions among the eligible and ineligible groups vary considerably. While in the first group, only 23% of the individuals able to work reported being emplyed in in the ineligible group, this rate increases to 36%. Furthermore, the non-poor group, on average, earns 15 pesos more than people in the poor group.

Finally, a number of variables describing "social background" are reported. The results suggest important differences between groups. In the poor sample, only 17.5% of the households have access to running water, while, in the second group, this variable is 37%. Finally, two variables define the wall characteristics of the houses, but the results suggest that there are not a lot of people living in houses with this type of structure: asbestos, metal.

Table 5 shows evidence of the effect of PROGRESA on health outcomes in different years. For sickness incidence, the values show that people who do not receive the treatment are sick more frequently than people who receive the cash transfer. Furthermore, this impact is not significant in the first year of the program (1998), but in later years, the effect becomes stronger and statistically significant. An important fact is that the results show a greater impact on the non-poor group than on the poor group. Additionally, people who belong to the treated group seem to stay sick for shorter periods in comparison to people who do not receive the program. In this case, however, the impact is greater for the eligible group.

Weight control improved for the eligible group. Although these results are not surprising given that people were entitled to regular health checks and control if they receive PROGRESA, it is important to point out that that people receiving cash transfers and belonging to the poor group, significantly increased the number of visits to the doctor to check the health and nutrition of their child; hence, the program is accomplishing its main goals, at least in the beginning.

Variable		Eligible =	igible = poor Ineligible = non-			non-poor
Receive treatment	Yes	No	Diff. ¹ (abs.z-Value)	Yes	No	Diff. ¹ (abs.z-Value)
Sickness incidence in 1998^2	0.109	0.113	-0.004 (0.059)	0.126	0.114	0.012 (0.000)
Sickness incidence in 1999^2	0.188	0.192	-0.005 (0.080)	0.133	0.145	-0.012 (0.001)
Sickness incidence in 2000^2	0.203	0.211	-0.008 (0.002)	0.264	0.290	-0.027 (0.000)
Sickness spell in 1998^3	7.231	7.627	0.415 (0.002)	9.655	9.269	0.386 (0.035)
Sickness spell in 1999^3	9.429	10.14	-0.707 (0.007)	10.94	11.56	0.623 (0.269)
Sickness spell in 2000^3	9.874	10.75	-0.879 (0.006)	11.48	11.83	-0.356 (0.586)
Weight control in 1998^4	0.868	0.782	-0.087 (0.000)	0.797	0.846	-0.049 (0.000)
Weight control in 1999^4	0.932	0.917	0.015 (0.026)	0.911	0.912	-0.001 (0.515)
Weight control in 2000^4	0.923	0.920	0.003 (0.447)	0.869	0.891	-0.021 (0.171)
Weight control (times) in 1998^5	4.123	4.178	-0.055 (0.000)	3.798	3.649	0.149 (0.572)
Weight control (times) in 1999^6	3.526	3.112	0.413 (0.000)	3.235	3.181	0.128 (0.253)
Weight control (times) in 2000^6	4.795	4.568	0.227 (0.561)	4.041	4.096	-0.055 (0.686)

Table 5: Descriptive Evidence of the Effect of Cash Subsidies on Health Outcomes

^{1.} Difference between the treated and control groups.

^{2.} Percentage of people who reported a sickness incidence during the previous 4 weeks.

^{3.} Days people reported being sick.

^{4.} Children younger than 5 years old who in the last six months had their weight were controlled.

^{5.} Number of times that children were weight-controlled in the previous year.

⁶. Number of times that children were weight-controlled in the previous 6 months.

4 Identification Framework

The topic of treatment evaluations concerns measuring the impact of interventions on outcomes of interest. A measure of causal impact is the average difference in the outcomes of the treated and non-treated (control) groups. Within a framework of a potential outcome model, which assumes that every individual in the population is potentially exposed to the treatment, the outcomes of the different groups form a basis for treatment evaluation.

With $D \in [0, 1]$ being the variable defining whether a person lives in a village that receives PROGRESA, and H_{1i} and H_{0i} being the outcomes if an individual receives or not the treatment respectively, we have:

$$H_i = \begin{cases} H_{1i} & \text{if } D_i = 1\\ H_{0i} & \text{if } D_i = 0 \end{cases}$$

Receiving and not receiving the treatment are mutually exclusive states: for any given individual, only one of the measures is available. The effect of the cause D on the health status of individual i is measured by: $(H_{1i} - H_{0i})$. The average causal effect of $D_i = 1$, relative to $D_i = 0$, is measured by the average treatment effect (ATE):

$$ATE = E[H|D = 1] - E[H|D = 0]$$

where the expectations are with respect to the probability distribution over the target population.

The descriptive evidence presented in section 3 shows that the randomization process that assigned individuals to the control and treatment groups was successfully implemented in PROGRESA. The random assignment implies that individuals exposed to the treatment are chosen randomly, so the treatment assignment does not depend on the outcome and is uncorrelated with the attributes of the treated subjects.

Due to comparability issues among the questions and structure, this paper includes data from two surveys: ENCEL99 and ENCEL00. Thus, the evaluation format allows the estimation of PROGRESA's impact on different age groups using, as explained, several health outcomes.

Let H_{iv} denote the individual health outcome per village and age group measured in 1999 and 2000. Let $P_i = 1$ if the individual lives in a household classified as poor, and $P_i = 0$ otherwise. The variable Pn_{ig} corresponds to the percentage of poor (=eligible) neighbors per village. Including this variable in the analysis allows us to determine the impact of the treated neighbors on their non-treated peers. It is obvious that not everybody is related or socialize with everybody in the PROGRESA villages. Therefore, to calculate this variable, we use the geographical references of each family. By determining each household localization, it is possible to establish the number of treated and non-treated neighbors to whom each household member might be related based on two conditions: age and closeness. Thus, the indirect impact of PROGRESA is measured by treated individuals interacting with their non-treated peers who might engage in the same activity: attending school, working in a specific area, and living in the same neighbourhood. The next figure explains how the model incorporate the measure of the spillover effect.

Figure 4: PROGRESA: Interations within villages



Let $D_v = 1$ if the individual lives in a village that receives PROGRESA and $D_v = 0$ otherwise. X_i contains background characteristic variables such as gender, maximum level of education in the family, state where the household is located, work status, and dummy variables for the different years in the sample.

$$H_{iv} = \varphi_0 + \varphi_1 D_v + \varphi_2 P n_{ig} + \varphi_3 (P n_{ig} * D_v) + \sum_i \gamma_i X_i + \mu_i$$

where $E(\mu_i|D_{iv}) = 0$ by virtue of the randomized assignment of PROGRESA to villages when estimating this model in the sub-groups of the non-poor and the poor. In this context, φ_1 defines the direct effect of the program, while φ_2 and φ_3 measure the

spillover effect when the last equation is estimated for the non-poor (=ineligible).

 H_{iv} takes different values defining each group's health status. For babies and children, the dependent variable corresponds to sickness incidence, sickness spell, specific illness faced by children, and weight control. For young individuals, people between 12 and 18 years old, H_{iv} defines whether an individual had difficulty with normal activities during the previous four weeks and sickness incidences. Finally, for adult and older individuals, H_{iv} defines whether individuals can deal with normal, hard, and moderate activities: the number of km that people can walk before getting tired, sickness incidence, and their capacity to dress and bathe themselves.

If $H_{iv} \in [0, 1]$, this paper uses probit and logit specifications in addition to a standard OLS estimation. When H_{iv} is a count variable, such as the number of days that an individual reported being sick, this document estimates a negative binomial model.⁴. This type of model is applicable when a Poisson estimation is inappropriate due to over-dispersion (which is the general behavior of the explained variables). In a poisson distribution the mean and variance are equal. If the variance is greater than the mean, the distribution is said to display over-dispersion.

A zero-inflated negative binomial regression generates two separate models and then combines them. First, a logit model is generated for the "certain zero" cases, predicting whether or not an individual would be in this group. Then, a negative binomial model is generated predicting the counts for those individuals who are not certain zeros.

$$Pr[y|x] = \frac{\Gamma(y+\alpha^{-1})}{y!\Gamma(\alpha^{-1})} (\frac{\alpha^{-1}}{\alpha^{-1}+\lambda})^{\alpha^{-1}} (\frac{\lambda}{\alpha^{-1}+\lambda})^y$$

In the context of count regression models, the negative binomial distribution can be thought of as a Poisson distribution with unobserved heterogeneity which, in turn, can be conceptualized as a mixture of two probability distributions, poisson and gamma.

When the count variables have an excessive number of zeros, the model that best fits the distribution of these variables is a zero-inflated Negative Binomial. This specification supplements a count density with a binary process. If the binary process takes a value 0, with probability $f_1(0)$, $H_{it} = 0$. If the binary process takes a value of 1, with a probability of $f_1(1)$, $H_{it} = 0, 1, 2...$ This definition lets zero counts occur in two ways: as a realization of the binary process and as a realization of the count process. The density for this specification can be defined as:

$$g(y) = \begin{cases} f_1(0) + (1 - f_1(0))f_2(0) & \text{if } H_{it} = 0\\ (1 - f_1(0))f_2(y) & \text{if } H_{it} \ge 1 \end{cases}$$

⁴The negative binomial distribution is given by:

5 Estimation Results

Table 6 and 7 summarize the estimation results of the direct and indirect effect for each of the health outcomes. The data included in the estimations corresponds to the 1999 and 2000 surveys. The table is divided into eligibility conditions (poor and non-poor) and age groups. In the Appendix, each estimated equation is reported, including all the covariates.

5.1 Direct Effect of PROGRESA

In general the variable, "treatment" has a positive effect on the health outcomes in the analysis. In fact, both groups, ineligible and eligible, seem to improve their health status due to the cash grants. In spite of the fact that few of the results are statistically significant, the significant ones show an important impact. It is important to stress the fact that the main variables enforced by the program such as sickness incidence and weight and height control, improved considerably among children.

Specifically, for babies, children, and young individuals, the program reduces their sickness spell length in addition to lowering the possibility that they will become ill. In fact, sickness such as cough, fever, and respiratory problems are less frequent among PROGRESA kids. Moreover, children immersed in the program increased their frequency of height and weight control.

PROGRESA has a significant impact on adult individuals, and there is no significant effect registered for older individuals, results consistent with related literature. In the case of adults, the program not only improved their capabilities to deal with normal, simple, and hard activities but also enabled them to walk at least 2 km before getting tired. Finally, adult individuals who received the treatment show decreased probabilities of getting sick.

5.2 Indirect Effect of PROGRESA: Spillover Results

An interesting fact about PROGRESA is that it seems to have a major impact on people who do not receive the cash grants (=ineligible). This result is explained by the interactions that these families have with people receiving the treatment (=eligible). Not surprisingly, the ineligible age group that show the greatest improvement in health

Age	Variable	OLS	3	\mathbf{Log}	it	\mathbf{Prob}	it
group		Non-Poor	Poor	Non-Poor	Poor	Non-Poor	Poor
	Sickness incidence	0.108^{*} (0.05)	-0.065^{*} (0.03)	0.113^{*} (0.05)	-0.058^{*} (0.03)	0.129^{*} (0.05)	-0.060^{*} (0.03)
	Days sick	$0.019 \\ (0.07)$	-0.019 (0.06)	$0.522 \\ 0.435$	-0.373 0.189		
Babies	Sickness: diahrrea	-0.057 (0.12)	-0.007 (0.06)	-0.095 (0.13)	-0.008 (0.07)	-0.092 (0.13)	-0.012 (0.07)
	Sickness: fever	-0.490 (0.41)	-0.490^{*} (0.23)	-0.267 (0.24)	-0.261^{*} (0.13)	-0.268 (0.23)	-0.255^{*} (0.12)
	Sickness: cough	0.656^{**} (0.22)	0.279^{**} (0.10)	0.196^{**} (0.07)	0.074^{**} (0.03)	0.205^{**} (0.07)	0.080^{**} (0.03)
	Sickness: respiratory prob.	-0.169 (0.18)	-0.062 (0.08)	-0.038 (0.04)	-0.010 (0.01)	-0.039 (0.04)	-0.008 (0.01)
	Sickness: other	0.445^{**} (0.17)	0.193^{*} (0.08)	0.075^{*} (0.03)	0.024^{*} (0.01)	0.082^{**} (0.03)	0.026^{*} (0.01)
	Control height & weight	-0.119 (0.07)	$0.051 \\ (0.04)$	-0.114 (0.06)	$0.047 \\ (0.04)$	-0.122 (0.07)	0.044 (0.04)
	Control weight (number times)	$0.434 \\ (0.40)$	-0.183 (0.25)	$0.302 \\ (0.371)$	-0.141 (0.228)		
Children	Sickness incidence	$0.017 \\ (0.05)$	-0.013 (0.02)	$0.026 \\ (0.03)$	-0.010 (0.02)	$\begin{array}{c} 0.031 \\ (0.04) \end{array}$	-0.008 (0.02)
	Days sick	$0.015 \\ (0.04)$	-0.014 (0.02)				
Young	Sickness incidence	$0.015 \\ (0.04)$	-0.014 (0.02)	0.001 (0.03)	-0.012 (0.02)	0.001 (0.03)	-0.016 (0.02)
	Days sjck	$0.015 \\ (0.04)$	-0.014 (0.02)				

Table 6: Marginal Probabilities: Impact of PROGRESA on Health

Age group	Variable	OL	S	Logi	t	Probit		
	, and the	No Pobre	Pobre	No Pobre	Pobre	No Pobre	Pobre	
	Problem: normal act.	-0.043^{*} (0.02)	-0.003 (0.01)	-0.031 (0.02)	-0.003 (0.01)	-0.037 (0.02)	-0.004 (0.01)	
	Problem: hard act.	0.029 (0.03)	-0.004 (0.02)	$0.025 \\ (0.03)$	-0.002 (0.02)	$0.025 \\ (0.03)$	-0.002 (0.02)	
	Problem: moderate act.	0.037 (0.03)	$0.009 \\ (0.02)$	$0.033 \\ (0.03)$	$0.011 \\ (0.01)$	$0.031 \\ (0.03)$	$0.010 \\ (0.02)$	
Adultos	Problem: simple act.	-0.036 (0.02)	-0.005 (0.01)	-0.029 (0.02)	-0.006 (0.01)	-0.029 (0.02)	-0.009 (0.01)	
	Problem: walk	-0.034 (0.02)	-0.004 (0.01)	-0.029 (0.02)	-0.006 (0.01)	-0.028 (0.02)	-0.007 (0.01)	
	Problem: dress	-0.057^{**} (0.02)	-0.002 (0.01)	-0.035^{*} (0.01)	-0.002 (0.01)	-0.048^{**} (0.02)	-0.004 (0.01)	
	Days sick	-0.406^{*} (0.17)	-0.161 (0.10)					
	No. km able to walk	-0.705 (0.37)	$\begin{array}{c} 0.743^{***} \\ (0.22) \end{array}$					
	Problem: normal act.	-0.046 (0.06)	$0.011 \\ (0.05)$	-0.043 (0.06)	$0.010 \\ (0.05)$	-0.042 (0.06)	$0.010 \\ (0.05)$	
	Problem: hard act.	0.019 (0.07)	-0.019 (0.06)	$0.023 \\ (0.08)$	-0.017 (0.06)	$0.026 \\ (0.08)$	-0.011 (0.06)	
	Problem: moderate act.	0.059 (0.07)	-0.018 (0.06)	$0.065 \\ (0.08)$	-0.017 (0.06)	$0.069 \\ (0.08)$	-0.012 (0.06)	
Ancianos	Problem: simple act.	0.098 (0.08)	-0.014 (0.06)	$0.105 \\ (0.08)$	-0.014 (0.06)	$0.103 \\ (0.08)$	-0.012 (0.06)	
	Problem: walk	0.012 (0.07)	-0.020 (0.06)	$0.016 \\ (0.08)$	-0.017 (0.06)	$0.015 \\ (0.08)$	-0.013 (0.06)	
	Problem: dress	-0.084^{***} (0.02)	-0.012 (0.01)	-0.015^{*} (0.01)	-0.005 (0.01)	-0.017^{*} (0.01)	-0.004 (0.01)	
	Days sick	-1.090 (1.04)	-0.516 (0.10)					
	No. Km able to walk	$0.995 \\ (0.94)$	-0.574 (0.83)					

Table 7: Marginal Probabilities: Impact of PROGRESA on Health

due to this situation is children between 0 and 12 years of age.

Specifically, children decrease not only their probabilities of getting sick but also their sickness spell. Figures 4-7 show in a detailed way the evolution of the main responsive variables for children. The graphs are constructed using an Epanechnikov-Kernel regression of the impact on health outcomes for children who are not eligible for cash transfers on the eligible fraction in their peer group (other children in the same village). The figures show that health conditions improve for ineligible individuals as the number of treated poor neighbors increases and compare the results with the same environment the the non-PROGRESA villages. Ineligible individuals living in treated villages who have neighbors that receive PROGRESA tend to have a lower probability of getting sick. Similarly, they decrease the number of days sick, and in the case of cough, they are less likely to get this type of illness.

For adults, the variables that have improved are their capacity to deal with normal activities: ineligible individuals living in treated villas have considerably decreased their problems when dealing with normal activities as the number of treated neighbors has increased. This last characteristic is shared with their capabilities to deal with hard activities, bathe and dress themselves, and walk a certain number of kilometers.



Figure 5: Spillover Effect: Babies

Source: Estimation Results



Figure 6: Spillover Effect: Sickness Spell

Source: Estimation Results

Figure 7: Spillover Effect: Adult Individuals



Source: Estimation Results



Figure 8: Spillover Effect: Old Individuals

Summary and Conclusions 6

This document addresses the impact of an anti poverty program on several health outcomes. The program combines a traditional cash transfer with financial incentives for families to invest in the human capital of children: nutrition, health, and education. To receive the cash transfers, families are required to participate in growth monitoring and nutrition supplement programs, as well as to regularly attend education programs about health and hygiene. The next table summarizes the significant effects presented by the estimation results to gain a clearer view of the program's impact.

As we can see, the direct effect of the program has a major impact on the enforced variables for the 14 targeted population with respect to sickness incidence and health checkups. In general, treated individuals exhibit better health status based on their sickness frequency as well as their sickness spell. Moreover, people receiving PRO-GRESA are more likely to deal with simple/moderate and strong activities. Finally, adult individuals and older individuals living in villages receiving the treatment not only seem to be able to walk more km. on average, but also feel that their capacity to walk at least 2 km. before getting tired has improved.

Regarding the spillover effect, the results are surprisingly large. People who live in treated villages but are ineligible for the program improve their health status, measured in their probabilities of being sick, the number of days that sickness spell lasts,

Source: Estimation Results

	Non-Poor	Poor
Babies	 Lower sickness incidence: diarrhea, fever, respiratory problems Higher incidence of sickness: cough Less frequency of controlling height and weight 	 Lower sickness incidence: diarrhea, fever, respiratory problems Higher incidence of sickness: cough Higher frequency controlling height and weight
Children	Higher probability of getting sickMore days sick	 Lower probability of getting sick Less days sick
Young	\bullet Lower probability of getting sick	\bullet Lower probability of getting sick
Adults	Fewer problems dealing with normal, simple, and daily activities; walkingFewer days sick	 Fewer problems dealing hard activities. Less days sick On average is able to walk more km.
Old	 Fewer problems dealing with normal, simple, and daily activities; walking Fewer days sick On average able to walk more km. 	 Fewer problems dealing with normal, simple, and daily activities; walking Less days sick

 Table 8: Direct Effect of PROGRESA

	Non-Poor	Poor
Babies	 Lower sickness incidence: cough and other illness Fewer days sick Higher frequency of controlling height and weight 	 Lower sickness incidence: cough and other illness Less days sick Higher frequency of controlling height and weight
Children	• Less days sick	• Lower probability of getting sick
Young	• Lower probability to get sick	• Less days sick
Adults		• Fewer problems dealing with normal, simple and daily activities
Old		• Fewer problems dealing with daily ac- tivities

and their capacity to deal with different daily activities. We attribute this result to the social interactions generated by the structure of the program.

Although PROGRESA benefits both eligible (=poor) and ineligible (=non-poor) individuals, we can say that, globally, the impact is stronger in poor families. The effects vary over age groups, being more significant for children (0-5 years old) and adult individuals (19-65 years old). In contrast, those people who seem to be less affected or have almost no benefits are young (6-18) and old (older 65 years) individuals. This result might occur because young individuals are generally healthier, so they are less exposed to diseases, while old individuals are more often exposed because of declining health.

The results presented correspond to the evaluation data of the PROGRESA program. It is important to note that there might be a problem in the data collection because one person in each family provides information. Therefore, in most cases, their answers are just beliefs instead of affirmations. In spite of this, the results show that PROGRESA seems to have achieved most of its goals, and, in general, the health condition of treated individuals has improved.

References

- Aguero J., & Carter M., The Impact of Unconditional Cash Transfers on Nutrition: The South African Child Support Grant. International Poverty Centre, United Nations Development Program.
- Behrman, J. R., 1993. Will Depeloping Coutry nutrition improve with income? A case study for rural South India. Journal of Political Economy.
- Bouis, H. E., and L. J. Haddad. 1992. Are estimates of calorie-income elasticities too high? A recalibration of the plausible range. Journal of Development Economics.
- 4. Coady P., (2001). An Evaluation of the Distributional Power of PROGRESA'S Cash Transfers in Mexico. International Food Policy Research Institute.
- Coordinacion Nacional de Programa de Desarrollo Humano Oportunidades, SEDESOL (2005). General Rural Methodology Note. Instituto Nacional de Salud Publica.
- 6. Don S., (2000). PROGRESA: An Overview. EGAT/PR.
- Gertler P. & Boyce S., (2000). An Experiment in Incentive-Based Welfare: The Impact of PROGRESA on Health. International Food Policy Research Institute.
- Jere R. Behrman & John Hoddinott, (2005). Programme Evaluation with Unobserved Heterogeneity and Selective Implementation: The Mexican "PROGRESA" Impact on Child Nutrition, Oxford Bulletin of Economics and Statistics, Department of Economics, University of Oxford, vol. 67(4), pages 547-569, 08.
- Cattaneo A. & Lalive R., Social Interactions and Schooling Decisions (September 2006). CEPR Discussion Paper No. 5816
- Lara Gisela, 2008. Conferencia Binacional de Promotoras Voluntarias de Salud, Programa IMSS-Oportunidades, Instituto Mexicano de Seguro Social.
- Loggia, Silvia (in progress), "Los programas de salud reproductiva en el contexto inter- cultural: Un acercamiento a las representaciones y prácticas de los agentes de la medi- cina institucional en una zona indígena", Doctoral thesis, CEDDU, El Colegio de México.

- Lozano, Rafael, et al. (1993), Desigualdad, pobreza y salud en México, Consejo Consultivo del Programa Nacional de Solidaridad, El Nacional, México.
- 13. Nanak Kakwani & Fabio Veras Soares & Hyun H. Son, 2005. "Conditional cash transfers in African countries," Working Papers 9, International Poverty Centre.
- 14. Nigenda, Gustavo, et al. (2001), Descentralización de los servicios de salud y los programas de salud reproductiva: Situación en cuatro estados de México, Instituto Nacional de Salud Públi- ca/ Fundación Mexicana para la Salud/Fundación Ford.
- 15. Rawlings L., & Rubio G., Evaluating the Impact of Conditional Cash Transfer Programs, (2005). The World Bank Research Observer, vol. 20, no.1
- 16. Rivera J. et al (2004). Impact of the Mexican Program for Education, health, and Nutrition (PROGRESA) on rates of Growth and Anemia in Infants and Young Children. A Randomized Effectiveness Study. American Medical Association.
- Sepulveda Jaime, ed. (1993), La salud de los pueblos indígenas en México, Secretaría de Salud/Instituto Nacional Indigenista, México.
- Scott John, 1999. Analisis del Programa de Educacion, Salud y Alimentacion (PROGRESA), Estudio RIMISP-FAO, CIDE. Programa IMSS-Oportunidades, In stituto Mexicano de Seguro Social.
- Scott John, 1999. Analisis del Programa de Educacion, Salud y Alimentacion (PRO- GRESA), Estudio RIMISP-FAO, CIDE. Programa IMSS-Oportunidades, In stituto Mexicano de Seguro Social.
- 20. Wooldrige J., Econometric Analysis of Cross Section and Panel Data, The MIT Press, Cambridge Massachusetts, London, England.
- 21. World Bank (2001a), Engendering Development Through Gender Equality in Rights, Resources, and Voice, World Bank, Oxford University.

7 Appendix

Ineligible Eligible Ineligible Eligible Ineligible Eligible Sickness inciderer 0.001 -0.008 -0.002 -0.054 -0.002 -0.029 poor share -0.156*** -0.133*** -0.849*** -0.874*** -0.484** -0.484** -0.484** -0.497*** -0.497*** poor share -0.066 (0.02) -0.013 (0.13) (0.16) (0.02) N 2434 20192 2434 20192 2434 20192 Diharrea - - -0.003 -0.003 -0.043 -0.006 poor share -0.003 0.000 -0.130 (0.13) (0.13) (0.13) poor share -0.042 -0.005 1.554 -0.315 0.713 -0.094 (0.03) (0.01) (0.89) (0.45) (0.41) (0.18) N 1902 16839 1902 16839 1902 16839 poor share -0.066 -0.069 0.379
Sickness incidence U <thu< th=""> U <thu< th=""></thu<></thu<>
treatment 0.001 -0.008 -0.002 -0.054 -0.002 -0.029 poor share -0.156** -0.133*** -0.849** -0.849** -0.484** -0.484** -0.497*** N 2434 20192 2434 20192 2434 20192 Diharrea - - - - - - - - - - - - 0.007 - 0.007 - 0.007 - 0.097 - 0.497*** - 0.497*** - 0.497*** - 0.497*** - 0.497*** - 0.497*** - 0.497*** - 0.497****
$\begin{array}{llllllllllllllllllllllllllllllllllll$
poor share -0.156^{**} (0.06) -0.133^{***} (0.02) -0.849^{***} (0.32) -0.484^{***} (0.13) -0.484^{***} (0.07)N243420192243420192243420192Diharrea treatment -0.003 (0.01)0.000 (0.00) -0.102 (0.03) -0.043 (0.13) -0.006 (0.03)poor share 0.042 (0.03) -0.005 (0.01) 1.554 (0.98) -0.315 (0.41) -0.094 (0.45)N190216839190216839190216839Pever treatment -0.003 (0.03) 0.011 (0.16) -0.021 (0.06) -0.003 (0.08) -0.003 (0.03)poor share 0.066 (0.03) -0.066 (0.03) 0.011 (0.16) -0.021 (0.06) -0.003 (0.03)poor share 0.066 (0.03) -0.064 (0.03) -0.042^* (0.20) 0.207 (0.20) -0.240^* (0.26)N204817934204817934204817934Cough treatment 0.052 (0.05) -0.064 (0.41) -0.069 (0.07) (0.07) (0.03) -0.300^* (0.04) -0.887^{***} (0.17) -0.587^{***} (0.24)N212518279212518279212518279Respiratory problems treatment -0.048 (0.04) -0.300 (0.04) -0.300 (0.44) -0.618 (0.17) -0.734^{***} (0.18)N212518279212518279212518279Respiratory problems treatment
(0.06) (0.02) (0.32) (0.13) (0.19) (0.07) N243420192243420192243420192Diharrea </td
N 2434 20192 2434 20192 2434 20192 Diharrea treatment -0.003 0.000 -0.102 -0.003 -0.043 -0.006 poor share 0.042 -0.005 1.554 -0.315 0.713 -0.094 N 1902 16839 1902 16839 1902 16839 Fever - - -0.003 0.011 -0.021 -0.003 -0.066 poor share 0.066 -0.003 0.011 -0.021 -0.003 -0.066 fever - - -0.066 0.379 -0.482* 0.207 -0.240* poor share 0.066 -0.069* 0.379 -0.482* 0.207 -0.240* N 2048 17934 2048 17934 2048 17934 Cough - - -0.004 0.133 -0.014 0.069 -0.079 poor share 0.052 -0.004 0.133 -0.014 0.069
Diharrea -0.003 0.000 -0.102 -0.003 -0.043 -0.006 poor share 0.042 -0.005 1.554 -0.315 0.713 -0.094 (0.03) (0.01) (0.98) (0.45) (0.41) (0.18) N 1902 16839 1902 16839 1902 16839 1902 16839 Fever -
$\begin{array}{cccccccccccccccccccccccccccccccccccc$
$\begin{array}{c c c c c c c c c c c c c c c c c c c $
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$
(0.03) (0.01) (0.98) (0.45) (0.41) (0.18) N190216839190216839190216839Fever </td
$\begin{array}{c c c c c c c c c c c c c c c c c c c $
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $
$\begin{array}{cccccccccccccccccccccccccccccccccccc$
$\begin{array}{c c c c c c c c c c c c c c c c c c c $
$\begin{array}{c c c c c c c c c c c c c c c c c c c $
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$
$\begin{array}{c c c c c c c c c c c c c c c c c c c $
$\begin{array}{c c c c c c c c c c c c c c c c c c c $
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$
$\begin{tabular}{ c c c c c c c c c c c c c c c c c c c$
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$
$\begin{array}{c c c c c c c c c c c c c c c c c c c $
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$
poor share -0.210 (0.12) -0.191^{***} (0.04) -1.389 (0.78) -1.681^{***} (0.32) -0.734^{***}
(0.12) (0.04) (0.78) (0.32) (0.36) (0.14) N 1932 17040 1932 17040 1932 17040
N 1932 17040 1932 17040 1932 17040
Other sickness
treatment -0.053 -0.007 -0.472 -0.108 -0.216 -0.049
$\begin{array}{cccccccccccccccccccccccccccccccccccc$
poor share -0.141 -0.175^{***} -1.262 -2.372^{***} -0.599 -0.942^{***}
(0.12) (0.03) (1.07) (0.43) (0.45) (0.18)
N 1894 16795 1894 16795 1894 16795
Weight control
treatment -0.022 0.019*** -0.426 0.394*** -0.206 0.179***
$\begin{array}{cccccccccccccccccccccccccccccccccccc$
poor share 0.094^* -0.007 1.740^* -0.140 0.852^* -0.082
N 1212 8365 1212 8365 1212 8365
familu's heal work Ves Ves Ves Ves Ves Ves
status Yes Yes Yes Yes Yes Yes
state effects Yes Yes Yes Yes Yes
year effects Yes Yes Yes Yes Yes Yes

Table 10: Baby 1: Health Status

* p<0.05, ** p<0.01, *** p<0.001 Var.=1 if baby was sick during the last 4 weeks; 0 otherwise

	OI	\mathbf{S}	Log	git	Probit		
	Ineligible	Eligible	Ineligible	Eligible	Ineligible	Eligible	
Normal activities							
treatment	-0.011*	-0.003	-0.106*	-0.044	-0.061*	-0.023	
	(0.01)	(0.00)	(0.05)	(0.03)	(0.03)	(0.02)	
poor share	0.008	-0.009	0.090	-0.092	0.050	-0.046	
	(0.02)	(0.01)	(0.16)	(0.10)	(0.09)	(0.05)	
Ν	16309	58932	16309	58932	16309	58932	
Hard activities							
treatment	-0.014*	0.003	-0.134*	0.033	-0.074^{**}	0.016	
	(0.01)	(0.00)	(0.05)	(0.03)	(0.03)	(0.02)	
poor share	-0.039*	-0.011	-0.375^{*}	-0.149	-0.201*	-0.071	
	(0.02)	(0.01)	(0.17)	(0.11)	(0.09)	(0.06)	
Ν	14763	54936	14763	54936	14763	54936	
Moderate activiti	es						
treatment	-0.003	0.006^{**}	-0.030	0.081^{*}	-0.018	0.040^{*}	
	(0.01)	(0.00)	(0.05)	(0.03)	(0.03)	(0.02)	
poor share	-0.033	-0.006	-0.331	-0.079	-0.177	-0.038	
	(0.02)	(0.01)	(0.17)	(0.11)	(0.09)	(0.06)	
Ν	14760	54923	14760	54923	14760	54923	
Simple activities							
treatment	0.015^{***}	0.001	0.227^{***}	0.026	0.116^{***}	0.012	
	(0.00)	(0.00)	(0.06)	(0.04)	(0.03)	(0.02)	
poor share	0.025	0.009	0.349	0.199	0.166	0.089	
	(0.01)	(0.01)	(0.21)	(0.14)	(0.10)	(0.07)	
Ν	14749	54915	14749	54915	14749	54915	
Walk more than 2	2 km befor	e getting	tired				
treatment	0.008	-0.003	0.096	-0.042	0.050	-0.020	
	(0.00)	(0.00)	(0.06)	(0.04)	(0.03)	(0.02)	
poor share	0.038^{*}	0.021**	0.445^{*}	0.352^{**}	0.224^{*}	0.167^{**}	
	(0.02)	(0.01)	(0.19)	(0.13)	(0.10)	(0.06)	
Ν	14726	54794	14726	54794	14726	54794	
Bath and dress							
treatment	0.005^{*}	0.001	0.291^{*}	0.072	0.125^{*}	0.032	
	(0.00)	(0.00)	(0.13)	(0.07)	(0.05)	(0.03)	
poor share	0.018*	0.004	0.943^{*}	0.282	0.371^{*}	0.105	
	(0.01)	(0.00)	(0.42)	(0.26)	(0.17)	(0.10)	
Ν	14729	54826	14729	54826	14729	54826	
Number of km a	person can	walk					
treatment	-0.003	0.121^{**}					
	(0.10)	(0.04)					
poor share	0.109	0.390^{*}					
	(0.32)	(0.15)					
family's head work	Yes	Yes	Yes	Yes	Yes	Yes	
status	Yes	Yes	Yes	Yes	Yes	Yes	
state effects	Yes	Yes	Yes	Yes	Yes	Yes	
year effects	Yes	Yes	Yes	Yes	Yes	Yes	

Table 11: Adult 1: Health Status

* p < 0.05, ** p < 0.01, *** p < 0.001Var.=1 if baby was sick during the last 4 weeks; 0 otherwise

	O	LS	Logit		Probit	
	Ineligible	Eligible	Ineligible	Eligible	Ineligible	Eligible
Normal activities						
treatment	0.015	-0.017	0.092	-0.099	0.050	-0.057
	(0.02)	(0.01)	(0.12)	(0.07)	(0.07)	(0.04)
poor share	0.093	0.015	0.591	0.088	0.330	0.042
	(0.07)	(0.04)	(0.41)	(0.24)	(0.24)	(0.14)
Ν	1806	5495	1806	5495	1806	5495
Hard activities						
treatment	-0.022	0.006	-0.099	0.027	-0.063	0.019
	(0.02)	(0.01)	(0.11)	(0.06)	(0.07)	(0.04)
poor share	-0.073	-0.041	-0.337	-0.203	-0.207	-0.137
	(0.08)	(0.05)	(0.36)	(0.23)	(0.22)	(0.14)
Ν	1729	5158	1729	5158	1729	5158
Moderate activitie	es					
treatment	-0.003	0.006^{**}	-0.030	0.081^{*}	-0.018	treatment
-0.023	0.013	-0.102	0.064	-0.063	0.041	
	(0.02)	(0.01)	(0.11)	(0.06)	(0.07)	(0.04)
poor share	-0.124	-0.108*	-0.572	-0.518*	-0.351	-0.326*
	(0.08)	(0.05)	(0.36)	(0.23)	(0.22)	(0.14)
Ν	1728	5155	1728	5155	1728	5155
Simple activities						
treatment	0.027	-0.002	0.111	-0.007	0.070	-0.005
	(0.02)	(0.01)	(0.10)	(0.06)	(0.06)	(0.04)
poor share	0.170^{*}	0.055	0.722^{*}	0.235	0.448^{*}	0.151
	(0.08)	(0.05)	(0.35)	(0.21)	(0.22)	(0.13)
Ν	1725	5146	1725	5146	1725	5146
Walk more than 2	km before	getting tir	ed			
treatment	0.044	-0.005	0.194	-0.024	0.120	-0.015
	(0.02)	(0.01)	(0.11)	(0.06)	(0.06)	(0.04)
poor share	0.108	0.035	0.479	0.168	0.290	0.120
	(0.08)	(0.05)	(0.36)	(0.22)	(0.22)	(0.14)
Ν	1724	5133	1724	5133	1724	5133
Bath and dress						
treatment	0.003	0.000	-0.069	-0.035	-0.035	-0.015
	(0.00)	(0.00)	(0.11)	(0.06)	(0.05)	(0.03)
poor share	0.000	-0.002	-0.901*	-0.327	-0.378*	-0.148
	(0.01)	(0.01)	(0.40)	(0.22)	(0.16)	(0.09)
Ν	16452.000	59966.000	16452.000	59966.000	16452.000	59966.000
Number of km a person can walk						
treatment	-0.138	0.313				
	(0.33)	(0.20)				
poor share	1.718	-1.412				
	(1.11)	(0.73)				
family's head work	Yes	Yes	Yes	Yes	Yes	Yes
status	Yes	Yes	Yes	Yes	Yes	Yes
state effects	Yes	Yes	Yes	Yes	Yes	Yes
year effects	Yes	Yes	Yes	Yes	Yes	Yes

Table 12: Old 1: Health Status

* p < 0.05, ** p < 0.01, *** p < 0.001Var.=1 if baby was sick during the last 4 weeks; 0 otherwise

	OLS		Zero-Inflated/Poisson	
	Ineligible	Eligible	Ineligible	Eligible
treatment	0.207**	0.240***	0.094*	0.108***
	(0.08)	(0.03)	(0.04)	(0.02)
poor share	0.195	0.093	0.106	0.046
	(0.26)	(0.11)	(0.14)	(0.05)
r2	0.785	0.784		
F	350.410	2434.378		
Ν	1164	8071	1164	8071
family's head work	Yes	Yes	Yes	Yes
status	Yes	Yes	Yes	Yes
state effects	Yes	Yes	Yes	Yes
year effects	Yes	Yes	Yes	Yes

Table 13: Baby: Weight Control Frequency

* p < 0.05, ** p < 0.01, *** p < 0.001Var.= number of days a baby was sick

	OLS		Lo	Logit		Probit	
	Ineligible	Eligible	Ineligible	Eligible	Ineligible	Eligible	
Children							
treatment	0.019^{*}	-0.002	0.370^{*}	-0.061	0.186^{*}	-0.024	
	(0.01)	(0.00)	(0.16)	(0.06)	(0.08)	(0.03)	
poor share	0.014	-0.012	0.203	-0.361	0.062	-0.175	
	(0.03)	(0.01)	(0.49)	(0.22)	(0.24)	(0.10)	
parents work	-0.027*	-0.021***	-0.460*	-0.509^{***}	-0.229^{*}	-0.226^{***}	
	(0.01)	(0.00)	(0.20)	(0.10)	(0.11)	(0.05)	
r2	0.113	0.044					
F	36.412	123.762					
Ν	3443.000	32420.000	3443.000	32420.000	3443.000	32420.000	
Young							
treatment	-0.024**	-0.004	-0.308**	-0.051	-0.160**	-0.027	
	(0.01)	(0.00)	(0.11)	(0.05)	(0.06)	(0.03)	
poor share	0.030	0.027^{*}	0.352	0.369*	0.181	0.175	
	(0.03)	(0.01)	(0.38)	(0.17)	(0.20)	(0.09)	
parents work	-0.017	-0.024***	-0.181	-0.289***	-0.099	-0.152^{***}	
	(0.01)	(0.01)	(0.15)	(0.07)	(0.08)	(0.04)	
r2	0.168	0.156	. ,	. ,	. ,	. ,	
F	73.966	370.971					
Ν	4421.000	24137.000	4421.000	24137.000	4421.000	24137.000	
status	Yes	Yes	Yes	Yes	Yes	Yes	
state effects	Yes	Yes	Yes	Yes	Yes	Yes	
year effects	Yes	Yes	Yes	Yes	Yes	Yes	

Table 14: Sickness Incidence

 $- \frac{1}{p} < 0.05, ** p < 0.01, *** p < 0.001$

	OLS		Zero-Inflated		
	Ineligible	Eligible	Ineligible	Eligible	
Baby					
treatment	-0.194	-0.036	-0.163*	-0.001	
	(0.15)	(0.04)	(0.07)	(0.03)	
poor snare	(0.50)	(0.14)	(0.157)	(0.10)	
inflate		. ,		. ,	
treatment			0.006	0.076	
			(0.10)	(0.04)	
poor share			1.006**	1.124***	
			(0.31)	(0.13)	
lnalpha			0 710***	0.005***	
cons			$-0.710^{+0.10}$	-0.995^{++++}	
			(0.10)	(0.00)	
N	2409	19912	2409	19912	
Children	0.677	0.000	0.010		
treatment	0.052	-0.001	0.243	-0.254	
poor chara	(0.03) 0.185	(0.01)	(0.42) 1 553	(0.26) 1 506	
poor snare	(0.10)	(0.04)	(1.11)	(0.85)	
inflato		. ,	. ,	. ,	
treatment			-0.522	-1.838	
			(0.36)	(1.09)	
poor share			-0.935	18.118	
			(1.00)	(10.45)	
lnalpha					
cons			-0.519	5.695^{***}	
			(0.48)	(0.11)	
Ν	3303	31834	3303	31834	
Young					
treatment	0.043	0.020	-0.613	0.545	
noor choro	(0.04)	(0.01)	(0.46)	(0.29)	
poor snare	(0.12)	(0.054)	(1.022)	-1.142	
parents work	-0.026	-0.013	1.110^{*}	-0.359	
parents work	(0.05)	(0.02)	(0.55)	(0.40)	
inflate					
treatment			-0.764	-0.071	
			(0.47)	(0.19)	
poor share			-1.407	0.814	
cons			(1.35) 6 477***	(0.60) 4 184***	
COIIS			(1.07)	(0.50)	
Inalpha			d	(/	
cons			u -0.996*	0.486	
			(0.44)	(0.28)	
N	4060.000	22154.000	4060.000	22154.000	

Table 15: Sickness Spell: Days reported as sick

* p < 0.05,** p < 0.01,**
*p < 0.001 Var.= number of days a baby was sick

	OLS		Zero-Inflated	
	Ineligible	Eligible	Ineligible	Eligible
Adult				
treatment	-0.037	0.001	-0.024	0.026
	(0.05)	(0.02)	(0.13)	(0.07)
poor share	0.045	0.159^{*}	0.052	0.686^{**}
	(0.15)	(0.07)	(0.40)	(0.26)
inflate				
treatment			0.141	0.068
			(0.11)	(0.06)
poor share			-0.017	-0.235
			(0.35)	(0.22)
Inalpha				
cons			0.179	0.217^{**}
			(0.13)	(0.08)
N	14781	55003	14781	55003
Old				
treatment	-0.333	-0.416*	-0.116	-0.064
	(0.35)	(0.20)	(0.14)	(0.07)
poor share	2.186	0.402	0.482	-0.049
	(1.17)	(0.73)	(0.45)	(0.29)
inflate				
treatment			0.107	0.255^{**}
			(0.16)	(0.09)
poor share			-0.454	-0.474
			(0.53)	(0.31)
lnalpha				
cons			-0.391**	-0.347***
			(0.14)	(0.08)
N	1725.000	5169.000	1725.000	5169.000
family's head work	Yes	Yes	Yes	Yes
status	Yes	Yes	Yes	Yes
state effects	Yes	Yes	Yes	Yes
year effects	Yes	Yes	Yes	Yes

Table 16: Sickness Spell: Days reported as sick