

A Service of

ZBW

Leibniz-Informationszentrum Wirtschaft Leibniz Information Centre for Economics

Briskorn, Dirk; Knust, Sigrid

## Working Paper — Digitized Version On Circular 2-Factorizations of the Complete Tripartite Graph

Manuskripte aus den Instituten für Betriebswirtschaftslehre der Universität Kiel, No. 636

**Provided in Cooperation with:** Christian-Albrechts-University of Kiel, Institute of Business Administration

*Suggested Citation:* Briskorn, Dirk; Knust, Sigrid (2008) : On Circular 2-Factorizations of the Complete Tripartite Graph, Manuskripte aus den Instituten für Betriebswirtschaftslehre der Universität Kiel, No. 636, Universität Kiel, Institut für Betriebswirtschaftslehre, Kiel

This Version is available at: https://hdl.handle.net/10419/201520

#### Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

#### Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.



# WWW.ECONSTOR.EU

# Manuskripte aus den Instituten für Betriebswirtschaftslehre der Universität Kiel

No. 636

### On Circular 2-Factorizations of the Complete Tripartite Graph

Dirk Briskorn, Sigrid Knust

April 2008

Dirk Briskorn Christian-Albrechts-Universität zu Kiel, Institut für Betriebswirtschaftslehre, Olshausenstr. 40, 24098 Kiel, Germany, http://www.bwl.uni-kiel.de/bwlinstitute/Prod briskorn@bwl.uni-kiel.de

Sigrid Knust Universität Osnabrück Fachbereich Mathematik/Informatik Albrechtstr. 28, 49069 Osnabrück, Germany http://www.informatik.uni-osnabrueck.de/knust/ sigrid@informatik.uni-osnabrueck.de

#### Abstract

We consider 2-factorizations of the complete tripartite graph K(n;3) where each 2-factor consists of cycles with even length. An additional requirement is related to pairs of edges being consecutive in an arbitrary cycle. For all pairs of edges we require that the three nodes incident to these two edges must be from different partite sets of the tripartite graph. We show that such a 2-factorizations cannot exist if n = 2 or n is odd. Furthermore, we show how to construct such a 2-factorization for all even n with  $n \notin \{2, 26, 34, 58, 74\}$ .

Keywords: Graph decomposition; complete tripartite graph; 2-factorization

### **1** Introduction

Decompositions of graphs have received a great deal of attention. In particular, this is true for edge-disjoint decompositions of complete multipartite graphs into cycles, see Lindner and Rodger [9] for a survey.

There are several papers considering special cycle structures. Billington [1] studies decompositions of complete tripartite graphs into cycles of lengths 3 and 4 while Cavenagh and Billington [4] consider decompositions of complete multipartite graphs into cycles with even length. Cavenagh [3] shows that each complete tripartite graph G can be partitioned into cycles of length k if the number of edges of G is a multiple of k. More generally, Billington and Cavenagh [2] prove that the edges of each complete tripartite graph can be partitioned into cycles of arbitrary lengths. Hilton and Rodger [5], Laskar [6], Laskar and Auerbach [7], and Leach and Rodger [8] investigate decompositions of complete multipartite graphs into Hamilton cycles.

In this paper we study special structures of 2-factorizations of the complete tripartite graph where all three partite sets have identical size. The graph having 3n nodes is denoted by K(n;3) in the following. The three subsets of nodes forming the partite sets of the tripartite graph are represented by  $V_0$ ,  $V_1$ , and  $V_2$ . Recall that a 2-factor of a graph G = (V, E) is a subset E' of edges such that each node  $v \in V$  is incident to exactly two edges  $e_0, e_1 \in E'$ ,  $e_0 \neq e_1$ . A 2-factorization of a graph is a partition of the edges into edge-disjoint 2-factors. Note that each 2-factor consists of a collection of node-disjoint cycles. We consider 2-factors meeting two additional requirements:

- (i) Each cycle contained in the 2-factor must have even length.
- (ii) In the 2-factor each node  $i \in V_s$ ,  $s \in \{0, 1, 2\}$ , must be adjacent to exactly one node  $j \in V_{(s-1) \mod 3}$  and to exactly one node  $j' \in V_{(s+1) \mod 3}$ .

A 2-factor fulfilling (i) and (ii) is referred to as circular even 2-factor (ce2f) below. Accordingly, a circular even 2-factorization (ce2F) of a graph G is defined as a decomposition of E into edge-disjoint ce2fs.

In the following section we study for which numbers  $n \in \mathbb{N}$  a ce2F of K(n;3) exists. We answer this question for all n with  $n \notin \{26, 34, 58, 74\}$ .

### 2 Results

It is easy to see that no ce2F of K(n; 3) exists if n is odd. Since then the number of nodes 3n is odd, there must be at least one cycle having odd length in each 2-factor. Therefore, we restrict ourselves to n being even in the following.

#### 2.1 Case n = 4k with $k \in \mathbb{N}$

In this subsection we provide a construction scheme for a ce2F of the graph K(n; 3) with n = 4k for all  $k \in \mathbb{N}$ . Note that the subgraph induced by two of the three partite sets is the complete bipartite graph  $K_{n,n}$  with 2n nodes. It is well-known that for this graph 1-factorizations exist, e.g. the bipartite 1-factorization  $F^{bip} = \left\{ F_0^{bip}, \ldots, F_{n-1}^{bip} \right\}$  with

$$F_l^{bip} = \{[m, n + (m+l) \mod n] \mid m = 0, \dots, n-1\}$$
 for  $l = 0, \dots, n-1$ .

The basic idea of our construction scheme is to combine certain 1-factors of  $F^{bip}$  between pairs of partite sets in each ce2f. First, we adress the question how to combine 1-factors of  $F^{bip}$  in order to obtain a ce2f of K(n; 3). For two different partite sets  $V_s, V_{s'}$  and  $l \in \{0, \ldots, n-1\}$ we denote by  $F_l^{bip}(s, s')$  the 1-factor  $F_l^{bip}$  between the partite sets  $V_s$  and  $V_{s'}$ .

**Lemma 1.** Three 1-factors  $F_{l_0}^{bip}(0,1)$ ,  $F_{l_1}^{bip}(1,2)$ , and  $F_{l_2}^{bip}(2,0)$  with  $l_0, l_1, l_2 \in \{0, \ldots, n-1\}$  form a ce2f of K(n;3) if  $l_0 + l_1 + l_2$  is odd.

Proof. Obviously,  $F_{l_0}^{bip}(0,1)$ ,  $F_{l_1}^{bip}(1,2)$ , and  $F_{l_2}^{bip}(2,0)$  form together a 2-factor of K(n;3) since exactly two 1-factors are chosen for each group. Moreover, the 2-factor fulfills condition (ii), i.e. it is circular. Let us consider an arbitrary path from a node  $i \in V_s$  to another node  $i' \in V_s$  induced by  $F_{l_0}^{bip}(0,1)$ ,  $F_{l_1}^{bip}(1,2)$ , and  $F_{l_2}^{bip}(2,0)$ . Obviously, the length of this path is a multiple of 3. Now consider an arbitrary path having length 3. Note that there are exactly two such paths for each  $i \in V_s$  and both connect i with another node  $i' \in V_s$ . Let i and i' be the start and end node of such a path. Since n is even and  $l_0 + l_1 + l_2$  is odd, |i - i'| is odd, as well. This implies that an even number of paths of length 3 must be concatenated to construct a circle and, hence, circles have even length.

In the following we will show that a ce2F of the graph K(n;3) with n = 4k for all  $k \in \mathbb{N}$  exists. According to Lemma 1 combining 1-factors  $F_{l_0}^{bip}(0,1)$ ,  $F_{l_1}^{bip}(1,2)$ , and  $F_{l_2}^{bip}(2,0)$  with indices  $l_0, l_1, l_2 \in \{0, \ldots, n-1\}$  yields a ce2f if

- exactly one index is odd, and the other two are even, or
- all three indices are odd.

According to Lemma 1 the specific combination of 1-factors can be chosen arbitrarily. However, we focus on a specific arrangement in the following.

At first we combine the odd 1-factors  $F_l^{bip}$  for  $l = 4\lambda + 1$  ( $\lambda = 0, \ldots, \frac{n}{4} - 1$ ) with the two even 1-factors  $F_{l-1}^{bip}$  and  $F_{l+1}^{bip}$ . As an example Figure 1 illustrates the case l = 1 for the graph K(4;3), where the 1-factors  $F_0^{bip}(0,1)$ ,  $F_1^{bip}(1,2)$  and  $F_2^{bip}(2,0)$  are shown. Here, the node on top of the column representing a part has the smallest index. We apply each combination three times by rotating the specific 1-factors such that each 1-factor is applied to each pair of partite sets exactly once. This means that each combination is arranged in three ce2fs:  $F_{l-1}^{bip}(0,1)$ ,  $F_l^{bip}(1,2)$ ,  $F_{l+1}^{bip}(2,0)$  in the first ce2f,  $F_{l-1}^{bip}(1,2)$ ,  $F_l^{bip}(2,0)$ ,  $F_{l+1}^{bip}(0,1)$  in the second ce2f, and  $F_{l-1}^{bip}(2,0)$ ,  $F_l^{bip}(0,1)$ ,  $F_{l+1}^{bip}(1,2)$  in the third ce2f. By arranging this configuration for all  $\lambda = 0, \ldots, \frac{n}{4} - 1$  we arrange edges corresponding to  $\frac{3n}{4}$  1-factors between each pair of partite sets in  $\frac{3n}{4}$  ce2fs. Note that we cover  $F_l^{bip}$  for each even index  $l \in \{0, \ldots, n-1\}$ .

It remains to arrange edges according to  $F_l^{bip}$  for all odd  $l = 4\mu + 3$  ( $\mu = 0, ..., \frac{n}{4} - 1$ ) in the  $\frac{n}{4}$  remaining ce2fs. This can be done by arranging  $F_l^{bip}$  between each pair of partite sets in



Figure 1:  $F_0^{bip}(0,1)$ ,  $F_1^{bip}(1,2)$ , and  $F_2^{bip}(2,0)$  for K(4;3)



Figure 2:  $F_3^{bip}(0,1)$ ,  $F_3^{bip}(1,2)$ , and  $F_3^{bip}(2,0)$  for K(4;3)

a single ce2f, i.e. by choosing  $F_l^{bip}(0,1)$ ,  $F_l^{bip}(1,2)$ , and  $F_l^{bip}(2,0)$ . Clearly, the prerequisite of Lemma 1 is fulfilled and, hence, we obtain a ce2f. As an example Figure 2 illustrates the case l = 3 for the graph K(4;3), where the 1-factors  $F_3^{bip}(0,1)$ ,  $F_3^{bip}(1,2)$  and  $F_3^{bip}(2,0)$  are shown. The preceding construction implies

**Theorem 1.** For the graph K(n; 3) with n = 4k a ce2F exists for all  $k \in \mathbb{N}$ .

### **2.2** Case $n \notin \{26, 34, 58, 74\}$

In this subsection we provide a construction scheme for a ce2F of the graph K(n;3) for all even  $n \notin \{2, 26, 34, 58, 74\}$ . If n is a multiple of 4, the construction method from the previous subsection can be applied. By using an integer programming formulation it can be shown that for n = 2 no ce2F exists. On the other hand, using Cplex we found ce2Fs  $F^6$  and  $F^{10}$  for K(6;3) and K(10;3). These ce2Fs are given in the appendix.

At first we will show that from ce2Fs for a graph K(l;3) with  $l \in \mathbb{N}$  ce2Fs for the larger graphs K(kl;3) can be constructed for all  $k \in \mathbb{N}$ .

**Lemma 2.** If a ce2F of K(l;3) for some integer  $l \in \mathbb{N}$  exists, then also a ce2F of K(kl;3) for all  $k \in \mathbb{N}$  exists.

*Proof.* The basic idea of this constructive proof is to cluster nodes and to use the structure known for the graph induced by the cluster. This technique has been applied in several works concerning decomposition of graphs, see e.g. Laskar [6] or Laskar and Auerbach [7]. Suppose we have a ce2F F of K(l; 3) for some  $l \in \mathbb{N}$ . Then we construct a ce2F F' of K(kl; 3) for all  $k \in \mathbb{N}$  as follows:

- 1. In the graph K(kl; 3) partition the nodes of each partite set  $V_s$ ,  $s \in \{0, 1, 2\}$ , into l subparts  $V_{s,m}$  for m = 0, ..., l 1, having size k each.
- Consider K(l; 3) as the complete tripartite graph induced by the 3l subparts. Each 2-factor in the 2-factorization F of K(l; 3) is expanded to k 2-factors in the 2-factorization F' of K(kl; 3). Note that for each pair of subparts (belonging to different partite sets) all edges between the nodes in these subparts can be represented by the complete bipartite graph K<sub>k,k</sub>. We may employ F<sup>bip</sup> to decompose them into k 1-factors. For this purpose we assign an arbitrary orientation to each circle from the 2-factors in F. Then, we combine the 1-factors F<sup>bip</sup><sub>λ</sub> for λ = 0,..., k 1 between each pair of subparts being consecutive in a circle.

It is easy to see that F' is a 2-factorization of K(kl; 3). Moreover, we have a circular structure due to the circular structure of F. It is obvious that circles of even length are constructed since each underlying circle in F is of even length q and circles in F' can only have lengths being multiples of q.

Thus, according to Lemma 2, we can construct ce2Fs of K(6k; 3) and K(10k; 3) for all  $k \in \mathbb{N}$  using the ce2Fs  $F^6$  and  $F^{10}$  as building blocks.

In a similar way the following lemma provides a basis for a more powerful recursive construction.

**Lemma 3.** If a ce2F of K(l; 3) for some  $l \in \mathbb{N}$  exists, then also a ce2F of K(kl + 4m; 3) for each pair of integers k, m with  $4m \ge kl \ge 3m$  exists.

*Proof.* The basic idea of this constructive proof is to divide the nodes into two parts inducing subgraphs  $G^{kl} := K(kl; 3)$  and  $G^{4m} := K(4m; 3)$ . Suppose we have a ce2F F of K(l; 3) for some  $l \in \mathbb{N}$ . Then we construct a ce2F of K(kl + 4m; 3) using structural elements of F and known ce2Fs of  $G^{4m}$  (cf. Subsection 2.1) as follows:

- 1. According to Lemma 2 from F we can construct a ce2F F' of  $G^{kl}$  for all  $k \in \mathbb{N}$ . We arrange F' of  $G^{kl}$  as part of the first kl ce2fs of K(kl + 4m; 3).
- 2. We complete the first kl ce2fs of K(kl+4m; 3) by ce2fs of  $G^{4m}$ . These ce2fs of  $G^{4m}$  are arranged by combining 1-factors according to  $F^{bip}$  of  $K_{4m,4m}$  as in the construction in Section 2.1. More specifically, we choose all 2m even 1-factors  $F_{2\lambda}^{bip}$  for  $\lambda = 0, \ldots, 2m-1$  and the kl-2m odd 1-factors  $F_{4m-kl+1+2\lambda}^{bip}$  for  $\lambda = 0, \ldots, kl-2m-1$ . Since  $kl \geq 3m$ , the number of used odd 1-factors is at least m and, therefore, at least half the number of used even 1-factors. Regarding Lemma 1, we can arrange 3m ce2fs of  $G^{4m}$  by combining two even 1-factors with one odd 1-factor in each ce2f. Furthermore, we can arrange kl 3m ce2fs of  $G^{4m}$  using the remaining odd 1-factors (only a single 1-factor in each ce2f, cf. Subsection 2.1).

Up to now, we have kl complete ce2fs of K(kl + 4m; 3). It remains to arrange all edges of  $G^{4m}$  according to  $F_{2\lambda+1}^{bip}$  and  $F_{4m-1-2\lambda}^{bip}$  for all  $\lambda = 0, \ldots, \frac{4m-kl}{2} - 1$  (these edges have been left out in Step 2) and all edges between  $G^{kl}$  and  $G^{4m}$  in the remaining 4m ce2fs of K(kl+4m; 3). In the following we represent node i by (s(i), i') where s(i) denotes the partite set of i and  $i' := i \mod n$  is the index of i within its partite set. W.l.o.g. we assume that the nodes of  $G^{4m}$  are numbered such that they satisfy  $0 \le i' \le 4m - 1$ .

3. For  $\lambda = 0, ..., 4m-1$  in the ce2f with index  $kl + \lambda$  for q = 0, 1, 2 and i' = 4m, ..., kl + 4m - 1 we arrange edges

$$[(q, i'), ((q+1) \mod 3, (i'-4m+\lambda) \mod 4m)] \text{ and } \\ [(q, i'), ((q-1) \mod 3, (i'-4m+\lambda) \mod 4m)].$$

In Step 3 we exclusively arrange circles of length 6. In the ce2f with index  $\mu \in \{kl, \ldots, kl + 4m-1\}$  of K(kl+4m; 3) the nodes  $(q, (\mu+i') \mod 4m)$  for q = 0, 1, 2 and  $i' = 0, \ldots, 4m-kl-1$  are not involved yet.

4. For  $\mu = kl, \ldots, kl + 4m - 1$  in the ce2f with index  $\mu$  for q = 0, 1, 2 and  $i' = 0, \ldots, 4m - kl - 1$  we arrange edges

In Step 4 we, again, exclusively arrange circles of length 6. Edges originating from Step 4 form ce2fs corresponding to  $F_{2\lambda+1}^{bip}$  and  $F_{4m-1-2\lambda}^{bip}$  for  $\lambda = 0, \ldots, \frac{4m-kl}{2} - 1$ . Summarizing, we obtain a ce2F of K(kl + 4m; 3).

In the following we want to apply Lemma 3 with k = 1. Thus, we study which even numbers  $n \in \mathbb{N}$  can be written in the form n = l + 4m with  $l, m \in \mathbb{N}$  and  $4m \ge l \ge 3m$ . We get

$$(l,m) = \begin{cases} \left(\frac{n}{2}, \frac{n}{8}\right) & \text{for } n \mod 8 = 0, \quad n \ge 8, \\ \left(\frac{n-6}{2}, \frac{n+6}{8}\right) & \text{for } n \mod 8 = 2, \quad n \ge 42, \\ \left(\frac{n-4}{2}, \frac{n+4}{8}\right) & \text{for } n \mod 8 = 4, \quad n \ge 28, \\ \left(\frac{n-2}{2}, \frac{n+2}{8}\right) & \text{for } n \mod 8 = 6, \quad n \ge 14. \end{cases}$$
(1)

Thus, each even number  $n \ge 36$  can be written in the form n = l + 4m with  $l, m \in \mathbb{N}$  and  $4m \ge l \ge 3m$ .

n	existence	argument
2	no	IP
6	yes	IP
10	yes	IP
14	yes	Lemma 3 ( $k = 1, l = 6, m = 2$ )
18	yes	Lemma 2 ( $k = 3, l = 6$ )
22	yes	Lemma 3 $(k = 1, l = 10, m = 3)$
26	?	
30	yes	Lemma 2 ( $k = 5, l = 6$ )
34	?	

Table 1: Existence results for a ce2F of K(n; 3)

**Theorem 2.** For the graph K(n; 3) a ce2F exists for all even  $n \notin \{2, 26, 34, 58, 74\}$ .

l	$D_m$	m	l+4m	argument
26	{7,8}	7	54	Lemma 2 $(k = 9, l = 6)$
		8	58	
34	{9,10,11}	9	70	Lemma 3 ( $k = 1, l = 30, m = 10$ )
		10	74	
		11	78	Lemma 2 ( $k = 13, l = 6$ )
58	{15,16,17,18,19}	15	118	Lemma 3 ( $k = 1, l = 54, m = 16$ )
		16	122	Lemma 3 $(k = 1, l = 54, m = 17)$
		17	126	Lemma 3 ( $k = 1, l = 54, m = 18$ )
		18	130	Lemma 3 ( $k = 1, l = 62, m = 17$ )
		19	134	Lemma 3 ( $k = 1, l = 62, m = 18$ )
74	{19,20,21,22,23,24}	19	150	Lemma 3 ( $k = 1, l = 70, m = 20$ )
		20	154	Lemma 3 ( $k = 1, l = 70, m = 21$ )
		21	158	Lemma 3 ( $k = 1, l = 70, m = 22$ )
		22	162	Lemma 3 ( $k = 1, l = 70, m = 23$ )
		23	166	Lemma 3 ( $k = 1, l = 78, m = 22$ )
		24	170	Lemma 3 ( $k = 1, l = 78, m = 23$ )

Table 2: Numbers l + 4m depending on  $l \in \{26, 34, 58, 74\}$ 

*Proof.* Due to Theorem 1 we already know a ce2F of K(n; 3) for all n which are a multiple of 4. For all other even n < 36 in Table 1 we summarize the results whether a ce2F is known for the graph K(n; 3) or not. Only for  $n \in \{2, 26, 34\}$  no ce2F is known. Note that for n = 26, 34 no numbers k, l, m exist which satisfy the conditions of Lemma 3.

Due to (1) each even  $n \ge 36$  can be represented as n = l + 4m with  $4m \ge l \ge 3m$ . Thus, if a ce2F for the specific graph K(l;3) is known, according to Lemma 3 also a ce2F of K(n;3) exists.

This construction covers all values n which do not depend on a (direct or indirect) representation with l = 26 or l = 34. The first part of Table 2 lists all numbers which directly depend on l = 26 or l = 34. The second column gives the feasible domain  $D_m$  consisting of all values  $m \in \mathbb{N}$  with  $4m \ge l \ge 3m$  for the specific value of l. For each element of  $D_m$  the third column contains the value l + 4m. If another argument not based on the specific value of lfor the existence of a ce2F of K(l + 4m; 3) is known, it is stated in the fourth column.

The only numbers which directly depend on  $l_1 = 26$  or  $l_2 = 34$  and for which no other argument is known are  $l_1 + 4 \cdot 8 = 58$  and  $l_2 + 4 \cdot 10 = 74$ . These numbers are treated in the same way in the second part of Table 2. It can be seen that all numbers which directly depend on  $l_3 = 58$  or  $l_4 = 74$  can be covered by other arguments. Thus, our construction only fails for all even  $n \in \{2, 26, 34, 58, 74\}$ . While for n = 2 it is proven that no ce2F exists, the other four cases remain open.

We can conclude that a ce2F of the graph K(n; 3) for all even  $n \notin \{2, 26, 34, 58, 74\}$  does exist. For n = 2 or n odd no ce2F can exist. Cases  $n \in \{26, 34, 58, 74\}$  remain open so far. However, we strongly conjecture ce2Fs to exist for these cases.

### References

- E. J. Billington. Decomposing complete tripartite graphs into cycles of lengths 3 and 4. Discrete Mathematics, 197/198:123–135, 1999.
- [2] E. J. Billington and N. J. Cavenagh. Decomposing Complete Tripartite Graphs Into Closed Trails Of Arbitrary Lenghts. *Czechoslovak Mathematical Journal*, 57(132):523–551, 2007.
- [3] N. J. Cavenagh. Decompositions of Complete Tripartite Graphs into k-cycles. Australasian Journal of Combinatorics, 18:193–200, 1998.
- [4] N. J. Cavenagh and E. J. Billington. Decompositions of Complete Multipartite Graphs into Cycles of Even Length. Graphs and Combinatorics, 16:49-65, 2000.
- [5] A. J. W. Hilton and C. A. Rodger. Hamilton decompositions of complete regular s-partite graphs. Discrete Mathematics, 58:63–78, 1986.
- [6] R. Laskar. Decomposition of some composite graphs into Hamiltonian cycles. In A. Hajnal and V. Sos, editors, *Proceedings of the Fifth Hungarian Coll. Keszthely 1976*, pages 705– 716. North-Holland, Amsterdam, 1978.
- [7] R. Laskar and B. Auerbach. On the decomposition of *r*-partite graphs into edge-disjoint hamilton circuits. *Discrete Mathematics*, 14:265–268, 1976.
- [8] C. D. Leach and C. A. Rodger. Fair Hamilton Decompositions of Complete Multipartite Graphs. *Journal of Combinatorial Theory, Series B*, 85:290–296, 2002.
- [9] C. Lindner and C. Rodger. Decomposition into cycles II: cycle systems. In J. Dinitz and D. Stinson, editors, *Contemporary Design Theory: A Collection of surveys*, pages 325–369. Wiley, New York, 1992.

# A Ce2Fs of K(6; 3) and K(10; 3)

In the following we represent the nodes  $i = 0, \ldots, n-1$  of the graph  $K(\frac{n}{3}; 3)$  by pairs (s(i), i') where  $i' = i \mod \frac{n}{3}$ . In the following a ce2F  $F^6 = \{F_f^6 \mid 0 \le f \le 5\}$  of K(6; 3) and a ce2F  $F^{10} = \{F_f^{10} \mid 0 \le f \le 9\}$  of K(10; 3) are shown. Each ce2f is given as collection of circles.

$$\begin{array}{rcl} F_{0}^{6} &=& \{ [(0,0),(1,3),(2,0),(0,3),(1,0),(2,3)], \\ && [(0,1),(2,4),(1,1),(0,4),(2,1),(1,4)], \\ && [(0,2),(1,5),(2,2),(0,5),(1,2),(2,5)] \} \\ F_{1}^{6} &=& \{ [(0,0),(1,1),(2,2),(0,3),(1,4),(2,5)], \\ && [(0,1),(1,2),(2,3),(0,4),(1,5),(2,0)], \\ && [(0,2),(1,3),(2,4),(0,5),(1,0),(2,1)] \} \\ F_{2}^{6} &=& \{ [(0,0),(2,2),(1,4),(0,2),(2,0),(1,2), \\ && (0,3),(2,1),(1,1),(0,5),(2,3),(1,3), \\ && (0,4),(2,4),(1,5),(0,1),(2,5),(1,0)], \} \\ F_{3}^{6} &=& \{ [(0,0),(1,4),(2,0),(0,4),(1,2),(2,1)], \\ && [(0,1),(2,2),(1,3),(0,5),(2,5),(1,5), \\ && (0,3),(2,3),(1,1),(0,2),(2,4),(1,0)] \} \\ F_{4}^{6} &=& \{ [(0,0),(2,0),(1,1),(0,3),(2,4),(1,2), \\ && (0,1),(2,3),(1,4),(0,5),(2,1),(1,5)] \} \\ F_{6}^{6} &=& \{ [(0,0),(1,2),(2,2),(0,4),(1,4),(2,4)], \end{array}$$

$$F_5 = \{[(0,0), (1,2), (2,2), (0,4), (1,4), (2,4)], \\[(0,1), (1,1), (2,5), (0,3), (1,3), (2,1)], \\[(0,2), (2,3), (1,5), (0,5), (2,0), (1,0)]\}$$

$$\begin{split} F_{1}^{10} &= \left\{ [(0,0),(1,7),(2,3),(0,4),(1,3),(2,1),(0,9),(1,9),(2,4),(0,3),(1,8),(2,0)], \\ &= ((0,1),(1,6),(2,7),(0,7),(1,0),(2,8),(0,5),(1,4),(2,6),(0,8),(1,5),(2,2), \\ &= (0,2),(1,2),(2,9),(0,6),(1,1),(2,5)] \right\} \\ F_{1}^{10} &= \left\{ [(0,0),(1,3),(2,0),(0,6),(1,5),(2,5),(0,4),(1,2),(2,4),(0,8), \\ &= (1,0),(2,6),(0,7),(1,7),(2,9),(0,5),(1,9),(2,3),(0,1),(1,4), \\ &= (2,7),(0,3),(1,1),(2,1),(0,2),(1,8),(2,2),(0,9),(1,6),(2,8)] \right\} \\ F_{2}^{10} &= \left\{ [(0,0),(1,8),(2,7),(0,9),(1,1),(2,3)],[(0,3),(1,2),(2,1),(0,6),(1,7),(2,5)], \\ &= ((0,0),(1,8),(2,7),(0,9),(1,1),(2,3)],(0,3),(1,2),(2,1),(0,6),(1,7),(2,5)], \\ &= ((0,0),(1,6),(2,9),(0,2),(1,5),(2,9),(0,7),(1,4),(2,0),(0,8),(1,9),(2,6)] \right\} \\ F_{3}^{10} &= \left\{ [(0,0),(1,6),(2,9),(0,2),(1,0),(2,0),(0,5),(1,2),(2,7),(0,4), \\ &= (1,9),(2,1),(0,3),(1,3),(2,5),(0,7),(1,1),(2,2),(0,6),(1,4), \\ &= (2,4),(0,1),(1,7),(2,8),(0,9),(1,5),(2,3),(0,8),(1,8),(2,6)] \right\} \\ F_{4}^{10} &= \left\{ [(0,0),(1,0),(2,7),(0,1),(1,9),(2,5),(0,6),(1,8),(2,1),(0,8), \\ &= (1,6),(2,3),(0,5),(1,5),(2,8),(0,4),(1,1),(2,4),(0,2),(1,7), \\ &= (2,0),(0,7),(1,3),(2,2)],[(0,3),(1,4),(2,9),(0,3),(1,2),(2,6)] \right\} \\ F_{5}^{10} &= \left\{ [(0,0),(1,2),(2,2),(0,3),(1,5),(2,0),(0,9),(1,4),(2,8),(0,1), \\ &= (1,6),(2,3),(0,5),(1,5),(2,8),(0,4),(1,1),(2,4),(0,2),(1,7), \\ &= (2,0),(0,7),(1,3),(2,2)],[(0,3),(1,4),(2,9),(0,3),(1,2),(2,6)] \right\} \\ F_{6}^{10} &= \left\{ [(0,0),(1,1),(2,2),(0,3),(1,5),(2,0),(0,9),(1,4),(2,8),(0,1), \\ &= (1,9),(2,2),(0,8),(1,2),(2,0),(0,2),(1,1),(2,7),(0,5),(1,0),(2,3)] \right\} \\ F_{7}^{10} &= \left\{ [(0,0),(1,1),(2,3),(0,3),(1,0),(2,3),(0,7),(1,6),(2,4)] \right\} \\ F_{8}^{10} &= \left\{ [(0,0),(1,4),(2,5),(0,8),(1,3),(2,3),(0,7),(1,6),(2,4)] \right\} \\ F_{8}^{10} &= \left\{ [(0,0),(1,4),(2,5),(0,8),(1,3),(2,4),(0,9),(1,7),(2,7),(0,2), \\ &= (1,9),(2,0),(0,1),(1,2),(2,8),(0,3),(1,7),(2,2),(0,3),(1,9),(2,7)] \right\} \\ F_{9}^{10} &= \left\{ [(0,0),(1,4),(2,5),(0,8),(1,3),(2,4),(0,9),(1,7),(2,7),(0,2), \\ &= (1,9),(2,0),(0,1),(1,2),(2,3),(0,3),(1,6),(2,1),(0,4),(1,8), \\ &= (2,8),(0,7),(1,5),(2,6),(0,6),(1,0),(2,2),(0,5),(1,1),(2,9)] \right\} \\ F_{9}^{10} &$$

# Manuskripte aus den Instituten für Betriebswirtschaftslehre der Universität Kiel

- 636 Briskorn, D., Knust, S., On circular 2-factorizations of the complete tripartite graph (2008)
- 635 Briskorn, D., Knust, S., Constructing fair sports league schedules (2008)
- 634 Büther, M., Beam search for the elastic generalized assignment problem (2008)
- 633 o.V., Jahresbericht 2007
- 632 Büther, M., Reducing the elastic generalized assignment problem to the standard generalized assignment problem (2007)
- 631 Veit, K.-R., Aktienrechtliches und übernahmerechtliches Squeeze out im Vergleich (2007)
- 630 Veit, K.-R., Die Verlustanzeige gemäß § 92 Abs. 1 AktG bzw. § 49 Abs. 3 GmbHG aus der Sicht des Abschlussprüfers (2007)
- 629 Büther, M., Briskorn, D., Reducing the 0-1 knapsack problem with a single continuous variable to the standard 0-1 knapsack problem (2007)
- 628 Friedl, B., Wertanalyse nach DIN EN 12973 als Instrument des produktorientierten Kostenmanagements (2007)
- 627 Drexl, A., Jørnsten, K., Pricing the generalized assignment problem (2007)
- 626 Drexl, A., Jørnsten, K., Pricing the multiple-choice nested knapsack problem (2007)
- 625 Drexl, A., Jørnsten, K., Knof, D., Non-linear anonymous pricing combinatorial auctions (2007)
- 624 Drexl, A., Jørnsten, K., Knof, D., Column aggregation-based pricing combinatorial auctions (2007)
- 623 Briskorn, D., Drexl, A., A branching scheme based on first-break-then-schedule decomposition (2007)
- 622 Rohrlack, C., Wolf, J., Koordination von länderübergreifenden Technologietransfers innerhalb multinationaler Unternehmen (2007)
- 621 Himme, A., Eine empirische Untersuchung zu Erfolgsfaktoren des Kostenmanagements – Befragungsergebnisse und abgeleitete Managementempfehlungen – (2007)
- 620 Meincke, S., Nippel, P., Der optimale Bezugspreis bei Kapitalerhöhungen mit Bezugsrechten (2007)
- 619 Nekat, K., Nippel, P., The impact of a firm's payout policy on stock prices and shareholders' wealth in an inefficient market (2007)
- 618 Himme, A., Eine kritische Analyse bestehender empirischer Arbeiten zum Kostenmanagement (2007)
- 617 Göthlich, S.E., Zur Ideengeschichte von Buchführung, Kostenrechnung und Controlling vor dem Hintergrund ihres historischen Kontexts (2007)