

Singh, Housila P.; Mehta, Vishal

Article

Improved Estimation of the Scale Parameter for Log-Logistic Distribution Using Balanced Ranked Set Sampling

Statistics in Transition New Series

Provided in Cooperation with:

Polish Statistical Association

Suggested Citation: Singh, Housila P.; Mehta, Vishal (2017) : Improved Estimation of the Scale Parameter for Log-Logistic Distribution Using Balanced Ranked Set Sampling, Statistics in Transition New Series, ISSN 2450-0291, Exeley, New York, NY, Vol. 18, Iss. 1, pp. 53-74, <https://doi.org/10.21307/stattrans-2016-057>

This Version is available at:

<https://hdl.handle.net/10419/207845>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.



<https://creativecommons.org/licenses/by/4.0/>

IMPROVED ESTIMATION OF THE SCALE PARAMETER FOR LOG-LOGISTIC DISTRIBUTION USING BALANCED RANKED SET SAMPLING

Housila P. Singh¹, Vishal Mehta²

ABSTRACT

In this article we have suggested some improved estimators of a scale parameter of log-logistic distribution (LLD) under a situation where the units in a sample can be ordered by judgement method without any error. We have also suggested some linear shrinkage estimator of a scale parameter of LLD. Efficiency comparisons are also made in this work.

Key words: minimum mean squared error estimator, shrinkage estimator, log-logistic distribution, best linear unbiased estimator, median ranked set sample.

AMS Subject Classification: 62G30; 62H12.

1. Introduction

Ranked set sampling (RSS) is a method of sampling that can be advantageous when quantification of all sampling units is costly but a small set of units can be easily ranked, according to the character under investigation, without actual quantification. The technique was first introduced by McIntyre (1952) for estimating means pasture and forage yields. The theory and application of ranked set sampling given by Chen *et al.* (2004).

A random variable X is said to have a log-logistic distribution with the scale parameter α and the shape parameter β if its cumulative distribution function (CDF) and probability density function (PDF) are respectively given as (see, Lesitha and Thomas (2012))

$$F(x; \alpha, \beta) = \frac{x^\beta}{\alpha^\beta + x^\beta}, x > 0, \alpha > 0, \beta > 1 \quad (1)$$

and

¹ School of Studies in Statistics, Vikram University, Ujjain - 456010, Madhya Pradesh, India.
E-mail: hpsujn@gmail.com.

² Indian Statistical Institute (ISI), North-East Centre, Tezpur - 784028, Assam, India.
E-mail: visdewas@gmail.com.

$$f(x; \alpha, \beta) = \frac{\beta \alpha^\beta x^{\beta-1}}{(\alpha^\beta + x^\beta)^2}, x > 0, \alpha > 0, \beta > 1. \quad (2)$$

Also, the k th moment of (2) exists only when $k < \beta$ and is given by

$$E(X^k) = \alpha^k B\left(1 - \frac{k}{\beta}, 1 + \frac{k}{\beta}\right), \quad (3)$$

where B denotes beta function.

The applications of log-logistic distribution are well known in a survival analysis of data sets such as survival times of cancer patients in which the hazard rate increases initially and decreases later (for example, see Bennett (1983)). In economic studies of distributions of wealth or income, it is known as Fisk distribution (see Fisk (1961)) and is considered as an equivalent alternative to a lognormal distribution. For further details on the importance and applications of a log-logistic distribution one may refer to Shoukri *et al.* (1988), Geskus (2001), Robson and Reed (1999) and Ahmad *et al.* (1988). For current reference in this context the reader is referred to Singh and Mehta (2013; 2014, a, b, 2015, 2016 a, b, c), Mehta and Singh (2014) and Mehta (2015).

If $X_{1:n}, X_{2:n}, \dots, X_{n:n}$ are the order statistics of a random sample of size n drawn from (1) then

$$Y_{r:n} = \frac{X_{r:n}}{\alpha}, r = 1, 2, \dots, n, \quad (4)$$

are distributed as order statistics of the same sample size drawn from a $LLD(1, \beta)$ with PDF given by

$$g(y, \beta) = \frac{\beta y^{\beta-1}}{(1 + y^\beta)^2}, y > 0, \beta > 1. \quad (5)$$

For a detailed description of various properties of order statistics arising from a $LLD(1, \beta)$ one may refer to Ragab and Green (1984). Balakrishnan and Malik (1987) have given some recurrence relations on the single and product moments of order statistics arising from a $LLD(1, \beta)$. Suppose

$$\gamma_{r:n} = E(Y_{r:n}), r = 1, 2, \dots, n, \quad (6)$$

$$\sigma_{r,s:n} = Cov(Y_{r:n}, Y_{s:n}), 1 \leq r < s \leq n \quad (7)$$

and

$$\sigma_{r,r:n} = Var(Y_{r:n}), 1 \leq r \leq n. \quad (8)$$

By using (4) in (6)-(8) we have

$$E(X_{r:n}) = \alpha \gamma_{r:n}, 1 \leq r \leq n \tag{9}$$

$$\text{Cov}(X_{r:n}, X_{s:n}) = \alpha^2 \sigma_{r,s:n}, 1 \leq r < s \leq n \tag{10}$$

$$\text{Var}(X_{r:n}) = \alpha^2 \sigma_{r,r:n}, 1 \leq r \leq n. \tag{11}$$

Lesitha and Thomas (2012) have computed the values of $\gamma_{r:n}$ and $\sigma_{r,s:n}, 1 \leq r, s \leq n$ independently for $n = 2(1)8$ and for $\beta = 2.5(0.5)5.0$ using Mathcad software so as to use those values for the computation of BLUE of α based on order statistics. If $\mathbf{X} = (X_{1:n}, X_{2:n}, \dots, X_{n:n})'$ then the mean vector $E(\mathbf{X})$ and dispersion matrix $D(\mathbf{X})$ of \mathbf{X} are

$$E(\mathbf{X}) = \gamma \alpha$$

and

$$D(\mathbf{X}) = \alpha^2 \mathbf{G},$$

where $\gamma = (\gamma_{1:n}, \gamma_{2:n}, \dots, \gamma_{n:n})'$ and $\mathbf{G} = ((\sigma_{r,s:n}))$.

Thus, by Gauss-Markov theorem Lesitha and Thomas (2012) gives the BLUE $\hat{\alpha}$ based on order statistics of a random sample of size n as:

$$t_1 = \hat{\alpha} = (\gamma' \mathbf{G}^{-1} \gamma)^{-1} \gamma' \mathbf{G}^{-1} \mathbf{X}$$

and

$$\text{Var}(t_1) = (\gamma' \mathbf{G}^{-1} \gamma)^{-1} \alpha^2 = \alpha^2 V_1, \tag{12}$$

where $V_1 = (\gamma' \mathbf{G}^{-1} \gamma)^{-1}$.

Lesitha and Thomas (2012) further estimate α based on the mean of unbiased estimators of α defined from each individual observations in the balanced ranked set sampling as:

$$t_2 = \alpha^* = \frac{1}{n} \sum_{r=1}^n \left[\frac{X_{(r:n)r}}{\gamma_{r:n}} \right],$$

with

$$\text{Var}(t_2) = \frac{1}{n^2} \sum_{r=1}^n \left[\frac{\sigma_{r,r:n}}{\gamma_{r:n}^2} \right] \alpha^2 = \alpha^2 V_2, \tag{13}$$

where $V_2 = \frac{1}{n^2} \sum_{r=1}^n \left[\frac{\sigma_{r,r:n}}{\gamma_{r:n}^2} \right]$.

Lesitha and Thomas (2012) also estimate α based on BLUE in the balanced ranked set sampling as:

$$t_3 = \alpha^{**} = (\gamma' G_1^{-1} \gamma)^{-1} \gamma' G_1^{-1} X_{rss}$$

and

$$\text{Var}(t_3) = (\gamma' G_1^{-1} \gamma)^{-1} \alpha^2 = \alpha^2 V_3. \quad (14)$$

where $\gamma = (\gamma_{1:n}, \gamma_{2:n}, \dots, \gamma_{n:n})'$, $G_1 = \text{diag}(\sigma_{1,1:n}, \sigma_{2,2:n}, \dots, \sigma_{n,n:n})$ and $V_3 = (\gamma' G_1^{-1} \gamma)^{-1}$.

When n is small the estimators α^* and α^{**} may not be acceptable for the expected level of precision. In such situations Lesitha and Thomas (2012) makes N cycles of RSS. For details see Chen *et al.* (2004). Suppose α_i^* and α_i^{**} denote the estimators of α corresponding to α^* and α^{**} respectively, based on the i th cycle. Then, estimators of α based on N cycles are given by:

$$\bar{\alpha}^* = \frac{1}{N} \sum_{i=1}^N \alpha_i^*,$$

and

$$\bar{\alpha}^{**} = \frac{1}{N} \sum_{i=1}^N \alpha_i^{**}$$

with

$$\text{Var}(\bar{\alpha}^*) = \frac{\alpha^2}{Nn^2} \sum_{r=1}^N \left[\frac{\sigma_{r,r:n}}{\gamma_{r:n}^2} \right],$$

and

$$\text{Var}(\bar{\alpha}^{**}) = \frac{(\gamma' G_1^{-1} \gamma)^{-1} \alpha^2}{N}.$$

Median ranked set sampling (MRSS) was first introduced by Muttlak (1997) to estimate the mean of a normal distribution. In general, MRSS is applied as a modification of RSS when one is interested in estimating a parameter associated with the central tendency of a distribution. The procedures of MRSS are given as: Select n independent samples each with n units as in the case of RSS. Then rank the units in each sample either by judgement method or by using some inexpensive means without having actual measurement on the unit. Lesitha and Thomas (2012) used MRSS method to estimate α as:

$$t_4 = \tilde{\alpha} = \begin{cases} \frac{1}{n} \sum_{r=1}^n \left[\frac{X_{(m:n)r}}{\gamma_{m:n}} \right] & ; \text{when } m = \frac{n+1}{2} \text{ and } n \text{ is odd} \\ (\gamma_1' G_2^{-1} \gamma_1)^{-1} \gamma_1' G_2^{-1} X_{mrss} & ; \text{when } m = \frac{n}{2} \text{ and } n \text{ is even} \end{cases}$$

with

$$Var(t_4) = \begin{cases} \frac{1}{n} \left[\frac{\sigma_{m,m:n}}{\gamma_{m:n}^2} \right] \alpha^2 & ; \text{ when } m = \frac{n+1}{2} \text{ and } n \text{ is odd} \\ \left(\gamma_1' G_2^{-1} \gamma_1 \right)^{-1} \alpha^2 & ; \text{ when } m = \frac{n}{2} \text{ and } n \text{ is even} \end{cases} = \alpha^2 V_4, \tag{15}$$

where

$$X_{mrss} = (X_{(m:n)1}, X_{(m+1:n)2}, X_{(m:n)3}, X_{(m+1:n)4}, \dots, X_{(m+1:n)n})$$

$$\gamma_1 = (\gamma_{m:n}, \gamma_{m+1:n}, \gamma_{m:n}, \gamma_{m+1:n}, \dots, \gamma_{m+1:n})$$

$$G_2 = \text{diag}(\sigma_{m,m:n}, \sigma_{m+1,m+1:n}, \sigma_{m,m:n}, \sigma_{m+1,m+1:n}, \dots, \sigma_{m+1,m+1:n}) \text{ and}$$

$$V_4 = \begin{cases} \frac{1}{n} \left[\frac{\sigma_{m,m:n}}{\gamma_{m:n}^2} \right] & ; \text{ when } m = \frac{n+1}{2} \text{ and } n \text{ is odd} \\ \left(\gamma_1' G_2^{-1} \gamma_1 \right)^{-1} & ; \text{ when } m = \frac{n}{2} \text{ and } n \text{ is even} \end{cases}.$$

2. Improved estimation of the scale parameter α

Let $t_i, i = 1, 2, 3, 4$ be an unbiased estimator of the parameter α , then we define a class of estimators for α as

$$T_i = A_i t_i, i = 1, 2, 3, 4,$$

where A_i 's, $i = 1, 2, 3, 4$ are suitably chosen constants such that mean squared error of the estimators T_i 's, $i = 1, 2, 3, 4$ is minimum.

The biases and mean squared errors (MSEs) of $T_i, i = 1, 2, 3, 4$ are respectively given by

$$B(T_i) = \alpha(A_i - 1),$$

and

$$MSE(T_i) = A_i^2 Var(t_i) + (A_i - 1)^2 \alpha^2 = \alpha^2 [A_i^2 (1 + V_i) - 2A_i + 1].$$

The $MSE(T_i), i = 1, 2, 3, 4$ is minimized for

$$A_i = (1 + V_i)^{-1}, i = 1, 2, 3, 4.$$

Thus, the resulting minimum MSE estimator of α is given by

$$T_{0i} = t_i (1 + V_i)^{-1}, i = 1, 2, 3, 4.$$

The biases and MSEs of $T_{0i}, i = 1, 2, 3, 4$ are respectively given as

$$B(T_{0i}) = -\alpha \left(\frac{V_i}{1+V_i} \right), \quad (16)$$

and

$$MSE(T_{0i}) = \alpha^2 \left(\frac{V_i}{1+V_i} \right). \quad (17)$$

We have from (12) - (15) and (17) that

$$Var(t_i) - MSE(T_{0i}) = \alpha^2 \frac{V_i^2}{(1+V_i)} > 0, i = 1, 2, 3, 4. \quad (18)$$

It follows from (18) that the proposed MMSE estimators $T_{0i}'s, i = 1, 2, 3, 4$ are better than the corresponding usual unbiased estimators $t_i's, i = 1, 2, 3, 4$.

3. Improved estimation of the scale parameter α with prior information

Let $t_i, i = 1, 2, 3, 4$ be an unbiased estimator of the parameter α , then we define a class of estimators of α using the prior point estimate α_0 of α as

$$T_{1i} = \alpha_0 + B_i t_i, i = 1, 2, 3, 4, \quad (19)$$

where $B_i's, i = 1, 2, 3, 4$ are suitably chosen constants such that mean squared error of the estimators $T_{1i}'s, i = 1, 2, 3, 4$ are minimum.

The biases and mean squared errors (MSEs) of $T_{1i}, i = 1, 2, 3, 4$ are respectively given by

$$B(T_{1i}) = \alpha(\phi + B_i),$$

and

$$MSE(T_{1i}) = \alpha^2 [\phi^2 + B_i^2(1+V_i) + 2\phi B_i].$$

where $\phi = \left(\frac{\alpha_0}{\alpha} - 1 \right) = (\lambda - 1)$ with $\lambda = \frac{\alpha_0}{\alpha}$.

The $MSE(T_{1i}), i = 1, 2, 3, 4$ is minimized for

$$B_i = -\phi(1+V_i)^{-1}, i = 1, 2, 3, 4. \quad (20)$$

The value of $B_i, i = 1,2,3,4$ at (20) depends on the unknown parameter α , so an estimate of $B_i, i = 1,2,3,4$ based on sample data is given by

$$B_i^* = -\frac{\phi^*}{(1+V_i)} = -\frac{(\theta_{20} - t_i)}{t_i(1+V_i)}, i = 1,2,3,4.$$

Putting $B_i^*, i = 1,2,3,4$ in (19), we get a shrinkage estimator of t_i 's, $i = 1,2,3,4$ as

$$T_{li}^* = \alpha_0 - (1+V_i)^{-1}(\alpha_0 - t_i), i = 1,2,3,4. \tag{21}$$

The biases and mean squared errors (MSEs) of the estimators T_{li}^* 's, $i = 1,2,3,4$ are respectively given by

$$B(T_{li}^*) = \alpha\phi\left(\frac{V_i}{1+V_i}\right), \tag{22}$$

and

$$MSE(T_{li}^*) = \alpha^2 \frac{V_i(\phi^2 V_i + 1)}{(1+V_i)^2}. \tag{23}$$

Comparisons of the proposed shrinkage estimators T_{li}^* 's, $i = 1,2,3,4$ with that of corresponding usual unbiased estimators t_i 's, $i = 1,2,3,4$ are given in the following Theorem 1.

Theorem 1: *The proposed shrinkage estimators T_{li}^* 's, $i = 1,2,3,4$ are better than the corresponding usual unbiased estimators t_i 's, $i = 1,2,3,4$ if*

$$\begin{aligned} &\lambda \in \left(0, \left(1 + \sqrt{(2+V_i)}\right)\right), \\ &\text{i.e. if } \alpha_0 \in \left(0, \alpha\left\{1 + \sqrt{(2+V_i)}\right\}\right), \\ &\text{i.e. if } \alpha \in \left(\frac{\alpha_0}{\left\{1 + \sqrt{(2+V_i)}\right\}}, \infty\right). \end{aligned}$$

Proof: From (12) - (15) and (23), we have that

$$MSE(T_{li}^*) < Var(t_i), i = 1,2,3,4 \text{ if}$$

$$\alpha^2 \frac{V_i(\phi^2 V_i + 1)}{(1+V_i)^2} < \alpha^2 V_i,$$

$$\text{i.e. if } \frac{(\phi^2 V_i + 1)}{(1+V_i)^2} < 1,$$

$$\begin{aligned}
& \text{i.e. if } \phi^2 < \frac{\left((1+V_i)^2 - 1\right)}{V_i}, \\
& \text{i.e. if } \phi^2 < \frac{V_i(2+V_i)}{V_i}, \\
& \text{i.e. if } \phi^2 < 2+V_i, \\
& \text{i.e. if } (\lambda-1)^2 < 2+V_i, \\
& \text{i.e. if } \left\{1 - \sqrt{(2+V_i)}\right\} < \lambda < \left\{1 + \sqrt{(2+V_i)}\right\}. \tag{24}
\end{aligned}$$

Since $\left\{1 - \sqrt{(2+V_i)}\right\} < 0$ and $\lambda (= \alpha_0 / \alpha)$ cannot be negative therefore (24) reduces to

$$0 < \lambda < \left\{1 + \sqrt{(2+V_i)}\right\},$$

or

$$0 < \alpha_0 < \alpha \left\{1 + \sqrt{(2+V_i)}\right\},$$

or

$$\left\{\frac{\alpha_0}{1 + \sqrt{(2+V_i)}}\right\} < \alpha < \infty.$$

Hence the theorem. ♦

Further, we have compared the proposed shrinkage estimators T_{1i}^* 's, $i = 1, 2, 3, 4$ with that of corresponding MMSE estimators T_{0i} 's, $i = 1, 2, 3, 4$ and the results are presented in Theorem 2.

Theorem 2: The proposed shrinkage estimators T_{1i}^* 's, $i = 1, 2, 3, 4$ are better than the corresponding MMSE estimators T_{0i} 's, $i = 1, 2, 3, 4$ if

$$\lambda \in (0, 2),$$

$$\text{i.e. if } \alpha_0 \in (0, 2\alpha),$$

$$\text{i.e. if } \alpha \in \left(\frac{\alpha_0}{2}, \infty\right).$$

Proof: From (17) and (23) we have that

$$MSE(T_{1i}^*) < MSE(T_{0i}), i = 1, 2, \dots, 7 \text{ if}$$

$$\alpha^2 \frac{V_i(\phi^2 V_i + 1)}{(1+V_i)^2} < \alpha^2 \frac{V_i}{1+V_i},$$

i.e. if $\frac{(\phi^2 V_i + 1)}{(1 + V_i)} < 1,$

i.e. if $\phi^2 < 1,$

i.e. if $-1 < \phi < 1,$

i.e. if $0 < \lambda < 2,$

or $0 < \alpha_0 < 2\alpha,$

or $\frac{\alpha_0}{2} < \alpha < \infty.$

Hence the theorem. ♦

4. Relative efficiencies

We have computed the relative efficiencies of various suggested estimators to usual estimators by using the formulae:

$$e_1 = RE(T_{01}, t_1) = 1 + V_1;$$

$$e_2 = RE(T_{02}, t_2) = 1 + V_2;$$

$$e_3 = RE(T_{03}, t_3) = 1 + V_3;$$

$$e_4 = RE(T_{04}, t_4) = 1 + V_4;$$

$$e_5 = RE(T_{04}, T_{01}) = \frac{V_1(1 + V_4)}{V_4(1 + V_1)};$$

$$e_6 = RE(T_{04}, T_{02}) = \frac{V_2(1 + V_4)}{V_4(1 + V_2)};$$

$$e_7 = RE(T_{04}, T_{03}) = \frac{V_3(1 + V_4)}{V_4(1 + V_3)};$$

$$e_8 = RE(T_{11}^*, t_1) = \frac{(1 + V_1)^2}{(\phi^2 V_1 + 1)};$$

$$e_9 = RE(T_{11}^*, T_{01}) = \frac{(1 + V_1)}{(\phi^2 V_1 + 1)};$$

$$e_{10} = RE(T_{12}^*, t_2) = \frac{(1 + V_2)^2}{(\phi^2 V_2 + 1)};$$

$$e_{11} = RE(T_{12}^*, T_{02}) = \frac{(1 + V_2)}{(\phi^2 V_2 + 1)};$$

$$e_{12} = RE(T_{13}^*, t_3) = \frac{(1 + V_3)^2}{(\phi^2 V_3 + 1)};$$

$$e_{13} = RE(T_{13}^*, T_{03}) = \frac{(1 + V_3)}{(\phi^2 V_3 + 1)};$$

$$e_{14} = RE(T_{14}^*, t_4) = \frac{(1 + V_4)^2}{(\phi^2 V_4 + 1)};$$

and

$$e_{15} = RE(T_{14}^*, T_{04}) = \frac{(1 + V_4)}{(\phi^2 V_4 + 1)}.$$

- The values of $e_i, i = 1, 2, \dots, 7$ are shown in Table 1 for $n = 2(1)8$ and $\beta = 2.5(0.5)5$.

- The values of $e_i, i = 8, 9, \dots, 15$ are shown in Tables 2 to 5 for $n = 2(1)8$;

$\beta = 2.5(0.5)5$ and different values of $\lambda = \frac{\alpha_0}{\alpha}$.

Table 1. The values of e_i 's, $i = 1, 2, \dots, 7$.

n	β	e_1	e_2	e_3	e_4	e_5	e_6	e_7
2	2.5	1.3371	1.4025	1.2799	1.2799	1.1530	1.3124	1.0000
	3.0	1.2158	1.1978	1.1674	1.1674	1.2381	1.1516	1.0000
	3.5	1.1514	1.1254	1.1135	1.1135	1.2901	1.0932	1.0000
	4.0	1.1126	1.0885	1.0827	1.0827	1.3242	1.0643	1.0000
	4.5	1.0872	1.0665	1.0633	1.0633	1.3476	1.0474	1.0000
	5.0	1.0697	1.0521	1.0501	1.0501	1.3644	1.0368	1.0000
3	2.5	1.2011	1.1904	1.1180	1.0832	2.1798	2.0828	1.3740
	3.0	1.1308	1.0970	1.0752	1.0543	2.2446	1.7159	1.3572
	3.5	1.0937	1.0626	1.0526	1.0385	2.3093	1.5893	1.3475
	4.0	1.0706	1.0447	1.0391	1.0288	2.3527	1.5272	1.3422
	4.5	1.0552	1.0338	1.0303	1.0224	2.3815	1.4910	1.3382
	5.0	1.0443	1.0266	1.0242	1.0180	2.4033	1.4675	1.3356
4	2.5	1.1392	1.1118	1.0648	1.0477	2.6846	2.2089	1.3376
	3.0	1.0939	1.0582	1.0424	1.0317	2.7929	1.7903	1.3246
	3.5	1.0678	1.0380	1.0301	1.0227	2.8598	1.6495	1.3176
	4.0	1.0514	1.0273	1.0226	1.0171	2.9034	1.5803	1.3126
	4.5	1.0413	1.0208	1.0176	1.0134	3.0062	1.5408	1.3102
	5.0	1.0324	1.0164	1.0141	1.0107	2.9570	1.5158	1.3085
5	2.5	1.1078	1.0740	1.0409	1.0278	3.5999	2.5482	1.4547
	3.0	1.0733	1.0391	1.0272	1.0187	3.7126	2.0476	1.4397
	3.5	1.0531	1.0257	1.0195	1.0135	3.7819	1.8805	1.4317
	4.0	1.0403	1.0186	1.0147	1.0101	3.8704	1.8188	1.4421
	4.5	1.0316	1.0141	1.0115	1.0080	3.8489	1.7503	1.4223
	5.0	1.0256	1.0112	1.0092	1.0065	3.8785	1.7199	1.4196
6	2.5	1.0880	1.0528	1.0282	1.0196	4.2168	2.6150	1.4288
	3.0	1.0600	1.0282	1.0189	1.0133	4.3313	2.0988	1.4170
	3.5	1.0437	1.0187	1.0136	1.0096	4.3995	1.9265	1.4101
	4.0	1.0332	1.0135	1.0103	1.0073	4.4473	1.8430	1.4065
	4.5	1.0261	1.0103	1.0080	1.0057	4.4756	1.7943	1.4024
	5.0	1.0211	1.0082	1.0065	1.0046	4.5010	1.7638	1.4009
7	2.5	1.0735	1.0397	1.0206	1.0138	5.0286	2.8038	1.4800
	3.0	1.0509	1.0214	1.0139	1.0094	5.1991	2.2498	1.4690
	3.5	1.0372	1.0147	1.0106	1.0068	5.2926	2.1298	1.5419
	4.0	1.0282	1.0103	1.0076	1.0052	5.3054	1.9708	1.4542
	4.5	1.0222	1.0079	1.0060	1.0041	5.3447	1.9217	1.4556
	5.0	1.0179	1.0062	1.0048	1.0033	5.3582	1.8854	1.4494
8	2.5	1.0643	1.0310	1.0157	1.0107	5.7315	2.8525	1.4632
	3.0	1.0444	1.0168	1.0106	1.0073	5.8758	2.2888	1.4526
	3.5	1.0329	1.0112	1.0077	1.0053	6.0530	2.1067	1.4484
	4.0	1.0245	1.0081	1.0058	1.0040	5.9651	2.0116	1.4441
	4.5	1.0190	1.0062	1.0046	1.0032	5.9067	1.9593	1.4396
	5.0	1.0156	1.0049	1.0037	1.0025	6.0905	1.9479	1.4568

Table 2. The values of e_i for $i = 8$ and 9

n	β	$e_8 = RE(t_{II}^*, t_I)$						Range of λ in which T_{II}^* is efficient to t_I
		$\lambda = 1.00$	$\lambda = 1.25$ and $\lambda = 0.75$	$\lambda = 1.50$ and $\lambda = 0.50$	$\lambda = 1.75$ and $\lambda = 0.25$	$\lambda = 2.00$	$\lambda = 2.25$	
2	2.5	1.7878	1.7509	1.6489	1.5028	1.3371	1.1710	(0,2.53)
	3.0	1.4782	1.4586	1.4026	1.3182	1.2158	1.1054	(0,2.49)
	3.5	1.3257	1.3133	1.2773	1.2217	1.1514	1.0721	(0,2.47)
	4.0	1.2378	1.2292	1.2039	1.1641	1.1126	1.0527	(0,2.45)
	4.5	1.1820	1.1756	1.1568	1.1268	1.0872	1.0403	(0,2.44)
	5.0	1.1442	1.1392	1.1246	1.1010	1.0697	1.0319	(0,2.44)
3	2.5	1.4426	1.4247	1.3735	1.2960	1.2011	1.0977	(0,2.48)
	3.0	1.2786	1.2683	1.2382	1.1910	1.1308	1.0617	(0,2.46)
	3.5	1.1961	1.1892	1.1688	1.1363	1.0937	1.0434	(0,2.45)
	4.0	1.1461	1.1411	1.1263	1.1024	1.0706	1.0323	(0,2.44)
	4.5	1.1133	1.1095	1.0982	1.0798	1.0552	1.0250	(0,2.43)
	5.0	1.0906	1.0876	1.0787	1.0641	1.0443	1.0200	(0,2.43)
4	2.5	1.2977	1.2865	1.2541	1.2035	1.1392	1.0659	(0,2.46)
	3.0	1.1966	1.1896	1.1692	1.1366	1.0939	1.0435	(0,2.45)
	3.5	1.1402	1.1354	1.1212	1.0983	1.0678	1.0310	(0,2.44)
	4.0	1.1053	1.1018	1.0913	1.0743	1.0514	1.0232	(0,2.43)
	4.5	1.0843	1.0815	1.0732	1.0597	1.0413	1.0186	(0,2.43)
	5.0	1.0659	1.0638	1.0574	1.0468	1.0324	1.0145	(0,2.43)
5	2.5	1.2272	1.2190	1.1950	1.1570	1.1078	1.0503	(0,2.45)
	3.0	1.1519	1.1466	1.1312	1.1063	1.0733	1.0336	(0,2.44)
	3.5	1.1091	1.1054	1.0945	1.0769	1.0531	1.0241	(0,2.43)
	4.0	1.0823	1.0796	1.0715	1.0583	1.0403	1.0181	(0,2.43)
	4.5	1.0643	1.0622	1.0559	1.0457	1.0316	1.0141	(0,2.43)
	5.0	1.0518	1.0501	1.0451	1.0369	1.0256	1.0114	(0,2.42)
6	2.5	1.1837	1.1772	1.1582	1.1279	1.0880	1.0406	(0,2.44)
	3.0	1.1237	1.1195	1.1071	1.0870	1.0600	1.0273	(0,2.44)
	3.5	1.0892	1.0863	1.0775	1.0631	1.0437	1.0197	(0,2.43)
	4.0	1.0675	1.0653	1.0587	1.0479	1.0332	1.0149	(0,2.43)
	4.5	1.0529	1.0512	1.0461	1.0377	1.0261	1.0116	(0,2.42)
	5.0	1.0426	1.0413	1.0372	1.0304	1.0211	1.0094	(0,2.42)
7	2.5	1.1524	1.1471	1.1316	1.1066	1.0735	1.0337	(0,2.44)
	3.0	1.1043	1.1008	1.0905	1.0736	1.0509	1.0230	(0,2.43)
	3.5	1.0759	1.0734	1.0659	1.0538	1.0372	1.0167	(0,2.43)
	4.0	1.0572	1.0554	1.0498	1.0407	1.0282	1.0126	(0,2.42)
	4.5	1.0449	1.0434	1.0391	1.0320	1.0222	1.0099	(0,2.42)
	5.0	1.0362	1.0350	1.0316	1.0258	1.0179	1.0079	(0,2.42)
8	2.5	1.1327	1.1282	1.1148	1.0932	1.0643	1.0293	(0,2.44)
	3.0	1.0907	1.0877	1.0787	1.0641	1.0444	1.0200	(0,2.43)
	3.5	1.0669	1.0647	1.0582	1.0475	1.0329	1.0147	(0,2.43)
	4.0	1.0497	1.0481	1.0433	1.0354	1.0245	1.0109	(0,2.42)
	4.5	1.0384	1.0372	1.0335	1.0274	1.0190	1.0084	(0,2.42)
	5.0	1.0315	1.0305	1.0275	1.0225	1.0156	1.0069	(0,2.42)

Table 2. The values of e_i for $i = 8$ and 9 (cont.)

n	β	$e_9 = RE(T_{11}^*, T_{01})$					Range of λ in which T_{11}^* is efficient to T_{01}
		$\lambda = 1.00$	$\lambda = 1.25$ and $\lambda = 0.75$	$\lambda = 1.50$ and $\lambda = 0.50$	$\lambda = 1.75$ and $\lambda = 0.25$	$\lambda = 2.00$ and $\lambda = 0.00$	
2	2.5	1.3371	1.3095	1.2332	1.1240	1.0000	[0,2]
	3.0	1.2158	1.1996	1.1536	1.0842	1.0000	[0,2]
	3.5	1.1514	1.1406	1.1094	1.0610	1.0000	[0,2]
	4.0	1.1126	1.1048	1.0821	1.0463	1.0000	[0,2]
	4.5	1.0872	1.0813	1.0640	1.0364	1.0000	[0,2]
	5.0	1.0697	1.0650	1.0514	1.0293	1.0000	[0,2]
3	2.5	1.2011	1.1862	1.1436	1.0790	1.0000	[0,2]
	3.0	1.1308	1.1216	1.0950	1.0533	1.0000	[0,2]
	3.5	1.0937	1.0873	1.0687	1.0389	1.0000	[0,2]
	4.0	1.0706	1.0659	1.0520	1.0297	1.0000	[0,2]
	4.5	1.0552	1.0515	1.0408	1.0234	1.0000	[0,2]
	5.0	1.0443	1.0414	1.0329	1.0189	1.0000	[0,2]
4	2.5	1.1392	1.1294	1.1009	1.0565	1.0000	[0,2]
	3.0	1.0939	1.0875	1.0688	1.0390	1.0000	[0,2]
	3.5	1.0678	1.0633	1.0500	1.0286	1.0000	[0,2]
	4.0	1.0514	1.0480	1.0380	1.0218	1.0000	[0,2]
	4.5	1.0413	1.0386	1.0307	1.0177	1.0000	[0,2]
	5.0	1.0324	1.0304	1.0241	1.0139	1.0000	[0,2]
5	2.5	1.1078	1.1004	1.0787	1.0445	1.0000	[0,2]
	3.0	1.0733	1.0684	1.0540	1.0308	1.0000	[0,2]
	3.5	1.0531	1.0496	1.0393	1.0226	1.0000	[0,2]
	4.0	1.0403	1.0377	1.0300	1.0173	1.0000	[0,2]
	4.5	1.0316	1.0296	1.0235	1.0136	1.0000	[0,2]
	5.0	1.0256	1.0239	1.0191	1.0110	1.0000	[0,2]
6	2.5	1.0880	1.0820	1.0646	1.0367	1.0000	[0,2]
	3.0	1.0600	1.0561	1.0444	1.0254	1.0000	[0,2]
	3.5	1.0437	1.0408	1.0324	1.0186	1.0000	[0,2]
	4.0	1.0332	1.0311	1.0247	1.0143	1.0000	[0,2]
	4.5	1.0261	1.0244	1.0195	1.0113	1.0000	[0,2]
	5.0	1.0211	1.0197	1.0157	1.0091	1.0000	[0,2]
7	2.5	1.0735	1.0686	1.0541	1.0309	1.0000	[0,2]
	3.0	1.0509	1.0475	1.0377	1.0216	1.0000	[0,2]
	3.5	1.0372	1.0348	1.0277	1.0160	1.0000	[0,2]
	4.0	1.0282	1.0264	1.0210	1.0122	1.0000	[0,2]
	4.5	1.0222	1.0208	1.0166	1.0096	1.0000	[0,2]
	5.0	1.0179	1.0168	1.0134	1.0078	1.0000	[0,2]
8	2.5	1.0643	1.0600	1.0474	1.0271	1.0000	[0,2]
	3.0	1.0444	1.0415	1.0329	1.0189	1.0000	[0,2]
	3.5	1.0329	1.0308	1.0245	1.0141	1.0000	[0,2]
	4.0	1.0245	1.0230	1.0183	1.0106	1.0000	[0,2]
	4.5	1.0190	1.0178	1.0142	1.0082	1.0000	[0,2]
	5.0	1.0156	1.0146	1.0117	1.0068	1.0000	[0,2]

Table 3. The values of e_i for $i = 10$ and 11

n	β	$e_{10} = RE(T_{12}^*, t_2)$						Range of λ in which T_{12}^* is efficient to t_2
		$\lambda = 1.00$	$\lambda = 1.25$ and $\lambda = 0.75$	$\lambda = 1.50$ and $\lambda = 0.50$	$\lambda = 1.75$ and $\lambda = 0.25$	$\lambda = 2.00$	$\lambda = 2.25$	
2	2.5	1.9669	1.9186	1.7871	1.6038	1.4025	1.2075	(0,2.55)
	3.0	1.4347	1.4171	1.3671	1.2910	1.1978	1.0960	(0,2.48)
	3.5	1.2665	1.2566	1.2280	1.1830	1.1254	1.0590	(0,2.46)
	4.0	1.1849	1.1784	1.1592	1.1287	1.0885	1.0409	(0,2.45)
	4.5	1.1374	1.1327	1.1188	1.0964	1.0665	1.0304	(0,2.44)
3	5.0	1.1069	1.1033	1.0926	1.0754	1.0521	1.0236	(0,2.43)
	2.5	1.4171	1.4004	1.3527	1.2800	1.1904	1.0922	(0,2.48)
	3.0	1.2034	1.1961	1.1749	1.1411	1.0970	1.0450	(0,2.45)
	3.5	1.1292	1.1248	1.1118	1.0908	1.0626	1.0285	(0,2.44)
	4.0	1.0914	1.0884	1.0794	1.0646	1.0447	1.0202	(0,2.43)
4	4.5	1.0688	1.0665	1.0598	1.0488	1.0338	1.0151	(0,2.43)
	5.0	1.0539	1.0522	1.0470	1.0384	1.0266	1.0119	(0,2.42)
	2.5	1.2360	1.2274	1.2024	1.1629	1.1118	1.0523	(0,2.45)
	3.0	1.1199	1.1158	1.1038	1.0843	1.0582	1.0265	(0,2.43)
	3.5	1.0775	1.0749	1.0673	1.0549	1.0380	1.0171	(0,2.43)
5	4.0	1.0554	1.0536	1.0482	1.0394	1.0273	1.0122	(0,2.42)
	4.5	1.0419	1.0406	1.0366	1.0299	1.0208	1.0092	(0,2.42)
	5.0	1.0330	1.0320	1.0288	1.0236	1.0164	1.0072	(0,2.42)
	2.5	1.1534	1.1481	1.1325	1.1073	1.0740	1.0339	(0,2.44)
	3.0	1.0798	1.0771	1.0693	1.0565	1.0391	1.0176	(0,2.43)
6	3.5	1.0521	1.0504	1.0454	1.0371	1.0257	1.0115	(0,2.42)
	4.0	1.0375	1.0363	1.0327	1.0267	1.0186	1.0082	(0,2.42)
	4.5	1.0285	1.0276	1.0249	1.0204	1.0141	1.0062	(0,2.42)
	5.0	1.0225	1.0218	1.0196	1.0161	1.0112	1.0049	(0,2.42)
	2.5	1.1084	1.1047	1.0939	1.0764	1.0528	1.0239	(0,2.43)
7	3.0	1.0572	1.0554	1.0498	1.0407	1.0282	1.0126	(0,2.42)
	3.5	1.0377	1.0365	1.0328	1.0269	1.0187	1.0083	(0,2.42)
	4.0	1.0272	1.0263	1.0237	1.0194	1.0135	1.0060	(0,2.42)
	4.5	1.0207	1.0201	1.0181	1.0148	1.0103	1.0045	(0,2.42)
	5.0	1.0164	1.0159	1.0143	1.0117	1.0082	1.0036	(0,2.42)
8	2.5	1.0809	1.0783	1.0703	1.0573	1.0397	1.0178	(0,2.43)
	3.0	1.0433	1.0419	1.0377	1.0308	1.0214	1.0095	(0,2.42)
	3.5	1.0295	1.0286	1.0258	1.0211	1.0147	1.0065	(0,2.42)
	4.0	1.0207	1.0200	1.0181	1.0148	1.0103	1.0045	(0,2.42)
	4.5	1.0158	1.0153	1.0138	1.0113	1.0079	1.0035	(0,2.42)
9	5.0	1.0125	1.0121	1.0109	1.0090	1.0062	1.0027	(0,2.42)
	2.5	1.0629	1.0609	1.0548	1.0447	1.0310	1.0138	(0,2.43)
	3.0	1.0339	1.0328	1.0296	1.0242	1.0168	1.0074	(0,2.42)
	3.5	1.0225	1.0218	1.0197	1.0161	1.0112	1.0049	(0,2.42)
	4.0	1.0163	1.0158	1.0143	1.0117	1.0081	1.0036	(0,2.42)
10	4.5	1.0125	1.0121	1.0109	1.0090	1.0062	1.0027	(0,2.42)
	5.0	1.0099	1.0096	1.0087	1.0071	1.0049	1.0022	(0,2.42)

Table 3. The values of e_i for $i = 10$ and 11 (cont.)

n	β	$e_{11} = RE(T_{12}^*, T_{02})$					Range of λ in which T_{12}^* is efficient to T_{02}
		$\lambda = 1.00$	$\lambda = 1.25$ and $\lambda = 0.75$	$\lambda = 1.50$ and $\lambda = 0.50$	$\lambda = 1.75$ and $\lambda = 0.25$	$\lambda = 2.00$ and $\lambda = 0.00$	
2	2.5	1.4025	1.3680	1.2743	1.1436	1.0000	[0,2]
	3.0	1.1978	1.1831	1.1413	1.0779	1.0000	[0,2]
	3.5	1.1254	1.1166	1.0912	1.0512	1.0000	[0,2]
	4.0	1.0885	1.0825	1.0650	1.0369	1.0000	[0,2]
	4.5	1.0665	1.0621	1.0491	1.0280	1.0000	[0,2]
3	5.0	1.0521	1.0487	1.0386	1.0221	1.0000	[0,2]
	2.5	1.1904	1.1764	1.1363	1.0752	1.0000	[0,2]
	3.0	1.0970	1.0904	1.0710	1.0402	1.0000	[0,2]
	3.5	1.0626	1.0585	1.0463	1.0265	1.0000	[0,2]
	4.0	1.0447	1.0418	1.0332	1.0191	1.0000	[0,2]
4	4.5	1.0338	1.0316	1.0252	1.0145	1.0000	[0,2]
	5.0	1.0266	1.0249	1.0198	1.0115	1.0000	[0,2]
	2.5	1.1118	1.1040	1.0815	1.0460	1.0000	[0,2]
	3.0	1.0582	1.0544	1.0430	1.0247	1.0000	[0,2]
	3.5	1.0380	1.0356	1.0282	1.0163	1.0000	[0,2]
5	4.0	1.0273	1.0256	1.0203	1.0118	1.0000	[0,2]
	4.5	1.0208	1.0194	1.0155	1.0090	1.0000	[0,2]
	5.0	1.0164	1.0153	1.0122	1.0071	1.0000	[0,2]
	2.5	1.0740	1.0690	1.0545	1.0311	1.0000	[0,2]
	3.0	1.0391	1.0366	1.0291	1.0167	1.0000	[0,2]
6	3.5	1.0257	1.0241	1.0192	1.0111	1.0000	[0,2]
	4.0	1.0186	1.0174	1.0139	1.0080	1.0000	[0,2]
	4.5	1.0141	1.0132	1.0106	1.0061	1.0000	[0,2]
	5.0	1.0112	1.0105	1.0084	1.0049	1.0000	[0,2]
	2.5	1.0528	1.0493	1.0391	1.0224	1.0000	[0,2]
7	3.0	1.0282	1.0264	1.0210	1.0122	1.0000	[0,2]
	3.5	1.0187	1.0175	1.0139	1.0081	1.0000	[0,2]
	4.0	1.0135	1.0126	1.0101	1.0059	1.0000	[0,2]
	4.5	1.0103	1.0097	1.0077	1.0045	1.0000	[0,2]
	5.0	1.0082	1.0076	1.0061	1.0036	1.0000	[0,2]
8	2.5	1.0397	1.0371	1.0295	1.0170	1.0000	[0,2]
	3.0	1.0214	1.0200	1.0160	1.0093	1.0000	[0,2]
	3.5	1.0147	1.0137	1.0110	1.0064	1.0000	[0,2]
	4.0	1.0103	1.0097	1.0077	1.0045	1.0000	[0,2]
	4.5	1.0079	1.0074	1.0059	1.0034	1.0000	[0,2]
9	5.0	1.0062	1.0058	1.0047	1.0027	1.0000	[0,2]
	2.5	1.0310	1.0290	1.0231	1.0133	1.0000	[0,2]
	3.0	1.0168	1.0158	1.0126	1.0073	1.0000	[0,2]
	3.5	1.0112	1.0105	1.0084	1.0049	1.0000	[0,2]
	4.0	1.0081	1.0076	1.0061	1.0035	1.0000	[0,2]
10	4.5	1.0062	1.0058	1.0047	1.0027	1.0000	[0,2]
	5.0	1.0049	1.0046	1.0037	1.0022	1.0000	[0,2]

Table 4. The values of e_i for $i = 12$ and 13

n	β	$e_{12} = RE(T_{13}^*, t_3)$						Range of λ in which T_{13}^* is efficient to t_3
		$\lambda = 1.00$	$\lambda = 1.25$ and $\lambda = 0.75$	$\lambda = 1.50$ and $\lambda = 0.50$	$\lambda = 1.75$ and $\lambda = 0.25$	$\lambda = 2.00$	$\lambda = 2.25$	
2	2.5	1.6380	1.6099	1.5309	1.4152	1.2799	1.1397	(0,2.51)
	3.0	1.3628	1.3487	1.3080	1.2455	1.1674	1.0803	(0,2.47)
	3.5	1.2398	1.2311	1.2056	1.1654	1.1135	1.0531	(0,2.45)
	4.0	1.1723	1.1663	1.1485	1.1202	1.0827	1.0381	(0,2.44)
	4.5	1.1306	1.1262	1.1130	1.0917	1.0633	1.0288	(0,2.44)
	5.0	1.1028	1.0993	1.0891	1.0725	1.0501	1.0227	(0,2.43)
3	2.5	1.2499	1.2407	1.2141	1.1721	1.1180	1.0553	(0,2.46)
	3.0	1.1560	1.1506	1.1347	1.1091	1.0752	1.0345	(0,2.44)
	3.5	1.1080	1.1044	1.0936	1.0761	1.0526	1.0238	(0,2.43)
	4.0	1.0797	1.0771	1.0692	1.0565	1.0391	1.0176	(0,2.43)
	4.5	1.0614	1.0594	1.0535	1.0437	1.0303	1.0135	(0,2.42)
	5.0	1.0489	1.0473	1.0426	1.0348	1.0242	1.0107	(0,2.42)
4	2.5	1.1338	1.1293	1.1158	1.0940	1.0648	1.0296	(0,2.44)
	3.0	1.0867	1.0838	1.0753	1.0613	1.0424	1.0191	(0,2.43)
	3.5	1.0612	1.0592	1.0533	1.0435	1.0301	1.0135	(0,2.42)
	4.0	1.0457	1.0442	1.0398	1.0326	1.0226	1.0100	(0,2.42)
	4.5	1.0355	1.0344	1.0310	1.0253	1.0176	1.0078	(0,2.42)
	5.0	1.0284	1.0275	1.0248	1.0203	1.0141	1.0062	(0,2.42)
5	2.5	1.0835	1.0808	1.0726	1.0592	1.0409	1.0184	(0,2.43)
	3.0	1.0551	1.0533	1.0480	1.0392	1.0272	1.0121	(0,2.42)
	3.5	1.0393	1.0381	1.0343	1.0281	1.0195	1.0086	(0,2.42)
	4.0	1.0295	1.0286	1.0258	1.0211	1.0147	1.0065	(0,2.42)
	4.5	1.0231	1.0223	1.0201	1.0165	1.0115	1.0051	(0,2.42)
	5.0	1.0185	1.0179	1.0162	1.0133	1.0092	1.0041	(0,2.42)
6	2.5	1.0571	1.0553	1.0497	1.0406	1.0282	1.0126	(0,2.42)
	3.0	1.0381	1.0369	1.0332	1.0272	1.0189	1.0084	(0,2.42)
	3.5	1.0274	1.0265	1.0239	1.0196	1.0136	1.0060	(0,2.42)
	4.0	1.0206	1.0200	1.0180	1.0148	1.0103	1.0045	(0,2.42)
	4.5	1.0161	1.0156	1.0141	1.0116	1.0080	1.0035	(0,2.42)
	5.0	1.0130	1.0126	1.0113	1.0093	1.0065	1.0028	(0,2.42)
7	2.5	1.0415	1.0402	1.0362	1.0296	1.0206	1.0091	(0,2.42)
	3.0	1.0279	1.0270	1.0244	1.0200	1.0139	1.0061	(0,2.42)
	3.5	1.0213	1.0206	1.0186	1.0152	1.0106	1.0047	(0,2.42)
	4.0	1.0152	1.0147	1.0133	1.0109	1.0076	1.0033	(0,2.42)
	4.5	1.0119	1.0116	1.0104	1.0086	1.0060	1.0026	(0,2.42)
	5.0	1.0096	1.0093	1.0084	1.0069	1.0048	1.0021	(0,2.42)
8	2.5	1.0316	1.0306	1.0275	1.0226	1.0157	1.0069	(0,2.42)
	3.0	1.0213	1.0207	1.0186	1.0153	1.0106	1.0047	(0,2.42)
	3.5	1.0154	1.0149	1.0135	1.0111	1.0077	1.0034	(0,2.42)
	4.0	1.0117	1.0113	1.0102	1.0084	1.0058	1.0026	(0,2.42)
	4.5	1.0092	1.0089	1.0080	1.0066	1.0046	1.0020	(0,2.42)
	5.0	1.0074	1.0072	1.0065	1.0053	1.0037	1.0016	(0,2.42)

Table 4. The values of e_i for $i = 12$ and 13 (cont.)

n	β	$e_{13} = RE(T_{13}^*, T_{03})$					Range of λ in which T_{13}^* is efficient to T_{03}
		$\lambda = 1.00$	$\lambda = 1.25$ and $\lambda = 0.75$	$\lambda = 1.50$ and $\lambda = 0.50$	$\lambda = 1.75$ and $\lambda = 0.25$	$\lambda = 2.00$ and $\lambda = 0.00$	
2	2.5	1.2799	1.2578	1.1962	1.1058	1.0000	[0,2]
	3.0	1.1674	1.1553	1.1205	1.0669	1.0000	[0,2]
	3.5	1.1135	1.1056	1.0828	1.0467	1.0000	[0,2]
	4.0	1.0827	1.0772	1.0608	1.0346	1.0000	[0,2]
	4.5	1.0633	1.0591	1.0467	1.0267	1.0000	[0,2]
3	5.0	1.0501	1.0469	1.0371	1.0213	1.0000	[0,2]
	2.5	1.1180	1.1098	1.0859	1.0484	1.0000	[0,2]
	3.0	1.0752	1.0702	1.0553	1.0316	1.0000	[0,2]
	3.5	1.0526	1.0492	1.0389	1.0224	1.0000	[0,2]
	4.0	1.0391	1.0365	1.0290	1.0167	1.0000	[0,2]
4	4.5	1.0303	1.0283	1.0225	1.0130	1.0000	[0,2]
	5.0	1.0242	1.0226	1.0180	1.0104	1.0000	[0,2]
	2.5	1.0648	1.0605	1.0478	1.0274	1.0000	[0,2]
	3.0	1.0424	1.0397	1.0315	1.0181	1.0000	[0,2]
	3.5	1.0301	1.0282	1.0224	1.0130	1.0000	[0,2]
5	4.0	1.0226	1.0211	1.0168	1.0098	1.0000	[0,2]
	4.5	1.0176	1.0165	1.0131	1.0076	1.0000	[0,2]
	5.0	1.0141	1.0132	1.0105	1.0061	1.0000	[0,2]
	2.5	1.0409	1.0383	1.0304	1.0175	1.0000	[0,2]
	3.0	1.0272	1.0254	1.0203	1.0117	1.0000	[0,2]
6	3.5	1.0195	1.0182	1.0145	1.0084	1.0000	[0,2]
	4.0	1.0147	1.0137	1.0110	1.0064	1.0000	[0,2]
	4.5	1.0115	1.0107	1.0086	1.0050	1.0000	[0,2]
	5.0	1.0092	1.0086	1.0069	1.0040	1.0000	[0,2]
	2.5	1.0282	1.0264	1.0210	1.0121	1.0000	[0,2]
7	3.0	1.0189	1.0177	1.0141	1.0082	1.0000	[0,2]
	3.5	1.0136	1.0127	1.0102	1.0059	1.0000	[0,2]
	4.0	1.0103	1.0096	1.0077	1.0045	1.0000	[0,2]
	4.5	1.0080	1.0075	1.0060	1.0035	1.0000	[0,2]
	5.0	1.0065	1.0061	1.0048	1.0028	1.0000	[0,2]
8	2.5	1.0206	1.0193	1.0153	1.0089	1.0000	[0,2]
	3.0	1.0139	1.0130	1.0104	1.0060	1.0000	[0,2]
	3.5	1.0106	1.0099	1.0079	1.0046	1.0000	[0,2]
	4.0	1.0076	1.0071	1.0057	1.0033	1.0000	[0,2]
	4.5	1.0060	1.0056	1.0045	1.0026	1.0000	[0,2]
9	5.0	1.0048	1.0045	1.0036	1.0021	1.0000	[0,2]
	2.5	1.0157	1.0147	1.0117	1.0068	1.0000	[0,2]
	3.0	1.0106	1.0099	1.0079	1.0046	1.0000	[0,2]
	3.5	1.0077	1.0072	1.0057	1.0033	1.0000	[0,2]
	4.0	1.0058	1.0055	1.0044	1.0025	1.0000	[0,2]
10	4.5	1.0046	1.0043	1.0034	1.0020	1.0000	[0,2]
	5.0	1.0037	1.0035	1.0028	1.0016	1.0000	[0,2]

Table 5. The values of e_i for $i = 14$ and 15

n	β	$e_{14} = RE(T_{14}^*, t_4)$						Range of λ in which T_{14}^* is efficient to t_4
		$\lambda = 1.00$	$\lambda = 1.25$ and $\lambda = 0.75$	$\lambda = 1.50$ and $\lambda = 0.50$	$\lambda = 1.75$ and $\lambda = 0.25$	$\lambda = 2.00$	$\lambda = 2.25$	
2	2.5	1.6380	1.6099	1.5309	1.4152	1.2799	1.1397	(0,2.51)
	3.0	1.3628	1.3487	1.3080	1.2455	1.1674	1.0803	(0,2.47)
	3.5	1.2398	1.2311	1.2056	1.1654	1.1135	1.0531	(0,2.45)
	4.0	1.1723	1.1663	1.1485	1.1202	1.0827	1.0381	(0,2.44)
	4.5	1.1306	1.1262	1.1130	1.0917	1.0633	1.0288	(0,2.44)
3	5.0	1.1028	1.0993	1.0891	1.0725	1.0501	1.0227	(0,2.43)
	2.5	1.1733	1.1672	1.1494	1.1209	1.0832	1.0383	(0,2.44)
	3.0	1.1116	1.1078	1.0967	1.0786	1.0543	1.0246	(0,2.43)
	3.5	1.0785	1.0759	1.0682	1.0557	1.0385	1.0173	(0,2.43)
	4.0	1.0585	1.0566	1.0509	1.0416	1.0288	1.0129	(0,2.42)
4	4.5	1.0454	1.0439	1.0396	1.0324	1.0224	1.0100	(0,2.42)
	5.0	1.0363	1.0351	1.0316	1.0259	1.0180	1.0080	(0,2.42)
	2.5	1.0976	1.0944	1.0847	1.0690	1.0477	1.0215	(0,2.43)
	3.0	1.0644	1.0623	1.0561	1.0458	1.0317	1.0142	(0,2.43)
	3.5	1.0459	1.0445	1.0400	1.0327	1.0227	1.0101	(0,2.42)
5	4.0	1.0345	1.0334	1.0301	1.0247	1.0171	1.0076	(0,2.42)
	4.5	1.0269	1.0261	1.0235	1.0193	1.0134	1.0059	(0,2.42)
	5.0	1.0216	1.0209	1.0189	1.0155	1.0107	1.0047	(0,2.42)
	2.5	1.0563	1.0545	1.0490	1.0401	1.0278	1.0124	(0,2.42)
	3.0	1.0378	1.0366	1.0330	1.0270	1.0187	1.0083	(0,2.42)
6	3.5	1.0272	1.0264	1.0238	1.0195	1.0135	1.0060	(0,2.42)
	4.0	1.0203	1.0197	1.0178	1.0146	1.0101	1.0045	(0,2.42)
	4.5	1.0161	1.0156	1.0141	1.0116	1.0080	1.0035	(0,2.42)
	5.0	1.0130	1.0126	1.0113	1.0093	1.0065	1.0028	(0,2.42)
	2.5	1.0395	1.0382	1.0344	1.0282	1.0196	1.0087	(0,2.42)
7	3.0	1.0267	1.0258	1.0233	1.0191	1.0133	1.0059	(0,2.42)
	3.5	1.0193	1.0187	1.0169	1.0138	1.0096	1.0042	(0,2.42)
	4.0	1.0146	1.0142	1.0128	1.0105	1.0073	1.0032	(0,2.42)
	4.5	1.0115	1.0111	1.0100	1.0082	1.0057	1.0025	(0,2.42)
	5.0	1.0092	1.0090	1.0081	1.0066	1.0046	1.0020	(0,2.42)
8	2.5	1.0278	1.0269	1.0243	1.0199	1.0138	1.0061	(0,2.42)
	3.0	1.0189	1.0183	1.0165	1.0135	1.0094	1.0041	(0,2.42)
	3.5	1.0137	1.0133	1.0120	1.0098	1.0068	1.0030	(0,2.42)
	4.0	1.0104	1.0101	1.0091	1.0075	1.0052	1.0023	(0,2.42)
	4.5	1.0082	1.0079	1.0071	1.0059	1.0041	1.0018	(0,2.42)
9	5.0	1.0066	1.0064	1.0058	1.0047	1.0033	1.0014	(0,2.42)
	2.5	1.0214	1.0207	1.0187	1.0153	1.0107	1.0047	(0,2.42)
	3.0	1.0146	1.0142	1.0128	1.0105	1.0073	1.0032	(0,2.42)
	3.5	1.0106	1.0103	1.0093	1.0076	1.0053	1.0023	(0,2.42)
	4.0	1.0081	1.0078	1.0071	1.0058	1.0040	1.0018	(0,2.42)
10	4.5	1.0064	1.0062	1.0056	1.0046	1.0032	1.0014	(0,2.42)
	5.0	1.0051	1.0049	1.0044	1.0036	1.0025	1.0011	(0,2.42)

Table 5. The values of e_i for $i = 14$ and 15 (cont.)

n	β	$e_{15} = RE(T_{14}^*, T_{04})$					Range of λ in which T_{14}^* is efficient to T_{04}
		$\lambda = 1.00$	$\lambda = 1.25$ and $\lambda = 0.75$	$\lambda = 1.50$ and $\lambda = 0.50$	$\lambda = 1.75$ and $\lambda = 0.25$	$\lambda = 2.00$ and $\lambda = 0.00$	
2	2.5	1.2799	1.2578	1.1962	1.1058	1.0000	[0,2]
	3.0	1.1674	1.1553	1.1205	1.0669	1.0000	[0,2]
	3.5	1.1135	1.1056	1.0828	1.0467	1.0000	[0,2]
	4.0	1.0827	1.0772	1.0608	1.0346	1.0000	[0,2]
	4.5	1.0633	1.0591	1.0467	1.0267	1.0000	[0,2]
	5.0	1.0501	1.0469	1.0371	1.0213	1.0000	[0,2]
3	2.5	1.0832	1.0776	1.0611	1.0348	1.0000	[0,2]
	3.0	1.0543	1.0508	1.0402	1.0231	1.0000	[0,2]
	3.5	1.0385	1.0360	1.0286	1.0165	1.0000	[0,2]
	4.0	1.0288	1.0270	1.0215	1.0124	1.0000	[0,2]
	4.5	1.0224	1.0210	1.0167	1.0097	1.0000	[0,2]
	5.0	1.0180	1.0168	1.0134	1.0078	1.0000	[0,2]
4	2.5	1.0477	1.0446	1.0353	1.0203	1.0000	[0,2]
	3.0	1.0317	1.0297	1.0236	1.0136	1.0000	[0,2]
	3.5	1.0227	1.0213	1.0169	1.0098	1.0000	[0,2]
	4.0	1.0171	1.0160	1.0128	1.0074	1.0000	[0,2]
	4.5	1.0134	1.0125	1.0100	1.0058	1.0000	[0,2]
	5.0	1.0107	1.0101	1.0080	1.0047	1.0000	[0,2]
5	2.5	1.0278	1.0260	1.0207	1.0120	1.0000	[0,2]
	3.0	1.0187	1.0175	1.0140	1.0081	1.0000	[0,2]
	3.5	1.0135	1.0127	1.0101	1.0059	1.0000	[0,2]
	4.0	1.0101	1.0095	1.0076	1.0044	1.0000	[0,2]
	4.5	1.0080	1.0075	1.0060	1.0035	1.0000	[0,2]
	5.0	1.0065	1.0061	1.0048	1.0028	1.0000	[0,2]
6	2.5	1.0196	1.0183	1.0146	1.0085	1.0000	[0,2]
	3.0	1.0133	1.0124	1.0099	1.0058	1.0000	[0,2]
	3.5	1.0096	1.0090	1.0072	1.0042	1.0000	[0,2]
	4.0	1.0073	1.0068	1.0055	1.0032	1.0000	[0,2]
	4.5	1.0057	1.0054	1.0043	1.0025	1.0000	[0,2]
	5.0	1.0046	1.0043	1.0035	1.0020	1.0000	[0,2]
7	2.5	1.0138	1.0129	1.0103	1.0060	1.0000	[0,2]
	3.0	1.0094	1.0088	1.0070	1.0041	1.0000	[0,2]
	3.5	1.0068	1.0064	1.0051	1.0030	1.0000	[0,2]
	4.0	1.0052	1.0049	1.0039	1.0023	1.0000	[0,2]
	4.5	1.0041	1.0038	1.0031	1.0018	1.0000	[0,2]
	5.0	1.0033	1.0031	1.0025	1.0014	1.0000	[0,2]
8	2.5	1.0107	1.0100	1.0080	1.0046	1.0000	[0,2]
	3.0	1.0073	1.0068	1.0055	1.0032	1.0000	[0,2]
	3.5	1.0053	1.0050	1.0040	1.0023	1.0000	[0,2]
	4.0	1.0040	1.0038	1.0030	1.0018	1.0000	[0,2]
	4.5	1.0032	1.0030	1.0024	1.0014	1.0000	[0,2]
	5.0	1.0025	1.0024	1.0019	1.0011	1.0000	[0,2]

5. Conclusion

It is observed from Table 1 that the values of the relative efficiencies $e_i' s, i = 1, 2, \dots, 7$ of the proposed minimum mean squared error (MMSE) estimators $T_{0i}, i = 1, 2, 3, 4$ with respect to Lesitha and Thomas (2012) estimators $t_i' s, i = 1, 2, 3, 4$ respectively are greater than 'unity'. Thus, the proposed estimators $T_{0i}, i = 1, 2, 3, 4$ are more efficient than the corresponding usual estimators $t_i' s, i = 1, 2, 3, 4$ respectively.

It is further observed that the values of the relative efficiency e_5 of T_{04} with respect to T_{01} are the largest among $e_i' s, i = 1, 2, \dots, 7$, from which follows that the proposed MMSE estimator T_{04} is the best estimator among Lesitha and Thomas (2012) estimators $t_i' s, i = 1, 2, 3, 4$ and MMSE estimators $T_{0i}, i = 1, 2, 3, 4$.

Tables 2 to 5 demonstrate that for fixed (n, β) the values of relative efficiencies $e_i' s, i = 8, 9, \dots, 15$ increase as λ increases up to 1, while it decreases if λ goes beyond 'unity'. When the value of λ is unity (i.e. the guessed value α_0 coincides with the true value α) a higher gain in efficiency is seen. For fixed values of (n, λ) the values of $e_i' s, i = 8, 9, \dots, 15$ decrease as β increases. When (β, λ) are fixed the values of $e_i' s, i = 8, 9, \dots, 15$ also decrease as sample size n increases. A higher gain in efficiency is obtained when the sample size n is small. In general, the estimators $T_{li}^*, i = 1, 2, 3, 4$ are more efficient than Lesitha and Thomas (2012) estimators $t_i' s, i = 1, 2, 3, 4$ and MMSE estimators $T_{0i}, i = 1, 2, 3, 4$ respectively when $\lambda \in (0, 2.42)$. It is further observed that $T_{li}^*, i = 1, 2, 3, 4$ are respectively better than MMSE estimators $T_{0i}, i = 1, 2, 3, 4$ when $\lambda \in [0, 2]$.

Acknowledgement

The authors are highly grateful to the referees for their constructive comments/ suggestions that helped in the improvement of the revised version of the paper.

REFERENCES

- AHMAD, M. I., SINCLAIR, C. D., WERRITTY, A., (1988). Log-logistic flood frequency analysis. *J Hydrol*, 98, pp. 205–224.
- BALAKRISHNAN, N., MALIK, H. J., (1987). Best linear unbiased estimation of location and scale parameter of the log-logistic distribution. *Commun Stat Theory Methods*, 16, pp. 3477–3495.
- BENNETT, S., (1983). Log-logistic regression models for survival data. *J R Stat Soc, Ser C* 32, pp. 165–171.
- CHEN, Z., BAI, Z., SINHA, B. K., (2004). Ranked set sampling, theory and applications. *Lecture Notes in Statistics*, Springer, New York.
- FISK, P. R., (1961). The graduation of income distributions. *Econometrica*, 29, pp. 171–185.
- GESKUS, R. B., (2001). Methods for estimating the AIDS incubation time distribution when data of seroconversion is censored. *Stat Med*, 20, 795–812.
- LESITHA, G., THOMAS, P. Y., (2012). Estimation of the scale parameter of A LOG-LOGISTICS DISTRIBUTION. *METRIKA*, DOI 10.1007/S00184-012-0397-5.
- MCINTYRE, G. A., (1952). A method for unbiased selective sampling using ranked sets. *Aust J Agric Res*, 3, pp. 385–390.
- MEHTA, V., (2015). Estimation in Morgenstern Type Bivariate Exponential Distribution with Known Coefficient of Variation by Ranked Set Sampling. Proceeding of the “30th M. P. Young Scientist Congress” (MPYSC-2015), M. P. Council of Science and Technology, Vigyan Bhawan, Nehru Nagar, Bhopal -462 003, Madhya Pradesh, India.

- MEHTA, V., SINGH, H. P., (2014). Shrinkage Estimators of Parameters of Morgenstern Type Bivariate Logistic Distribution Using Ranked Set Sampling. *Journal of Basic and Applied Engineering Research (JBAER)*, 1 (13), pp. 1–6.
- MUTTLAK, H. A., (1997). Median ranked set sampling. *J Appl Stat Sci.* 6, pp. 245–255.
- RAGAB, A., GREEN, J., (1984). On order statistics from the log-logistic distribution and their properties. *Commun Stat Theory Methods*, 13, pp. 2713–2724.
- ROBSON, A., REED, D., (1999). Flood estimation handbook, 3. Statistical procedures for flood frequency estimation. Institute of Hydrology, Wallingford, UK.
- SHOUKRI, M. M., MIAN, I. U. M., TRACY, D., (1988). Sampling properties of estimators of log-logistic distribution with application to Canadian precipitation data. *Can J Stat*, 16, pp. 223–236.
- SINGH, H. P., MEHTA, V., (2013). An Improved Estimation of Parameters of Morgenstern Type Bivariate Logistic Distribution Using Ranked Set Sampling. *STATISTICA*, 73 (4), pp. 437–461.
- SINGH, H. P., MEHTA, V., (2016a). Improved Estimation of Scale Parameter of Morgenstern Type Bivariate Uniform Distribution Using Ranked Set Sampling. *Communications in Statistics – Theory and Methods*, 45 (5), pp. 1466–1476.
- SINGH, H. P., MEHTA, V., (2014a). Linear shrinkage estimator of scale parameter of Morgenstern type bivariate logistic distribution using ranked set sampling. *Model Assisted Statistics and Applications (MASA)*, 9, pp. 295–307.
- SINGH, H. P., MEHTA, V., (2014b). An Alternative Estimation of the Scale Parameter for Morgenstern Type Bivariate Log-Logistic Distribution Using Ranked Set Sampling. *Journal of Reliability and Statistical Studies*, 7 (1), pp. 19–29.
- SINGH, H. P., MEHTA, V., (2015). Estimation of Scale Parameter of a Morgenstern Type Bivariate Uniform Distribution Using Censored Ranked Set Samples. *Model Assisted Statistics and Applications (MASA)*, 10, pp. 139–153.

SINGH, H. P., MEHTA, V., (2016b). Some Classes of Shrinkage Estimators in the Morgenstern Type Bivariate Exponential Distribution Using Ranked Set Sampling. Hacettepe Journal of Mathematics and Statistics, 45 (2), pp. 575–591.

SINGH, H. P., MEHTA, V., (2016c). A Class of Shrinkage Estimators of Scale Parameter of Uniform Distribution Based on K- Record Values. National Academy Science Letters, 39, pp. 221–227.