

A Service of



Leibniz-Informationszentrum Wirtschaft Leibniz Information Centre for Economics

Garegnani, María Lorena; Gómez Aguirre, Maximiliano

Working Paper Forecasting inflation in Argentina

IDB Working Paper Series, No. IDB-WP-891

Provided in Cooperation with: Inter-American Development Bank (IDB), Washington, DC

Suggested Citation: Garegnani, María Lorena; Gómez Aguirre, Maximiliano (2018) : Forecasting inflation in Argentina, IDB Working Paper Series, No. IDB-WP-891, Inter-American Development Bank (IDB), Washington, DC, https://doi.org/10.18235/0001160

This Version is available at: https://hdl.handle.net/10419/208111

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.



ND https://creativecommons.org/licenses/by-nc-nd/3.0/igo/legalcode







IDB WORKING PAPER SERIES Nº IDB-WP-891

Forecasting Inflation in Argentina

Lorena Garegnani Maximiliano Gómez Aguirre

Inter-American Development Bank Department of Research and Chief Economist

June 2018



Forecasting Inflation in Argentina

Lorena Garegnani Maximiliano Gómez Aguirre

Central Bank of Argentina

Cataloging-in-Publication data provided by the Inter-American Development Bank Felipe Herrera Library

Garegnani, María Lorena.

Forecasting inflation in Argentina / Lorena Garegnani, Maximiliano Gómez Aguirre.

p. cm. — (IDB Working Paper Series ; 891)

Includes bibliographic references.

1. Inflation (Finance)-Argentina-Forecasting. 2. Inflation (Finance)-Argentina-Econometric models. I. Gómez Aguirre, Maximiliano. II. Inter-American Development Bank. Department of Research and Chief Economist. III. Title. IV. Series. IDB-WP-891

http://www.iadb.org

Copyright © 2018 Inter-American Development Bank. This work is licensed under a Creative Commons IGO 3.0 Attribution-NonCommercial-NoDerivatives (CC-IGO BY-NC-ND 3.0 IGO) license (<u>http://creativecommons.org/licenses/by-nc-nd/3.0/igo/</u> <u>legalcode</u>) and may be reproduced with attribution to the IDB and for any non-commercial purpose, as provided below. No derivative work is allowed.

Any dispute related to the use of the works of the IDB that cannot be settled amicably shall be submitted to arbitration pursuant to the UNCITRAL rules. The use of the IDB's name for any purpose other than for attribution, and the use of IDB's logo shall be subject to a separate written license agreement between the IDB and the user and is not authorized as part of this CC-IGO license.

Following a peer review process, and with previous written consent by the Inter-American Development Bank (IDB), a revised version of this work may also be reproduced in any academic journal, including those indexed by the American Economic Association's EconLit, provided that the IDB is credited and that the author(s) receive no income from the publication. Therefore, the restriction to receive income from such publication shall only extend to the publication's author(s). With regard to such restriction, in case of any inconsistency between the Creative Commons IGO 3.0 Attribution-NonCommercial-NoDerivatives license and these statements, the latter shall prevail.

Note that link provided above includes additional terms and conditions of the license.

The opinions expressed in this publication are those of the authors and do not necessarily reflect the views of the Inter-American Development Bank, its Board of Directors, or the countries they represent.



Abstract*

In 2016 the Central Bank of Argentina began to announce inflation targets. In this context, providing authorities with good estimates of relevant macroeconomic variables is crucial for making pertinent corrections in order to reach the desired policy goals. This paper develops a group of models to forecast inflation for Argentina, which includes autoregressive models and different scale Bayesian VARs (BVAR), and compares their relative accuracy. The results show that the BVAR model can improve the forecast ability of the univariate autoregressive benchmark's model of inflation. The Giacomini-White test indicates that a BVAR performs better than the benchmark in all forecast horizons. Statistical differences between the two BVAR model specifications (small and large-scale) are not found. However, looking at the RMSEs, one can see that the larger model seems to perform better for longer forecast horizons.

JEL classifications: C11, C13, C32, C53

Keywords: Bayesian Vector Autoregressive, Forecasting, Prior specification, Marginal likelihood, Small-scale and large-scale models

^{*} This study was undertaken within the framework of CEMLA's Joint Research Program 2017 coordinated by the Central Bank of Colombia. The authors thank counseling and technical advisory provided by the Financial Stability and Development (FSD) Group of the Inter-American Development Bank in the process of writing this document. The opinions expressed in this publication are those of the authors and do not reflect the views of CEMLA, the FSD group, the Inter-American Development Bank or the Central Bank of Argentina.

1. Introduction

Long-term nominal commitments such as labor contracts, mortgages and other debt are widespread features of modern economies. Forecasting how the general price level will evolve over the life of a commitment is an essential part of private sector decision-making.

The existence of long-term nominal obligations is also among the primary reasons economists generally believe that monetary policy is not neutral, at least over moderate horizons.

Central banks aim is to keep inflation stable, and perhaps also to keep output near an efficient level. With these objectives, the New Keynesian model makes explicit that optimal policy will depend on optimal forecasts (e.g., Svensson, 2005), and further, that policy will be most effective when it is well understood by the public.

Under "inflation targeting," central banks have generally released forecasts in quarterly "Inflation Reports" in order to be more transparent in their actions. The costs and benefits of transparency are widely debated, but the need for a central bank to be concerned with inflation forecasting is broadly agreed. In short, inflation forecasting is of the foremost importance to households, businesses, and policymakers.

During the year 2016, the Central Bank of Argentina has begun to announce inflation targets. In this context, providing the authorities with good estimates of relevant macroeconomic variables turns out to be crucial to making the pertinent corrections in order to reach the desired policy goals.

A standard tool in macroeconomics that is widely employed in forecasting is Vector Autoregressive (VAR) analysis. VARs are flexible time series models that can capture complex dynamic relationships among macroeconomic aggregates. However, their dense parameterization often leads to unstable inference and inaccurate out-of-sample forecasts, particularly for models with many variables, due to the estimation uncertainty of the parameters.

Litterman (1980) and Doan, Litterman, and Sims (1984) have proposed to combine the likelihood function (the data) with some informative prior distributions (the researcher's belief about the values of coefficients) to improve the forecasting performance of VAR models, introducing a Bayesian approach into VAR modeling.

In any Bayesian inference, a fundamental yet challenging step is prior specification, which influences posterior distributions of the unknown parameters and, consequently, the forecasts (Geweke, 2005). Fortunately, the literature has proposed some methodologies to set how informative the prior distributions should be.

Regarding prior selection, Litterman (1980) and Doan, Litterman, and Sims (1984) set the tightness of the prior by maximizing the out-of-sample forecasting performance of a small-scale model. Many authors follow this strategy, such as Robertson and Tallman (1999) and Wright (2009), and Giannone et. al (2014), who minimize the Root Mean Square Error (RMSE) of the forecasts.

On the other hand, Banbura, Giannone and Reichlin (2008) propose controlling the overfitting caused by the considerable number of variables in the model by selecting the shrinkage of the coefficients in such a way as to provide appropriate fitting "in-sample." Within this second selection strategy, we can find authors such as Giannone, Lenza and Primiceri (2012), Bloor and Mathenson (2009), Carriero, Clark and Marcellino (2015) and Koop (2011).

Banbura, Giannone and Reichlin (2008) showed that by applying Bayesian VAR methodology, they were able to handle large unrestricted VARs models and therefore they demonstrated that VAR framework can be applied to empirical problems that require the analysis of more than a few sets of time series. The authors showed that a Bayesian VAR is a viable alternative to factor models or panel VARs for analysis of large dynamic systems.

This paper develops a group of models to forecast inflation for Argentina, which includes autoregressive models, and different scale Bayesian VARs (BVAR), and compares their relative accuracy.

The paper is organized as follows. Section 2 presents the methodological aspects related to the application of Bayesian analysis in a VAR framework, and Section 3 presents a brief description of the data. Section 4 goes through the empirical results, and finally, Section 5 concludes.

2. Bayesian VAR Methodology

A VAR model has the following structure

$$\mathbf{y}_{t} = \mathbf{c} + \mathbf{B}_{1}\mathbf{y}_{t-1} + \ldots + \mathbf{B}_{p}\mathbf{y}_{t-p} + \mathbf{\varepsilon}_{t}, \tag{1}$$

where \mathbf{y}_t is a n × 1 vector of endogenous variables, $\boldsymbol{\varepsilon}_t \sim N(\mathbf{0}, \boldsymbol{\Sigma})$ is a n × 1 vector of exogenous shocks, **c** is a n × 1 vector of constants, \mathbf{B}_1 to \mathbf{B}_p are n × n matrices and $\boldsymbol{\Sigma}$ is n × n covariance matrix.

The BVAR coefficients are a weighted average of the prior mean (researcher's belief) and the maximum likelihood (ML) estimators (inferred from the data), with the inverse covariance of the prior and the ML estimators as weights.

Consider the following posterior distribution for the VAR coefficients

$$\boldsymbol{\beta} | \boldsymbol{\Omega} \sim N(\boldsymbol{\beta}_0, \boldsymbol{\Omega}^{-1} \boldsymbol{\xi}) \tag{2}$$

where the vector $\boldsymbol{\beta}_0$ is the prior mean (whose elements will represent the coefficient in equation (1), the matrix $\boldsymbol{\Omega}$ is the known variance of the prior and $\boldsymbol{\xi}$ is a scalar parameter controlling the tightness of the prior information. Even though $\boldsymbol{\Omega}$ could have many shapes, gamma and Wishart distributions are frequently used in the literature, since they ensure a normally distributed posterior.²

The conditional posterior of β can be obtained by multiplying the prior by the likelihood function. The posterior takes the form

$$\boldsymbol{\beta}|\boldsymbol{\Omega}, \mathbf{y} \sim N\left(\widehat{\boldsymbol{\beta}}(\boldsymbol{\xi}), \widehat{\mathbf{V}}(\boldsymbol{\xi})\right), \tag{3}$$

where

$$\widehat{\mathbf{\beta}}(\xi) \equiv \operatorname{vec}\left(\widehat{\mathbf{B}}(\xi)\right),$$
(4)

and

$$\widehat{\mathbf{B}}(\xi) \equiv (\mathbf{x}'\mathbf{x}\,\boldsymbol{\Sigma}^{-1} + (\boldsymbol{\Omega}\xi)^{-1})^{-1}(\mathbf{x}'\mathbf{y}\boldsymbol{\Sigma}^{-1} + (\boldsymbol{\Omega}\xi)^{-1}\boldsymbol{\beta}_0),\tag{5}$$

$$\widehat{\mathbf{V}}(\xi) \equiv (\mathbf{x}'\mathbf{x}\,\boldsymbol{\Sigma}^{-1} + (\boldsymbol{\Omega}\xi)^{-1})^{-1}.$$
(6)

Vectors **y** and **x** represent observed data while β_0 is a matrix where each column corresponds to the prior mean of each equation.

It is important to note that if we choose a large value for ξ , the prior will have little weight into the posterior. This translates to a large volatility of the prior and not enough information coming from the prior. On the other hand, if the ξ is set to a small value (i.e., close to zero), the

 $^{^{2}}$ If the posterior distributions are in the same family as the prior probability distribution, the prior and posterior are then called conjugate distributions.

prior becomes more informative and the posterior mean moves towards the prior mean. To see this point, we can express (5) as follows:

$$\widehat{\mathbf{B}}(\xi) \equiv \widehat{\mathbf{\Omega}}[\mathbf{\Omega}_0^{-1} \boldsymbol{\beta}_0 + (\boldsymbol{\Sigma}^{-1} \otimes \mathbf{x}') \mathbf{y}]$$
(7)

and

$$\widehat{\mathbf{\Omega}} = [\mathbf{\Omega}_0^{-1} + \mathbf{\Sigma}^{-1} \otimes \mathbf{x}' \mathbf{x}]^{-1}$$
(8)

If the second element between brackets in equation (7) is multiplied by $(\mathbf{x}'\mathbf{x})^{-1}(\mathbf{x}'\mathbf{x})$, we obtain the following equation

$$\widehat{\mathbf{B}}(\xi) \equiv \widehat{\mathbf{\Omega}} \big[\mathbf{\Omega}_0^{-1} \mathbf{\beta}_0 \big] + \widehat{\mathbf{\Omega}} \big[\mathbf{\Sigma}^{-1} \otimes \mathbf{x}' \mathbf{x} \, (\mathbf{x}' \mathbf{x})^{-1} \mathbf{x}' \mathbf{y} \, \big] \tag{9}$$

$$\widehat{\mathbf{B}}(\xi) \equiv \widehat{\mathbf{\Omega}} [\mathbf{\Omega}_0^{-1} \mathbf{\beta}_0] + \widehat{\mathbf{\Omega}} [\mathbf{\Sigma}^{-1} \otimes \mathbf{x}' \mathbf{x} \, \mathbf{\beta}_{\text{ols}}]$$
(10)

As can be seen, the posterior is a weighted average between the prior and the Ordinary Least Square (OLS) estimators,³ where the weights are the reciprocal of the prior covariance matrix and the reciprocal of the OLS covariance matrix respectively. As a result, if the information contained in the data is good enough to describe the process behind it, the posterior will move towards the OLS estimators. However, it is important to underscore that, even if the available series are adequate to describe the data generating process, the researcher could still formulate hypothesis about the distribution of the parameters based on his own beliefs. That would imply ignoring the information contained in the data, and usually that kind of decisions are based on strong beliefs.

The issue mentioned in the last paragraph demonstrates the need to be cautious about choosing the prior mean and the hyperpriors. In the following subsections, these aspects are discussed in more detail.

2.1 Level or Growth Rate

It is unclear a priori whether transforming variables into their growth rates can enhance the forecast performance of a BVAR model. On one hand, the level specification can better accommodate the existence of long-run (cointegrating) relationships across the variables, which would be omitted in a VAR in differences. On the other hand, Clements and Hendry (1996) have shown that in a

³ The OLS estimators of a VAR coincide exactly with the ML estimators conditional on the initial values.

classical framework, differencing can improve forecasting performance in the presence of instability.

There has been little effort in the BVAR literature to compare specifications in levels versus differences. Carriero, Clark and Marcellino (2015) work with this specific topic and found that models in growth rates generally yield more accurate forecasts than those obtained from the models in levels. However, we can find both approaches in the literature. Following the Litterman (1986) tradition, some authors considered BVARs with variables in levels (e.g., Banbura et al., 2008; Giannone et al., 2014; and Giannone, Lenza and Primiceri, 2012). Other authors used BVARs with variables in differences or growth rates (e.g., Clark and McCracken, 2007, and Del Negro et al. (2004).

As mentioned above, there is no apparent reason to opt for series in levels or in differences to work with; nevertheless, choosing a representation ex ante gives us information about the characteristics of the prior distribution (values of the mean prior). For example, working with variables in differences implies that the persistence of those variables should be low, and that one should impose a number close to zero as a prior mean of the first lag, denoting low persistence in the series.

Since it is a good practice to start with some idea about the value that the prior could take and encouraged by the evidence found by Carriero, Clark and Marcellino (2015), we have opted to work with variables in differences.

In the next subsection, we will treat the variance of the prior as another aspect of prior distribution.

2.2 Choice of Hyperparameters and Lag Length Strategy

To select the hyperparameters and the lag length we will follow the strategy suggested by Banbura, Giannone and Reichlin (2008), Carriero, Clark and Marcellino (2015) and Giannone, Lenza and Primiceri (2012). Suppose, that a model is described by a likelihood function $p(\mathbf{y}|\mathbf{\theta})$ and a prior distribution $p_{\mathbf{y}}(\mathbf{\theta})$, where $\mathbf{\theta}$ is a vector parameter of the model and \mathbf{y} is a vector of hyperparameters affecting the distribution of all the priors of the model. It is natural to choose these hyperparameters by interpreting the model as a hierarchical one, i.e. replacing $p_{\mathbf{y}}(\mathbf{\theta})$ with $p(\mathbf{\theta}|\mathbf{y})$ and evaluating their posterior (Berger, 1985; Koop, 2003). In this way, the posterior can be obtained by applying Bayes' law

$$p(\mathbf{\gamma}|\mathbf{y}) \approx p(\mathbf{y}|\mathbf{\gamma})p(\mathbf{\gamma}),$$
 (11)

where $p(\mathbf{\gamma})$ is the density of the hyperparameters and $p(\mathbf{y}|\mathbf{\gamma})$ is the marginal likelihood. In turn, the marginal likelihood is the density that comes from the data when the hyperparameters change, in other words, the marginal likelihood can be obtained after integrating out the uncertainty about the parameters in the model,

$$p(\boldsymbol{\gamma}|\boldsymbol{y}) = \int p(\boldsymbol{y}|\boldsymbol{\theta},\boldsymbol{\gamma}) p(\boldsymbol{\theta}|\boldsymbol{\gamma}) d\boldsymbol{\theta}.$$
 (12)

For every conjugate prior, the density $p(\mathbf{\gamma}|\mathbf{y})$ can be computed in closed form. To obtain the Bayesian hierarchical structure, it is necessary to obtain the distribution of $p(\mathbf{\theta})$ by integrating out the hyperparameters

$$p(\boldsymbol{\theta}) = \int p(\boldsymbol{\theta}, \boldsymbol{\gamma}) p(\boldsymbol{\gamma}) d\boldsymbol{\gamma}.$$
 (13)

More precisely, we can find different values of the prior distribution from different hyperparameter values and, in this way, we can represent the posterior as:

$$p(\boldsymbol{\theta}, \boldsymbol{\gamma} | \boldsymbol{y}) = p(\boldsymbol{y} | \boldsymbol{\theta}, \boldsymbol{\gamma}) p(\boldsymbol{\theta}, \boldsymbol{\gamma}) p(\boldsymbol{\gamma}).$$
(14)

The marginal likelihood should be sufficient to discriminate among models; in this sense, we can choose models with different hyperparameters and different likelihood specification (more precisely, lags length structure). To make this point operational, we estimate different models, following Giannone, Lenza and Primiceri (2012) who introduce a procedure allowing to optimize the values of the hyperparameters that maximize the value of the marginal likelihood of the model. This implies that the hyperparameter values are not set a priori but are estimated.

Then the marginal likelihood can be estimated for every combination of hyperparameter values within specified ranges and for different lag length structures, and the optimal combination is retained as the one that maximizes that value.

2.3 Comparison Strategy

In this subsection we present some details about our strategy for model comparison. We will mention the steps that we will follow to do it and then give more details about the predictive ability tests used for comparison:

- a) Estimate a univariate AR model.
- b) Compute the relative RMSE to the AR from a).
- c) Compute the relative RMSE to the BVAR.⁴
- d) Run the test of Giacomini and White (2006) to compare both models.

Our benchmark is a univariate model. This means that we have at hand different statistical measures that cover both the frequentist and the Bayesian approaches. While frequentist literature tends to compare the forecasts with actual values, Bayesian literature compares the realized values with the whole posterior predictive density.

The testing methodology of Giacomini and White (2006) consists on evaluating relative forecast accuracy with a Diebold-Mariano (1995) like test, but with one central difference: the size of the in-sample estimation window is kept fixed, instead of expanding. Using the sample observations available at time "t", forecasts of $\mathbf{y}_{t+\tau}$ are produced for different "t" for given " τ " periods into the future, with rolling windows of estimation with the two models that are being compared. The sequences of forecasts are then evaluated according to some loss function and then the difference of forecast losses is computed. This way, a time series of differences of forecast losses $\Delta L_{t+\tau}(\hat{\theta})$ that depends on the estimated parameters is constructed. The test then consists of a Wald test on the coefficients of the regression of that series against a constant, the unconditional version of the test in equation (15), or against other explanatory variables, the conditional version in equation (16),

$$\Delta \mathcal{L}_{t+\tau}(\hat{\theta}) = \mu + \varepsilon_t, \tag{15}$$

$$\Delta \mathcal{L}_{t+\tau}(\hat{\theta}) = \beta' X_t + \varepsilon_t. \tag{16}$$

Standard errors may be calculated using the Newey-West covariances estimator, controlling for heteroskedasticity and autocorrelation. In this paper, the unconditional version is used.

The Giacomini-White test⁵ has many advantages: it captures the effect of estimation uncertainty on relative forecast performance, it allows for comparison between either nested or non-nested models, and, finally, it is quite easy to compute.

⁴ The mean of the predictive density is considered.

⁵ See chapter 17 of Hashimzade and Thornton (2013) for a detailed discussion of this test.

2.4 Model Specification

We follow Banbura et al. (2008) and analyze two VAR models that incorporate variables of special interest, including indicators of real economic activity, consumer prices and monetary variables. We consider the following two alternative models:

Small-Scale Model. This is a small monetary VAR including three key variables:

- a) Prices: we used the consumer price index constructed by the National Institute of Statistics and Census of Argentina (INDEC). After December 2006 until July 2012, the previous series is linked with the evolution of the consumer price index provided Provincial Institute of Statistics and Census of the Province of San Luis and, after July 2012, series is again linked with the evolution of the consumer price index of the City of Buenos Aires.⁶
- b) Economic Activity: we used a monthly economic activity indicator known as EMAE (*Estimador Mensual de Actividad Económica*) published by INDEC. The EMAE is based on the value added for each activity at a base price plus net taxes (without subsidies), and it uses weights provided by Argentina's National Accounts (2004). It tries to replicate quarterly GDP at a monthly frequency.
- c) Interest Rate: we used data from the Central Bank of Argentina (BCRA) on 30 to 59-day fixed-term deposit rates.

Large-Scale Model. In addition to the variables included in the small-scale model, this version also includes the rest of variables in the data set. These are detailed in the next section.

In September 2016, Argentina transitioned to an inflation targeting regime. This could generate a structural break in the mean or/and variance. To account for this possible change in the mean of the process, we incorporate a dummy variable in both specifications (Marcelino and Mizon, 2000).⁷

As we compare models of different sizes, we need a strategy on how to choose the shrinkage hyperparameter as models become larger. As the dimension increases, we want more

⁶ From December 2006 to October 2015, the index by INDEC presented severe discrepancies with provincial and private price indexes and was therefore discard for that period.

⁷ In the Appendix, we show the posterior estimation of the whole sample to see the effect of this. We controlled the change in the mean due the transition to an inflation targeting regime and indeed obtained a significant coefficient in both models.

shrinkage, as suggested by the analysis in De Mol, Giannone and Reichlin (2008) to control for overfitting. We set the tightness of the prior for the model to have better in-sample fit; in this way, we are shrinking more in a larger dimension model.

3. Data

Our data set is composed of a group of 16 monthly macroeconomic variables for Argentina.

	Source	Source Description		Charae	cteristics
1	INDEC	EMAE	log	SA	Unit-Root
2	INDEC	CPI Inflation	-	-	Trend
3	INDEC	Core CPI Inflation (ex. seasonal and	-	-	Trend
4	INDEC	Industrial Employment	log	SA	Unit-Root
5	INDEC	Construction Employment	log	SA	Unit-Root
6	INDEC	Retail Trade Employment	log	SA	Stationary
7	BCRA	M2 Monetary Aggregate	log	SA	Unit-Root
8	BCRA	Multilateral Nominal Exchange Rate	log	-	Unit-Root
9	BCRA	30 to 59-day Deposit Rate	-	-	Unit-Root
10	INDEC	Imports of Intermediate Goods	log	SA	Unit-Root
11	INDEC	Total Exports	log	SA	Unit-Root
12	UTDT	Consumer Confidence Index		-	Unit-Root
13	INDEC	Monthly Supermarket Sales	log	SA	Unit-Root
14	AFCP	Cement Sales	log	SA	Unit-Root
15	MINEM	Asphalt Sales	log	-	Stationary
16	MERVAL	AL Stock Market Index		-	Unit-Root

 Table 1. List of Endogenous Variables

4. Results

4.1 Estimation of the BVAR Model

4.1.1 The Optimal Hyperparameters

We work with a Normal-Wishart BVAR specification. In this type of specification there are two hyperparameters and two parameters. We estimate the overall tightness λ_1 , lag decay λ_3 and the lag length as we have described in Section 2.2, and then we impose the value of the prior mean (the autoregressive coefficient) equal to zero as discussed earlier.

The hyperparameter of the overall tightness λ_1 is the standard deviation of the prior of all the coefficients in the system different from the constant. In other words, it determines how all the coefficients are concentrated around their prior means.

The term λ_3 is a decay factor, and $1/(L^{\lambda_3})$ controls the tightness on lag "L" relative to the first lag. Since the coefficients of higher order lags are more likely to be close to zero than those of lower order lags, the prior for the standard deviations of the coefficients decrease as the lag length increases. The values usually used in the literature are 1 or 2, so we settle for $\lambda_3 = 2$.

The prior variance of the parameters of $\hat{\beta}(\xi)$ is set according to:

$$\sigma_{ij}^2 = \left(\frac{1}{\sigma_j^2}\right) \left(\frac{\lambda_1}{L^{\lambda_3}}\right)^2,\tag{17}$$

where σ_j^2 denotes the OLS residual variance of the autoregressive coefficient for variable "j", λ_1 is an overall tightness parameter, "L" is the current lag, and λ_3 is a scaling coefficient controlling the speed at which coefficients for lags greater than 1 converge to 0.

For exogenous variables, we define the variances as:

$$\sigma^2 = (\lambda_1 \lambda_4)^2 \tag{18}$$

The results for the hyperparameters and prior means of the small and the large-scale model are shown in Table 2; as can be seen, all the hyperparameters are equal for both type of models excepted for the hyperparameter λ_1 .

The characteristics of our hyperparameters after the optimization procedure are as follows:

Hyperparameters Values	Large-Scale Model	Small-Scale Model
Autoregressive Coefficient:	0	0
Overall Tightness (λ ₁):	0.05	0.23
Lag Decay (λ ₃):	2	2
Exogenous Variable Tightness	1	1
Lag Length	1	1

Table 2. List of Hyperparameter Values

The hyperparameter λ_1 is equal to 0.05 for the large-scale model while the hyperparameter λ_1 for the small-scale is 0.23. From a practical point of view, this means that the "true" value of the coefficients estimated (posterior) is probably farther from the prior mean in the small-scale model than in the large-scale one.

Another aspect to consider about λ_1 , is the fact that this hyperparameter impacts on the distribution of the parameters of lagged endogenous and exogenous variables of each equation in the system. In this sense, with more shrinkage for example, it is less probable that the posterior coefficients of the lagged endogenous and exogenous variables depart from the prior.

As can be seen in Table 2, the posterior coefficients of the variables in the large-scale model are less probable to depart from the prior than the small-scale ones. Models with many variables will tend to have a better in-sample fit even when λ_1 is set to a "loose" value.

The posteriors obtained for the small- and the large-scale model of the inflation equation in each type of model are shown in the Appendix.

4.1.2 Forecasting Exercise

Our forecasting exercise is conducted in the following way. We estimate the hyperparameters considering the whole sample, through the maximization of the marginal likelihood; and then, we compute the forecasts.

As mentioned above, the data set goes from January 2004 to July 2017. We compute one, three and six-step-ahead forecasts with rolling windows. The size of the estimation sample is the same for each forecast horizon. Out-of-sample forecast accuracy is measured in terms of RMSE of the forecasts. Therefore, we obtained three RMSEs for each model.

Relative forecast accuracy is analyzed in Table 3, by computed the different combinations of RMSE ratios. On average, the BVAR presents better accuracy than the benchmark independently of the forecast horizon. For immediate horizons, the small-scale model slightly outperforms the large-scale model, but the large-scale model outperforms the small-scale model for longer forecast horizons.

Table 3. Relative Forecast Accuracy

One Step Ahead			Three Step Ahead		Six Step Ahead			
Ratio Small Model- Benchmark	Ratio Large Model- Benchmark	Ratio Large Model-Small Model	Ratio Small Model- Benchmark	Ratio Large Model- Benchmark	Ratio Large Model-Small Model	Ratio Small Model- Benchmark	Ratio Large Model- Benchmark	Ratio Large Model- Small Model
0.77	0.90	1.69	0.78	0.77	1.02	0.87	0.82	0.94

In the next subsection, we analyze these results with a Giacomini-White test.

4.2 Forecast Evaluation

To evaluate the predictive performance of the different models, we used the tests described earlier. Each column of Table 4 contains the probability value of Giacomini-White test statistic for the different models.

Forecast Horizon	Large BVAR vs. Benchmark	Small BVAR vs. Benchmark	Difference Between BVAR Models
One-Step-Ahead	0.03	0.01	0.29
Three-Steps- Ahead	0.00	0.00	0.49
Six-Steps-Ahead	0.09	0.05	0.41

Table 4. Giacomini-White Test

The result of the Giacomini-White test show that at a 5% significance level, the large BVAR model outperforms the benchmark for one step and three-step-ahead forecast horizon, while the small BVAR outperforms the benchmark at a 5% significance level for all forecast horizons. The last column of the table shows the Giacomini-White test applied to the differences in predictive ability between the small- and large-scale BVAR models, but in this case the differences are not significant for all forecast horizons.

5. Conclusions

This paper assesses the performance of Bayesian VAR to forecast inflation in Argentina. We considered a Normal Wishart BVAR specification for a small- and a large-scale model of differentiated variables setting the prior mean according to standard recommendations in previous studies. The overall tightness hyperprior and the lag length of the different models were set by optimization of the marginal likelihood. We found that large-scale models have narrower priors, giving more weight to the priors mean than small-scale models.

Overall, the results show that the BVAR model can improve the forecast ability of the univariate autoregressive benchmark's model of inflation. The Giacomini-White test indicates that a BVAR performes better than the benchmark in all forecast horizons. Statistical differences between the two BVAR model specifications (small and large-scale) are not found. However, looking at the RMSEs, one can see that the larger model seems to perform better for longer forecast horizons.

References

- Banbura M., D. Giannone and L. Reichlin. 2008. "Large Bayesian VARs." Working Paper 996. Frankfurt, Germany: European Central Bank.
- Berger, J. 1985. *Statistical Decision Theory and Bayesian Analysis*. New York, United States: Springer Series in Statistics.
- Blake, A., and H. Mumtaz. 2012. Applied Bayesian Econometrics for Central Bankers. Centre for Central Banking Studies Technical Handbook 4. London, United Kingdom: Bank of England.
- Bloor, C., and T. Matheson. 2009. "Real-Time Conditional Forecasts with Bayesian VARs: An Application to New Zealand." Reserve Bank of New Zealand Discussion Paper DP2009/02. Wellington, New Zealand: Reserve Bank of New Zealand.
- Carriero, A., T. Clark and M. Marcellino. 2015. "Bayesian VARs: Specification Choices and Forecast Accuracy." *Journal of Applied Econometrics* 30(1): 46–73.
- Clark, T., and M. McCracken. 2007. "Forecasting with Small Macroeconomic VARs in the Presence of Instabilities." Finance and Economics Discussion Series 2007-41. Washington, DC, United States: Board of Governors of the Federal Reserve System.
- Clements, M.P., and D.F. Hendry. 1996. "Multi-Step Estimation for Forecasting." *Oxford Bulletin* of Economics and Statistics 58(4): 657–684.
- De Mol, C., D. Giannone and L. Reichlin. 2008. "Forecasting Using a Large Number of Predictors: Is Bayesian Regression a Valid Alternative to Principal Components?" *Journal of Econometrics* 146(2): 318-328.
- Del Negro, M. et al. 2004. "On the Fit and Forecasting Performance of New Keynesian Models." FRB Atlanta Working Paper 2004-37. Atlanta, United States: Federal Reserve Bank of Atlanta.
- Diebold, F.X., and R.S. Mariano. 1995. "Comparing Predictive Accuracy." *Journal of Business* and Economic Statistics 13: 253-63.
- Doan, T., R. Litterman and C.A. Sims. 1984. "Forecasting and Conditional Projection Using Realistic Prior Distributions." *Econometric Reviews* 3: 1–100.
- Gelman, A. 2006. "Prior Distributions for Variance Parameters in Hierarchical Models." *Bayesian Analysis* 1(3): 515–534.
- Geweke, J. 2005. Contemporary Bayesian Econometrics and Statistics. Hoboken: Wiley.

- Giacomini, R., and H. White. 2006. "Test of Conditional Predictive Ability." *Econometrica* 74: 1545-1578.
- Giannone, D., M. Lenza and G.E. Primiceri. 2012. "Prior Selection for Vector Autoregressions." Working Paper ECARES 2012-002. Brussels, Belgium: Université Libre de Bruxelles (ULB).
- Giannone, D. et al. 2014. "Short-Term Inflation Projections: A Bayesian Vector Autoregressive Approach." *International Journal of Forecasting* 30(3): 635–644.
- Gneiting, T., and A. Raftery. 2007. "Strict Proper Scoring Rules, Prediction, and Estimation." Journal of the American Statistical Association 102(477): 359-378.
- Gneiting, T. 2011. "Making and Evaluating Point Forecasts." *Journal of the American Statistical Association* 106: 746–762.
- Hashimzade, N., and M.A. Thornton. 2013. *Handbook of Research Methods and Applications in Empirical Macroeconomics*. Cheltenham, United Kingdom: Edward Elgar.
- Koop, G. 2003. Bayesian Econometrics. Hoboken, United States: Wiley.
- Koop, G. 2011. "Forecasting with Medium and Large Bayesian VARs." SIRE Discussion Papers 2011-38. Edinburgh, United Kingdom: Scottish Institute for Research in Economics (SIRE).
- Litterman, R.B. 1979. "Techniques of Forecasting Using Vector Autoregressions." Minneapolis, United States: University of Minnesota. Doctoral dissertation.
- Litterman, R.B. 1980. "A Bayesian Procedure for Forecasting with Vector Autoregression." Working paper. Cambridge, United States: Massachusetts Institute of Technology, Department of Economics.
- Litterman, R.B. 1986. "Forecasting with Bayesian Vector Autoregressions: Five Years of Experience." *Journal of Business and Economic Statistics* 4(1): 25–38.
- Mao, W. 2010. "Bayesian Multivariate Predictions." Iowa City, United States: University of Iowa. Doctoral dissertation.
- Marcellino, M., and G. Mizon. 2000. "Modelling Shifts in the Wage-Price and Unemployment-Inflation Relationships in Italy, Poland and the UK." *Economic Modelling* 17: 387-413.
- Robertson, J.C., and E.W. Tallman. 1999. "Vector Autoregressions: Forecasting and Reality." *Federal Reserve Bank of Atlanta Economic Review* (Q1): 4–18.

- Svensson, L.E.O. 2005. "Monetary Policy with Judgment: Forecast Targeting." *International Journal of Central Banking* 1: 1–54.
- Waggoner, D.F., and T. Zha. 1999. "Conditional Forecasts in Dynamic Multivariate Models." *Review of Economics and Statistics* 81(4):639–651.
- Warne, A., G. Coenen and K. Christoffel. 2013. "Marginalised Predictive Likelihood Comparisons with Application to DSGE, DSGE-VAR, and VAR Models." CFS Working Paper Series, 478. Frankfurt, Germany: Goethe University, Center for Financial Studies.
- Wright, J.H. 2009. "Forecasting US Inflation by Bayesian Model Averaging." *Journal of Forecasting* 28(2): 131–144.

Appendix

Endogenous variables:	Inflation, Interest Rate, Real Activity
Exogenous variables:	Constant, Dummy 2016-11
Estimation sample:	July 2004 to July 2017
Sample size (omitting initial conditions):	156
Number of lags included in regression:	1
Prior:	Normal-Wishart
Autoregressive coefficient:	0
Overall tightness:	0.23
Lag decay:	2
Exogenous variable tightness:	1

Table 1A. Small BVAR Characteristics

Table 2A. Small BVAR Inflation Equation Coefficient Values

	Median	SD	LB	UB
INF(-1)	0.468	0.066	0.338	0.598
I(-1)	0.901	0.640	-0.356	2.157
Y(-1)	2.631	3.500	-4.237	9.499
Constant	0.280	0.071	0.140	0.420
d112016	-0.197	0.144	-0.479	0.086

Sum of squared residuals: 91.05 R-squared: 0.291 Adj. R-squared: 0.272

Endogenous variables	Inflation, Interest Rate, Real Activity, Multilateral Exchange Rate, Industrial Employment, Cement Sales, Asphalts Sales, Imports of Intermediate Goods, Total Exports, M2, Core Inflation, Construction Employment, Consumer Confidence Index, Supermarket Sales, Stock Market Index
Exogenous variables:	Constant, Dummy 2016-11
Estimation sample:	July 2004 to July 2017
Sample size:	156
Number of lags:	1
Prior:	Normal-Wishart
Autoregressive coefficient:	0
Overall tightness:	0.05
Lag decay:	2
Exogenous variable tightness:	1

Table 3A. Large BVAR Characteristics

	Median	SD	LB	UB
INF(-1)	0.145	0.045	0.057	0.234
I(-1)	0.436	0.407	-0.362	1.235
Y(-1)	1.177	2.131	-3.005	5.359
E(-1)	7.261	3.431	0.528	13.994
EMPI(-1)	16.644	11.611	-6.143	39.431
CEM(-1)	-0.680	0.556	-1.771	0.410
ASPH(-1)	0.083	0.411	-0.723	0.888
IMP(-1)	0.125	0.477	-0.810	1.061
EXP(-1)	0.091	0.491	-0.873	1.055
M2(-1)	4.093	2.410	-0.637	8.823
INFC(-1)	0.183	0.047	0.091	0.275
EMPC(-1)	-1.452	2.933	-7.207	4.303
ICC(-1)	-0.011	0.013	-0.036	0.013
SUP(-1)	2.243	1.322	-0.351	4.837
STK(-1)	0.133	1.110	-2.045	2.310
Constant	0.056	0.039	-0.021	0.132
d112016	-0.014	0.042	-0.096	0.067

 Table 4A. Large BVAR Inflation Equation Coefficient Values

Sum of squared residuals: 89.33 R-squared: 0.304 Adj. R-squared: 0.224