

Myerson, Roger B.

**Working Paper**

## An Extension of the Kakutani Fixed Point Theorem

Discussion Paper, No. 245

**Provided in Cooperation with:**

Kellogg School of Management - Center for Mathematical Studies in Economics and Management Science, Northwestern University

*Suggested Citation:* Myerson, Roger B. (1976) : An Extension of the Kakutani Fixed Point Theorem, Discussion Paper, No. 245, Northwestern University, Kellogg School of Management, Center for Mathematical Studies in Economics and Management Science, Evanston, IL

This Version is available at:

<http://hdl.handle.net/10419/220605>

**Standard-Nutzungsbedingungen:**

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

**Terms of use:**

*Documents in EconStor may be saved and copied for your personal and scholarly purposes.*

*You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.*

*If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.*

Discussion Paper No. 245

AN EXTENSION OF THE KAKUTANI  
FIXED POINT THEOREM

by

Roger B. Myerson

September 1976

# AN EXTENSION OF THE KAKUTANI FIXED POINT THEOREM

by Roger B. Myerson

Northwestern University, Evanston, IL 60201

In arguments using the Kakutani Fixed Point Theorem, special difficulties often arise for points on the boundary of the domain. In this paper we present an extension of the Kakutani Theorem which relaxes the restrictions on the map at such points.

Let  $S$  be a nonempty, convex, closed and bounded subset of a finite dimensional Euclidean space. Let  $S^*$  be the set of all subsets of  $S$ . Let  $S^{**}$  be the set of all subsets of  $S$  which are nonempty, convex, and closed. Let  $S^0$  be a nonempty subset of  $S$  such that

$$(1) \quad \forall x \in S, \forall y \in S^0, \frac{1}{2}(x+y) \in S^0.$$

For example,  $S^0$  may be the relative interior of  $S$ , which will always satisfy (1).

A point-to-set map  $F:S \rightarrow S^*$  is upper-semicontinuous iff  $x_n \rightarrow \bar{x}$ ,  $y_n \in F(x_n)$ , and  $y_n \rightarrow \bar{y}$  imply that  $\bar{y} \in F(\bar{x})$ .

KAKUTANI [1941] showed that, if  $F:S \rightarrow S^{**}$  is upper-semicontinuous, then there exists some  $\bar{x} \in S$  such that  $\bar{x} \in F(\bar{x})$ . Our extension of his theorem weakens the convexity assumptions on  $F(x)$ .

Theorem. If  $F:S \rightarrow S^*$  is upper-semicontinuous, and if  $F(y) \in S^{**}$  for all  $y \in S^0$ , then there exists an  $\bar{x} \in S$  such that  $\bar{x} \in F(\bar{x})$ .

Proof. Select any  $\bar{y} \in S^0$ . For any positive integer  $n$ ,

define  $F_n: S \rightarrow S^*$  by  $F_n(x) = F\left(\frac{2^n-1}{2^n}x + \frac{1}{2^n}\bar{y}\right)$ .

$F_n$  is clearly upper-semicontinuous. For any  $x \in S$ ,

$$\frac{2^n-1}{2^n}x + \frac{1}{2^n}\bar{y} = \frac{1}{2}(x + \frac{1}{2}(x + \dots + \frac{1}{2}(x + \bar{y}) \dots))$$

must be in  $S^0$ , so  $F_n(x) \in S^{**}$ . Hence  $F_n$  satisfies Kakutani's conditions; and so for each  $n$  there exists some  $x_n \in S$  such that  $x_n \in F_n(x_n)$ .

Since  $S$  is compact, we can find some  $\bar{x}$  and a convergent subsequence such that  $N(i) \rightarrow \infty$  and  $x_{N(i)} \rightarrow \bar{x}$  as  $i \rightarrow \infty$ .

Thus,  $x_{N(i)} \rightarrow \bar{x}$ ,  $x_{N(i)} \in F\left(\frac{2^{N(i)}-1}{2^{N(i)}}x_n + \frac{1}{2^{N(i)}}\bar{y}\right)$ , and  $\left(\frac{2^{N(i)}-1}{2^{N(i)}}x_n + \frac{1}{2^{N(i)}}\bar{y}\right) \rightarrow \bar{x}$ . By upper-semicontinuity of  $F$ , we

conclude that  $\bar{x} \in F(\bar{x})$ . Q.E.D.

Suppose  $F^0: S^0 \rightarrow S^{**}$  is an upper-semicontinuous map.

$F^0$  has a unique minimal upper-semicontinuous extension  $F: S \rightarrow S^*$  defined by:

$$(2) \quad F(x) = \left\{ z \mid \begin{array}{l} \text{there exist sequences } \{x_n\} \text{ and } \{z_n\} \text{ such} \\ \text{that } x_n \in S^0, x_n \rightarrow x, z_n \in F^0(x_n), \text{ and } z_n \rightarrow z \end{array} \right\}.$$

(Upper-semicontinuity assures us that  $F(x) = F^0(x)$  if  $x \in S^0$ .)

Then  $F$  may not be convex-valued outside  $S^0$ , but it will satisfy the hypotheses of our theorem. A fixed point of  $F$  can be interpreted as a limit of almost-fixed points of  $F^0$ .

Corollary For any upper-semicontinuous map  $F^0: S^0 \rightarrow S^{**}$ , there exist sequences  $\{x_n\}$  and  $\{z_n\}$  and an  $\bar{x} \in S$  such that  $x_n \in S^0$ ,  $z_n \in F^0(x_n)$ ,  $x_n \rightarrow \bar{x}$ , and  $z_n \rightarrow \bar{x}$ .

## REFERENCES

Kakutani, S., "A Generalization of Brouwer's Fixed Point Theorem,"  
Duke Mathematical Journal 8 (1941), pages 457 - 459.

## ABSTRACT

An extension of the Kakutani Fixed Point Theorem is presented, in which convexity restrictions are relaxed for points on the boundary of the domain.