

Kojima, M.; Saigal, R.

Working Paper

A Property of Matrices with Positive Determinants

Discussion Paper, No. 332

Provided in Cooperation with:

Kellogg School of Management - Center for Mathematical Studies in Economics and Management Science, Northwestern University

Suggested Citation: Kojima, M.; Saigal, R. (1978) : A Property of Matrices with Positive Determinants, Discussion Paper, No. 332, Northwestern University, Kellogg School of Management, Center for Mathematical Studies in Economics and Management Science, Evanston, IL

This Version is available at:

<https://hdl.handle.net/10419/220692>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.

DISCUSSION PAPER NO. 332

A PROPERTY OF MATRICES WITH POSITIVE DETERMINANTS

by

M. Kojima ^{*/}

R. Saigal

June 1978

The work of both authors was supported by Grant MSC77-03472 from the National Science Foundation.

^{*/} Department of Information Sciences, Tokyo Institute of Technology, Tokyo, Japan. The work of this author was done while visiting Northwestern University.

A Property of Matrices with Positive Determinants

M. Kojima*

R. Saigal

ABSTRACT

Let \mathcal{P} be the set of all $n \times n$ real matrices which have a positive determinant. We show here that at least 2^{n-1} matrices are needed to "see" each matrix in \mathcal{P} . Also, any finite subset of \mathcal{P} can be "seen" from a class of at most 2^{n-1} matrices in \mathcal{P} .

The work of both authors was supported by Grant MCS77-03472 from the National Science Foundation.

*Department of Information Sciences, Tokyo Institute of Technology, Tokyo, Japan. The work of this author was done while visiting Northwestern University.

A Property of Matrices with Positive Determinants

M. Kojima

R. Saigal

§1. Introduction

Let \mathcal{P} be the set of all $n \times n$ real matrices which have a positive determinant. Given two matrices A and B in \mathcal{P} , we say A can be seen from B , and vice-versa, if

$$(1-t)A + tB \text{ is nonsingular for every } t \text{ in } [0,1] \quad (1.1)$$

i.e., the line joining A and B lies inside \mathcal{P} . Also, we say a subset \mathcal{Q} can see B if for some A in \mathcal{Q} , (1.1) holds. In this note we consider the following question: What is the smallest set of matrices \mathcal{Q} from which each matrix in \mathcal{P} can be seen? It follows readily that $\mathcal{Q} \subset \mathcal{P}$. We give here a partial answer to the above question. We first show that $|\mathcal{Q}| \geq 2^{n-1}$, and also that any finite set of matrices in \mathcal{P} can be seen from a subset of cardinality at most 2^{n-1} .

Certain other properties of \mathcal{P} are known. Eaves [1] showed that \mathcal{P} is path connected, and Todd [7] showed that any two matrices in \mathcal{P} can be seen from a third (and thus considerably strengthened the result of [1]). Todd's work has implications for understanding the produce of the fixed point algorithms [1],[2],[5],[8]. This work is motivated by the recent use of property (1.1) by Kojima and Saigal [3],[4] in establishing conditions when PL mappings are homeomorphisms, and the use of property (1.1) in establishing PL approximations to diffeomorphisms, Saigal [6]. Since the set of all matrices with negative determinants is a nonsingular linear transformation of \mathcal{P} , all our methods are also valid for this set.

§2. Notation, Definitions and Preliminary Results

In this section, we present our notation and establish some preliminary results.

For an $n \times n$ matrix A , by $A(k)$ we represent the $k \times k$ submatrix consisting of the first k rows and k columns of A , for each $k = 1, \dots, n$, and call these the leading principal submatrices of A ; and their determinants the leading principal minors of A . A simple fact about these leading principal minors of A is the following

Lemma 2.1: Let $\mathcal{P}^0 = \{A: \det A(k) \neq 0 \text{ for each } k\}$. Then, \mathcal{P}^0 is open and dense in the space of all $n \times n$ matrices.

Proof: \mathcal{P}^0 is clearly open. Now, for $A \notin \mathcal{P}^0$, we note that for all sufficiently small $\epsilon > 0$, $A + \epsilon I \in \mathcal{P}^0$, and thus the denseness follows.

We now state a result, without proof, from the work of Saigal [5]:

Theorem 2.2 (Saigal): Let A and B be $n \times n$ matrices in \mathcal{P} . Then

$$\min_{0 \leq t \leq 1} \det[(1-t)A + tB] > 0 \quad (2.1)$$

if and only if

$$\inf_{\lambda \geq 0} \det[B^{-1}A + \lambda I] > 0. \quad (2.2)$$

Proof: See [5, Lemma 3.1.1].

(2.1) characterizes the situation when matrix A can see the matrix B in \mathcal{P} , and the theorem states that this is so if and only if $B^{-1}A$ has no negative real eigenvalues. Now, for $\delta > 0$, define an $n \times n$ diagonal matrix:

$$E(\delta) = \begin{bmatrix} \delta & & & \\ & \delta^2 & & \\ & & \ddots & \\ & & & \delta^{2n-1} \end{bmatrix} \quad (2.3)$$

and a polynomial

$$\begin{aligned}\phi(\lambda, \delta) &= \lambda^n + a_1(\delta)\lambda^{n-1} + \dots + a_{n-1}(\delta)\lambda + a_n(\delta) \\ &= \det(E(\delta)A + \lambda I).\end{aligned}\tag{2.4}$$

Then, we can prove:

Lemma 2.3: Let $\det(E(\delta)A + \lambda I) = \phi(\lambda, \delta)$, and

$$\text{sign } \det A(k) = \varepsilon_k \quad \text{for } k = 1, \dots, n,$$

where $\varepsilon_k \in \{-1, 1\}$. Then, there exist positive constants δ^* , b_k , c_k , $k = 1, \dots, n$, such that

$$\varepsilon_k b_k \delta^{2^k-1} \leq a_k(\delta) \leq \varepsilon_k c_k \delta^{2^k-1}.\tag{2.5}$$

Proof: The proof is by induction. For $n = 1$, we see that

$$\phi(\lambda, \delta) = \lambda + \delta \det A$$

and thus (2.5) holds with $b_1 = c_1 = |\det A|$. Now, assume that the result is true for $n = 1, \dots, r$, and consider the case when A is an $(r+1) \times (r+1)$ matrix. Let

$$A = \begin{bmatrix} \bar{A} & \begin{smallmatrix} 1 & b \\ - & - \end{smallmatrix} \\ a & \begin{smallmatrix} 1 & \gamma \\ - & - \end{smallmatrix} \end{bmatrix}.$$

Then

$$\begin{aligned}\phi(\lambda, \delta) &= \det(E(\delta)A + \lambda I) \\ &= \det \begin{bmatrix} \bar{E}(\delta)\bar{A} + \lambda I & \begin{smallmatrix} 1 & \bar{E}(\delta)b \\ - & - \end{smallmatrix} \\ \delta^{2^r}a & \begin{smallmatrix} 1 & \delta^{2^r}\gamma + \lambda \\ - & - \end{smallmatrix} \end{bmatrix} \\ &= \lambda \det(\bar{E}(\delta)\bar{A} + \lambda I) + \delta^{2^r} \det \begin{bmatrix} \bar{E}(\delta)\bar{A} + \lambda I & \begin{smallmatrix} 1 & \bar{E}(\delta)b \\ - & - \end{smallmatrix} \\ a & \begin{smallmatrix} 1 & \gamma \\ - & - \end{smallmatrix} \end{bmatrix}\end{aligned}$$

Denoting the first term by $\lambda \bar{\phi}(\lambda, \delta)$ and the second term by $\chi(\lambda, \delta)$, we note, from the induction hypothesis, that there exist $\bar{\delta}^*$, \bar{b}_k , \bar{c}_k , $k = 1, \dots, r$ such that

$$\bar{\phi}(\lambda, \delta) = \lambda^r + \bar{a}_1(\delta)\lambda^{r-1} + \dots + \bar{a}_r(\delta)\tag{2.6}$$

and $\varepsilon_k \bar{b}_k \delta^{2^{k-1}} \leq \bar{a}_k(\delta) \leq \varepsilon_k \bar{c}_k \delta^{2^{k-1}}$ for $k = 1, \dots, r$, and δ in $(0, \delta^*)$.

Also

$$\chi(\lambda, \delta) = \delta^{2^{r+1}-1} \varepsilon_{r+1} |\det A| + \delta^{2^r} \omega(\lambda, \delta) \quad (2.7)$$

where $\omega(\lambda, \delta)$ is a polynomial of degree r in λ , whose coefficients are polynomials in the variable δ , $\omega(0, \delta) = 0$ for all δ . Now, noting that $\phi(\lambda, \delta)$ is the sum of λ times (2.6) and (2.7), we have our result.

A consequence of Lemma 2.3 is the following:

Theorem 2.4: Let A be an $n \times n$ matrix in \mathcal{P} , and let

$$\text{sign } \det A(k) = \varepsilon_k, \quad k = 1, \dots, n,$$

where $\varepsilon_k \in \{-1, +1\}$. Then

(i) for some positive number $\delta^* > 0$ and all δ in $(0, \delta^*)$

$$\inf_{\lambda \geq 0} \phi(\lambda, \delta) > 0$$

if and only if $\varepsilon_k = +1$ for all $k = 1, \dots, n-1$.

(ii) If $\varepsilon_p = -1$, then there exists a $\delta^* > 0$ such that for all δ in $(0, \delta^*)$,

$$\inf_{\lambda \geq 0} \phi(\lambda, \delta) < 0.$$

Proof: Now, if $\varepsilon_k = +1$ for every k , then from Lemma 2.3, $\phi(\lambda, \delta) \geq \lambda^n$

+ $\sum_{k=1}^n b_k \delta^{2^{k-1}} \lambda^{n-k}$ for some $b_k > 0$, all δ in $(0, \delta^*)$, and $\lambda \geq 0$. Hence, for

$\lambda \geq 0$, $\phi(\lambda, \delta) > 0$. Thus, the if part of (i) follows. Now, let $\varepsilon_p = -1$ for some $p < n$, $\varepsilon_n = +1$. From Lemma 2.3, there exist positive numbers c_k and δ^*

such that $\phi(\lambda, \delta) \leq \lambda^n + \sum_{k=1}^n \varepsilon_k c_k \delta^{2^{k-1}} \lambda^{n-k}$, for all δ in $(0, \delta^*)$. Now, consider the above for $\lambda(\delta) = \delta^{2^{(p-1)} + \frac{1}{2}}$, for δ in $(0, \delta^*)$. Then

$$\phi(\lambda(\delta), \delta) = \sum_{k=0}^n \varepsilon_k c_k \delta^{2^{k-1}} \{\delta^{2^{p-1} + \frac{1}{2}}\}^{n-k} = \sum_{k=0}^n \varepsilon_k c_k \delta^{\omega(k)}$$

where $\omega(x) = (2^x - 1) + (n-x)(2^{(p-1)} + \frac{1}{2})$ for all $x \geq 0$, and $\epsilon_0 = c_0 = +1$. We note that $\omega(x)$ is convex, and that $\omega(p+1) - \omega(p) = 2^{p-1} - \frac{1}{2} \geq \frac{1}{2}$ and $\omega(p-1) - \omega(p) = \frac{1}{2}$. Hence $\omega(k) - \omega(p) \geq \frac{1}{2}$ for all $k \neq p$. Hence, we have

$$\phi(\lambda(\delta), \delta) = \delta^{\omega(p)} \left\{ \epsilon_p c_p + \sum_{k \neq p} \epsilon_k c_k \delta^{\omega(k) - \omega(p)} \right\}.$$

Since $\epsilon_p c_p < 0$, we note that $\phi(\lambda(\delta), \delta)$ is negative for all sufficiently small $\delta > 0$, and thus (ii) follows. Also, the only if part of (i) follows from the above.

§3. The Main Theorems:

In this section we prove our main results that if a class \mathcal{Q} of matrices can see any matrix in \mathcal{P} , then $|\mathcal{Q}| \geq 2^{n-1}$, and also that any finite subset of \mathcal{P} can be seen from a subset \mathcal{Q} of cardinality at most 2^{n-1} .

For this purpose, consider the class of diagonal matrices

$$\mathcal{D} = \{D: D_{ii} = +1 \text{ or } -1, D_{ij} = 0, i \neq j\} \subset \mathcal{P}.$$

Then, as can be readily confirmed, $|\mathcal{D}| = 2^{n-1}$.

We are now ready to prove our main theorem:

Theorem 3.1: Let $\mathcal{Q} \subset \mathcal{P}$ be a subset of m $n \times n$ matrices. Assume that $m < 2^{n-1}$. Then, there exists a matrix B in \mathcal{P} such that

$$\min_{0 \leq t \leq 1} \det[(1-t)A + tB] < 0 \text{ for all } A \text{ in } \mathcal{Q}.$$

Proof: From Lemma 2.1, since \mathcal{P}^0 is open and dense, there exists a matrix C such that $CA \in \mathcal{P}^0$ for each $A \in \mathcal{Q}$. Now, since $m < 2^{n-1}$, there exists a matrix $D \in \mathcal{D}$ such that if $A \in \mathcal{Q}$ then DCA has at least one negative leading principal minor. Hence, by Theorem 2.4(ii), for each $A \in \mathcal{Q}$, there exists a positive number δ_A such that for every $\delta \in (0, \delta_A]$,

$$\min_{\lambda \geq 0} \det[E(\delta)DCA + \lambda I] < 0.$$

Letting $\delta^* = \min \delta_A > 0$, and $B = (E(\delta^*)DC)^{-1}$, we obtain the desired result from Theorem 2.2.

We now prove our other main result:

Theorem 3.2: Let $\mathcal{Q} \subset \mathcal{P}$ be a finite set of matrices. Then, there exist $m \leq 2^{n-1}$ matrices B^1, \dots, B^m in \mathcal{P} such that if $A \in \mathcal{Q}$, then

$$\min_{0 \leq t \leq 1} \det[(1-t)A + tB^j] > 0 \text{ for some } j.$$

Proof: From Lemma 2.1, there exists a matrix C such that $CA \in \mathcal{P}^0$ for each A in \mathcal{Q} . Now, classify the matrices in $\{CA: A \in \mathcal{Q}\}$ by the equivalence relation $A \sim B$ if and only if $\text{sign det } A(k) = \text{sign det } B(k)$ for each $k = 1, \dots, n-1$. There are at most 2^{n-1} classes generated by the equivalence relation. Now, let $\mathcal{Q}_1, \dots, \mathcal{Q}_m$ be these classes. Consider \mathcal{Q}_r . For $A \in \mathcal{Q}_r$, $k = 1, \dots, n-1$, define

$$\varepsilon_k = \text{sign det } A(k)$$

and the matrix $D \in \mathcal{D}$ such that $D_{ii} = \varepsilon_{k-1} \varepsilon_k$ with $\varepsilon_0 = 1$. Then, $\text{sign det } [DA](k) = \varepsilon_k^2 > 0$, for $k = 1, \dots, n$, and A in \mathcal{Q}_r . From Theorem 2.4, there exists a δ_A such that for all δ in $(0, \delta_A]$, $\min_{\lambda \geq 0} \det[E(\delta)DA + \lambda I] > 0$. Now, define $\delta^* = \min_{A \in \mathcal{Q}_r} \delta_A$, and we note that all matrices in \mathcal{Q}_r can be seen from $B_r = [E(\delta^*)D]^{-1}$ (from Theorem 2.2), and we have our result.

As is evident from the proof of Theorem 3.2, the "sign matrices" in \mathcal{D} may not be sufficient, since some scaling matrix $E(\delta)$ is also involved. It can be readily verified that the matrices $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ and $\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$ can see all 2×2 matrices $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ in \mathcal{P} . The former can see any matrix for which $a+d \geq 0$, and the latter can see any matrix with $(a+d) < 0$. But we conjecture

that 2^{n-1} matrices are sufficient to see all $n \times n$ matrices in \mathcal{P} . Our approach appears to fail for this case. In the space of 3×3 matrices, for example, the 3×3 sign matrices in \mathcal{D} , namely

$$\begin{bmatrix} 1 & & \\ & 1 & \\ & & 1 \end{bmatrix}, \begin{bmatrix} -1 & & \\ & -1 & \\ & & 1 \end{bmatrix}, \begin{bmatrix} -1 & & \\ & 1 & \\ & & -1 \end{bmatrix}, \begin{bmatrix} 1 & & \\ & -1 & \\ & & -1 \end{bmatrix}$$

cannot see the matrix $\begin{bmatrix} -39 & 20 & 20 \\ 20 & -39 & 20 \\ 20 & 20 & -39 \end{bmatrix}$. (We are grateful to Todd [9] for this example.) Thus, in our approach, scaling is important.

References

- [1] Eaves, B. C., "A Short Course in Solving Equations with PL Homotopies," SIAM-AMS Proceedings, 9 (1976), 73-143.
- [2] Eaves, B. C. and R. Saigal, "Homotopies for Computation of Fixed Points in Unbounded Regions," Mathematical Programming, 3 (1972), 225-237.
- [3] Kojima, M. and R. Saigal, "A Study of PC^1 Homeomorphisms on Subdivided Polyhedrons," Submitted to SIAM Math. Analysis, April 1978.
- [4] Kojima, M. and R. Saigal, "On the Relationship between Conditions that Insure a Piecewise Linear Mapping is a Homeomorphism," Submitted to Math. of O.R., May, 1978.
- [5] Saigal, R., "On Paths Generated by Fixed Point Algorithms," Mathematics of Operations Research, 1 (1976), 359-380.
- [6] Saigal, R., "On Piecewise Linear Approximations to Smooth Mappings," Submitted to Math. of O.R., Dec. 1977.
- [7] Todd, M. J., "On the Jacobian of a Function at a Zero Computed by a Fixed Point Algorithm," To appear Math. of O.R.
- [8] Todd, M. J., The Computation of Fixed Points and Applications, Springer-Verlag, New York, 1976.
- [9] Todd, M. J., Private Communication, June 1, 1978.