

Yamamoto, Taku; Kunitomo, Naoto

Working Paper

Asymptotic Bias of Ordinary Least Squares Estimator for Multivariate Autoregressive Models

Discussion Paper, No. 504

Provided in Cooperation with:

Kellogg School of Management - Center for Mathematical Studies in Economics and Management Science, Northwestern University

Suggested Citation: Yamamoto, Taku; Kunitomo, Naoto (1981) : Asymptotic Bias of Ordinary Least Squares Estimator for Multivariate Autoregressive Models, Discussion Paper, No. 504, Northwestern University, Kellogg School of Management, Center for Mathematical Studies in Economics and Management Science, Evanston, IL

This Version is available at:

<https://hdl.handle.net/10419/220864>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.

DISCUSSION PAPER NO. 504
ASYMPTOTIC BIAS OF ORDINARY LEAST SQUARES
ESTIMATOR FOR MULTIVARIATE AUTOREGRESSIVE
MODELS

by

Taku Yamamoto */

and

Naoto Kunitomo **/

October 1981

*/

Department of Economics
Soka University
1-236 Tangi-cho
Hachioji-shi, Tokyo 192
Japan

**/

Department of Economics
Northwestern University
Evanston, Illinois 60201
U.S.A.

Asymptotic Bias of Ordinary Least Squares Estimator
for Multivariate Autoregressive Models

by

Taku Yamamoto

and

Naoto Kunitomo

Department of Economics
Soka University
1-236 Tangi-cho
Hachioji-shi, Tokyo 192
Japan

Department of Economics
Northwestern University
Evanston, Illinois 60201
U.S.A.

August 1981

Summary

Asymptotic bias is derived, as a relatively simple matrix function of the true parameters, for the ordinary least squares estimator in multivariate autoregressive models. The result is obtained through a convenient asymptotic expansion of the estimator.

Some key words: Asymptotic bias; Asymptotic expansion; Autoregressive model; Ordinary least squares estimator.

1. Introduction and Model

Kendall (1954) obtained the bias of the ordinary least squares estimator to order n^{-1} for the first order autoregressive model with a constant term, while White (1961) to orders higher than n^{-1} for the one without a constant term. In the present note, we derive the asymptotic bias to order n^{-1} for the ordinary least squares estimator of a multivariate autoregressive model with a constant term. It can be reduced to a model without a constant term as a special case, which corrects an error in Groenewald and de Waal (1979). While the asymptotic bias, whose precise definition is given in Section 2, is conceptually different from the approximation of the finite bias by Kendall and White, they are shown to be identical to order n^{-1} for both types of the first order multivariate autoregressive models.

Let us consider the p -variate first order autoregressive process

$$\underline{y}_t = \underline{B}\underline{y}_{t-1} + \underline{b}_0 + \underline{v}_t, \quad (1)$$

where $\underline{y}_t = [y_{t1}, y_{t2}, \dots, y_{tp}]'$, $\underline{v}_t = [v_{t1}, v_{t2}, \dots, v_{tp}]'$,

$\underline{B} = [\beta_{ij}]$ is the $p \times p$ coefficient matrix and $\underline{b}_0 = [\beta_{i0}]$ is the $p \times 1$ coefficient vector. We assume that \underline{v}_t is independently identically distributed as $N(0, \underline{\Omega}_v)$ and all characteristic roots of \underline{B} are less than unity in absolute value. Following Anderson (1959), the process is alternatively expressed as

$$\underline{z}_t = \underline{A}\underline{z}_{t-1} + \underline{u}_t, \quad (2)$$

where $\underline{z}_t = [y_t', 1]'$, $\underline{u}_t = [v_t', 0]$ with $\underline{\Omega}_u = E(\underline{u}_t \underline{u}_t')$, and \underline{A}

is the $(p+1) \times (p+1)$ matrix

$$\tilde{A} = \begin{bmatrix} \tilde{B} & \tilde{b}_0 \\ \tilde{0} & 1 \end{bmatrix}.$$

We can express z_t and $E(z_t z_t')$ as

$$z_t = \tilde{d} + \sum_{i=0}^{\infty} \tilde{A}_{p+1}^i u_{t-i}, \quad \tilde{\Gamma} = E(z_t z_t') = \tilde{d} \tilde{d}' + \sum_{i=0}^{\infty} \tilde{A}_{p+1}^i \Omega_u \tilde{A}_{p+1}^{i'}$$

where $\tilde{A}^0 = \tilde{I}_{p+1}, \tilde{I}_k$ is the identity matrix of rank k , and \tilde{d} is the $(p+1)$ th column of \tilde{A}^∞ , i.e., $\tilde{d} = \tilde{A}^\infty z_{-\infty} [((\tilde{I}_p - \tilde{B})^{-1} \tilde{b}_0), 1]'$

2. Asymptotic Bias of Ordinary Least Squares Estimator

For given observations y_0, y_1, \dots, y_n , the ordinary least squares estimator of \tilde{A} is given by

$$\hat{\tilde{A}} = \left(\sum_{t=1}^n z_t z_{t-1}' \right) \left(\sum_{t=1}^n z_{t-1} z_{t-1}' \right)^{-1}. \quad (5)$$

Here we assume that the initial observation y_0 obeys the same multivariate normal distribution as y_t for $t \geq 1$. As a generalization of Akahira (1979), we can expand $\hat{\tilde{A}} - \tilde{A}$ as

$$\begin{aligned} \hat{\tilde{A}} - \tilde{A} &= \left(\sum_{t=1}^n u_t z_{t-1}' \right) \left(\sum_{t=1}^n z_{t-1} z_{t-1}' \right)^{-1} \\ &= W \left(n^{-\frac{1}{2}} \tilde{I}_{p+1} + \tilde{Y} \right) \tilde{\Gamma}^{-1} \\ &= W \tilde{\Gamma}^{-1} \left(n^{-\frac{1}{2}} \tilde{I}_{p+1} - n^{-1} \tilde{Y} \right) + o_p(n^{-1}) \end{aligned} \quad (6)$$

where $W = n^{-\frac{1}{2}} \sum_{t=1}^n u_t z_{t-1}'$

$$\tilde{Y} = n^{\frac{1}{2}} (n^{-1} \sum_{t=1}^n z_{t-1} z_{t-1}' - \tilde{\Gamma}) \tilde{\Gamma}^{-1} = n^{-\frac{1}{2}} \left(\sum_{t=1}^n z_{t-1} z_{t-1}' \right) \tilde{\Gamma}^{-1} - n^{\frac{1}{2}} \tilde{I}_{p+1}.$$

Let us define the asymptotic bias of $\hat{\tilde{A}}$, denoted by $ABIAS(\hat{\tilde{A}})$, as follows:

$$\begin{aligned} \text{ABIAS}(\hat{\underline{A}}) &= \text{AE}(\hat{\underline{A}} - \underline{A}) \\ &= E\{\underline{W}\Gamma^{-1}(n^{-\frac{1}{2}}\underline{I}_{p+1} - n^{-1}\underline{v})\} + o(n^{-1}), \end{aligned} \quad (7)$$

where AE is the expectation of the asymptotic expansion of the distribution function of $\hat{\underline{A}}$.

Theorem: The asymptotic bias of the ordinary least squares estimator (5) for the first order multivariate autoregressive model with a constant term (1) is given by

$$\begin{aligned} \text{ABIAS}(\hat{\underline{A}}) &= -n^{-1}\underline{\Omega}_{\underline{u}}\left[\sum_{k=0}^{\infty} \underline{A}^k \Gamma^{-1}\{\underline{d}'\Gamma^{-1}\underline{d} + \underline{d}\underline{d}'\Gamma^{-1}\right. \\ &\quad \left.+ \left(\sum_{i=0}^{\infty} \underline{A}^i \underline{\Omega}_{\underline{u}} \underline{A}^{-i}\right)\underline{A}^{k+1}\Gamma^{-1}\right. \\ &\quad \left.+ \text{tr}(\underline{A}^{k+1}\left(\sum_{i=0}^{\infty} \underline{A}^i \underline{\Omega}_{\underline{u}} \underline{A}^{-i}\right)\Gamma^{-1})\}\right] + o(n^{-1}), \end{aligned} \quad (8)$$

Proof: See Appendix.

As easily seen, the above result includes a univariate p-th order autoregressive model as a special case. Specifically, for the univariate first order autoregressive model, we have

$$\underline{A} = \begin{bmatrix} \beta_1 & \beta_0 \\ 0 & 1 \end{bmatrix}, \quad \underline{\Gamma} = \begin{bmatrix} \omega^2(1-\beta_1^2)^{-1} + c^2 & c \\ c & 1 \end{bmatrix},$$

and $\underline{d} = [c, 1]'$, $c = \beta_0/(1-\beta_1)$, $\sum_{i=0}^{\infty} \underline{A}^i \underline{\Omega}_{\underline{u}} \underline{A}^{-i} = \omega^2(1-\beta_1^2)^{-1}\underline{M}$, $\omega^2 = E(u_t^2)$,

$\underline{M} = \underline{e}\underline{e}'$ and $\underline{e} = [1, 0]'$. Then, the first row of the first term in (8) reduces to $(1+\beta_1)[1, -c]$, while that of the second term vanishes.

The first rows of the third and fourth terms are simplified to the same vector $\beta_1[1, -c]'$. Thus, for the univariate first order autoregressive model with a constant term, we get

$$\text{ABIAS}(\hat{\beta}_1, \hat{\beta}_0) = n^{-1}(1+3\beta_1)[-1, \beta_0/(1-\beta_1)] + o(n^{-1}).$$

The result of $\text{ABIAS}(\hat{\beta}_1)$ conforms with Kendall (1954), while, as far as we know, $\text{ABIAS}(\hat{\beta}_0)$ has not previously been obtained.

The result (8) is also simplified for the case of no constant term.

Corollary: Let \hat{B} be the ordinary least squares estimator for the first order multivariate autoregressive model in (1) with a priori knowledge of $b_0 = 0$. Then, the asymptotic bias of \hat{B} is given by

$$\begin{aligned} \text{ABIAS}(\hat{B}) = & -n^{-1} \Omega_v^{-1} \left\{ \sum_{k=0}^{\infty} B^{*2k+1} \Gamma^{-1} + \sum_{k=0}^{\infty} B^{*k} \Gamma^{-1} \text{tr}(B^{*k+1}) \right\} \\ & + o(n^{-1}), \end{aligned}$$

$$\text{where } \hat{B} = \left(\sum_{t=1}^n y_t y_{t-1}' \right) \left(\sum_{t=1}^n y_{t-1} y_{t-1}' \right)^{-1}, \quad \Gamma = E(y_t y_t') = \sum_{i=0}^{\infty} B^i \Omega_v B^{*i}.$$

Proof: Since $d = 0$ in this case, the first two terms of (8) are dropped.

Further, replacing A and Ω_u by B and Ω_v in (8), and noting that

$$\Gamma = \sum_{i=0}^{\infty} B^i \Omega_v B^{*i} \text{ in this case, the proof is completed.}$$

We note that, while the second term of (9) is the same as the second term of equation (3.13) in Groenewald and de Waal (1979), the first term is different from their first term.

For the case of the univariate first order autoregressive model without a constant term, we have $B = B^* = \beta_1$ and $\Gamma = \sigma^2/(1-\beta_1^2)$. Thus the above result is reduced to

$$\text{ABIAS}(\hat{\beta}_1) = -(2\beta_1)/n,$$

which conforms with White (1961) to order n^{-1} .

Finally, it is easily seen from the proof in Appendix that all

5.

the results remain to hold, even if we replace the assumption on \underline{y}_0 by $\underline{y}_0 = 0$.

Appendix: Proof of Theorem

By the structure of \tilde{Y} , we can rewrite (6) as

$$\hat{\tilde{A}} - \tilde{A} = n^{-\frac{1}{2}}\{2\tilde{W}\Gamma^{-1}\} - n^{-1}\tilde{W}\Gamma^{-1}\{n^{-\frac{1}{2}}(\sum_{t=0}^{n-1} \tilde{z}_t \tilde{z}'_t)\Gamma^{-1}\} + o_p(n^{-1}).$$

Since $E(\tilde{w}) = 0$, $ABIAS(\hat{\tilde{A}})$ is reduced to

$$ABIAS(\hat{\tilde{A}}) = -n^{-1}E\{n^{-1}\sum_{t=1}^n \tilde{u}_t \tilde{z}'_{t-1}\Gamma^{-1}(\sum_{t=0}^{n-1} \tilde{z}_t \tilde{z}'_t)\Gamma^{-1}\} + o(n^{-1}). \quad (A.1)$$

Using the expression of \tilde{z}_t in (4), we can express $\sum_{t=1}^n \tilde{u}_t \tilde{z}'_{t-1}$ and $\sum_{t=0}^{n-1} \tilde{z}_t \tilde{z}'_t$ as

$$\sum_{t=1}^n \tilde{u}_t \tilde{z}'_{t-1} = \sum_{t=1}^n (\tilde{g}_t + \tilde{h}_t), \quad (A.2)$$

$$\sum_{t=0}^{n-1} \tilde{z}_t \tilde{z}'_t = \sum_{t=0}^{n-1} (\tilde{d}\tilde{d}' + \tilde{F}_t + \tilde{G}_t + \tilde{G}'_t + \tilde{H}_t + \tilde{H}'_t),$$

where $\tilde{g}_t = \tilde{u}_t \tilde{d}'$

$$\tilde{h}_t = \sum_{i=1}^{\infty} \tilde{h}_{t,i} = \sum_{i=1}^{\infty} \tilde{u}_t \tilde{u}'_{t-i} \tilde{A}^{i-1},$$

$$\tilde{F}_t = \sum_{i=0}^{\infty} \tilde{A}^i \tilde{u}_{t-i} \tilde{u}'_{t-i} \tilde{A}^{-i},$$

$$\tilde{G}_t = \sum_{i=0}^{\infty} \tilde{G}_{t,i} = \sum_{i=0}^{\infty} \tilde{d} \tilde{u}'_{t-i} \tilde{A}^{-i},$$

$$\tilde{H}_t = \sum_{i=0}^{\infty} \tilde{H}_{t,i} = \sum_{i=0}^{\infty} \tilde{A}^i \tilde{u}_{t-i} (\sum_{j=1}^{\infty} \tilde{u}'_{t-i-j} \tilde{A}^{i+j}).$$

Noting that $E\{(\tilde{u}_s \tilde{z}'_{s-1})(\tilde{z}_t \tilde{z}'_t)\} = 0$ for $s > t$, the expectation for given s is given by

$$\begin{aligned} & E\{\tilde{u}_s \tilde{z}'_{s-1}\Gamma^{-1}(\sum_{t=0}^{n-1} \tilde{z}_t \tilde{z}'_t)\Gamma^{-1}\} \\ &= E\{\tilde{u}_s \tilde{z}'_{s-1}\Gamma^{-1}(\sum_{t=s}^{\infty} \tilde{z}_t \tilde{z}'_t)\Gamma^{-1}\} - E\{\tilde{u}_s \tilde{z}'_{s-1}\Gamma^{-1}(\sum_{t=n}^{\infty} \tilde{z}_t \tilde{z}'_t)\Gamma^{-1}\}, \\ & \quad s=1, 2, \dots, n. \end{aligned} \quad (A.3)$$

We first evaluate the first term of the above. Since the expectation exists only when the time indices of u_t 's are equal or pairwise equal, it can be expressed by (A.2) as

$$\begin{aligned} & E\{u_{\underline{s}} z'_{\underline{s}-1} \Gamma^{-1} (\sum_{t=\underline{s}}^{\infty} z_t z'_t) \Gamma^{-1}\} \\ &= E\{g_{\underline{s}} \Gamma^{-1} (\sum_{t=\underline{s}}^{\infty} G_t) \Gamma^{-1} + g_{\underline{s}} \Gamma^{-1} (\sum_{t=\underline{s}}^{\infty} G'_t) \Gamma^{-1} \\ &+ h_{\underline{s}} \Gamma^{-1} (\sum_{t=\underline{s}}^{\infty} H_t) \Gamma^{-1} + h_{\underline{s}} \Gamma^{-1} (\sum_{t=\underline{s}}^{\infty} H'_t) \Gamma^{-1}\}. \end{aligned} \quad (A.4)$$

The first term of the above can be reduced to

$$\begin{aligned} E\{g_{\underline{s}} \Gamma^{-1} (\sum_{t=\underline{s}}^{\infty} G_t) \Gamma^{-1}\} &= E\{g_{\underline{s}} \Gamma^{-1} (\sum_{t=\underline{s}}^{\infty} \sum_{i=0}^{\infty} G_{t,i}) \Gamma^{-1}\} \\ &= E\{g_{\underline{s}} \Gamma^{-1} (\sum_{k=0}^{\infty} G_{\underline{s}+k,k}) \Gamma^{-1}\} \\ &= E\{u_{\underline{s}} d' \Gamma^{-1} \sum_{k=0}^{\infty} d u_{\underline{s}}' A^k \Gamma^{-1}\}. \end{aligned}$$

Since $d' \Gamma^{-1} d$ is a scalar, we get

$$E\{g_{\underline{s}} \Gamma^{-1} (\sum_{t=\underline{s}}^{\infty} G_t) \Gamma^{-1}\} = \Omega_u (\sum_{k=0}^{\infty} A^k) \Gamma^{-1} (d' \Gamma^{-1} d). \quad (A.5)$$

Similarly, the second term of (A.4) is reduced to

$$\begin{aligned} E\{g_{\underline{s}} \Gamma^{-1} (\sum_{t=\underline{s}}^{\infty} G'_t) \Gamma^{-1}\} &= E\{g_{\underline{s}} \Gamma^{-1} (\sum_{k=0}^{\infty} G'_{\underline{s}+k,k}) \Gamma^{-1}\} \\ &= E\{u_{\underline{s}} d' \Gamma^{-1} \sum_{k=0}^{\infty} A^k u_{\underline{s}}' d' \Gamma^{-1}\}. \end{aligned}$$

Since $d' \Gamma^{-1} \sum_{k=0}^{\infty} A^k u_{\underline{s}}$ is a scalar, first transposing it and then taking the expectation, we get

$$E\{g_{\underline{s}} \Gamma^{-1} (\sum_{t=\underline{s}}^{\infty} G'_t) \Gamma^{-1}\} = \Omega_u (\sum_{k=0}^{\infty} A^k) \Gamma^{-1} d d' \Gamma^{-1}. \quad (A.6)$$

The third term of (A.4) can be written as

$$\begin{aligned}
E\{h_s \Gamma^{-1} (\sum_{t=s}^{\infty} H_t) \Gamma^{-1}\} &= E\{h_s \Gamma^{-1} (\sum_{t=s}^{\infty} \sum_{i=0}^{\infty} H_{t,i}) \Gamma^{-1}\} \\
&= E\{\sum_{k=0}^{\infty} \sum_{i=1}^{\infty} h_{s,i} \Gamma^{-1} H_{s+k,k} \Gamma^{-1}\} \\
&= E\{\sum_{k=0}^{\infty} \sum_{i=1}^{\infty} u_s u'_{s-i} A^{i-1} \Gamma^{-1} A^k u_s u'_{s-i} A^{k+i} \Gamma^{-1}\}.
\end{aligned}$$

Since $u'_{s-i} A^{i-1} \Gamma^{-1} A^k u_s$ is a scalar, first transposing it and then

taking the expectation, we get

$$E\{h_s \Gamma^{-1} (\sum_{t=s}^{\infty} H_t) \Gamma^{-1}\} = \Omega_u \{ \sum_{k=0}^{\infty} A^k \Gamma^{-1} (\sum_{i=0}^{\infty} A^i \Omega_u A^{-i}) A^{k+1} \} \Gamma^{-1}. \quad (A.7)$$

The fourth term of (A.4) can be similarly written as

$$\begin{aligned}
E\{h_s \Gamma^{-1} (\sum_{t=s}^{\infty} H'_t) \Gamma^{-1}\} &= E\{\sum_{k=0}^{\infty} \sum_{i=1}^{\infty} h_{s,i} \Gamma^{-1} H'_{s+k,k} \Gamma^{-1}\} \\
&= E\{\sum_{k=0}^{\infty} \sum_{i=1}^{\infty} u_s u'_{s-i} A^{i-1} \Gamma^{-1} A^{k+i} u_{s-i} u'_s A^k \Gamma^{-1}\}.
\end{aligned}$$

Since $u'_{s-i} A^{i-1} \Gamma^{-1} A^{k+i} u_{s-i}$ is a scalar, moving it to the end and then taking the expectation, we get

$$E\{h_s \Gamma^{-1} (\sum_{t=s}^{\infty} H'_t) \Gamma^{-1}\} = \Omega_u [\sum_{k=0}^{\infty} A^k \Gamma^{-1} \text{tr}\{A^{k+1} (\sum_{i=0}^{\infty} A^i \Omega_u A^{-i}) \Gamma\}]. \quad (A.8)$$

Thus, the first term of (A.3) is

$$\begin{aligned}
&E\{u_s z'_{s-1} \Gamma^{-1} (\sum_{t=0}^{\infty} z_t z'_t) \Gamma^{-1}\} \\
&= \Omega_u [\sum_{k=0}^{\infty} A^k \Gamma^{-1} \{d \bar{\Gamma}^{-1} d + d d' \Gamma^{-1} + (\sum_{i=0}^{\infty} A^i \Omega_u A^{-i}) A^{k+1} \Gamma^{-1} \\
&\quad + \text{tr}(A^{k+1} (\sum_{i=0}^{\infty} A^i \Omega_u A^{-i}) \Gamma^{-1})\}], \quad s = 1, 2, \dots, n. \quad (A.9)
\end{aligned}$$

It is easily seen that the second term of (A.3) is $O(B^{n-s})$ for $s=1, 2, \dots, n$,

because $A^i u_t = [(B^i v_t), 0]'$. Since $O(B^{n-s}) = O(\rho^{n-s})$ where $0 \leq \rho < 1$ by the assumption on the characteristic roots, we have $\sum_{s=1}^n O(B^{n-s}) = O(1)$. Thus,

$$\begin{aligned}
& E\{n^{-1} \sum_{t=1}^n \tilde{u}_t \tilde{z}'_{t-1} \Gamma^{-1} (\sum_{t=0}^{n-1} \tilde{z}_t \tilde{z}'_t) \Gamma^{-1}\} \\
& = E\{\tilde{u}_s \tilde{z}'_{s-1} \Gamma^{-1} (\sum_{t=0}^{\infty} \tilde{z}_t \tilde{z}'_t) \Gamma^{-1}\} + o(n^{-1})
\end{aligned}$$

From (A.1) and (A.9), the proof is completed.

References

- Akahira, M.(1979). On the second order asymptotic optimality of estimators in an autoregressive process. Rep.Univ. Electro-Comm., 29, 213-8.
- Anderson, T.W. (1959). On asymptotic distributions of estimates of parameters of stochastic difference equations, Ann. Math. Statist., 30, 676-687.
- Groenewald, P.C.N. and de Waal, D.J. (1979). An asymptotic distribution for the coefficient in a multiple time series, South African Statis. J., 13, 15-28.
- Kendall, M.G. (1954). Note on bias in the estimation of autocorrelation. Biometrika, 41, 403-4.
- White, J.S. (1961). Asymptotic expansion for the mean and variance of the serial correlation coefficient. Biometrika, 48, 85-95.