

Saari, Donald G.

Working Paper

The Borda Dictionary

Discussion Paper, No. 821

Provided in Cooperation with:

Kellogg School of Management - Center for Mathematical Studies in Economics and Management Science, Northwestern University

Suggested Citation: Saari, Donald G. (1989) : The Borda Dictionary, Discussion Paper, No. 821, Northwestern University, Kellogg School of Management, Center for Mathematical Studies in Economics and Management Science, Evanston, IL

This Version is available at:

<https://hdl.handle.net/10419/221180>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.

Discussion Paper No. 821

THE BORDA DICTIONARY

by

Donald G. Saari*
Department of Mathematics
Northwestern University

January 1989

Abstract

For n candidates, a profile of voters defines a unique Borda election ranking for each of the $2^n - (n+1)$ subsets of two or more candidates. The Borda Dictionary is the set of all of these election listings that occur for any choice of a profile. As such, the dictionary contains all positive features, all flaws, and all paradoxes that can occur with sincere Borda elections. After the Borda Dictionary is characterized, it is used to show in what ways the Borda Count (BC) is an improvement over other positional voting methods and to derive several new BC properties. These properties include several new characterizations of the BC expressed in terms of axiomatic representations of social choice functions, as well as showing, for example, that the BC ranking of n candidates can be uniquely determined by the BC rankings of all sets of $k < n$ candidates for any choice of k between 2 and n .

*This research was supported, in part, by NSF grants IRI8415348, IRI-8803505 and a Fellowship from the Guggenheim Memorial Foundation.

The Borda Count (BC) is the simple method used to tabulate ballots where, for n candidates, $n-i$ points are assigned to a voter's i^{th} ranked candidate; the candidate with the most points wins. While the BC has both attractive features and (as a corollary of Arrow's Theorem) flaws, only some of them are known. To redress this situation, I characterize *everything* that could possibly happen when the BC is used. To understand this assertion, note that a profile determines a unique ordinal BC election ranking for each subset of candidates: i.e., associated with each profile is a unique listing of election rankings. In this paper I characterize all possible BC election listings that ever occur with some profile. This collection of all possible BC election listings is called a *Borda Dictionary*. By construction, the Borda Dictionary contains everything that could possibly occur with a BC election, so it catalogues all of the BC flaws, all of the BC paradoxes, and all of the BC positive features.

This goal of characterizing the Borda Dictionary continues my program, initiated in [3] (also see [4,5]), to characterize everything that could possibly occur with positional election procedures.¹ A conclusion of [3], repeated in Section 1.2, is that *the Borda method is the unique positional voting method to minimize the kinds and number of paradoxes that can occur*. This statement explains the partial results in the literature suggesting that the BC enjoys a favored status. It also follows from this conclusion that, in some sense, the BC plays a critical role within positional voting methods. As part of my development of the Borda Dictionary, I'll outline the mathematical reasons for this. Also, I show in what ways the BC is an improvement over other positional voting methods, and (although I defer a detailed discussion of the applications of the Borda dictionary for elsewhere) I indicate in Section 1.3 how these results apply to procedures such as runoff elections, etc.

The Borda Dictionary lists all of the possible listings of BC election outcomes, so one might correctly surmise that it can be used to derive and extend known BC conclusions. To illustrate this, I rederive Smith's conclusion [10] that the BC ranking of the candidates is related to the majority vote tallies of all pairs of candidates. As I show, Smith's result turns out to be just one of many possibilities: e.g., I show that the BC ranking of all n candidates is related to the BC rankings of all sets of m candidates if m satisfies $n > m \geq 2$.

1. This paper can be read independent of [3]. On the other hand, the proofs of several of my main results strongly depend upon arguments developed in [3].

as another example, if m satisfies $2 \leq m \leq n$, then there are relationships among the BC rankings of the set of all n candidates and all sets of m candidates. (There are other situations, and, elsewhere [8], I completely characterize them.) To illustrate a different kind of application of the Borda Dictionary, I show how to use it to significantly extend in many different directions those characterizations of the BC based on axiomatic properties of social choice functions. (These results, which start in Section 3.2, can be read independent of the technical Section 2.) Indeed, by using BC Dictionary, it turns out that a surprisingly large number of new BC conclusions can be found. Because it is impossible to provide an exhaustive listing of possible new results, I adopt the strategy of emphasizing how the BC Dictionary can be used to obtain new statements. As such, several of my examples not only illustrate certain points, but also offer new BC conclusions.

1.1. Notation and dictionaries

To motivate the notation and the dictionary, I start with two paradoxes.

Example 1. a. One of the oldest voting paradoxes, the *Condorcet Cycle*, occurs when, say, 5 voters have the rankings $c_1 \succ c_2 \succ c_3$, 5 have $c_2 \succ c_3 \succ c_1$, and 5 have $c_3 \succ c_1 \succ c_2$. By majority votes of 10 to 5, these voters prefer $c_1 \succ c_2$, and $c_2 \succ c_3$. Therefore, one might suspect that the voters prefer c_1 to c_3 . The paradox is that, by a vote of 10 to 5, they prefer $c_3 \succ c_1$; the voters' election rankings create a cycle.

b. A second example [3] has 6 voters with the ranking $c_3 \succ c_1 \succ c_2$, 5 with $c_2 \succ c_1 \succ c_3$, and 4 with $c_1 \succ c_2 \succ c_3$. The plurality ranking is $c_3 \succ c_2 \succ c_1$. If candidate c_1 were to withdraw, then it is common practice to assume that the electing group's ranking now is given by the truncated ranking of $c_3 \succ c_2$. However, by majority votes of at least 9 to 6, the majority vote ranking of each pair is the exact opposite of its relative ranking in the plurality outcome. By majority votes, these same, sincere voters prefer $c_1 \succ c_2$, $c_2 \succ c_3$, and $c_1 \succ c_3$.

Thus, a "paradox" is where a profile of voters (i.e., choices of rankings for each of the sincere voters) determines election rankings among the subsets of candidates that unveil an unexpected, counter-intuitive outcome. As indicated in (b), paradoxes illustrate that the way we use and interpret election rankings may be incorrect. So, to understand what actually can occur (i.e., to find all possible paradoxes of this kind), we need to determine all listings of election

rankings over all possible subsets of candidates that are associated with a profile. To start, list all feasible subsets of candidates. With the $n \geq 2$ candidates $C^n = \{c_1, \dots, c_n\}$, there are $2^n - (n+1)$ subsets with enough candidates (at least two) to permit an election. List these subsets as $\{S_1, \dots, S_{2^n - (n+1)}\}$ where, for convenience, the first $n(n-1)/2$ subsets are the pairs of candidates, the next $n!/3!(n-3)!$ are the sets of three candidates, etc. Also, in each S_i , list the candidates in the order determined by the subscripts. For example, with $C^3 = \{c_1, c_2, c_3\}$, the sets could be $S_1 = \{c_1, c_2\}$, $S_2 = \{c_1, c_3\}$, $S_3 = \{c_2, c_3\}$, and $S_4 = \{c_1, c_2, c_3\}$.

For each S_i , let R_i be the set of all complete, binary, reflexive, transitive rankings of S_i . Thus R_i is the listing of all possible ordinal election rankings associated with the S_i candidates. For instance with $S_4 = \{c_2, c_3, c_1\}$, $R_4 = \{c_2 > c_3, c_2 = c_3, c_3 > c_2\}$, while R_4 contains all $3!$ linear rankings of the three candidates in S_4 along with the 7 rankings that have a tie among the candidates. The *universal space*, $U^n = R_1 \times R_2 \times \dots \times R_{2^n - (n+1)}$, is a product space, so an element of U^n is a listing of $2^n - (n+1)$ rankings -- there is a ranking assigned to each subset of candidates. (For instance, $\{c_2 > c_1, c_1 > c_3, c_3 > c_2, c_1 = c_3 > c_2\}$ specifies a ranking for each subset of candidates, so it is an element of U^3 .) Thus U^n is the space of all possible, as well as all impossible coordinated (ordinal) election outcomes over the subsets of candidates.

A positional voting system for a set of candidates is where specified weights, $\{w_i\}$, are used to tally the ballots. In tabulating the ballots, w_i points are assigned to a voter's i^{th} ranked candidate, and the election ranking of each candidate is determined by the total number of points she receives. For k candidates, the assigned weights define a *voting vector* $W = (w_1, w_2, \dots, w_k)$ where $w_i \geq w_{i+1}$ and $w_1 > w_k$. Let W_i designate the voting vector assigned to tally the ballots for the candidates in S_i . The listing of the $2^n - (n+1)$ selected voting vectors, the *system voting vector* is

$$(1) \quad W^n = (W_1, W_2, \dots, W_{2^n - (n+1)}).$$

The obvious equivalence relationship among the voting vectors can be described with the vector E_k where, with k candidates, $E_k = (1, \dots, 1)$.

2. A *reversed positional voting system* is where $w_i \leq w_{i+1}$ and where the winning candidate is the one with the lowest total. With only minor changes, the conclusions of [3] and here hold for such systems. Indeed, the main difference is to permit negative values for "a" in Proposition 1. See [6,7] for more discussion.

Proposition 1. Let $\mathbf{W}$ be a voting vector for a set of k candidates. If a and b are scalars, then the ordinal election rankings of an election tallied with the voting vector $\mathbf{W}$ and with $a\mathbf{W} + b\mathbf{E}_k$ must always be the same.

This proposition holds because the factor $a\mathbf{W}$ just scales the final tally while $b\mathbf{E}_k$ just adds the same value to the tally of each candidate. Throughout this paper, I always use equivalence classes of voting vectors. For instance, for $k = 4$, both $(21, 18, 14, 10) = 4(3, 2, 1, 0) + 10(1, 1, 1, 1)$ and $(3, 1, -1, -3) = 2(3, 2, 1, 0) + 3(1, 0, 1, 1)$ are Borda vectors.

Definition. A *Borda voting vector* for $k \geq 3$ candidates is any voting vector equivalent to an $\mathbf{E}_k$ vector $(k-1, k-2, \dots, 0)$. A *Borda system voting vector*, denoted by $\mathbf{B}^k$, is where a Borda voting vector is used to tally the ballots for all subsets with three or more candidates, and where $(1, 0)$ is used for pairs of candidates.

In the obvious fashion, once a system voting vector, $\mathbf{W}^k$, is specified, then a profile of voters, $\mathbf{p}$, uniquely determines a listing of rankings. This listing, denoted by $f(\mathbf{p}; \mathbf{W}^k)$, consists of the election ranking for each subset of candidates.

Example 2. List the subsets of $\mathcal{C}^3$ as $\{(c_1, c_2), (c_1, c_3), (c_2, c_3), \{c_1, c_2, c_3\}\}$.

a. If the system voting vector is $\mathbf{W}^3 = (1, 0; 1, 0; 1, 0; 1, 0, 0)$, (i.e., a majority election is used for the first three subsets of candidates and a plurality election for S_3) and if $\mathbf{p}_a$ is the profile from Example 1.a, then $f(\mathbf{p}_a, \mathbf{W}^3) = (c_1 \succ c_2, c_3 \succ c_1, c_2 \succ c_3, c_1 = c_2 = c_3) \in \mathcal{U}^3$.

b. If $\mathbf{p}_b$ be the profile in Example 1b, then $f(\mathbf{p}_b, \mathbf{W}^3) = (c_1 \succ c_2, c_1 \succ c_3, c_2 \succ c_3, c_3 \succ c_2 \succ c_1) \in \mathcal{U}^3$.

c. $f(\mathbf{p}_a, \mathbf{B}^3) = (c_1 \succ c_2, c_3 \succ c_1, c_2 \succ c_3, c_1 = c_2 = c_3)$, while $f(\mathbf{p}_b, \mathbf{B}^3) = (c_1 \succ c_2, c_1 \succ c_3, c_2 \succ c_3, c_1 \succ c_2 \succ c_3)$.

The paradox in Example 1b disappears when the $\mathbf{B}^k$, rather than the plurality vote, is used to tally the ballots. As shown in Section 3, this is no coincidence.

Definition. Let $n \geq 2$ candidates be given. For a given $\mathbf{W}^0$, the *dictionary generated by $\mathbf{W}^0$* is the set

$$1.2 \quad D(\mathbf{W}^0) = \{ f(\mathbf{p}, \mathbf{W}^0) \mid \mathbf{p} \text{ is a profile with a finite number of voters} \}.$$

An entry in the dictionary – the sequence of election rankings over the various subsets of candidates – is called a *word*. Each ranking within a word is called a *symbol*.³

1.2. Characterization of Dictionaries

There is an important difference between a word in $D(\mathbf{W}^0)$ and an *element* of U^n . A word is a listing of election outcomes realized by a profile whereas an element of U^n is a sequence of rankings that may, or may not be achieved through an election. Consequently, if an element of U^n is not in $D(\mathbf{W}^0)$, it corresponds to a listing of rankings that never can be attained in an election tailored with the system vector $\mathbf{W}^0$. (For instance, the sequence from Example 1.b., $\{c_1 \succ c_2, c_1 \succ c_3, c_2 \succ c_3, c_3 \succ c_2 \succ c_1\}$, is in U^3 , but, as I show in Corollary 3.1, it is not in $D(\mathbf{B}^3)$. This means that such an election outcome never can occur with the BC.) Thus, a dictionary, $D(\mathbf{W}^0)$, catalogues everything that can happen with $\mathbf{W}^0$, while $U^n \setminus D(\mathbf{W}^0)$ specifies what the listings of rankings that never can occur with an election with $\mathbf{W}^0$.

By construction, $D(\mathbf{W}^0)$ is a subset of U^n . If a system voting vector $\mathbf{W}^0$ admits only a small number of inconsistencies and potentially undesired outcomes, then $D(\mathbf{W}^0)$ is a small subset of U^n . But, this is not the general situation. The following theorem summarizes those results from [3] basic for my current discussion. Recall that $\mathbf{W}^0$ is a vector in a Euclidean space and that an algebraic set is a lower dimensional subset of this Euclidean space determined by the zeros of a given set of polynomials.

Theorem 1 [3] a. Let $n \geq 3$ candidates be given. With the exception of an algebraic subset, a^n , of possible choices for system voting vectors,

$$1.3 \quad D(\mathbf{W}^0) = U^n,$$

b. for $n = 3$, the only system voting vector in a^3 is $\mathbf{B}^3$. Namely, $D(\mathbf{B}^3)$

3. This useful terminology correctly suggests that a dictionary serves as a starting point for the analysis of election procedures. However, I adopted this term, and others, as a natural extension of the fact that the initial motivation for this direction of research came from "Chaos" and "Symbolic dynamics" in dynamical systems. See [5,6] for an exposition of the connection.

is a proper subset of U^0 , and if $W \neq B^0$, then $D(W^0) = U^0$.

c. For all $n \geq 3$, if $W^0 \neq B^0$, then $D(B^0)$ is a proper subset of $D(W^0)$.

d. Let $n \geq 3$. Let w be a word from $D(B^0)$. There exists a profile of voters, p , so that for any choice of a system voting vector, W^0 , $f(p, W^0) = w$.

Part a asserts that for almost all system voting vectors, anything can happen. To appreciate the implications of this statement, remember that while many election paradoxes in the literature compare election outcomes over different subsets of candidates, most of them involve only a few of the symbols of a word. Part a asserts that far more startling paradoxes exist. To create one, just fill in the remaining symbols of the word in *any desired manner*, and Theorem 1 ensures there is a profile to support this conclusion. Thus, by using Theorem 1 it becomes trivial to extend all such paradoxes from the literature in all possible ways. (See [3].) Indeed, part a guarantees that the wildest imaginable situations actually occur. As described in [5], one could even use a random number generator to determine a ranking for each subset of candidates, and, for almost all system voting vectors, there is a profile so that the randomly selected rankings are the sincere election outcomes of these voters.

This conclusion is disturbing: it is difficult to accept that a voting procedure reflects the voters' true wishes when the outcomes can radically change with even minor changes in which group of candidates just happens to be considered. Since part a asserts this can be the case for almost all voting vectors, the crucial issue is to find a method that permits some consistency among the election rankings. Part b asserts that with three candidates, the only possible way relief can be attained is with the BC: only the BC offers protection from all imaginable inconsistencies and paradoxes. This is illustrated in Example Three where the plurality ranking is in direct conflict with the same voters' majority vote rankings of the pairs of the candidates while, with the same profile, the BC ranking is consistent with the pairwise rankings.

As I will show elsewhere, part b of Theorem 1 does not extend: for all $n \geq 4$, there are choices of $W^0 \neq B^0$ where $D(W^0)$ is a proper subset of U^0 . However, even if $D(W^0)$ is a proper subset of U^0 , part c asserts that $D(B^0)$ always is a proper subset of $D(W^0)$. In other words, B^0 is the unique choice of a positional voting method to minimize not only the number, but also the kinds of paradoxes. Thus, for instance, a situation (a word) illustrating a negative feature of the BC also must occur with all other choices of positional voting methods. Part d

strengthens this statement by asserting that associated with each word in the Borda Dictionary is a profile whereby this word (this same feature) is realized for *all possible positional voting methods*.

1.2 Social Choice Procedures and Borda

To illustrate the central role played by the BC, I discuss an undesirable property of some of those procedures, such as tournaments, various kinds of runoff elections, and agendas, that involve the rankings of several subsets of candidates. These *social choice* procedures (i.e., instead of ranking the candidates, the procedure finds the "best" candidate or candidates) start by ranking specified initial subsets of candidates. Then, the rankings and the rules of the procedures determine which candidates are ranked at the next stage. For instance, bottom ranked candidates may be dropped from further consideration, as in a runoff election; replaced with other candidates, as with an agenda; or matched against other "winning" candidates, as in certain kinds of tournaments. This process continues until a final set of candidates is ranked. The chosen candidate(s) is determined by the rankings of the final set.

The typical way these social choice procedures are analyzed is to find a profile to prove that certain suspected properties occur. However, constructing an appropriate profile is a difficult combinatoric task. (This explains why many results use only small values of n and the plurality vote.) Note, the purpose of finding such a profile is to establish the existence of a particular word. But as the dictionaries catalogue all possible words, this complicated combinatoric step of profile design now can be eliminated. Indeed, by treating such social choice procedures as a *mapping* from a dictionary to the non-empty subsets of C^n , denoted by EP^n , the analysis becomes both more complete and much simpler. In other words, the dictionary -- the domain of the mapping -- replaces the the profiles as the primitive in the analysis.

To illustrate an advantage of taking this approach, I offer an important new result. Namely, *the possible outcomes of a given social choice procedure based on the Borda rankings are more restrictive than those based on any other positional voting ranking*. This statement, which is difficult to prove if profiles are the primitives, is an immediate consequence of the assertion that the Borda Dictionary is a proper subset of any other positional voting dictionary. Now, the social choice procedure is restricted to a smaller domain,

What this assertion implies is that one can expect more consistence among social choice procedures if their outcomes are based on the BC rankings rather than the rankings of any other positional voting method.

This new assertion about the BC is illustrated below with Corollary 1.1, which extends aspects of Corollary 6.1 in [3]. The goal is to understand which social choice procedures admit the troubling "abstention paradox" whereby, by abstaining, a voter forces the final result to be personally more favorable than had he voted. By using a dictionary, this issue can be analyzed for a large class of social choice procedures -- including runoff elections, tournaments, and agendas -- rather than just individual procedures as typically is the case when profiles are the primitive.

Definition A social choice method is an assignment of profile to a set in $P(C^n)$; that is, a nonempty subset of candidates from $C^n = \{c_1, c_2, \dots, c_n\}$. A social choice method based on the W^n rankings of subsets of the candidates is a mapping

1.4 $f: D(W^n) \rightarrow P(C^n)$.

A social choice method is *disjoint* if

- i) the method is based on the positional voting rankings, W^n , of the candidates,
- ii) there is a subset of candidates, called the *swing set*, and a ranking of this set whereby the reversal of two adjacently ranked candidates changes the choice of the final set of candidates to be ranked,
- iii) the selected candidate is based on the ranking of the final set of candidates, and
- iv) the image of f contains at least two different outcomes.

In a runoff election, any set of candidates, other than the final one, is a "swing set." This is because the reversal of the relative rankings of two candidates, one just above the cutoff and the other just below, changes which set of candidates are advanced.

In [1] I show *for almost all choices of W^n* (i.e., those that are not in $\emptyset^n$), and *all disjoint social choice methods admit the abstention paradox*.

Namely, there is a profile p and two additional voters with identical rankings whereby if the two voters vote, the outcome will be personally less favorable

4. I don't know the history of this paradox, but at least the flavor of it is described in Smith's paper [10]. It may have been known much earlier.

them all they abstain. The next question is to determine whether a disjoint method can avoid this conclusion with a $\mathbf{W}^0 \in \mathcal{W}^0$.

Corollary 1.1. For $n \geq 3$, if a disjoint social choice method based on the B^0 rankings admits the abstention paradox, then the abstention property holds for all choices $\mathbf{W}^0$. In particular, all runoff elections have this property.

This corollary underscores a basic consequence of Theorem 1: if a “negative” election phenomena occurs with the BC, then it probably occurs with all positional voting methods. Thus, to understand a social welfare or social choice procedure based on positional voting methods, the analysis must start with the Borda Dictionary to determine what can and cannot occur.

Outline of proof for Corollary 1.1. (The proof is in Section 4.) To outline the ideas, I show that for $n = 3$, the BC runoff election has the abstention property. The basic ideas extend to any BC runoff election and for $n \geq 3$. By using results in Section 3, it follows that $(c_1 \succ c_2, c_3 \succ c_1, c_2 \succ c_3, c_1 \succ c_3 = c_2) \in D(B^0)$. This means there is a profile that yields this word, and, of course, any fixed scalar multiple of the number of voters with the same ranking yields the same outcome. As shown in Theorem 6 of [3], this profile can be selected so that at least one of these voters has the ranking $c_2 \succ c_3 \succ c_1$. Add another voter with the identical ranking. Choose the scalar multiple determining the replicated number of voters to be large enough so that the outcomes of the pairwise elections are not affected whether or not these two voters vote. On the other hand, if these two voters abstain, then the ranking of the swing set, the set of all candidates, is $c_1 \succ c_3 \succ c_2$, and c_3 wins the runoff between c_1 and c_2 . But, should these two voters vote, the outcome is $c_1 \succ c_2 \succ c_3$, and the undesired c_1 wins the runoff. This completes the proof. The general proof, in Section 4 is similar. It just involves using those words in words in a dictionary (those points in the domain of the social choice procedure) where, by changing from one to the other, the final outcome changes.

2. How much of an improvement, and what representation?

Based on the results stated above, the BC appears to be superior to any other positional voting method -- at least with respect to the kinds of issues addressed by Theorem 1. (See [3] for a more detailed discussion.) A shortcoming

of Theorem 1 is that it does not indicate whether Borda's method provides only a marginal, or a significant improvement over any other positional voting system. If the difference between $D(W^0)$ and $D(B^0)$ is only a small number of words, then the advantage achieved by using Borda's method may not be of any importance. After all, this would imply that the BC merely avoids a small number of paradoxes.

The BC is a significant improvement; I show in this section that the Borda Dictionary has far fewer words than any other dictionary. For instance, for 5 candidates the general situation is that

$$2.1 \quad 10^{11} \leq |D(B^5)| \leq 5 \leq |D(W^5)|,$$

and for six candidates

$$2.2 \quad 10^{14} \leq |D(B^6)| \leq 5 \leq |D(W^6)|.$$

For instance, suppose for six candidates that each of the $2^6 - 1 = 57$ possible subsets of two or more candidates are plurality ranked. It follows from Eq. 2.2 that, on the average, *each* Borda word must be replaced with at least 10^{14} different words to complete the plurality election dictionary. To appreciate this number, recall that the projected supercomputers will perform about 10^{12} operations per second, and that there have been about 10^{47} seconds of time since the "Big Bang". Thus, if such a supercomputer started at the Big Bang to list the words that replace just *one* Borda word from $D(B^6)$, then, at the very best, the computer would only be $1/10^{33}$ through.

The large multiplies in these inequalities not only underscore the point that the BC avoids a shockingly large number of paradoxes², but also why it is impossible to list the entries in a dictionary. To circumvent the listing problem, I introduce a geometric approach to characterize the words in a dictionary, where I concentrate on the Borda Dictionary. This vector space representation characterizes all possible tallies for ballots, so it describes all cardinal relationships relating the election tallies among different subsets of candidates. Consequently, this approach allows one to address issues such as: what profiles define certain specified words in $D(B^0)$, what percentage of the R points cast does a candidate need to acquire in order to avoid (or to achieve) certain W properties, etc., etc. This is illustrated, in part, in Section 3.

This geometric representation for election tallies, partially developed

5. On the other hand, because of the large values, one might wonder whether most of the words in a dictionary require profiles with more voters than admitted by the population of the world. As indicated in [4], small numbers of voters suffice.

in 15.6.71, is critical for what follows. To simplify the exposition, first consider the set $S_2^{n-(n+1)} = \{c_1, \dots, c_n\}$. In the n -dimensional Euclidean space, E^n , identify the k^{th} component, x_k , with the k^{th} candidate, c_k , in the following manner. For $x = (x_1, \dots, x_n)$, let larger values of x_i denote a "stronger" preference for c_i . With this identification, the hyperplane $x_k = x_i$ divides E^n into three regions: the two half spaces are identified with the strict ordinal rankings (e.g., $x \in E^n$ satisfying $x_k > x_i$ corresponds to the ordinal ranking $c_i > c_j$), and the hyperplane is identified with indifference between the two candidates. By allowing the choices for i and j to vary over all pairs of indices, the resulting $n(n-1)/2$ hyperplanes divide E^n into cones that represent all possible ordinal rankings of the n candidates. Call each of these regions a *ranking region*. In this way, each ranking of the n candidates corresponds to a unique ranking region. For instance, $\{x \in E^n \mid x_2 > x_1 = x_3 > x_3 > x_1\}$ is identified with the ranking $c_2 > c_1 = c_3 > c_3 > c_1$. The line passing through the origin of E^n and $(1,1,\dots,1)$ represents complete indifference among the candidates: this line is the intersection of the $n(n-1)/2$ "indifference" hyperplanes.

For what follows, let A be the ranking $c_1 > c_2 > \dots > c_n$. If $W_2^{n-(n+1)}$ is the voting vector for $S_2^{n-(n+1)}$, then $W_2^{n-(n+1)}$ is in the closure of the ranking region identified with A . (If two or more of the components, w_i , agree, then $W_2^{n-(n+1)}$ is on the boundary of the ranking region; otherwise it is in the interior.) This vector also is the tally of a ballot with the ranking A . When used as a tally, denote it as $W_{A,2}^{n-(n+1)}$. Any other strict ranking of the n alternatives is a permutation of A , where i denote the generic representation of a permutation of A as $\pi(A)$. The tally for the ranking $\pi(A)$ is the appropriate permutation of $W_2^{n-(n+1)}$, denoted by $W_{\pi(A),2}^{n-(n+1)}$. Again, $W_{\pi(A),2}^{n-(n+1)}$ is in the closure of the ranking region associated with $\pi(A)$.

Example 3. For $n = 3$ and the voting vector (w_1, w_2, w_3) , the ranking $c_2 > c_1 > c_3$ is tallied with (w_2, w_1, w_3) to represent that w_2, w_1, w_3 points are assigned, respectively, to c_1, c_2, c_3 .

Let $f_{\pi(A)}$ denote the fraction of all the voters with the ranking $\pi(A)$. The tally of an election is given by

$$2.3 \quad F(\Pi_{\pi(A)}, W_2^{n-(n+1)}) = \sum_{\pi(A)} f_{\pi(A)} W_{\pi(A),2}^{n-(n+1)}$$

where the summation index varies over all $n!$ permutations of A . The election outcome is determined by the ranking region that contains this vector sum.

The non-negative variables $\{f_{\pi(A)}\}$ sum to unity. Thus the associated profile, p , is identified with a vector in the unit simplex in the positive

orthant of E^n : i.e., $\sum x_i = 1, x_i \geq 0$. Denote this simplex by $\text{SIM}(n)$. Consequently the point defined by Eq. 2.3 is in the convex hull of $\{W_{n(x),2}^0\}_{n(x) \in \pi(A)}$, where, as always, $\pi(A)$ varies over all $n!$ permutations. This means that the set of all possible election outcomes is in this convex hull. (Conversely, any ranking region in this convex hull is an election outcome for some profile of voters.) In turn, this hull is in the affine plane defined by $\{W_{n(x),2}^0\}_{n(x) \in \pi(A)}$ and $\pi(1, \dots, 1)$ where $\pi = \sum x_i$. The analysis is considerably simplified when this plane is a linear subspace of E^n . This motivates the first of two assumptions I impose on the voting vectors. This first one just specifies a value for "b" in Proposition 1.

Vector Normalization. The sum of the components of a voting vector equals zero.

Example 4. 1. The voting vector for a plurality election is $(1, 0, \dots, 0)$, so a normalized form is $n(1, 0, \dots, 0) - (1, \dots, 1) = (n-1, -1, \dots, -1)$. If $n = 3$, all election outcomes are in the convex hull determined by $(1, 0, 0)$, $(0, 1, 0)$, and $(0, 0, 1)$. For the normalized vector, the convex hull is defined by $(2, -1, -1)$, $(-1, 2, -1)$, and $(-1, -1, 2)$.

2. A vector normalized form for the BC vector $(n-1, n-2, \dots, 1, 0)$ is $2, 1, \dots, (n-1), \dots, (n+1-2i), \dots, 1-n$. If $n = 3$, this vector is $(2, 0, -2)$; if $n = 4$, it is $(3, 1, -1, -3)$.

The vector normalization assumption forces each of the normalized vectors, $W_{n(x),2}^0 - \pi(1, \dots, 1)$, to be orthogonal to $(1, \dots, 1)$, so the vote tally, given by Eq. 2.3, also is orthogonal to $(1, \dots, 1)$. Let E^{0*} be the linear subspace of E^n defined by the normal vector $(1, \dots, 1)$. The subspace E^{0*} contains the vote tally vectors, so it is the space of interest. For instance, if $n=3$, then E^{0*} is the two dimensional space $x+y+z=0$.

Now consider all $2^n - (n+1)$ subsets of candidates. Corresponding to the set S_j is the division of an Euclidean space of dimension $|S_j|$ into ranking regions. Denote the coordinate functions of this space by $x_{k,j}$ where the first subscript identifies the candidate, c_k , while the second identifies the subset S_j . For example, for $S_j = \{c_1, c_4, c_7\}$ the coordinates of the corresponding E^j are $(x_{1,j}, x_{4,j}, x_{7,j})$, and the ranking regions are in the two dimensional subspace, E^{0*} , defined by $x_{1,j} + x_{4,j} + x_{7,j} = 0$, with $(1, 1, 1)$ as a normal vector.

Let W^n be the cartesian product of the $2^n - (n+1)$ linear subspaces E^{k*} . A

ranking region in Ω^n is given by the product of ranking regions of the component spaces. For instance, Ω^5 is a five dimensional space where the ranking region $(N_{1,1} \times N_{2,1} \times N_{2,2} \times N_{3,1} \times N_{3,2} \times N_{3,3} \times N_{4,1} \times N_{4,2} \times N_{4,3} \times N_{4,4} \times N_{5,1} \times N_{5,2} \times N_{5,3} \times N_{5,4} \times N_{5,5})$ corresponds to the ranking $(c_1 \succ c_2, c_2 \succ c_3, c_3 \succ c_1, c_2 \succ c_3 \succ c_1)$. Note that there is a one to one correspondence between the ranking regions of Ω^n and the entries of Ω^n .

Using the obvious restriction, ranking A defines a ranking for each subset of candidates. If $\mathbf{W}^0$ is a system voting vector, then $\mathbf{W}^0$ is in the closure of the ranking region of Ω^n where each ranking is determined by A . Thus, this system voting vector represents how a voter with ranking A has his ballot tallied over each of the $2^n - (n+1)$ subsets of candidates. When treated as a tally, denote this vector as $\mathbf{W}^0_A$. Any other ranking is a permutation of A , $\pi(A)$, so the tally of the ballot for each subset of candidates is given by the appropriate permutation of the vector components of $\mathbf{W}^0$. This permutation, denoted by $\mathbf{W}^0_{\pi(A)}$, is in the closure of the $\pi(A)$ ranking region of Ω^n . For a profile $\{t_{\pi(A)}\}$, the simultaneous voting tally for all subsets of candidates is

$$2.5 \quad \mathbf{F}(\{t_{\pi(A)}\}, \mathbf{W}^0) = \sum_{\pi(A)} t_{\pi(A)} \mathbf{W}^0_{\pi(A)}.$$

This summation has the same interpretation as Eq. 2.3: it defines a point in the convex hull of $\{\mathbf{W}^0_{\pi(A)}\}_{\pi(A)}$. The point is in a ranking region of Ω^n , and the associated ranking is the word in $D(\mathbf{W}^0)$ defined by this profile.

Example 5. Suppose the elections over the sets $\{c_1, c_3, c_4\}$ and $\{c_1, c_2, c_3, c_4\}$ are tallied with the voting vectors $(2, 0, -2; 3, 1, -1, -3)$; i. e., both are FC elections. Consider the profile where five people have the ranking $c_1 \succ c_2 \succ c_3 \succ c_4$, three have $c_1 \succ c_4 \succ c_2 \succ c_3$, and two have $c_1 \succ c_4 \succ c_3 \succ c_2$. The tally is $(1/2)(2, 0, -2; 3, 1, -1, -3) + (3/10)(0, -2, 1; 1, -1, -3, 3) + (1/5)(0, -2, 2; 3, -1, -1, 1) = (1, -1, 0; 2, 1, -1, 0.4, -1.6, -0.4)$, or $(c_1 \succ c_4 \succ c_3; c_1 \succ c_2 \succ c_4 \succ c_3)$.

The key observation used to characterize the dictionaries is that a word, w , is in $D(\mathbf{W}^0)$ iff the product regions associated with w intersects the convex hull of the vectors $\{\mathbf{W}^0_{\pi(A)}\}_{\pi(A)}$. This convex hull is in the linear space, $\mathcal{L}(\mathbf{W}^0)$, spanned by $\{\mathbf{W}^0_{\pi(A)}\}_{\pi(A)}$. It is easier to analyze a linear space than a convex hull.

Proposition 2. For $n \geq 2$, let $\mathbf{W}^0$ be given. A word is in $D(\mathbf{W}^0)$ iff the product ranking regions in Ω^n associated with this word has a non-empty intersection with $\mathcal{L}(\mathbf{W}^0)$.

This proposition transfers the emphasis of characterizing $D(\mathbf{W}^0)$ to the

simpler task of characterizing $V(W^n)$: (e.g., $V(W^n) = \Omega^n$ iff $D(W^n) = \{0\}$). For instance, critical parts of Theorem 1, restated in this framework, assert for $n \geq 3$ candidate that with the exception of an algebraic subset, $\mathcal{A}^n$, of possible choices for system voting vectors,

$$(2.6) \quad V(W^n) = \Omega^n.$$

Moreover, for $n = 3$, if $W^3 \neq B^3$, then $V(W^3) = \Omega^3$. However, $V(B^3)$ is a proper linear subspace of Ω^3 . Indeed, for all $n \geq 3$, $V(B^n)$ is a proper linear subspace of Ω^n .

The added structure obtained by emphasizing $V(W^n)$, rather than $D(W^n)$, can be exploited in many ways. The first involves the dimension of $V(W^n)$. To develop insight, consider the simplified setting of two sets of candidates (c_1, c_2) and (c_1, c_3) . The coordinate representation for each pair is in $\mathbb{R}^{2*}$. If $x = (x_1, x_2)$ and $y = (y_1, y_3)$ are the coordinates, then Ω is $\{(x, y) : x_1 = -x_2, y_1 = -y_3\}$. Thus, Ω is a two dimensional vector space that contains 3^2 ranking regions. A one dimensional linear subspace of Ω meets only three regions. So if Λ_1 is a proper linear subspace of Λ_2 in Ω , then Λ_2 meets at least three times as many ranking regions as Λ_1 . This argument generalizes to prove the next statement.

Proposition 3. Let $W_1^n, n = 1, 2$, be two system voting vectors where $V(W_1^n)$ is a proper linear subspace of $V(W_2^n)$ and the difference in the dimension of the spaces is d . Then

$$(2.7) \quad 3^d \leq |D(W_1^n)| \leq |D(W_2^n)|.$$

The multiple 3^d is overly conservative because it is based on the assumption that the vector spaces differ in dimension only in those component spaces corresponding to pairs of candidates. This never happens. A more appropriate multiple is obtained by analyzing the geometry of the ranking regions. In particular, this multiple always is larger than 4^d . The estimates in the introductory comments of this section are based on the conservative Eq. (2.7), so that (2.7) actually provides even stronger relief from paradoxes.

It follows from the proposition that $\dim(V(W^n))$ serves as a crude measure of $|D(W^n)|$. The next statement imposes a lower bound on the dimension of any voting vector space.

Lemma 1.

or. An atom in $V(W^n)$ is a vector whose coefficients are uniquely determined by the profile $\Pi_{\text{at}(V)}$; thus, information about which profiles cause what kind of behavior are contained the $V(W^n)$ representation.

Proposition 4. For any $n \geq 2$

$$2.8 \quad \dim(V(W^0)) \geq n(n-1)/2.$$

An immediate corollary of Theorem 1 is that all possible rankings can occur with the majority vote rankings of the $n(n-1)/2$ pairs of candidates. The vector space representation for a pair of candidates is one dimensional, so the vector space corresponding to the rankings of all pairs of candidates has dimension $n(n-1)/2$. Because the space representing the rankings of the pairs of candidates is a subspace of $V(W^0)$, inequality 2.8 follows immediately. In other words, it is the subspace of the rankings of the pairs of candidates that forces this lower bound on $\dim(V(W^0))$.

It is clear that if $V(W_1^0)$ is a proper subspace of $V(W_2^0)$, then $D(W_1^0)$ must be a proper subset of $D(W_2^0)$. However, without imposing additional assumptions on the voting vectors, the converse is false. In fact, even if two system voting vectors are equivalent (so their dictionaries are identical), the vector spaces need not agree. This can be seen with the two Borda vectors $B_1^3 = (1,0;1,0;1,0;2,0,-2)$ and $B_2^3 = (1,0;1,0;1,0;6,0,-6)$ where the scaling difference in the voting vectors force $V(B_1^3) \neq V(B_2^3)$.

This example isolates the difficulty. If a dictionary of one system voting vector is properly contained in the dictionary of another, then the voting vector subspaces can have this same relationship only with an appropriate scaling of the voting vectors. This second normalization specifies the value of "a" from Proposition 1.

Definition. a. Let the system voting vector $W^0 = (W_1^0, \dots, W_{2^{n-(n+1)}}^0)$ be given. A *scalar normalization* of W^0 is a choice of $2^n-(n+1)$ scalars $\{s_j\}$ used to define the equivalent system voting vector $(s_1W_1, s_2W_2, \dots)$.

b. The *standard scalar normalization* for the Borda system voting vector is where the Borda vectors are given by Eq. 2.4 and the voting vectors for sets of two candidates is $(1,-1)$.

The next theorem is a much stronger version of Theorem 1. To appreciate the dimension assertions, note that $\dim(L^n) = k(n) = \sum_k (k-1)n/(n-k+1)$.

Theorem 2. a. For $n \geq 3$, $\dim(V(B^0)) = n(n-1)/2$.

- b. For $n \geq 3$ and $\mathbf{W}^0 = \mathbf{B}^0$, there is a scalar normalization of $\mathbf{W}^0$ so that $V(\mathbf{B}^0)$ is a proper linear subspace of $V(\mathbf{W}^0)$.
- c. With the exception of a lower dimensional algebraic subset $\mathcal{D}^0$ of system voting vectors, $V(\mathbf{W}^0) = \mathcal{Q}^0$.
- d. If $n = 3$, and if $\mathbf{W}^1 = \mathbf{B}^1$, then $V(\mathbf{W}^1) = \mathcal{Q}^1$.
- e. The assertions of parts b, c, d hold for *any* vector, $\mathbf{W}^0$, where the vector components for the pairs of candidates are (1,-1) and those for each of the remaining subspaces of $\mathcal{Q}^0$ are non-zero and satisfy the vector normalization assumption.

An amazing assertion of this theorem is that $\dim(V(\mathbf{B}^0)) = n(n-1)/2$; it agrees with the lower bound given in Proposition 4. By use of Proposition 3, this means that $b(\mathbf{B}^0)$ is a very small subset of $\mathcal{U}^0$ relative to the size of the dictionaries for most other methods. For instance, if $\mathbf{W}^0$ corresponds to where the each subset of candidates is plurality ranked, then the difference in dimension between $V(\mathbf{B}^0)$ and $V(\mathbf{W}^0)$ is $K(n) - n(n-1)/2$. Thus $\dim(V(\mathbf{B}^0)) = 10$ while $\dim(V(\mathbf{W}^1)) = 49$ and the dimensional difference is 39; $\dim(V(\mathbf{B}^0)) = 17$ while $\dim(V(\mathbf{W}^0)) = 129$ and the dimensional difference is 114. With Proposition 5, one can appreciate the effect the dimensional differences make in the comparative sizes of the dictionaries.

Another surprising assertion is part e. For instance, it follows from part c that the favorable BC properties do not depend on the monotonicity associated with the voting vectors (e.g., $w_i \geq w_{i+1}$), so, they must be a consequence of a deeper fundamental property of the BC.⁷ For instance, it follows from part e that $V(\mathbf{W}^0) = \mathcal{Q}^0$ for $\mathbf{W}^0 = (1,-1;1,-1;1,-1;5,-4,-1)$. $\mathbf{W}^0$ is not a system voting vector because (5,-4,-1) is equivalent to (10,1,1) which requires giving 10 points to a top ranked candidate, 4 points to a bottom ranked candidate, but only 1 point to a second ranked candidate.

Part b geometrically extends the assertion that if a word is in $b(\mathbf{B}^0)$, then it is in $b(\mathbf{W}^0)$. Moreover, as it will become clear with the techniques developed in Section 3, it also means that if $\mathbf{W}^0 \in \mathcal{D}^0$, then it must be constructed in terms of the properties of the BC.

From Theorem 2 it follows that the BC is, in certain important ways, a

7. This property is due to the fact the BC creates a singularity in the orbit of a certain wreath product of permutation groups. This singularity is based on the symmetry derived from the requirement that $w_i = w_{i+1}$ is the same constant for all choices of i .

significant improvement over the other positional voting methods. But, what about other kinds of social welfare procedures. For instance, positional voting methods can be viewed as determining certain weighted means over a profile. One could design other voting procedures based on the nonlinear variational methods from statistics. In such a manner, or with the use of other techniques, wouldn't it be possible to find a method much better than the BCC? Namely, how does the BCC fare within this larger class of voting procedures? ⁵

Definition. A *smooth, majority preserving social welfare mechanism* is a mapping $\Psi: S^{n^2} \rightarrow \mathbb{R}^n$ where the rankings of the pairs of candidates is determined by majority vote. Let $I(G^n)$ be the image set of G^n .

The intersection of $I(G^n)$ with the ranking regions of $\mathbb{R}^n$ determines what rankings and what paradoxes this social welfare mechanisms admits. With the above argument, it follows that a crude measure is $\dim(I(G^n))$, where smaller values indicate fewer paradoxes. Here, if $I(G^n)$ is a smooth manifold, then $\dim(I(G^n))$ is the dimension of the manifold. If it is not a smooth manifold, then let the smallest dimension of a manifold that contains $I(G^n)$.

Corollary 2.1. If Ψ^n is a smooth, majority preserving social welfare mechanism, then $\dim(I(G^n)) \geq \dim(V(B^n))$.

In other words, *for the class of smooth, majority preserving, social welfare mechanisms, a class which includes all positional voting procedures and many others, one cannot do better than Borda's method with respect to the dimensional measure.* For related comments concerning social choice mechanisms, see Corollary 3.2. By using the ideas in [8], the smoothness requirement can be dropped.

3. The Borda Dictionary and Vector Space.

To characterize $D(B^n)$ it suffices to characterize $V(B^n)$. But $V(B^n)$ is a linear subspace of $\mathbb{R}^n$, so it is uniquely determined by its normal bundle. (This is the set of all vectors orthogonal to $V(B^n)$.) Thus $V(B^n)$ is characterized once

⁵See [10] for more.

3. As one might suspect, the ideas developed in [4] and here can be used to analyze certain classes of statistical procedures.

a basis for its normal bundle in $\mathbb{Q}^n$ is determined. To do this, the sets $\{S_1, S_2, \dots, S_{2^{n(n-1)/2}}\}$ are used to identify the component spaces of $\mathbb{Q}^n$. Recall that the first $n(n-1)/2$ sets S_i are the pairs of alternatives, that the candidates in S_i are listed according to the subscripts, and that $\mathbb{Q}^n$ is a cartesian product of $\mathbb{R}^k$ spaces. Let $\text{ind}(S_i)$ be the set of the indices (the subscripts) of the candidates in S_i .

Theorem 3, given below, asserts that a basis for the normal bundle of $V(B^n)$ is given by the vectors $\{Z_{k,i}\}$, $j = 1+n(n-1)/2, 2+n(n-1)/2, \dots, 2^n-(n+1)$, $k \in \text{ind}(S_j)$, where the first subscript identifies the candidate and the second identifies the subset S_j . The vector $Z_{k,i}$ is defined in the following manner.

1. In the component subspaces of $\mathbb{Q}^n$ corresponding to S_i , $i > n(n-1)/2$, $Z_{k,i}$ has only one non-zero vector component, $Y_{k,i}$. Vector $Y_{k,i}$, in the component space of $\mathbb{Q}^n$ corresponding to S_i , has $-(|S_i|-1)/|S_i|$ in the $S_{k,i}$ coordinate, and $1/|S_i|$ in all others.

2. For each pair of candidates, S_i , $i \leq n(n-1)/2$, the S_i vector component of $Z_{k,i}$ is

- i. $0 = (0,0)$ if either $c_k \notin S_i$, or S_i is not a subset of S_i .
- ii. $(1/2, -1/2)$ if c_k is the first listed candidate in S_i .
- $(-1/2, 1/2)$ if c_k is the second listed candidate in S_i .

Example 6. 1. For $n = 3$ and $\{c_1, c_2\}$, $\{c_2, c_3\}$, $\{c_1, c_3\}$, $\{c_1, c_2, c_3\}$, the vector $Z_{2,4} = (-1/2, 1/2; 1/2, -1/2; 0, 0; 1/3, -2/3, 1/3)$ is identified with c_2 in S_4 . This vector corresponds to the S_4 ranking where c_2 is bottom ranked and c_1, c_3, c_1 are tied for first, but where c_2 is top ranked for each of the pairs, S_1, S_2 . (The S_3 component is 0 because $c_2 \notin S_3$.) Because $Z_{2,4}$ is a normal vector for $V(B^3)$, this is an impossible outcome for a Borda election.

2. For $n = 4$, let $S_1 = \{c_1, c_2\}$, $S_2 = \{c_2, c_3\}$, $S_3 = \{c_4, c_1\}$, $S_7 = \{c_2, c_3, c_1\}$, and $S_{11} = \{c_1, c_2, c_3, c_1\}$. The vector $Z_{3,7}$, which is identified with c_3 in S_7 , has $(-1/2, 1/2)$ as the S_2 component, $(1/2, -1/2)$ as the S_3 component, $(1/3, -2/3, 1/3)$ as the S_7 component, and 0 for all other components. By being identified with c_3 in S_7 , this vector corresponds to the S_7 ranking where c_3 is bottom ranked and the other S_7 candidates are tied for first place, but c_3 is top ranked in the pairwise comparisons with either of these candidates from S_7 (i.e., in the S_2 and S_3 rankings). Thus, such an sequence of election rankings is not a word in BtB^n ; it corresponds to an impossible outcome for a Borda election. The vector $Z_{1,11}$ also is identified with c_3 , but now it relates how c_1 fares among

the candidates in S_{j+1} rather than in S_j . This vector has $(-1/2, 1/2)$ for the S_j and S_{j+1} components, $(1/2, -1/2)$ for the S_j component, $(1/4, 1/4, -2/4, 1/4)$ for the S_{j+1} component, and 0 for all other components. Again, this normal vector is in a winning region that can never be attained by a Borda election. This winning region has a similar interpretation where c_j is S_{j+1} bottom ranked and top ranked in pairwise comparisons with any candidate from S_{j+1} .

4.1 The basic result

Theorem 3. a. For $n \geq 3$, $V(B^n)$ is the $n(n-1)/2$ dimensional linear subspace of 2^n defined by the normal vectors $\{Z_{k,j+1}\}$, $j = 1 + \lfloor n(n-1)/2 \rfloor, \dots, 2^n - (n+1)$, $k \in S_j$.

b. Suppose $V(W^n)$ admits a normal vector where the only non-zero components are in those component spaces corresponding to pairs of alternatives tied to S_j where $|S_j| \geq 3$. The voting vector component of W^n used to tally the ballots for S_j is a Borda vector.

c. Consider a vector space $V(W^n)$ defined by $\{v_i(W^n)\}$ where W^n is a vector admitted by part a of Theorem 2. The assertion in part b holds true for $V(W^n)$, where the S_j component is either the BC or the reversed BC.

This simple theorem can be used to completely specify the properties of the Borda count. As I stated, my emphasis is to show how to use this theorem, rather than to provide an extensive listing of new results. As a first step, I show how to use Theorem 3 to recover and extend, in a simple, elementary fashion, several well known conclusions about the BC. To state the first result, recall that a *Condorcet* or *unifurto* winner is a candidate that wins all pairwise comparisons (i.e. majority votes), while the *anti-majority* candidate is a candidate that loses all of the pairwise comparisons. (In Example 1.b, c_1 is the Condorcet winner, while c_4 is the anti-majority candidate.) Smith [10] discovered the important relationship, described in Corollary 3.1.a, that a Condorcet winner cannot be BC ranked last. Smith also showed that this particular property is satisfied by no other positional voting method. Combining Smith's assertion with some of Young's results [11], Fishburn and Gehrlein [2] strengthened Smith's statement by asserting that the BC is the only positional voting method whereby the winner can be determined "...solely on the basis of the outcome of pairwise votes between candidates." These two statements are special cases of a more encompassing question; namely, *for what choices of positional*

voting methods are there any relationship, whatsoever, among the rankings of the positional voting method and those of the pairwise comparisons with a majority rule?⁹ For instance, for a positional voting method other than the BC are there pairwise rankings (such as when each pairwise elections ends in a tie, or when both a Condorcet winner and an anti-majority candidate exist, or when the pairwise election rankings define a transitive, binary relationship) that preclude the possibility of at least *one* positional election rankings from occurring? This more general issue is answered in Corollary 3.1 b.

Corollary 3.1 a. Consider $n \geq 3$ candidates. A Condorcet winner can never be BC bottom ranked, and an anti-majority candidate can never be BC top-ranked.¹⁰

b. The BC is the only positional voting method that admits any relationship among the rankings of the positional voting procedure and those of the pairwise votes. Indeed, the same assertion holds for any choice of $(w_1, \dots, w_n)$ used to tally the ballots, where at least two choices of the w_i 's differ in value.

With Theorem 3, the proof of Corollary 3.1 reduces to a simple computation. This computation is carried out in detail to demonstrate the ideas.

Proof. Suppose a profile $p = \{P_{n \times n}\}$ admits c_1 as the Condorcet winner, but c_1 is BC bottom rank-1. Let $F(p, B^n) = (X_1, \dots, X_{2^n - (n+1)})$, so X_k is the normalized BC vote tally for S_k . The assumption that c_1 is BC bottom ranked requires the corresponding component of $X_{2^n - (n+1)}$ to be algebraically smallest: namely, $x_{1,k} < x_{1,k}$, $j = 2, \dots, n$, $k = 2^n - (n+1)$. Because $\sum_1 x_{1,2^n - (n+1)} = 0$ (by the vector normalization assumption), it follows that $x_{1,2^n - (n+1)} < 0$. It now follows from a direct computation, using the vector normalization, that

$$\begin{aligned} 3.1 \quad (Y_{1,2^n - (n+1)}, X_{2^n - (n+1)}) &= [-(n-1)x_{1,2^n - (n+1)} + (\sum_{j=2}^n x_{j,2^n - (n+1)})]/n \\ &= -x_{1,2^n - (n+1)} > 0. \end{aligned}$$

a. There always are some relationships among the *tallies*: i.e., a trivial one is if a candidate wins all pairwise elections by unanimous votes, then she is top ranked with any positional method.

10. A nice, related statement about the BC, by Fishburn and Gehrlein (2), asserts, for $n = 3$ candidates and for a specified probabilistic measure over the profiles that the BC is the unique positional voting method to maximize the *likelihood* that a Condorcet winner is top ranked. Using the mathematical structures introduced in Section 2 and in (3), Jill van Newenhuizen (13) significantly extends this statement in many different ways. In part, she shows that the same conclusion holds for all $n \geq 3$ and for a much larger class of probabilistic Distributions.

The assumption that c_1 is a Condorcet winner means that, for each k where $|S_k| = 2$ and $c_1 \in S_k$, the $x_{1,k}$ component of X_k is positive. For each such choice of k , it follows that

$$(3.1) \quad (1/2, -1/2) \cdot X_k = x_{1,k} > 0.$$

With Eqs. 3.1, 3.2, the computation of the scalar product is

$$(3.2) \quad (F(p, B^0), Z_{1/2^{0, \dots, 0, 1, \dots, 1}}) = -x_{1,2^{0, \dots, 0, 1, \dots, 1}} + \sum_k x_{1,k} > 0$$

where the summation is over the values of k selected for Eq. 3.2. But, Theorem 2 requires this scalar product to be zero. This contradiction proves that such a ranking is not a Borda outcome.

The proof that an anti-majority candidate cannot be BC top ranked is essentially the same.

Proof of part (c). F is a linear mapping, so if a system voting vector W^0 imposes an (arbitrary) relationship, whatsoever, among the outcomes of the pairs and the ranking of a set of candidates, then the image of F lies in a lower dimensional linear subspace of the specified coordinate subspaces of Ω^0 . This forces $V(W^0)$ to have a normal vector with its only non-zero vector components in those coordinate subspaces of Ω^0 . According to Theorem 3, the voting vector is a Borda vector.

3.2 Some axiomatic characterizations of the B⁰

To illustrate Corollary 3.1 (c), I use it to extend Young's insightful axiomatic characterization of the BC that is based on properties of *social choice functions* [11]. Say that two *social choice functions*, f, g , are *equivalent* if $f(p) = g(p)$ for all p . To state Young's result, recall that two standard assumptions on social choice functions are that f is *anonymous* if its outcome depends only on the numbers of voters with each preference, and that f is *neutral* if when α is a permutation of the indices of the candidates, then $f(\alpha(p)) = \alpha(f(p))$. (That is, both the different candidates and the different voters are treated equally; the outcome does not depend on their names.) The next assumption is that f is *consistent*: namely, if p and p' are profiles for distinct voter sets, then $f(p) \cap f(p') \neq \emptyset$ implies that $f(p) \cap f(p') = f(p \cup p')$. (Suppose

a group is subdivided into two subcommittees represented by p and p' . (In the subcommittees agrees in that $f(p) \cap f(p') \neq \emptyset$, then consistency requires the common set $f(p) \cap f(p')$ is the choice of the full group $p+p'$.) The function f is *faithful* if for a profile of a single voter with c_j as his top ranked candidate, $f(p) = c_j$. Finally, Young states that f has the *cancellation property* if when a profile p causes all $n(n-1)/2$ pairwise comparisons to result in a tie vote, then $f(p) = \emptyset$. Young proved for $n \geq 3$ that if a social choice function is anonymous, neutral, consistent, faithful, and has the cancellation property, then it is equivalent to choosing the top ranked candidates from the BC ranking of C^0 .¹¹

Definition A social choice procedure for C^0 is a *general scoring method* based on the vector $W^1 = (w_1^1, \dots, w_n^1)$ with tie breaker methods $W^i = (w_1^i, \dots, w_n^i)$, $i = 2, \dots, n$, if the following conditions are satisfied.

1. Not all of the w_j^1 's in W^1 have the same value.
2. A voter's i th ranked candidate receives w_j^i points.
3. The candidate, or candidates with the largest point total are selected. If a tie breaker is used to determine among several candidates with the largest point total, then, inductively, at the i th stage *all* candidates are re-ranked with W^i . Those candidates with the largest point total that also are selected at the $(i-1)$ th stage are selected for the i th stage.

The weights for a scoring vector need not satisfy any monotonicity condition, e.g., the vector $(-1, -1, 4, -3)$, where the third ranked candidate gets 4 points, the first and second get -1 points, and the last ranked candidate gets -3 points, is a scoring method. Young [12] found an interesting characterization of general scoring methods.

Proposition 5. [12] For $n \geq 3$, if a social choice function is anonymous, neutral, consistent, and does not have a fixed value, then it is equivalent to a general scoring method.

Footnotes

- [1]. The equivalence classes are necessary because it is easy to create an f that is not the BC. Indeed, if the BC tallies are composed with any monotonic function, even a discontinuous one, the result is an equivalent version of the BC. Moreover, any such mapping can be further composed with certain transformations of the space of profiles. These equivalent choices can have representations that appear to differ significantly from the BC.

To understand the tie-breaking scheme, suppose for $n = 3$ that the set is initially ranked with the BC, and that the two successive tie-breakers are $(1,0,0,0,0)$ and $(1,1,1,1,0)$. So, first BC rank the candidates. If there is a tie, choose the ones with the most first place votes; if there still is a tie, choose the ones with the least number of last place votes. As an illustration, there is a profile P so that the rankings of these three procedures are, respectively, $c_1=c_2=c_3>c_1>c_3$, $c_3>c_1=c_3=c_2>c_1$, and $c_1>c_2>c_1>c_3>c_2$; this means c_3 wins. One might wish to streamline the tie-breaking procedure by using a run-off among only $\{c_1, c_2, c_3\}$, rather than reconsidering c_1 and c_3 ; after all, c_1 and c_3 are eliminated from further consideration. However, *any such procedure violates consistency*. To see why, consider the symbols $\{c_1=c_2=c_3>c_1, c_1=c_2>c_3, c_1=c_2, c_1>c_2\}$ and $\{c_2=c_3=c_1>c_1, c_2=c_1>c_3, c_2=c_1, c_3>c_2\}$. Now, it follows from theorem 2 that for any choice of scoring methods, satisfying part 2, there are profiles $\mathbf{p}$ and $\mathbf{p}'$ defining the above two listings of election outcomes. Thus with a tie-breaker for a set of three and a set of two candidates, $f(\mathbf{p}) = \{c_1, c_2\}$ while $f(\mathbf{p}') = \{c_2>c_1\}$, so $f(\mathbf{p})\cap f(\mathbf{p}') = \{c_2\}$. However, $f(\mathbf{p}+\mathbf{p}') = \{c_3\}$. (This is because the linearity of summing tallies forces the top ranked candidates for $\mathbf{p}+\mathbf{p}'$ to be $\{c_2, c_3\}$.) Again, from the linearity of summation processes, c_3 is the winner of the run-off. Thus, while consistency appears to be a natural requirement, it imposes a strong condition on the social choice procedure.

"Faithfulness" in Young's Theorem plays two roles. The first is to impose a monotonicity on the choice of a scoring method; it requires $w_1>w_i$ for $i \geq 2$. As such, this condition prohibits W^1 from being $(1,1,0,0,0)$, or any other voting vector where the weights assigned to a voter's first and second ranked candidates agree. I use a weaker condition that includes "faithfulness" as a special case. The new condition admits all positional voting methods because it only requires w_n , the weight assigned to the bottom ranked candidate, to have a smaller value than at least one other weight. The second role of "faithfulness" is to ensure that the image of f contains singletons; i.e., there are situations where only one candidate is selected. It turns out that this property is not needed for what follows. On the other hand, this property does simplify the statements and the proofs of the results, so, for convenience, I include it as part of the following definition.

Definition. A social choice function, f , is *somewhat faithful* if the image of f contains a singleton and if for a profile $\mathbf{p}$ of a single voter, his bottom ranked

candidate is not in the set $f(p)$.

In light of Proposition 3, the role of the cancellation property in Young's result is to impose a pairwise ranking restriction on the choice of the quasi-positional voting method. It now follows immediately from Corollary 3.1 that the only possible choice is the BC. Indeed, in light of Corollary 3.1 b., it follows that *Young's result can be extended in many different directions simply by replacing the cancellation property with any other BC property*. For instance, the same conclusion holds if the cancellation property is replaced with the weaker requirement of *non-determinacy* whereby if p creates a tie vote in all pairwise elections, then $f(p)$ is not a singleton. Another natural choice is to replace cancellation with the requirement that if c_j is an anti-minority candidate, then $c_j \notin f(p)$; here the conclusion admits tie breakers. Either of these appealing, substitute axiomatic representations of the BC appears to be difficult to prove directly, but they are immediate consequences of Corollaries 3.2 and 3.4.a. Indeed, most of the BC properties derived here serve as substitute conditions.

Corollary 3.2. a. For $n \geq 3$, suppose an anonymous, neutral, consistent, and somewhat faithful social choice function, f , satisfies another specified condition whereby the pairwise rankings of the candidates imposes a (non-trivial) constraint on the image of f . If this specified condition is not satisfied by the BC, then no such f exists. If the condition is satisfied by the BC, then f is equivalent to first choosing the BC top ranked candidates of C^0 where possible ties may be broken by tie breakers.

b. For $n \geq 3$, there does not exist an anonymous, neutral, consistent, somewhat faithful social choice function that always selects the Condorcet winner.

c. For $n \geq 3$, if an anonymous, neutral, consistent, and somewhat faithful social choice function satisfies the non-determinacy property, then it is equivalent to the BC with no tie breakers.

By "non-trivial", I mean that when the designated pairwise rankings occur, then there is a subset of $P(C^0)$ that is not an image value for f . Part b, which is an immediate example of a condition that leads to an impossibility theorem, holds because the Condorcet winner need not be BC top ranked. Part c is

a direct extension of Young's statement, but it is not obvious that his techniques could prove this stronger statement. One purpose for including this assertion is to indicate in the proof why tie-breakers are not admitted. Finally, the only role played by the "somewhat faithful" condition is to impose enough monotonicity to outline the reversed Borda count. There are many substitutes and ancillary conditions.

3.2 The Borda tally

To extend Corollary 3.1a, note that the proof just uses the fact that $\text{Proj}_{\mathbf{Z}_{k+1}}(\mathbf{I}, \mathbf{B}^0)$ is orthogonal to each $\mathbf{Z}_{k+1}$. In the proof I separately computed the contribution of the scalar product due to the outcomes of the pairwise elections and the contribution due to the Borda tally. This same computation generalizes Corollary 3.1a from an assertion about the ranking of a Condorcet winner to a statement about any candidate who fares well in pairwise comparisons. More precisely, *if the pairwise election outcomes for candidate c_i satisfy $\sum_k x_{i+1,k} > 0$, where the summation index k is over the pairs of candidates S_k that include c_i , then c_i cannot be BC bottom ranked. Equivalently, if this summation is negative, then c_i cannot be BC top ranked.* This assertion uses all of the vectors $\mathbf{Z}_{i+2}^0, \dots, \mathbf{Z}_{n+1}^0$; it holds because, in light of Theorem 3, the assumption about the pairwise elections forces the scalar product $(\mathbf{Y}_{1,2}^0, \dots, \mathbf{Y}_{n+1}^0, \mathbf{X}_2^0, \dots, \mathbf{X}_{n+1}^0)$ to be negative. In turn, this forces the angle between $\mathbf{X}_2^0, \dots, \mathbf{X}_{n+1}^0$ and $\mathbf{Y}_{1,2}^0, \dots, \mathbf{Y}_{n+1}^0$ to be greater than $\pi/2$. It now follows from the position of $\mathbf{Y}_{1,2}^0, \dots, \mathbf{Y}_{n+1}^0$ and the geometry of the ranking regions that, not only is it impossible for c_i to be BC bottom ranked, but *it is impossible for c_i to even be BC tied for last.*

To appreciate the assumption $\sum_k x_{i+1,k} > 0$, note that if $S_i = \{c_i, c_n\}$, then $(x_{i+1,i} + 1)/2$ is the fraction of the voters that prefer c_i to c_n . Thus, the above generalization of Corollary 3.1a asserts if a candidate receives a surplus of votes over all of the pairwise comparisons (i.e., if the sum of the fractions of votes from each of the $n-1$ pairwise elections exceeds $(n-1)/2$), then she cannot be BC ranked last.

Example 7. (a) What words are admitted in $\text{D}(\mathbf{B}^0)$? To illustrate both the geometry and how algebraic relationships are found, I'll show that $(c_1 \geq c_2; c_1 \geq c_3; c_2 \geq c_3; c_2 \geq c_4; c_3 \geq c_4) \in \text{D}(\mathbf{B}^0)$. This word corresponds to the components $(x_{1,1}, -x_{1,1}; x_{1,2}, -x_{1,2}; x_{2,3}, -x_{2,3}; x_{1,1}, -x_{1,1}; -2x_{1,1})$ where $x_{1,1} > 0$ for $i = 1, 2, 3$, and $x_{1,4} = 0$. The orthogonal scalar product of this vector with $\mathbf{Z}_{1,1}$ yields $x_{1,1} +$

$x_{1,2} = x_{1,4}$, while with $Z_{2,3}$ it is $-x_{1,2} + x_{2,3} = x_{1,3}$. ($Z_{3,4} = -(Z_{1,4} + Z_{2,3})$, so it provides no new information.) Thus, the word is admissible in $\mathbf{B}^3$ because it is possible to satisfy the equation

$$(3.4) \quad x_{1,1} + 2x_{1,2} = x_{2,3}.$$

Therefore, not only is such a word in $\mathbf{D}(\mathbf{B}^3)$, but *Eq. 3.4 is a necessary and sufficient condition for a Condorcet winner to be BC tied for first place.* A necessary and sufficient condition for c_2 to be BC top ranked is if $x_{2,3}$ has a larger value. Notice that this imposes a significant burden on her winning margin over c_1 , particularly if c_1 does not just barely win both pairwise elections. Alternatively, a sufficient condition for c_1 , the Condorcet winner, to be BC top ranked is $x_{1,1} + 2x_{1,2} \geq 1/2$ ($\geq x_{2,3}$). As extreme examples, this happens if she just barely beats c_2 ($x_{1,1} > 0$) but gets at least 62.5% of the vote against c_3 ($x_{1,2} \geq 1/4$), or if she beats c_2 with a 75% vote ($x_{1,1} = 1/2$, $x_{1,2} > 0$). The main purpose for these new conclusions is to indicate how similar supporting relationships for any word in $\mathbf{D}(\mathbf{B}^n)$ and for any $n \geq 3$ can be found.

b. What words are in the dictionary for the plurality ranking, but not in $\mathbf{D}(\mathbf{B}^3)$? If $S_1 = \{c_1, c_2\}$, $S_2 = \{c_1, c_3\}$ and $S_3 = \{c_1, c_2, c_3\}$, then, by using $Z_{1,2}$, it follows that a word with the three symbols $c_1 > c_2$, $c_1 > c_3$, $c_2 > c_3 > c_1$ is not in $\mathbf{B}^3$. All such words are in $\mathbf{D}(\mathbf{W}^3)$ if the system vector is based on plurality votes. So, by permuting the indices, by filling in the other symbols in an arbitrary fashion, etc., this simple assertion accounts for 172,974,204 words in the plurality dictionary that are not in $\mathbf{D}(\mathbf{B}^3)$.

In his nice paper, Smith reminds the reader that Black questioned the naturalness of the Condorcet winner. For instance, the Condorcet winner, c_1 , may barely win each pairwise election, while c_2 barely loses to c_1 and then wins all other pairwise elections by substantial margins. It is reasonable to feel that c_2 should be the winner. Smith notes that "it would be interesting to try to formulate this feeling as a precise property: is may be that a suitable formulation is a necessary and sufficient condition for a positional voting system." To continue this line of thought, note that a Condorcet winner may achieve her status not by excellence, but through mediocrity by being most voters' compromise, or second choice candidate. For instance, suppose three voters have the ranking $c_1 > c_2 > c_3$, 50 have $c_3 > c_1 > c_2$, 50 have $c_2 > c_1 > c_3$, and two have $c_1 > c_3 > c_2$. Here, because 52 voters have c_2 as top choice, 50 have c_3 , and only three have c_1 , it is reasonable to believe that the true choice is between

c_1 and c_4 . Nevertheless, because 100 of the voters rank c_1 in second place, c_1 is the Condorcet winner, and c_4 is the *anti-majority candidate*.

Examples and criticisms of this kind do raise concern about the virtue of the Condorcet winner, so once again emphasis to Smith's question. Yet again, his question is easy to answer because, according to Proposition 5, the imposition of rational assumptions force the social choice function to be equivalent to a scoring method. If the social choice procedure is to be related in any manner whatsoever with the pairwise election outcomes -- and this is mandatory if one is to follow Smith's suggestion -- then, according to Corollary 3.1 a/c, *the procedure can only be the BC*. Thus, to avoid an impossibility conclusion, the appropriate condition must be based on BC properties. Example 7a illustrates the sensitivity of the BC to the pairwise tallies, while Ex. 3.4 illustrates that when the BC does allow a non-Condorcet winner to be BC top ranked, she can do so only by overcoming a significant burden. For instance, in the example of the previous paragraph, c_1 is the Borda winner with 107 points, while the Condorcet winner, c_4 , is Borda second ranked with 106 points.

3.4 Other subsets and consistent sets

The above argument just uses the vectors $\{Z_{1,2}^0, \dots, Z_{n+1,1}^0\}$. Many other conclusions follow by using all of the vectors $\{Z_{i,j}^0\}$. A first step is Corollary 3.3 given below. Part a is included to handle an obvious gap of Corollary 3.1a: namely, "is it possible for a Condorcet winner be BC ranked below an anti-majority candidate?"

Corollary 3.3. a. Let $n \geq 3$, and consider the subset S_1 where $1S, 1 \geq 3$. Each component of the BC tally for S_1 , X_j is given by

$$X_j = \sum_{i=1}^n N_{i,j,1} = \sum_k N_{1,k,1}$$

where the summation is over all pairs of candidates, c_j , where $c_j \in S_1$ and S_k is a subset of S_1 .

b. Suppose, for a set S_1 , $\sum_k N_{1,k,1} > 0$, where the summation is as described in part a. Then, candidate c_1 cannot be BC ranked or tied for last in S_1 . If this summation equals zero, then c_1 can be BC bottom ranked or top ranked iff the election ranking has all candidates in a tie.

c. If c_1 is the Condorcet winner and c_n is the anti-majority candidate, then the BC ranking for any subset of candidates that includes c_1 and c_n ranks c_1

strictly above c_n . The BC is the only positional voting method for which this is true.

Example 8. *a.* To illustrate Eq. 3.5, return to the introductory beverage example where c_1 corresponds to water, c_2 to beer, and c_3 to wine. This means that $x_{1,1} = -x_{2,1} = (6/15) - (9/15) = -3/15$, $x_{1,2} = -x_{3,2} = -3/15$, and $x_{2,3} = -x_{3,3} = (5/15) - (9/15) = -4/15$. Therefore, according to Eq. 3.5, $x_{1,1} = -6/15$, $x_{2,1} = -4/15$, and $x_{3,1} = 7/15$. This leads to the BC ranking $c_3/c_2/c_1$, or wine/beer/water.

b. Suppose a runoff election is being designed so that if a Condorcet winner exists, she will be elected. If the elimination procedure is based on ordinal rankings, then, as Smith noted, each set of candidates must be BC ranked and, at each stage, only the bottom ranked candidate is dropped. This can be accomplished by using the normalized BC tally. At each stage, drop all candidates with a negative (rather normalized) BC tally. As the Condorcet winner always has a positive normalized tally, she is advanced to all stages. A similar procedure holds for other social choice methods, such as a "generalized Borda" where the candidates are listed in some order and then the first $k/2$ candidates are ranked. The idea is to replace those candidates dropped from further consideration by the next listed candidates. Again, to ensure a Condorcet winner, at each stage, the candidates with a negative normalized BC tally should be dropped.

Equation 3.5 follows from the requirement that $(F(\{c_n(c_{-1}), B^0\}), Z_{1,1}) = 0$ for all c and 1 . Part a follows because if $c_1 \in S_1$, then, according to Eq. 3.5, $x_{1,1} > 0$. Similarly, if $c_n \in S_1$, $x_{n,1} < 0$. Part b is a simple illustration. Corollary 3.3b significantly extends Corollary 3.1 to all candidates (not just the Condorcet winner) and to all subsets of candidates (not just the set of all candidates). Based on the approach used by Borda, it wouldn't surprise me to discover that he already knew of a result of the general nature of Eq. 3.5 for $n = 2$. Versions of it do appear in Smith [10], and then later in Young [11], but only for the set of all candidates. By using this relationship for all sets of candidates, one can address issues such as whether there is a relationship between the BC ranking of a candidate in the set of all candidates and her BC ranking on the set where the bottom ranked candidate is removed. This question is easily answered below: the basis for the solution is stated for all subsets in Corollary 3.3a (by use of Eq. 3.5), and it is illustrated, with one simple

example, in Corollary 2.4b, Corollary 2.4a also shows that only the admissible such relationships. As such, Corollary 2.4 extends Corollary 2.1, and it can be used, in the obvious fashion, to extend Corollary 2.2, and, hence, the Corollary is stated in related issues coming from social choice theory, as introduced.

An important theme from social choice is to understand the appropriate restrictions on the “feasible” sets of candidates that leads to a possibility theorem. Social choice functions are extended to subsets of $\mathcal{C}$ by asserting that for S_1 , $(f(p, S_1)$ is a subset of S_1). A natural condition, introduced by Jones [11], is based on the idea that if a candidate is top ranked in S_1 , she should be top ranked in any subset of S_1 . Thus, if $(f(p, S_1)$ is a subset of S_1 , where S_0 is a subset of S_1 , then $(f(p, S_0) = f(p, S_1)$. However, when combined with some other natural assumptions, this condition turns out to be too strong to permit the existence of any social choice function. (For a geometric explanation, see [8].) Therefore, it is worth considering alternative conditions.

An arrow type condition leads to an impossibility conclusion, so it is reasonable to suspect that the selection of the “best” candidate depends on how she compares with other available candidates. For instance, it may be that part of the c_j ’s appeal is derived from her favorable comparison with c_2 . So, if c_2 is not an option, then c_1 loses some of her comparative advantage.

Correspondingly, if c_j suffers by comparison to c_2 , then her comparative advantage improves in the absence of c_2 . This motivates the following condition: a condition used in part c of Corollary 2.4 as one of many possible alternatives for Arrow’s condition that leads to a possibility conclusion.

Definition. a. A social choice function satisfies the *comparative advantage* property if for any subset of candidates, S_j , the following hold: suppose $c_2 \notin S_j$ and that $c_j \in S_j$ beats c_2 in a pairwise election. If there is a candidate $c_1 \in f(p, S_j \cup \{c_2\})$ that loses to c_2 in a pairwise election, then, $c_1 \notin f(p, S_j)$.

b. A social choice function satisfies the condition of *independence of irrelevant alternatives* (IIA) if the following condition holds for every possible set S_1 . If p and p' are two profiles where the relative rankings of the candidates in S_1 are the same, then $(f(p, S_1) = f(p', S_1)$.

The comparative advantage property determines who cannot be selected from a smaller feasible set, rather than who is, one might prefer to require $c_1 \in$

$f(p, S_j)$, but, such a requirement can lead to an impossibility conclusion.

Corollary 3.4. (a). Let S_j be a set of candidates and S_k a proper subset of at least three of the candidates in S_j . The BC is the only positional voting method that admits a relationship among the tallies (or rankings) of S_j , S_k , and the pairwise comparisons of the candidates in S_j . If $S_a = S_j - S_k$, then the BC relationship for each candidate $c_i \in S_k$ is

$$x_{i,j,a} = x_{i,j,k} + \sum_{c_l \in S_a} x_{i,j,l},$$

where the summation is over all pairs $S_l = \{c_l, c_l\}$ where $l \in \text{Index}_a$.

(b). Suppose $|S_j| \geq 3$ and that $S_l = S_j/\{c_a\}$. For $c_i \in S_k$, denote the pair $\{c_i, c_a\}$ by S_a . Then, $x_{i,j,a} = x_{i,j,k} + x_{i,j,a}$. In particular, if c_i has over half of the BC total vote tally in S_j and if c_a beats c_i in a pairwise election, then c_i cannot be bottom ranked, nor tied for bottom ranked in S_k . Such an assertion holds only if the BC is used to rank both sets.

(c). If for $n \geq 3$, an anonymous, neutral, consistent, somewhat faithful social choice function f that satisfies the IIA and the comparative advantage properties, then f is equivalent to the BC where $f(p, S_j)$ is the top ranked candidate from the BC ranking of S_j .

Example 9. If c_1 is BC bottom ranked in $S_j = \{c_1, c_2, c_3, c_4, c_5, c_6\}$ (so $x_{1,j,j} < 0$) and BC top ranked in $S_k = \{c_1, c_2, c_3\}$ (so $x_{1,k,k} > 0$), then it must be that c_1 lost in at least one of the pairwise comparisons with c_4 , c_5 , and/or c_6 . This assertion may seem to be obvious, so it is surprising that only the BC always satisfies it.

3.5. A sufficient condition

A difficulty with Eq. 3.5 is that it is useful only *if* the tallies for all of the pairwise elections are known. So to demonstrate various BC properties, examples need to be constructed so that the right hand side has the appropriate value. For normative or theoretical studies we need more. We need to know what values of $x_{i,j,k}$, $k \leq n(n-1)/2$, are admissible. With the vector space representation used here, the answer to this seemingly difficult combinatoric problem is immediate. Thus, Corollary 3.5 completes this direction of investigation. To state the results, notice that the vector $\mathbf{M}c_j = (1/2, -1/2, -1/2, -1/2, \dots, -1/2, -1/2)$ with $n(n-1)/2$ vector components corresponds to the tally

of a subset of the pairwise comparisons for a voter with the remaining 1. Similarly, $\mathbf{MP}_{a, S_1}^b$ corresponds to the entry for a voter with the ranking $a, 1$.

Corollary 3.5. The set for $\lambda = (\lambda_1, \dots, \lambda_{n(n-1)/2})$ can be realized by a profile of voters iff $\lambda \in \text{int}(\mathbf{MP}_{a, S_1}^b)$, the convex hull of $\{\mathbf{MP}_{a, S_1}^b\}$, and the components of λ are non-negative.

Example 10. a. By use of Corollary 3.5 and Eq. 3.5, one can show, for example, that there are profiles where c_1 is the Condorcet winner, but is ranked in the 10^{th} place, c_2 only loses to c_1 in the pairwise comparisons, but she is 1st ranked in the 10^{th} position, ..., c_j loses a pairwise comparisons to c_1 iff $j < i$ but she is i th ranked in the 10^{th} position, and c_n , the anti-majority candidate is i th bottom ranked. To prove this, note that 0 is an interior point of the convex hull specified in Corollary 3.5. Thus the $x_{1,j}$'s, for the pairwise comparisons, can be chosen to have arbitrarily small positive values. For candidate c_1 , only the value for pairwise comparison with c_j is specified, so the remaining $x_{1,j}$'s can be selected as arbitrarily small positive values that satisfy the requirement $0 < x_{1,2}^{b, \text{int}(\lambda)} < x_{2,2}^{b, \text{int}(\lambda)}$. Continuing, for candidate c_j , only the values for the pairwise comparisons with c_1 , $j < i$, have been specified. The remaining values of $x_{1,j}$, for the pairwise comparisons, can be selected so that they are positive and so that $x_{1,1}^{b, \text{int}(\lambda)} < x_{1,2}^{b, \text{int}(\lambda)}$. All of the values for the pairwise comparisons with c_n have been specified, and they are all negative. Thus, $x_{n,2}^{b, \text{int}(\lambda)} < 0$, and the conclusion holds. (I used the fact that all of the hyperplanes associated with a tie majority vote pass through 0. It is immediate that the appropriate values are admitted because the relevant hyperplanes obviously have independent normal vectors.)

With a similar argument, it is easy to show that if we only know that c_1 is the Condorcet winner and c_n is the anti-majority candidate, then the only restrictions on the B' ranking is that c_1 is strictly ranked above c_n (Corollary 3.3c): all rankings satisfying this condition are admitted. Related statements hold for the various subsets S_1 .

b. By use of Theorem 1.d and using a special case of the above, it follows that there is a profile of voters, $\mathbf{p}$, so that for all choices of positional voting methods, the election ranking is $c_{n-1} > c_{n-2} > \dots > c_2 > c_1, c_n > c_2$ while the pairwise rankings of the pairwise votes are $c_i > c_j$ iff $i < j$. Therefore, c_1 is the Condorcet winner, c_2 almost is the Condorcet winner (it only loses to c_1).

and c_k is the anti-majority winner.

6. Because I show in [31] how to extend voting paradoxes from the literature by use of the dictionaries, I do not emphasize this approach here. However, when this approach is used with the BC, complications arise because $\text{li}(\mathcal{B})$ is a subset of $\mathcal{Q}^n$. This means that the existence of certain symbols precludes the possibility that other symbols can occur. To find what symbols are admitted, note that Theorem 3 and Corollary 3.5 imply that they are determined by the image of σ mapping from $\mathcal{Q}^n$ to the various component spaces of $\mathcal{Q}^n$; the components of σ mapping are given by Eq. 3.5.

To illustrate this with a simple example, consider all possible words in $\text{li}(\mathcal{B})$ that have admit the Condorcet cycle $c_1 > c_2$, $c_3 > c_1$, and $c_2 > c_3$. This corresponds to the region in $\mathcal{CH}^3$ defined by positive values for $x_{1,1}$, $x_{3,2}$, and $x_{2,3}$. The easiest way is to find whether this region admits any symbols in S_1 with a tie vote. But, according to Eq. 3.5, if all three variables are equal to $\epsilon > 0$, then the S_1 ranking is $x_{1,1}=x_{2,1}=x_{3,1}$, or $c_1=c_2=c_3$. Because 0 is an interior point of $\mathcal{CH}^3$, the value of ϵ can be chosen so that $x_{1,2} = x_{3,2} = x_{2,3} = \epsilon$ is an interior point. Thus, by perturbing these values, it follows that the S_1 symbol can be anything. This same approach holds for all values of n .

3.6. More general EC characterizations.

So far I have emphasized the relationship among the BC ranking of a set S_1 and the rankings of the pairs of candidates in S_1 . This is not necessary; as already suggested by Eq. 3.6, there are relationships among the BC rankings of other subsets of candidates. Some natural ones are described below in Corollary 3.6; a complete listing is in [9].

The above characterizations of the BC are based on election outcomes of the minimal pairs of candidates. It is natural to wonder whether the BC rankings also can be characterized strictly in terms of the BC rankings of all triplets, or of all sets of 4 candidates. It can. Corollary 3.6 asserts that the BC ranking for the $\{S_1\} = k$ candidates in S_1 can be determined by the BC rankings for all possible subsets of m candidates from S_1 for $2 \leq m \leq k$. The explicit relationships are easily determined with elementary linear algebra techniques where the final result is of the form of Eq. 3.7 or of the form described in Example 11a.

For a different kind of motivation for Corollary 3.6, recall the Theorem

"pump" argument occasionally used by economists to discuss intransitivities). The argument is that a person with intransitive preferences soon would change his rankings to avoid being exploited. To see the idea, suppose a person has the rankings $a_1 > a_2$, $a_1 > a_3$, $a_2 > a_3$. If he is to choose a candidate from $\{a_1, a_2\}$, he selects a_1 . Presumably, he would be willing to pay (a bribe?) to change to set $\{a_2, a_3\}$ so he could select the personally more desirable outcome of a_2 . For similar reasons, after paying, presumably he will pay another sum to choose from the set $\{a_2, a_3\}$ rather than from $\{a_3, a_1\}$. Now, faced with selecting from $\{a_2, a_3\}$, presumably he will pay to obtain the personally more favorable situation of choosing from $\{a_1, a_2\}$. So, after paying, our victim returns to the original situation unless he changes his rankings; his cyclic preferences provide a never-ending opportunity to pump money out of him.

Can a money pump be applied to an organization? The same argument applies if an organization's top choices form a cycle. For instance, suppose a group's election rankings over triplets from $\{a, a_2, c_1, c_3\}$, $\{a_2, c_1, c_3, c_4\}$, $\{c_1, c_3, c_4, a_1\}$ and $\{c_3, c_4, a_1, a\}$. In such a situation, presumably, when faced with selecting from a specified set of three candidates, the organization would pay a sum of money for the privilege of selecting from the next triplet of candidates in order to obtain a "significantly better" candidate. (The first set of candidates "follows" the last set, etc. After all, the top ranked candidate in one set is bottom ranked in the following set of candidates. Thus, such an organization is a "money pump" target. In light of this example, it is reasonable to question if profit and voting methods could such cyclic rankings. It turns out that no positional voting method, even the BC, avoids all of them. However, only the BC gives partial relief because it avoids many of them. For instance, it follows as a consequence of Corollary 3.6 that the BC does not admit the above example of rankings. (This follows from the first specified normal vector. The normal vectors stated in Corollary 3.6 are in the subspace of 2^n identified with the subsets of candidates.)

Corollary 3.6 (a). Let $n = 4$. The BC is the only positional voting method that admits relationships among the tallies of the four triplets of candidates. If the sets are listed as $\{a_1, a_2, c_1\}$, $\{a_1, a_2, c_1\}$, $\{a_1, a_2, c_1\}$, $\{a_2, c_1, c_3\}$, then all possible relationships among the BC tallies for these four sets of three candidates are characterized by the normal bundle spanned by the two normal vectors $N_1 = (1, -2, 1, -2, 1, 1, 1, 1, -2, 1, -2, 1)$ and $N_2 = (-2, 1, 1, 1, -2, 1, 1, -2, 1, 1, 1, -2)$.

b. Let S_1 be a subset of $k \geq 3$ candidates, and consider all subsets of m candidates, $k \leq m \leq 2k$. If all sets are BC ranked, then there is a relationship among the rankings that is uniquely determined by a set of normal vectors of co-dimension $(k-1)/2$.

c. Let S_1 be a subset of candidates with $k \geq 3$ candidates, and let m be an integer such that $2 \leq m \leq k$. The BC imposes a relationship among the S_1 rankings and the $(k/m)(k-m)!$ subsets of m candidates from S_1 . These relationships are uniquely determined by the vectors in a normal bundle. A basis of this normal bundle of co-dimension $k(k-1)/2$ is given by the normal vectors determining the BC relationships among the sets of m candidates and by $k-1$ additional vectors of the following form: in the following manner: For $c_1 \in S_1$, the S_1 vector component is $[(k-2)!/(m-2)!(k-m)!]Y_{a_1,1}$, while the S_1 component, for $(S_1) = m$, $c_a \in S_1$, $-Y_{a,1}$. All other components are zero. If $k = 4$, $m=3$, and none of the sets are BC ranked, then there is no relationship among the admitted rankings.

In part c, the new normal vector associated with c_a is where she is top-ranked in all sets of m candidates that she belongs to while the remaining candidates are tied for bottom ranked, yet she is bottom ranked in the set of k candidates while the rest of the candidates are tied for top rank. This is a normal vector so such a ranking is not an admissible BC outcome. On the other hand, this ranking is an admissible election outcome for any other choice of a social ordering vector. Along with Proposition 5, the dimension of the normal bundle indicates the number of cycles and other phenomena avoided by the BC. The uniqueness assertion from part a extends to part c for $k = 4$, $m = 3$, and to other values of k and m . But, the approach used to prove part a becomes overly complicated to prove part c. Therefore, a different kind of argument (this is not a step-function) is required. In effect, this uniqueness assertion of part c is strong: if even one of the sets is not BC ranked while the other three are, then there are no relationships among the set of all possible election outcomes.

Proof of part a. Suppose the sets listed in part a are S_1 , S_2 , S_3 , and S_4 . To prove the conclusion, it suffices to show that the vectors $(Z_{k-1,i})$, $i = 1, 2, 3, \dots, k \in \text{lin}(S_1)$, determine the specified normal bundle over the specified linear subspace of $\mathbb{R}^k$. This is a straightforward computation. The uniqueness assertion is proved in the next section.

Example 11. (a) Because $\text{Eip}(\mathcal{B})$ satisfies the neutrality condition, other choices of normal vectors can be determined from the first specified one by using the permutation group theoretic structure. A quick way to find other normal vectors is to list the candidates in a repeating *chain*, such as $---c_1---c_2---c_1---c_1---c_1---c_2---$. Each candidate c_i is the *central candidate* in that subset consisting of c_i and the two candidates on either side of her in the chain. A normal vector for Corollary 3.6a is determined in the following manner: for each subset, assign the value -2 for the central element and 1 for the other two candidates. The first listed normal vector in the corollary comes from the above chain, while the second one comes from the chain $---c_1---c_2---c_1---c_2---c_1---c_2---$. Another normal vector (but linearly dependent) comes from the chain $---c_1---c_1---c_2---c_1---c_2---c_1---$, etc. One immediate consequence is that *it is impossible for all four of the central candidates to be BC bottom ranked, or to be BC top ranked*. The selection of normal vectors for other sets can be determined by a group-theoretic argument.

(b) The two specified normal vectors determine the relationships

$$3.7 \quad x_{1,7} + x_{1,8} + x_{1,9} + x_{2,10} = 0$$

$$3.8 \quad x_{1,7} + x_{2,8} + x_{3,9} + x_{4,10} = 0.$$

These are the only (independent) restrictions imposed on the heron outcomes.

Consequently, in either expression, it is impossible for the four variables all to be of the same sign. This implies, as already noted in part (a), that the central candidates cannot all be top (or bottom) ranked. In particular, it implies that the example demonstrating an organizational "money pump" never can occur with the $k=1$. On the other hand, these expressions do permit the rankings $c_1 \succ c_2 \succ c_3 \succ c_4$, $c_1 \succ c_1 \succ c_2$, $c_1 \succ c_1 \succ c_1$, and $c_2 \succ c_3 \succ c_4$; rankings that admit cycles. This is because these rankings require $x_{1,7}$, $x_{1,8}$, $x_{3,9}$, and $x_{2,10}$ to be positive. A simple way to show these choices are compatible with Eqs. 3.7, 3.8 is to set all of the variables in Eq. 3.7 equal to zero. This forces $x_{2,8}$ and $x_{4,10}$ to be negative. So, values exist to satisfy Eq. 3.8.

(c) These results provide another axiomatic representation of the rd. For this case, for $n=4$ candidates, *If a social choice function f satisfying the P is defined over all subsets of three candidates, if over each set it is unanimous, neutral, consistent, somewhat faithful, and if it never selects the*

candidate *not* being for an *elect*. Then *f* is equivalent to the *B* without the *elect* test. This serves as another alternative to Arrow's condition discussed in Corollary 3.4.

It is then asserted that the BC admits a relationship among the rankings of the four candidates and those of the four subsets of three candidates. The relationship is given by the specified normal vector n . Therefore, the new conditions relating the outcomes of the four sets are of the form

$$(3.9) \quad x_{123} + 6x_{134} + 6x_{234} + x_{124} = 2x_{1234}.$$

From this equation, all sorts of new conclusions, such as the impossibility of x_1 being BC bottom ranked in the three subsets of three candidates (which forces the left-hand side of Eq. (3.9) to be negative), yet BC top ranked in S_{123} (so $x_{123} > 0$) is immediate.

In a similar manner, it now is easy to show for n candidates and for any m where $n - m \geq 2$ that x_1 cannot be BC top (bottom) ranked in all possible sets of m candidates while being BC bottom (top) ranked in the set of all n candidates. The BC is the only positional voting vector with this property. (If another voting vector had this same property, then the linear space it defines would have the same dimension as the one for the BC.) As the vector space for the BC is contained in the new vector space, they must agree. Thus the new vector is a BC. When $m = 2$, this becomes a restatement of Corollary 3.1c. Moreover, by following the earlier arguments given in this paper, all of the results given in terms of x and the ranking of n candidates extend to this new situation.

4. Proofs.

The proofs of most of the assertions either are contained in the bulk of this paper, or they are immediate. The proofs of the certain of the remaining statements follow directly from arguments developed in [3]. For instance, Proposition 2 and Theorem 1 are proved in [3]. The proof of Theorem 2 also is given in the last section of [3]; indeed, while Theorem 2 is not formally stated in [3], the proof of this statement is the manner in which Theorem 1 is proved. As an outline, recall that a basis for $V(BP)$ is determined in the following manner. Starting with a ranking $\pi(A)$, a new ranking $\pi(A)'$ is determined by

transposing some (or, equivalently, ranked) candidates, the basis is found by computing

$$(4.1) \quad \mathbf{B}_{\text{out}(V)}^0 - \mathbf{B}_{\text{in}(V)}^0,$$

where the system vector $\mathbf{B}^0$ is expressed in a vector normalized form. The resulting vector has a fixed vector form in each subset of candidates containing these two candidates. By carefully choosing the two candidates that are to be transposed, it is shown that $\dim(V(\mathbf{B}^0)) = n(n-1)/2$. Moreover, as also shown in [3], each of the above basis vectors are in $V(\mathbf{W}^0)$ for an appropriately scalar normalized of $\mathbf{W}^0$. To prove this, the vector from $V(\mathbf{W}^0)$ involves the tally from more than one voter, and then an expression of the form given in (4.1) is used. This computation uses nothing more than the fact that when $\mathbf{W}^0$ is expressed in a vector normalized form, each voting vector component is non-zero. Thus, the first part of Theorem 2 holds.

Proof of Corollary 4.1. In the proof (Section 4.2) that the B^0 run-off election has the abstention paradox, the key step involved using three words that differed only in one symbol. The proof for the general case is much the same: again I use three words from $B(B^0)$ that differ only in a specified symbol. One choice for this symbol has the relative rankings of two candidates fixed; a second choice has one candidate ranked above the other, and the third choice reverses this relative ranking. To do this, I first show that such rankings can be found: namely, if two words in $B(B^0)$ differ only in one symbol and if the only difference in that symbol has two adjacent ranked candidates in reversed positions, then the voter whose this symbol has a tie vote between these candidates also is in $B(B^0)$. For instance, if $(c_1 \succ c_2, c_1 \succ c_3, c_2 \succ c_4, c_1 \succ c_2 \succ c_3)$ and $(c_1 \succ c_2, c_1 \succ c_4, c_2 \succ c_3, c_2 \succ c_1 \succ c_4)$ are in $B(B^0)$, then so is $(c_1 \succ c_2, c_1 \succ c_3, c_2 \succ c_4, c_1 \succ c_2 \succ c_3)$.

This assertion is immediate. There are several vectors in $V(\mathbf{W}^0)$ corresponding to each of the first two words: let v_1 be a vector corresponding to the first word, v_2 to the second, and $v_t = tv_1 + (1-t)v_2$ to a point on the line segment between them. Of course, because $V(\mathbf{W}^0)$ is a vector space, $v_t \in V(\mathbf{W}^0)$.

From Theorem 6 in [3], the region of profiles corresponding to a fixed symbol is a convex region, therefore all but one of the symbols of v_t, v_1, v_2 must vary. The remaining symbol is where the rankings of the two candidates are reversed. Assume without loss of generality that this is for the ranking of the set S_1 and that the candidates are $\{c_1, c_2\}$. This means that in the appropriate

common to some of $\mathcal{W}^0$, one ranking has $x_{1,j}, x_{2,j}$, and the other has the reversed inequality. For all other values of $x_{k,j}$, the same ordering applies. It now is immediate that there is a value of i so that the c_1, c_2 coordinates for v_i are $x_{1,j}, x_{2,j}$, while the remaining coordinate orderings are preserved. This completes the proof of the assertion.

If a social choice method based on B^0 admits the abstention property, then there are three words in $B(B^0)$, based on three candidates, c_1, c_2, c_3 where c_1 and c_3 are possible outcomes depending on the rankings of two possible final sets. The choice of which final set occurs depends on whether c_2 can be advanced one position in the ranking of a swing set. Start with the ranking of the swing set where c_2 is tied in the swing position. Choose the rest of the symbols of the word in $B(B^0)$ so that if c_2 is lowered from the tie position, then c_2 is the outcome, but if c_2 is advanced from the tie, then c_1 is the final outcome. (again, this is possible because the disjoint method has the abstention property.) This identifies the words that changes the outcomes.

The selected profile will be associated with the word with the tie vote in the swing set. Construct the rankings for the two voters with c_2 as top ranked, c_1 is bottom ranked, and c_3 is ranked somewhere in the middle. By use of Theorem 6 of [31], p can be selected so that it includes one of these voters. So, of both vote, the outcome is the undesired c_1 , if they abstain, the outcome is the more desired c_3 . Because this argument is based on a particular word being in $B(B^0)$ and because this same word is in all choices of $B(W^0)$, the same argument proves that this phenomenon occurs holds for all choices of positional voting vectors. (A slightly more complicated argument extends the conclusion to a much wider class of procedures.)

Proof of Theorem 4.a. This is a simple computation using the basis for $W(B^0)$ derived in [4].

Proof of part b. This follows immediately from Theorem 3.1 in [31]. An alternative proof can be derived by using the scheme developed to prove Corollary 3.6 on which is given below.

Proof of part c. This follows from Theorem 2 and the fact, pointed out above, that the main component of the proofs for other choices of W^0 is that when W^0 is expressed in a vector-normalized form, each voting vector component is non-zero.

Proof of Corollary 3.2.4. First I show that a somewhat faithful social choice function f cannot be single-valued. If it could be, then $f(p)$ is the same set from $PC(p)$ for all choices of p . Let $c_1 \in f(p)$, and let p_1 be a profile of a single voter where c_1 is bottom ranked. Then, according to the assumption of being somewhat faithful, $c_1 \notin f(p_1)$. This contradicts the assumption that f has only one value and proves the assertion.

According to assumption and the above, f is an anonymous, neutral, consistent social choice function that takes more than one value. Thus, according to Proposition 5, f is equivalent to a general scoring method. It now follows from Corollary 3.1b that the added assumption about f satisfying a condition involving pairwise rankings forces f to be equivalent either to the W or to the reversed PC . However, because f is somewhat faithful, the weight assigned to the bottom ranked candidate must be less than that for some other candidate, so f must be equivalent to choosing the top ranked candidates from the PC ranking. This completes the proof.

Incidentally, this assertion does not preclude the possibility of tie-breakers. For instance, suppose the binary condition is that the anti-majority candidate is not elected and that a single winner is being selected, then the tie-breaker is admitted. This is because, at the end of the first stage, the anti-majority candidate cannot be tied for first place. Thus, as the tie breaker is applied to those candidates that are top ranked, this does not involve the anti-majority candidate. $\square$

Proof of part 1. The PC need not have the Condorcet winner as top ranked, so, by part a, an impossibility assertion follows.

Proof of part a. According to part a, the method at the first stage must be the PC . Thus, the first stage is equivalent to selecting the top ranked candidates from the PC ranking. Now, suppose a tie breaker W_2 is admitted where W_2 is not a Borda Vector. According to Theorem 2, a profile can be found so that the pairwise votes are all ties (which forces the PC outcome to have all candidates tied for first place), but c_1 is the top ranked candidate of the W_2 tally. This means that the social choice mechanism selects c_1 . This choice of a simulation violates the assumption of non-determinacy. This completes the proof.

Incidentally, this is one of the few places I use the fact that the social choice function has a singleton in its range. On the other hand, if f does not admit any singleton sets in the range, only sets of at least two candidates, then the social choice method is equivalent to selecting the k top

number candidates from the electoral ranking. For instance, this may correspond to selecting a committee of k candidates. In this case the Borda ranking of $\mathcal{C}^n$ can be reduced by *a tie-breaker* to the top k candidates without coming into conflict with non-determinacy or consistency. To regain the same conclusion at a Borda election without use of *a tie-breaker* one just strengthens the non-determinacy condition to require more than k candidates in the set of f whenever all of the pairwise elections result in tie votes.

Proof of Corollary 3.6b. Let $k=n$ and let $\mathcal{W}_m$ be the subspace of $\mathcal{W}^n$ corresponding to the entries in the sets of m candidates. Let $\pi_m: \mathcal{W}^n \rightarrow \mathcal{W}_m$ be the obvious projection mapping. Part b holds iff the dimension of the vector space $\text{Pr}V(B^n)$ is $n(n-1)/2$. Because $\dim(V(B^n)) = n(n-1)/2$, we have that $\dim(\text{Pr}V(B^n)) \leq n(n-1)/2$. It remains to show that the dimension cannot be smaller than $n(n-1)/2$.

The space $\text{Pr}V(B^n)$ can be viewed (Corollary 3.5) as the image of a mapping $f: \mathcal{W}^n \rightarrow \mathcal{W}_m^m$ where each coordinate of f is given by Eq. (3.5). If the dimension of the image is less than $n(n-1)/2$, then, by the linearity of f , there is a point $\mathbf{p} \in \mathcal{W}^n$ and a direction $\mathbf{a}$ so that $f(\mathbf{p}+\mathbf{a}t) = f(\mathbf{p})$. By the Taylor series, this means that DF does not have rank $n(n-1)/2$.

The matrix DF is $\dim(\mathcal{W}_m^m) \times n(n-1)/2$ where each row corresponds to the value of $x_{i,j,k}$ while each column corresponds to a pair of candidates. It follows from Eq. (3.5) that the entries in this matrix are either unity or zero. To show that the rank of DF is $n(n-1)/2$, it suffices to show that the span of the row vectors includes the $n(n-1)/2$ coordinate vectors (in $\mathbb{R}^{n(n-1)/2}$), $\mathbf{e}_j$, with unity in the j^{th} component and zero in all others. To show this, first consider the coordinates associated with x_1 (i.e., $x_{1,j,k}$) and assume that the j -th columns of DF represents the pair $\{c_j, c_j\}$, $j = 2, \dots, n$. List all candidates other than c_1 according to subscript as $c_2, c_n, \dots, c_n, c_2, c_1, \dots$. From this list form the sets of candidates S_k , $k = 1, \dots, n-1$, where c_j is in S_k along with c_{k+1} and the $m-2$ candidates listed to the right of her. For instance, $S_1 = \{c_1, c_2, \dots, c_m\}$, $S_3 = \{c_1, c_4, \dots, c_{m+2}\}$, and $S_{n-1} = \{c_1, c_n, c_2, \dots, c_{m-1}\}$. Assume that the first $n-1$ rows of DF are listed in the order $x_{1,j,k}$ where j corresponds to the sets S_j defined above. With this notation, the first row of DF (for $x_{1,1,1}$) has unity in the first and columns, and zero in all others. In general, the j^{th} row has unity on the main diagonal, and unity in the next $m-2$ columns. If there are not $m-2$ columns remaining in this $(n-1) \times (n-1)$ block, then the remaining number of rows are filled

starting from the right-hand side. All other entries in this row are zero.

It can be noticed that this matrix is a circulant matrix (see [17]) and the determinant of such a square determinant is non-zero, or by elementary row reduction, it is clear that the resulting $(n-1)/(n-1)$ block has non-zero determinant. This means that the row vectors with unity in a component corresponding to a pair (c_j, c_k) and zero in all others is in the span of the row vectors of (18). The same construction is continued with all other choices of c_j and the conclusion follows.

Part c. Again, let $k = n$. For this part, we are considering the projection of $V(B^n)$ into the product space of W^n_m with the vector space representing the outcomes of $S_{2^1, 2^2, \dots, 2^{n-1}}$. As this projection has a subspace of dimension $n(n-1)/2$ (by part a), the dimension of the projection is no less than $n(n-1)/2$. As $\dim(V(B^n)) = n(n-1)/2$, the projection has exactly this dimension. Thus, it remains to find the $n-1$ normal vectors that were not in Part b.

Start with candidate c_1 . For each term in the sum $\sum_j Z_{1,j} c_j$, where the summation is over the $(n-1)/(m-1)/(n-m)$ of the sets S_j containing c_1 and $m-1$ other candidates, the vector component for each pair is either $(1/2, -1/2)$ (c_1 is top ranked) or 0. Each pair gets counted in a non-zero fraction $(n-2)/(m-2)/(n-m)$ times. Thus, if $-1/(n-2)/(m-2)/(n-m) \sum_j Z_{1,j} 2^{n-1-j}$ is added to the sum, all of the components for pairs become zero. This is the vector described in the statement of the corollary. Clearly, by changing the choice of the candidate from c_1 to c_n , $n-1$ normal vectors are defined.

What remains is to show the uniqueness assertion of part a. Namely, I prove that if any of the four sets of three candidates is not ranked with a BC, then all possible election rankings occur. Let Q' denote the product space of the vector ranking regions for the four sets. Suppose there exist choices of voting vectors, W_j , that admit a relationship among the ordinal election rankings for the four sets of three candidates specified in part a. This means that the span of the election tables define a lower dimensional, linear subspace of Q' , so this linear subspace has a non-zero normal vector, N , in Q' .

My first assertion is that N must be a linear combination of the two normal vectors specified in part a of the corollary. Extend the above choices of W_j 's in an arbitrary manner to obtain a W^n . Recall from Theorem 2 that $V(B^n)$ is a proper subset of $V(W^n)$ for any W^n that is not a Borda system vector. (Of course, this requires the proper scalar normalization of the components of W^n .)

Therefore, a normal vector for $V(W)$ must be in the span of the normal vectors for $V(B)$. Thus, N must be in the span of the two vectors specified in part a of the corollary, namely, $N = a_1N_1 + a_2N_2$. Because we assume that $N \neq 0$, either a_1 or a_2 is non-zero. Without loss of generality, assume that $a_1 \neq 0$. Thus, N can be expressed as $N_1 + bN_2$.

A scalar normalization of the W_i 's that suffices for Theorem 2 is to assume that $W_i = (2, x_i, -2, x_i)$, $i = 1, 2, 3, 4$, where $-2 \leq x_i \leq 2$. Thus, each W becomes a point (unit if $x_i = 0$). As N is a normal vector, there are 4! = 24 resulting equations corresponding to the the scalar product of N with the vote tallies assigned to a profile consisting of a single voter where the voter's preferences vary over each of the possible permutations of $A = (v_1, v_2, v_3, v_4)$. This gives rise to a system of 24 equations with the four unknowns x_i . Because N is a linear combination of the normal vectors for the B tallies, each equation has at least one solution -- the B . It remains to show that this is the unique solution. This requires finding four equations where the coefficients, determined by the entries of N , are linearly independent.

Looking at the equations for the cyclic permutations of the ranking (v_1, v_2, v_3, v_4) , (v_2, v_3, v_4, v_1) , (v_3, v_4, v_1, v_2) , and (v_4, v_1, v_2, v_3) , the equations for $X = (x_1, x_2, x_3, x_4)$ becomes $BX = 0$, where B is a 4×4 matrix with entries that are scalars or scalar multiples of b (the multiple of N_2 in the definition of N). The unique solution for this system is the B , as long as the matrix B is non-singular. The determinant of B equals $(-5+5b+b^2)(1-b-b^2)$, so for $b \neq -5$ and $b \neq -2$ are the only real values of b that cause this matrix to be singular. On the other hand, if one looks at the equations resulting from the cyclic permutations of the ranking (v_1, v_2, v_3, v_4) , a similar matrix equation results where the determinant of the coefficient matrix has the value $(-1-b-b^2)(-1-b^2+7b)$. This new polynomial has its real roots at the values $b = (-5 \pm \sqrt{17})/2 \approx -0.79$. Thus, no matter what is the value of b , there exists some selection of four independent equations that force the the solution of the system to be the B . This completes the proof.

References

1. Ales, J. (1994). "Public Social Choice and Individual Values." *Journal of Economic Theory*, 69, 1-24.
2. Fishburn, P. and S. Gehrlein. "Borda's rule, positional voting, and Condorcet's simple majority principle." *Public Choice* 70-82.

3. Smith, D. W. (1987) A foundation for Young's method. To appear in *J. Econ. Theory*.
4. Smith, D. W. (1988) Symmetric voting and social choice. *Math. Econ.* **10**: 101-121.
5. Smith, D. W. (1987) Union and intersection of decision mechanisms: systems of W. J. Crammink and K. Sigmund. *Lecture Notes in Economics and Mathematical Systems* **287**. Springer-Verlag, New York.
6. Smith, D. W. (1989) The arithmetic of choice resulting from sequential voting systems. *Journal of Applied Mathematics* 61:261-266.
7. Smith, D. W. (1990) The mathematics of weighted summation voting systems. *Math. Econ. Research* **7**: 475-490.
8. Smith, D. W. (1990) Conditions and extensions of Arrow's theorem. *U. Wisconsin paper*.
9. Smith, D. W. (1990) Generalizations admitting families of simple rules. *U. Wisconsin paper*.
10. Smith, D. W. (1976) Aggregation of preferences with variable and variable parameters. **41**: 1027-1041.
11. Young, H. P. (1974) An axiomatization of form's function. *Econ. Theory* **3**: 1-71.
12. Young, H. P. (1977) Social choice scoring functions. *SIAM J. Appl. Math.* **28**: 544-548.
13. Van Woudenberg, G. (1988) A geometric investigation of voting techniques: a comparison of Arrow's Voting postulate, Young's techniques and the rank count. Northeastern University. Ph.D. dissertation, August, 1988.