

Gilboa, Itzhak; Monderer, Dov

Working Paper

Quasi-Values on Subspaces

Discussion Paper, No. 855

Provided in Cooperation with:

Kellogg School of Management - Center for Mathematical Studies in Economics and Management Science, Northwestern University

Suggested Citation: Gilboa, Itzhak; Monderer, Dov (1989) : Quasi-Values on Subspaces, Discussion Paper, No. 855, Northwestern University, Kellogg School of Management, Center for Mathematical Studies in Economics and Management Science, Evanston, IL

This Version is available at:

<https://hdl.handle.net/10419/221214>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.

Discussion Paper No. 855

QUASI-VALUES ON SUBSPACES

by

Itzhak Gilboa^{*}

and

Dov Monderer^{**}

September 1989

^{*} Department of Managerial Economics and Decision Sciences, J.L. Kellogg Graduate School of Management, Northwestern University, Evanston, IL 60208.

^{**} Department of Industrial Engineering and Management. The Technion, Haifa 32000, Israel, visiting Department of Managerial Economics and Decision Sciences, J.L. Kellogg Graduate School of Management, Northwestern University, Evanston, IL 60208.

Abstract

Quasi-values are operators satisfying all axioms of the Shapley value with the possible exception of symmetry. We introduce the characterization and extendability problems for quasi-values on linear subspaces of games, provide equivalence theorems for these problems, and show that a quasi-value on a subspace Q is extendable to the space of all games iff it is extendable to $Q + \text{Sp}\{u\}$ for every game u .

Finally, we characterize restrictable subspaces and solve the characterization problem for those which are also monotone.

1. Introduction

Quasi-values are solutions of TU cooperative games that satisfy all the axioms of a value, except possibly the symmetry axiom. A complete characterization of quasi-values was given in Weber (1988): " ψ is a quasi-value iff there exists a probability distribution over the set of orders of the players such that $\psi v(i)$ is the expected marginal distribution of player i to its predecessors."

Here we deal with quasi-values defined on subspaces of games. Two main questions are discussed:

- (i) The characterization problem. Given a subspace of games Q , find a finite number of linear conditions on operators $\psi: Q \rightarrow \mathbb{R}^n$ (where n is the number of players) that are necessary and sufficient to ensure that ψ is a quasi-value.
- (ii) The extension problem. Given a quasi-value on a subspace of games, find a finite number of linear conditions that are necessary and sufficient for its extendability to a quasi-value on the space of all games.

This work was motivated by our attempt to solve the binary stochastic choice problem, which has drawn the attention of many economists and psychologists in the last five decades. For a description of this problem, many of its interpretations, and additional references, the reader is referred to Fishburn (1988). The technical description of the problem is the following.

Find a finite number of linear conditions on vectors $\beta = (p_j^i)_{i \neq j}$ that are necessary and sufficient for the existence of a probability distribution Pr on the set of all orders of $N = \{1, 2, \dots, n\}$ such that

$$p_j^1 = \Pr(i > j) \quad \forall i \neq j,$$

where $>$ denotes a generic order on N .

Monderer (1989) pointed out that the binary stochastic choice problem is equivalent to the extension problem for quasi-values on Q_2 , where Q_2 is the space spanned by all unanimity games on at most two-element sets. He also showed that the characterization problem for quasi-values on the space of all games turns out to be equivalent to another problem arising in social choice and psychology, the stochastic choice problem. This problem was formulated by Block and Marschak (1960), and was solved by Falmagne (1978). Monderer (1989) gave an independent game theory-based proof utilizing Weber's characterization mentioned above. Lately, Gilboa and Monderer (1989) used the game theoretic approach to obtain additional partial progress on the binary stochastic choice problem.

Our main results are the following: we formulate a condition $c(u)$ for every game u . It turns out that a linear operator ψ on a subspace Q is a quasi-value iff it satisfies $c(u)$ for all $u \in Q$. Furthermore, every quasi-value ψ is extendable to a quasi-value over the space of all games iff it satisfies $c(u)$ for all $u \notin Q$.

Using the above, we also prove that a quasi-value on Q can be extended to a quasi-value on the space of all games iff it can be extended to a quasi-value on $Q + u$ for all $u \notin Q$, where $Q + u$ is the linear space spanned by Q and u . These results are to be found in Section 2.

Next we solve the characterization problem for monotone restrictable subspaces. (Restrictable subspaces were studied by Neyman (1989) and Gilboa

(1989); monotonicity of a restrictable subspace is defined in Section 3. (The definition makes sense only in view of Lemma 3.1, which characterizes restrictable subspaces.)

2. General Subspaces

We start with some preliminaries. Let $N = \{1, 2, \dots, n\}$ be the set of players. The set of all TU cooperative games on N will be denoted by G . The subspace of all additive games will be denoted by A . A will be identified with the Euclidean space \mathbb{R}^n in the usual way. For $x, y \in \mathbb{R}^n$ we will write $x \geq y$, if $x_i \geq y_i \ \forall i \in N$.

The set of all one-to-one functions $\theta: N \rightarrow N$ (i.e., the set of all permutations) will be denoted by R .

For each $v \in G$ we define two vectors, v_* and v^* in \mathbb{R}^n as follows:

$$v_*(i) = \min_{s \subseteq N \setminus i} (v(s \cup i) - v(s)), \ \forall i \in N,$$

and

$$v^*(i) = \max_{s \subseteq N \setminus i} (v(s \cup i) - v(s)), \ \forall i \in N.$$

Throughout this paper, Q will denote a linear subspace of G . An operator, $\psi: Q \rightarrow A$ is called a Milnor operator if

$$v_* \leq \psi v \leq v^*, \ \forall v \in Q.$$

A quasi-value on Q is a linear efficient Milnor operator $\psi: Q \rightarrow A$. If Q contains the additive games, then by Monderer (1988), a linear operator $\psi: Q \rightarrow A$ is a Milnor operator iff it is a positive projection. So, for

subspaces $Q \supseteq A$, and in particular for $Q = G$, our definition coincides with the usual definition of quasi-values.¹

Let Pr be a probability distribution over the set, R , of permutations. Following Weber (1988), define the random-order value ψ_{Pr} on G as follows:

$$\psi_{Pr} v(i) = \sum_{\theta \in R} [v(S_{\theta}^i \cup i) - v(S_{\theta}^i)] Pr(\theta),$$

where

$$S_{\theta}^i = \{j \in N: \theta j < \theta i\}.$$

Weber (1988) proved that $\psi: G \rightarrow A$ is a quasi-value iff it is a random-order value.

Given a function $\psi: Q \rightarrow A$ and $u \in G$, we say that the condition $c(u)$ is satisfied by ψ if

$$c(u): \sum_{i=1}^n \psi v^i(i) \leq \sum_{i=1}^n (v^i - u)^*(i) + u(N), \quad \forall v^1, \dots, v^n \in G.$$

We now state the following:

Theorem A: Let $\psi: Q \rightarrow A$ be a linear operator. Then ψ is a quasi-value iff $c(u)$ holds for all $u \in Q$. //

Theorem B: Let $\psi: Q \rightarrow A$ be a quasi-value. Then ψ can be extended to a quasi-value on G iff $c(u)$ holds for all $u \notin Q$. //

Theorem C: Let $\psi: Q \rightarrow A$ be a quasi-value. Then ψ can be extended to a

¹Quasi-values are called also nonsymmetric values.

quasi-value on G iff ψ can be extended to a quasi-value on $Q + u$ for all $u \notin Q$. //

Note that Theorem A cannot be considered as a solution to our characterization problem because $c(u)$ involves infinitely many linear inequalities. Similarly, Theorem B does not solve the extension problem.

Proof of Theorem A: Let $\psi: Q \rightarrow A$ be a quasi-value, and let $u \in Q$. As ψ is a Milnor operator, we have:

$$(2.1) \quad \psi(v^i - u)(i) \leq (v^i - u)^*(i) \quad \forall i \in N.$$

Summation of (2.1) over $i \in N$ yields:

$$(2.2) \quad \sum_{i=1}^n \psi v^i(i) \leq \sum_{i=1}^n (v^i - u)^*(i) + \sum_{i=1}^n \psi u(i).$$

By the efficiency axiom,

$$\sum_{i=1}^n \psi u(i) = \psi u(N) = u(N),$$

and whence $c(u)$ holds.

As for the converse, let $\psi: Q \rightarrow A$ be a linear operator s.t. $c(u)$ holds for all $u \in Q$. We have to show that ψ is a Milnor and efficient operator. Substitute in $c(u)$: $v^i = v$, $v^j = 0 \quad \forall j \neq i$, and $u = 0$. This gives us:

$$\psi v(i) \leq v^*(i) \quad \forall v \in Q \text{ and } \forall i \in N.$$

Applying the same arguments to $-v$ instead of v , while observing that $(-v)^* = -v_*$, yields $\psi v(i) \geq v_*(i) \forall v \in Q$ and $\forall i \in N$. Therefore, ψ is a Milnor operator. To establish efficiency, substitute in $c(u)$: $v^i = v \forall i \in N$, and $u = v$. This yields:

$$\psi v(N) = \sum_{i=1}^n \psi v(i) \leq v(N).$$

Applying the same arguments to $-v$ instead of v proves that ψ is efficient, which completes the proof of Theorem A.

The following lemma will be used in the proof of Theorem B. It is, however, interesting in its own right.

Lemma 2.1: Let $\psi: Q \rightarrow A$ be a quasi-value, and let $u \notin Q$. Then ψ can be extended to a quasi-value on the space $Q + u$ spanned by Q and u iff $c(u)$ and $c(-u)$ hold.

Proof: The "only if" part follows from Theorem A. Suppose now that $c(u)$ and $c(-u)$ hold. Define $y = (y_1, y_2, \dots, y_n)$ and $z = (z_1, z_2, \dots, z_n)$ as follows:

$$y_i = \sup_{v \in Q} (\psi v(i) - (v - u)^*(i)) \forall i \in N,$$

and

$$z_i = \inf_{v \in Q} ((v + u)^*(i) - \psi v(i)) \forall i \in N.$$

Claim: y and z are well defined, $y \leq z$, and

$$\sum_{i=1}^n y_i \leq u(N) \leq \sum_{i=1}^n z_i.$$

Suppose we have already proved the claim. Choose any $x \in \mathbb{R}^n$ that satisfies $y \leq x \leq z$ and $x(N) = u(N)$, and define $\psi_u: Q + u \rightarrow A$ by:

$$\psi_u(v + \alpha u) = \psi v + \alpha x \quad \forall v \in Q, \text{ and } \forall \text{ real number } \alpha.$$

We will show that ψ_u is a quasi-value. Obviously, ψ_u is linear and efficient. To prove that ψ_u is a Milnor operator we have to show that

$$(2.4) \quad (v + \alpha u)_* \leq \psi v + \alpha x \leq (v + \alpha u)^* \quad \forall \alpha, v.$$

Obviously, for all $w \in G$: $(-w)^* = w_*$ and $(\alpha w)^* = \alpha w^*$ for $\alpha > 0$. Therefore, (2.4) is equivalent to the following two sets of inequalities:

$$(2.5) \quad \psi v - x \leq (v - u)^* \quad \forall v \in Q,$$

and

$$(2.6) \quad \psi v + x \leq (v + u)^* \quad \forall v \in Q.$$

Equation (2.5) holds because $x \geq y$, and (2.6) holds because $x \leq z$.

Therefore, ψu is a quasi-value.

We still have to prove the claim.

Proof of Claim: For each $v \in Q$ substitute in $c(u)$: $v^i = v$, and $v^j = 0$

$\forall j \neq i$. Therefore, $\psi v(i) = (v - u)^*(i) \leq \sum_{j=1; j \neq i}^n (-u)^*(j) + u(N)$. Hence, y is well defined. Similarly, z is proved to be well defined (using the condition $c(-u)$). To prove that $y \leq z$ it suffices to show that for every $v^1, v^2 \in Q$

$$(2.7) \quad \psi v^1(i) - (v^1 - u)^*(i) \leq (v^2 + u)^*(i) - \psi v^2(i).$$

Indeed, because ψ is a Milnor operator,

$$(2.8) \quad \begin{aligned} \psi(v^1 + v^2)(i) &\leq (v^1 + v^2)^*(i) \\ &= ((v^1 - u) + (v^2 + u))^*(i) \leq (v^1 - u)^*(i) + (v^2 + u)^*(i), \end{aligned}$$

by the obvious property of the $*$ operator: $(v + w)^* \leq v^* + w^* \forall v, w \in G$.

Rearranging terms in (2.8) yields (2.7). Finally, substituting $v^i = v$ for all i in $c(u)$ and $c(-u)$ yields $\sum_i [\psi v(i) - (v - u)^*(i)] \leq u(N) \leq \sum_i [(v + u)^*(i) - \psi v(i)]$ for all v , whence $\sum_i y_i \leq u(N) \leq \sum_i z_i$ follows. This completes the proof of Lemma 2.1.

Proof of Theorem B: The "only if" part follows from Lemma 2.1. As for the converse, suppose $c(u)$ holds for all $u \in Q$. For $\theta \in R$, denote by ψ_θ the quasi-value defined on Q by the permutation θ . that is,

$$\psi_\theta v(i) = v(\{j \in N: \theta j \leq \theta i\}) - v(\{j \in N: \theta j < \theta i\}).$$

By the characterization theorem of quasi-values on G (Weber (1988)), it suffices to show that ψ belongs to the convex hull of $\{\psi_\theta: \theta \in R\}$. By

standard convex analysis arguments, ψ belongs to the convex hull of $\{\psi_\theta: \theta \in R\}$ iff it satisfies every linear inequality satisfied by $\{\psi_\theta: \theta \in R\}$. These may be summarized in:

$$(2.9) \quad \sum_{i=1}^n \psi v^i(i) \leq \max_{\theta \in R} \sum_{i=1}^n \psi_\theta v^i(i) \quad \forall v^1, v^2, \dots, v^n \in Q.$$

Let then $v^1, v^2, \dots, v^n \in Q$. We will show that (2.9) holds. Define $u \in G$ by:

$$u(S) = \max_{\theta \in R} \sum_{i \in S} \psi_\theta v_S^i(i), \quad \forall S \subseteq N,$$

where $v_S^i(T) = v^i(S \cap T) \quad \forall T \subseteq N$.

If $u \notin Q$, then $c(u)$ holds because of our assumption. If $u \in Q$, then $c(u)$ holds by Theorem A. Thus, we may use $c(u)$ to prove (2.9). Note that it suffices to show

$$(v^i - u)^*(i) \leq 0 \quad \forall i \in N,$$

or, equivalently, to show that

$$(2.10) \quad u(S \cup i) - u(S) \geq v^i(S \cup i) - v^i(S), \quad \forall i \in N, \quad \forall S \subseteq N \setminus i.$$

Let then $i \in N$ and $S \subseteq N \setminus i$. Denote $s = \#S$. Let $\theta \in R$ satisfy:

$$u(S) = \sum_{j \in S} \psi_\theta v_S^j(j).$$

Denote by τ the element of R for which

$$u(S \cup i) = \sum_{j \in S \cup i} \psi_{\tau} v_{S \cup i}^i(j)$$

and let $\bar{\theta}$ be a permutation such that

$$\theta_k > \bar{\theta}_i, \forall k \notin S \cup i$$

$$\bar{\theta}_i > \bar{\theta}_j, \forall j \in S$$

and

$$\bar{\theta}_j > \bar{\theta}_k \Leftrightarrow \theta_j > \theta_k, \forall j, k \in S.$$

Then

$$\begin{aligned} u(S \cup i) &= \sum_{j \in S \cup i} \psi_{\tau} v_{S \cup i}^j(j) \geq \sum_{j \in S \cup i} \psi_{\bar{\theta}} v_{S \cup i}^j(j) \\ &= \sum_{j \in S} \psi_{\bar{\theta}} v_{S \cup i}^j(j) + \psi_{\bar{\theta}} v_{S \cup i}^i(i). \end{aligned}$$

However, for $j \in S$, $\psi_{\bar{\theta}} v_{S \cup i}^j(j) = \psi_{\bar{\theta}} v_S^j(j)$, so that

$$\sum_{j \in S} \psi_{\bar{\theta}} v_{S \cup i}^j(j) = u(S).$$

Noting that

$$\psi_{\bar{\theta}} v_{S \cup i}^i(i) = v(S \cup i) - v(S)$$

completes the proof. //

Note that Theorem B and Lemma 2.1 prove Theorem C.

3. Restrictable Subspaces

Recall that for each $v \in G$ and $S \subseteq N$ the game v_S is defined by $v_S(T) = v(S \cap T) \forall T \subseteq N$. Q is restrictable if $v_S \in Q \forall S \subseteq N$ whenever $v \in Q$.

Let $\{w^T: T \neq \emptyset\}$ be the linear base of G consisting of all unanimity games. That is:

$$w^T(S) = \begin{cases} 1, & S \supseteq T \\ 0, & \text{otherwise.} \end{cases}$$

For each $\Sigma \subseteq 2^N \setminus \{\emptyset\}$ let Q_Σ be the linear space spanned by $\{w^T: T \in \Sigma\}$. Note that

$$(3.1) \quad w_{T_0}^T = \begin{cases} w^T, & \text{if } T_0 \supseteq T \\ 0, & \text{if } T_0 \not\supseteq T. \end{cases}$$

Therefore, Q_Σ is a restrictable space.

Lemma 3.1: Q is restrictable iff $Q = Q_\Sigma$ for some $\Sigma \subseteq 2^N \setminus \{\emptyset\}$.

Proof: The "if" part is true by (3.1). As for the converse, denote

$$\Sigma = \{T: w^T \in Q\}.$$

We will show that $Q = Q_\Sigma$. For this purpose, it suffices to show that for all $v \in Q$, if the representation of v in the base $\{w^T\}_{T \subseteq N}$ is $v = \sum \alpha_T w^T$, then $w^T \in Q$ whenever $\alpha_T = 0$.

We will prove the latter claim by an induction on the number k of sets

T for which $\alpha_T \neq 0$. For $k = 1$ the claim is trivial. Suppose the claim holds for $k < p$, and let $v = \sum \alpha_T w^T$ satisfy $v \in Q$ and $\#\{T: \alpha_T \neq 0\} = P$. Let T_0 be a minimal set (w.r.t. inclusion) such that $\alpha_{T_0} \neq 0$. By (3.1) $v_{T_0} = \alpha_{T_0} w^{T_0}$, and hence $w^{T_0} \in Q$. Denote $v_0 = v - \alpha_{T_0} w^{T_0}$. Then $v_0 \in Q$, $v_0 = \sum_{T \neq T_0} \alpha_T w^T$, and $\#\{T \neq T_0: \alpha_T \neq 0\} = P - 1$. Therefore, by the induction hypothesis $w^T \in Q$, $\forall T \neq T_0$ for which $\alpha_T \neq 0$. //

We now turn to formulate the characterization problem for restrictable spaces, and to solve it for the class of monotone restrictable spaces. Let $\Sigma \subseteq 2^N \setminus \{\emptyset\}$. Let $\beta = (\beta^i(T))_{T \in \Sigma; i \in N}$ be a vector of real numbers. Define a linear operator $\psi_\beta: Q_\Sigma \rightarrow A$ by defining it on the base $\{w^T: T \in \Sigma\}$ as follows:

$$\psi_\beta w^T(i) = \beta^i(T), \quad \forall i \in N.$$

The Characterization Problem on Q_Σ

Find a finite number of linear inequalities on β that are necessary and sufficient for ψ_β to be a quasi-value on Q_Σ .

Obvious necessary conditions are:

$$(3.2) \quad \beta^i(T) = 0, \quad \forall i \notin T \in \Sigma,$$

$$(3.3) \quad \beta^i(T) \geq 0, \quad \forall i \in T \in \Sigma,$$

and

$$(3.4) \quad \sum_{i \in T} \beta^i(T) = 1, \quad \forall T \in \Sigma.$$

In some trivial cases the conditions (3.2)-(3.4) are also sufficient. E.g., if for all $S \neq T \in \Sigma$, $S \not\subseteq T$ and $T \not\subseteq S$. A class $\Sigma \subseteq 2^N \setminus \{\emptyset\}$ is monotone if $T \in \Sigma$ and $S \supseteq T$ imply $S \in \Sigma$. A restrictable subspace Q_Σ is said to be monotone if Σ is such. Let Σ be a monotone class. For each $i \in T \in \Sigma$ define:

$$(3.5) \quad c_{\beta}^i(T) = \sum_{k=t}^n (-1)^{k-t} \sum_{A \supseteq T; |A|=k} \beta^i(A),$$

where $t = \#T$.

Theorem D: Let Σ be a monotone class, and let $\beta = (\beta^i(T))_{T \in \Sigma; i \in N}$. Then ψ_{β} is a quasi-value on Q_Σ iff β satisfies (3.2)-(3.4), and the following conditions:

$$(3.6) \quad c_{\beta}^i(T) \geq 0, \quad \forall i \in T \in \Sigma,$$

and

$$(3.7) \quad \sum_{\{T \in \Sigma: i \in T\}} c_{\beta}^i(T) \leq 1, \text{ for every } i \in N.$$

Proof

Necessity

Suppose ψ_{β} is a quasi-value. We have to show that (3.6) and (3.7) hold.

For each $T \neq N$ define the game v^T as follows:

$$v^T(S) = \begin{cases} 1, & T \subset S \\ 0, & T \not\subseteq S \end{cases}$$

(where \subset denotes strict inclusion).

It is well known, and also easily verified, that for every $B \subset N$

$$(3.8) \quad v^B = \sum_{k=t+1}^n (-1)^{k-t+1} \sum_{|A|=k; A \supset B} w^A.$$

Fix $T \in \Sigma$ and $i \in T$. Then by (3.8) $v^{T \setminus i} = v_1^{T, i} + v_2^{T, i}$, where

$$v_1^{T, i} = \sum_{k=t+1}^n (-1)^{k-t+1} \sum_{\{A \mid |A|=k; A \supset T\}} w^A$$

and

$$v_2^{T, i} = \sum_{k=t+1}^n (-1)^{k-t+1} \sum_{\{A \mid |A|=k; i \notin A \supset T \setminus i\}} w^A.$$

Note that

$$v_1^{T, i}(S \cup i) - v_1^{T, i}(S) = v^{T \setminus i}(S \cup i) - v^{T \setminus i}(S)$$

for all $S \subseteq N \setminus i$, whence $(v_1)_*(i) \geq 0$. Since Q is monotone, $v_1^{T, i} \in Q$ and the Milnor condition yields $\psi_{\beta} v_1^{T, i}(i) \geq 0$. However, $\psi_{\beta} v_1^{T, i}(i) = c_{\beta}^i(T)$, and (3.6) follows.

To prove (3.7), let there be given $i \in N$. Denote

$$v = \sum_{\{T \in \Sigma: i \in T\}} v_1^{T, i}.$$

By linearity,

$$(3.9) \quad \psi_{\beta} v(i) = \sum_{\{T \in \Sigma: i \in T\}} c^i(T).$$

On the other hand, for each $i \in S \in \Sigma$, $v(S) - v(S \setminus i) = 1$ (because $v_1^{T,i}(S) - v_1^{T,i}(S \setminus i) = 1$ iff $S = T$), and by monotonicity of Q for each $i \in S \notin \Sigma$, $v(S) - v(S \setminus i) = 0$. Therefore, $v^*(i) \leq 1$. Combining this with (3.9) and the Milnor condition yields (3.7).

Sufficiency

ψ is obviously linear and efficient. We proceed to prove that it is a Milnor operator. Fix $i \in N$. Denote

$$\Sigma_i = \{T \in \Sigma: i \in T\}$$

and

$$\bar{\Sigma}_i = \{T \in \Sigma: i \notin T\}.$$

Obviously,

$$Q_{\Sigma} = Q_{\Sigma_i} + Q_{\bar{\Sigma}_i}.$$

As $\psi_{\beta} v(i) = v^*(i) = 0$ for all $v \in Q_{\bar{\Sigma}_i}$ (by (3.2)), it suffices to show that

$$\psi_{\beta} v(i) \leq v^*(i), \quad \forall v \in Q_{\Sigma_i}.$$

It is easily verifiable that $\{v_1^{T,i}: i \in T \in \Sigma\}$ is a linear base for Q_{Σ_i} .

Let $v = \sum_{i \in T \in \Sigma} d_T v_1^{T,i}$. Then

$$v^*(i) = \max\{0, \max_{i \in T \in \Sigma} d_T\}.$$

On the other hand,

$$\psi_{\beta} v(i) = \sum_{i \in T \in \Sigma} d_T c^i(T) \leq \max\{0, \max_{i \in T \in \Sigma} d_T\} \sum_{i \in T \in \Sigma} c^i(T) \leq v^*(i)$$

by (3.6) and (3.7).

Therefore, ψ is a Milnor operator. //

References

- Block, H. D. and J. Marshack (1960), "Random Orderings and Stochastic Theories of Responses," Cowles Foundation Paper No. 147.
- Falmagne, J. C. (1978), "A Representation Theorem for Finite Random Scale Systems," J. Math. Psych., 18, pp. 52-72.
- Fishburn, P. C. (1988), "Binary Probabilities Induced by Ranking," mimeo, AT&T Bell Laboratories, Murray Hill, New Jersey, to appear in J. Math. Psych.
- Gilboa, I. (1989), "Additivizations of Non-Additive Measures," Mathematics of Operations Research, 14, 1-17.
- Gilboa, I. and D. Monderer (1989), "A Game-Theoretic Approach to the Binary Stochastic Choice Problem," mimeo.
- Monderer, D., (1988), "Values and Semi-Values on Subspaces of Finite Games," Int. J. of Game Theory, 17, 4, pp. 301-310.
- Monderer, D. (1989), "The Stochastic Choice Problem: A Game Theoretic Approach," mimeo.
- Neyman, A. (1989), "Uniqueness of the Shapley Value," Games and Economic Behavior, 1, 1, pp. 116-118.
- Weber, R. J. (1988), "Probabilistic Values for Games," in The Shapley Value, A. E. Roth (ed.), Cambridge University Press, New York, pp. 101-119.