

A Service of

ZBW

Leibniz-Informationszentrum Wirtschaft Leibniz Information Centre for Economics

Caporale, Guglielmo Maria; Gil-Alana, Luis A.; Carmona-González, Nieves

Working Paper Particulate Matter 10 (PM10): Persistence and Trends in Eight European Capitals

CESifo Working Paper, No. 8402

Provided in Cooperation with: Ifo Institute – Leibniz Institute for Economic Research at the University of Munich

Suggested Citation: Caporale, Guglielmo Maria; Gil-Alana, Luis A.; Carmona-González, Nieves (2020) : Particulate Matter 10 (PM10): Persistence and Trends in Eight European Capitals, CESifo Working Paper, No. 8402, Center for Economic Studies and Ifo Institute (CESifo), Munich

This Version is available at: https://hdl.handle.net/10419/223474

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.



WWW.ECONSTOR.EU



Particulate Matter 10 (PM10): Persistence and Trends in Eight European Capitals

Guglielmo Maria Caporale, Luis A. Gil-Alana, Nieves Carmona-González



Impressum:

CESifo Working Papers ISSN 2364-1428 (electronic version) Publisher and distributor: Munich Society for the Promotion of Economic Research - CESifo GmbH The international platform of Ludwigs-Maximilians University's Center for Economic Studies and the ifo Institute Poschingerstr. 5, 81679 Munich, Germany Telephone +49 (0)89 2180-2740, Telefax +49 (0)89 2180-17845, email office@cesifo.de Editor: Clemens Fuest https://www.cesifo.org/en/wp An electronic version of the paper may be downloaded • from the SSRN website: www.SSRN.com

- from the RePEc website: <u>www.RePEc.org</u>
- from the CESifo website: <u>https://www.cesifo.org/en/wp</u>

Particulate Matter 10 (PM10): Persistence and Trends in Eight European Capitals

Abstract

This paper examines the statistical properties of daily PM10 in eight European capitals (Amsterdam, Berlin, Brussels, Helsinki, London, Luxembourg, Madrid and Paris) over the period 2014-2020 by applying a fractional integration framework; this is more general than the standard approach based on the classical dichotomy between I(0) stationary and I(1) non-stationary series used in most other studies on air pollutants. All series are found to be characterised by long memory and fractional integration, with orders of integration in the range (0, 1), which implies that mean reversion occurs and shocks do not have permanent effects. Persistence is highest in the case of Brussels, Amsterdam and London. The presence of negative trends in Brussels, Paris and Berlin indicates some degree of success in reducing pollution in these capitals.

JEL-Codes: C220, Q530.

Keywords: fractional integration, long memory, persistence, trends, air pollutants, PM₁₀.

Guglielmo Maria Caporale* Department of Economics and Finance Brunel University London United Kingdom – London, UB8 3PH Guglielmo-Maria.Caporale@brunel.ac.uk

Luis A. Gil-Alana University of Navarra Pamplona / Spain alana@unav.es Nieves Carmona-González Universidad Francisco de Vitoria Madrid / Spain

*corresponding author

June 2020

Prof. Luis A. Gil-Alana gratefully acknowledges financial support from the MINEIC-AEI-FEDER ECO2017-85503-R project from 'Ministerio de Economía, Industria y Competitividad' (MINEIC), 'Agencia Estatal de Investigación' (AEI) Spain and 'Fondo Europeo de Desarrollo Regional' (FEDER). He also acknowledges financial support from internal Projects of the Universidad Francisco de Vitoria.

1. Introduction

Particulate matter (PM) are microscopic particles of solid or liquid matter suspended in the air. Its sources can be natural or anthropogenic. It has a significant impact on both climate and precipitation, and therefore has both health and social costs. In particular, it affects the amount of incoming solar radiation and outgoing terrestrial radiation. The coarse particles can have a diameter between 2.5 and 10 micrometers (μ m) (PM₁₀) and are known to be a very harmful form of air pollution given their ability to penetrate into the lungs and blood streams and cause respiratory and heart diseases as well as premature death. Various countries have therefore set limits for particulars in the air, which are emitted during the combustion of vehicle engine fuels, braking and tyre wearing. In particular, the European Union has defined in a series of directives the acceptable limits for exhaust emissions of new vehicles sold in the European Union and EEA member states.

Numerous studies have analysed the connection between pollution and harmful health effects (e.g., Schwartz and Marcus, 1990; Anderson et al., 1996; Atkinson et al., 1999; Gardner and Dorling, 1999). The present study contributes to another branch of the literature which focuses instead on modelling various pollutants such as sulphur dioxide (SO₂), nitrogen dioxide (NO₂), carbon monoxide (CO), ozone (O₃), PM_{2.5} and PM₁₀. For instance, Zamri et al. (2009) applied the Box-Jenkins ARIMA approach to model CO and NO₂ in Malaysia and found an upward trend. Li et al. (2017) analysed air quality in Beijing from 2014 to 2016 using the Spatio-temporal Deep Learning (StDL) model, the Time Delay Neural Network (TDNN) model, the ARMA model, the Support Vector Regression (SVR) model, and the Long Short-Term Memory Neural Network Extended (LSTME) model, and concluded that the LSTME model is the most suitable one for time series characterised by long-term dependence with optimal time delays.

Naveen and Anu (2017) studied air quality in India using ARIMA, seasonal ARIMA (SARIMA) and other models. Pan and Chen (2008) is one of the few studies using longmemory AutoRegressive Fractional Integrated Moving Average (ARFIMA) models for air pollution data (in Taiwan) and concluding that these are more accurate than AutoRegressive Integrated Moving Average (ARIMA) models.

It is clearly important to investigate the dynamics of air pollution to develop suitable models for prediction purposes and design policies to manage air quality. This paper examines the statistical properties of daily PM_{10} in eight European capitals (Amsterdam, Berlin, Brussels, Helsinki, London, Luxembourg, Madrid and Paris) over the period 2014-2020 by applying a fractional integration framework that is more general than the standard approach based on the classical dichotomy between I(0) stationary and I(1) non-stationary series used in the vast majority of previous studies on air pollutants, since it allows for fractional as well as integer degrees of differentiation and thus for a much wider set of stochastic behaviours. In particular, it enables the researcher to analyse the long-memory properties of the series of interest and the possible presence of trends, to test for mean reversion, and to measure the degree of persistence and the speed of adjustment to the long-run equilibrium level. Therefore, it provides information about whether the effects of shocks are transitory or permanent, which is a crucial piece of information for adopting appropriate policy measures.

The remainder of the paper is structured as follows: Section 2 outlines the methodology used for the analysis; Section 3 describes the data; Section 4 presents the empirical results; Section 5 offers some concluding remarks.

2. Methodology

As mentioned above, we adopt a long-memory approach based on fractional integration. Long memory is a feature of time series that are characterised by a high degree of dependence between observations which are far apart in time. It has been found to be displayed by many time series in different fields such as climatology (Gil-Alana, 2005; 2008; 2017; Vyushin and Kushner, 2009; Franzke, 2012; Ludescher et al., 2016; Bunde, 2017; Yuan et al., 2019); environmental sciences (Barros et al., 2016; Gil-Alana et al., 2016; Tiwari et al., 2016; Gil-Alana and Solarin, 2018); economics and finance (Gil-Alana and Moreno, 2012; Abritti et al., 2017); etc.

There exist a variety of statistical models that can describe this type of behaviour; a very popular one among time series analysts is based on the concept of fractional integration, which occurs when the number of differences required to make a series stationary I(0) is a fractional value. More precisely, a time series is said to be integrated of order d or I(d) if it can be expressed as:

$$(1 - B)x_t = u_t, t = 1, 2, ..., (1)$$

where B is the backshift operator ($Bx_t = x_{t-1}$), the differencing parameter d can be any real value, and u_t is I(0) defined as a covariance stationary process with a spectral density function that is positive and bounded at all frequencies in the spectrum. This framework encompasses different cases such as short memory (d = 0), stationary long memory (0 < d < 0.5), non-stationary though mean-reverting processes ($0.5 \le d < 1$), unit roots (d = 1) and explosive patterns (d ≥ 1).

3. Data

The series analysed is the daily average air quality taken from the World Air Quality Index (WAQI) at https://aqicn.org/map/world/es/. All data have been converted using the US EPA standard (United States Environmental Protection Agency). Specifically, we use daily data for the past 7 years (2014-2020) concerning eight European capitals: Amsterdam, Berlin, Brussels, Helsinki, London, Luxembourg, Madrid and Paris. The series represents the daily level of air quality (PM10) measured in micrograms per cubic meter of air (ug/m3). The WAQI data come from the following original sources: Madrid: http://www.mambiente.madrid.es/opencms/opencms/calaire/ (Ayuntamiento de Madrid); Paris: http://www.airparif.asso.fr/ (AirParif - Association de surveillance de la qualité de l'air en Île-de-France); Amsterdan: https://www.luchtmeetnet.nl/ (RIVM); Luxembourg: https://environnement.public.lu/fr.html (Portail de l'Environnement du Grand-duché de Luxembourg); London: https://uk-air.defra.gov.uk/ (UK-AIR, air quality information resource Defra. UK); Helsinki: https://www.ilmatieteenlaitos.fi/ilmanlaatu (Ilmanlaatu Suomessa); Brussels: https://www.irceline.be/en/ (Belgian Interregional Environment Agency); Berlin: https://www.berlin.de/senuvk/umwelt/luftqualitaet/ (Luftqualität).

[TABLE 1 ABOUT HERE]

Table 1 reports the sample periods for each capital and provides some descriptive statistics for each series. It can be seen that Paris exhibits the highest mean value, while Helsinki has the lowest. Paris also has the most volatile series, whilst Luxembourg has the least volatile.

4. Empirical Results

We estimate the following model:

$$y_t = \alpha + \beta t + x_t, \quad (1 - B)^{a_0} x_t = u_t, \quad t = 1, 2, ...,$$
 (2)

where y_t stands for PM₁₀ in each European capital in turn; B is the backshift operator, and x_t is an I(d) process such that the error term u_t is I(0); the disturbances are assumed to follow a white noise (see Tables 2 and 3) and an autocorrelated process (see Tables 4 and 5) in turn, where the latter is modelled using the exponential spectral framework of Bloomfield (1973). In all cases we display the estimated values of d (and their associated 95% confidence bands) for three different specifications: (i) no deterministic terms in (1), i.e., we impose the restriction $\alpha = \beta = 0$ (the results for this case are reported in the second column in Tables 2 and 4); (ii) an intercept only (see the third column in both tables); (iii) an intercept and a linear time trend (see the fourth column in both tables). The estimated values reported in bold in these tables are those corresponding to our preferred specification, which has been selected on the basis of the statistical significance of the regressors.

[TABLES 2 AND 3 ABOUT HERE]

When assuming that u_t is a white noise, the intercept is found to be the only significant deterministic term in all cases; the estimated values of d are in the interval (0, 1), which implies long memory and fractional integration. They range between 0.39 (Amsterdam) and 0.62 (Madrid). For Amsterdam, the values are all within the stationary range (d < 0.5); for Brussels, London, Paris and London, they are around the stationary boundary (d = 0.5), while non-stationarity (d \geq 0.5) is found in the case of Helsinki, Berlin and Madrid.

[TABLES 4 AND 5 ABOUT HERE]

When allowing for autocorrelation (Tables 4 and 5) the time trend appears to be negative and statistically significant in the case of Brussels, Berlin and Paris, which might reflect the anti-pollution policies adopted in these three capitals. In particular, a low emission zone (LEZ) was established in the Brussels region with the aim of meeting the European air quality standards and emission ceilings; in Berlin the German Climate Action Plan 2050 is being implemented to control air pollution by laying down environmental quality standards and emission reduction requirements; a LEZ based on Euro norm vehicle classification has also been introduced in Paris.

The estimated values of d are once more in the interval (0, 1), though they are now significantly smaller than in the previous case. In fact, they are all within the stationary range, specifically between 0.22 (Luxembourg) and 0.33 (Helsinki and Paris). These lower estimates are likely to reflect the competition between the fractional integration and Bloomfield parameters in describing time dependence between the observations. Both sets of estimates, under the assumption of white noise and autocorrelated errors respectively, indicate that the degree of persistence is highest in the case of Brussels, Amsterdam and London, and lowest in the case of Helsinki, Berlin, and Madrid; thus, the effects of shocks are more long-lived in the former capitals.

5. Conclusions

This paper has used fractional integration methods to obtain evidence on persistence and time trends in PM_{10} in eight European capitals (Amsterdam, Berlin, Brussels, Helsinki, London, Luxembourg, Madrid and Paris). This approach is more general than the standard ones used in most of the literature on air pollutants and thus is more informative about the time series properties of the series of interest. The results indicate that all of them display fractional integration with orders of integration in the range

(0,1); this implies that mean reversion occurs and shocks do not have permanent effects. However, the degree of persistence is different in the eight capitals examined; in particular, the effects of shocks take longer to die away in the case of Brussels, Amsterdam and London. Such evidence should be taken into account by policy makers aiming to design effective measures to reduce pollution.

The estimated values of d are lower under the assumption of autocorrelated errors; in this case three of the capitals examined (Brussels, Paris and Berlin) exhibit statistically significant negative time trends, which suggests that the policies they have adopted to reduce pollution (such as the establishment of LEZs) have been successful, at least to some extent.

Other statistical properties of PM_{10} that could be investigated are seasonality, non-linearities and structural breaks; the forecasting properties of rival models could also be examined; all these issues are left for future work.

8

References

Abritti, M., L.A. Gil-Alana, Y. Lovcha and A. Moreno (2017) Term structure persistence, Journal of Financial Econometrics 14, 331-352.

Anderson HR, Ponce de Leon A, Martin-Bland J, Bower J.S., Strachan DP (1996) Air pollution and daily mortality in London: 1987-92, BMJ 312:665-669. http://doi:10.1136/bmj.312.7032.665

Atkinson RW, Anderson HR, Strachan DP, Bland JM, Bremner SA, Ponce de Leon A (1999) Short-term associations between outdoor air pollution and visits to accident and emergency departments in London for respiratory complaints Eur Respir J 13(2):257–265. http://doi: 10.1183/09031936.99.1322579

Barros, C.P., L.A. Gil-Alana and F. Perez de Gracia (2016), Stationarity and long range dependence of carbon dioxide emissions. Evidence from disaggregated data, Environmental and Resources Economics 63, 1, 45-56.

Bunde, A. (2017), Long-term memory in climate: Detection, extreme events and significance of trends, Chapter 11 in Nonlinear and Stochastic Climate Dynamics, edited by Christian L.E. Franzke and Terence O'Kane, Cambridge University Press

Franzke, C. (2012), Nonlinear trends, long-range dependence, and climate noise properties of surface temperature, *Journal of Climate*, 25(12), 4172–4183.

Gardner MW, Dorling SR (1999) Neural network modelling and prediction of hourly NOx and NO₂ concentrations in urban air in London. Atmos Environ 33:709-719. https://doi.org/10.1016/S1352-2310(98)00230-1

Gil-Alana, L.A. (2005), Statistical modeling of the temperatures in the Northern hemisphere using fractional integration techniques, Journal of Climate 27, 10, 3477-3491.

Gil-Alana, L.A. (2008), Time trend estimation with breaks in temperature series, Climatic Change 89, 3, 325-337.

Gil-Alana, L.A,. (2017) Alternative modeling approaches for the ENSO time series. Persistence and seasonality, International Journal of Climatology 37, 5, 2354-2363.

Gil-Alana, L.A., R. Gupta and F. Perez de Gracia (2016), Modelling persistence of carbon emissions allowance prices, Renewable and Sustainable Energy Review 55, 221-226.

Gil-Alana, L.A. and A. Moreno (2012), Uncovering the US term premium. An alternative route, Journal of Banking and Finance 36, 4, 1181-1193.

Gil-Alana, L. A., and Solarin, S. A. (2018). Have US environmental policies been effective in the reduction of US emissions? A new approach using fractional integration. Atmospheric Pollution Research, 9(1), 53-60.

Li X, Peng L, Yao X, Cui S, Hi Y, You C, Chi T (2017). Long short-term memory neural network for air pollutant concentration predictions: Method development and evaluation. Environ Pollut 231:1997-1004. <u>https://doi.org/10.1016/j.envpol.2017.08.114</u>

Ludescher, J., A. Bunde, C.L. Franzke and H. J. Schellnhuber (2016). Long-term persistence enhances uncertainty about anthropogenic warming of Antarctica, *Climate Dynamics*, 46(1–2), 263–271, doi: 10.1007/s00382-015-2582-5

Naveen V, Anu N (2017) Time series analysis to forecast air quality indices in Thiruvananthapuram District, Kerala, India. J Eng Res Appl 7(6): 66-84. https://doi: 10.9790/9622-0706036684

Pan JN, Chen ST (2008) Monitoring long-memory air quality data using ARFIMA model, Environmetrics 19 (2): 209–219. https://doi.org/10.1002/env.88

Schwartz O, Marcus A (1990) Mortality and air pollution in London: A time series analysis. Am J Epidemiol 131(1):185–194. https:// doi: 10.1136/jech.54.10.750

Tiwari, A.K., Kyophilavong, P. and Albulescu, C.T., (2016), Testing the stationarity of CO_2 emissions series in Sub-Saharan African countries by incorporating nonlinearity and smooth breaks, Research in International Business and Finance, 37, 527-540.

Vyushin, D. I. and P.J. Kushner (2009), Power-law and long-memory characteristics of the atmospheric general circulation, *Journal of Climate*, 22(11), 2890–2904.

Yuan, N., Y. Huang, J. Duan, C. Zhu, E. Xoplaki and J. Luterbacher (2019), On climate prediction; how much can we expect from climate memory?, Climate Dynamics 52, 1-2, 855-864.

Zamri IM, Roziah Z, Marzuki I, Muhd SL (2009) Forecasting and Time Series Analysis of Air Pollutants in Several Area of Malaysia. Am J Environ Sci 5:625-632. https://doi:10.3844/ajessp.2009.625.632

Table 1: Descriptive statistics

AMST	ERDAM	BERLIN		
Start date	End date		Start date	End date
01/01/2014	13/06/2020		20/08/2014	13/06/2020
No. of observati	ons: 2145		No. of observation	ons: 2114
M	ean: 25.7		Me	ean: 28.0
Standard deviat	ion: 11.1		Standard deviat	ion: 11.9
Varia	nce: 124.1		Varia	nce: 140.6
Min./M	lax.: 8 / 263		Min./M	ax.: 8 / 95
Rat	nge: 255		Rai	1ge: 87
BRUS	SSELS		HELS	SINKI
Start date	End date		Start date	End date
31/12/2013	13/06/2020		02/05/2014	13/06/2020
No. of observati	ons: 2334		No. of observation	ons: 2214
M	ean: 24.0	<i>Mean:</i> 18.1		ean: 18.1
Standard deviat	<i>ion:</i> 11.8	Standard deviation: 10.0		ion: 10.0
Varia	nce: 140.4	40.4 <i>Variance:</i> 99.9		
Min./Max.: 1 / 100 Min./Max.: 3 / 90		ax.: 3/90		
Rat	nge: 99	Range: 87		
LON	DON		LUXEM	IBOURG
LON Start date	DON End date		LUXEM Start date	BOURG End date
LON Start date 31/12/2013	End date 13/06/2020		LUXEM Start date 19/06/2015	IBOURG End date 13/06/2020
LON Start date 31/12/2013 No. of observation	End date 13/06/2020 ons: 2353		LUXEM Start date 19/06/2015 No. of observation	End date 13/06/2020 ons: 1583
LON Start date 31/12/2013 No. of observation Mo	End date 13/06/2020 ons: 2353 ean: 26.5		LUXEM Start date 19/06/2015 No. of observation Me	End date 13/06/2020 ons: 1583 ean: 19.5
LON Start date 31/12/2013 No. of observation Mathematical Standard deviat	End date 13/06/2020 ons: 2353 ean: 26.5 ion: 9.9		LUXEM Start date 19/06/2015 No. of observatio Me Standard deviat	End date 13/06/2020 ons: 1583 ean: 19.5 ion: 6.8
LON Start date 31/12/2013 No. of observation Mo Standard deviat Varia	End date 13/06/2020 ons: 2353 ean: 26.5 ion: 9.9 nce: 97.1		LUXEM Start date 19/06/2015 No. of observation Me Standard deviat Varian	End date 13/06/2020 ons: 1583 ean: 19.5 ion: 6.8 nce: 46.2
LON Start date 31/12/2013 No. of observation Mo Standard deviat Varian Min./M	End date 13/06/2020 ons: 2353 ean: 26.5 ion: 9.9 nce: 97.1 vax.: 5 / 89		LUXEM Start date 19/06/2015 No. of observation Ma Standard deviat Varian Min./M	End date 13/06/2020 ons: 1583 ean: 19.5 ion: 6.8 nce: 46.2 fax:: 2 / 52
LON Start date 31/12/2013 No. of observation Min Standard deviate Varian Min./Min Rate	End date 13/06/2020 ons: 2353 ean: 26.5 ion: 9.9 nce: 97.1 ax.: 5 / 89 nge: 84		LUXEM Start date 19/06/2015 No. of observation Ma Standard deviat Varian Min./M Ran	End date 13/06/2020 ons: 1583 ean: 19.5 ion: 6.8 nce: 46.2 ax.: 2 / 52 nge: 50
LON Start date 31/12/2013 No. of observation Min Standard deviat Varian Min./M Rat MA	End date 13/06/2020 ons: 2353 ean: 26.5 ion: 9.9 nce: 97.1 Jax.: 5 / 89 nge: 84		LUXEM Start date 19/06/2015 No. of observation Ma Standard deviat Varian Min./M Ran PA	End date 13/06/2020 ons: 1583 can: 19.5 ion: 6.8 nce: 46.2 cax.: 2 / 52 nge: 50
LON Start date 31/12/2013 No. of observation Ma Standard deviate Varian Min./M Rate MAN Start date	End date 13/06/2020 ons: 2353 ean: 26.5 ion: 9.9 nce: 97.1 Jax.: 5 / 89 nge: 84		LUXEM Start date 19/06/2015 No. of observation Ma Standard deviat Varian Min./M Ran PA Start date	End date 13/06/2020 ons: 1583 can: 19.5 ion: 6.8 nce: 46.2 ax.: 2 / 52 nge: 50
LON Start date 31/12/2013 No. of observation Mainer Min./M Rate MAI Start date 31/12/2013	End date 13/06/2020 ons: 2353 ean: 26.5 ion: 9.9 nce: 97.1 Jax.: 5 / 89 nge: 84 DRID End date 13/06/2020 13/06/2020		LUXEM Start date 19/06/2015 No. of observation Ma Standard deviat Varian Min./M Ran Start date 31/12/2013	End date End date 13/06/2020 ons: 1583 can: 19.5 ion: 6.8 nce: 46.2 ax.: 2 / 52 nge: 50 RIS End date 13/06/2020 13/06/2020
LON Start date 31/12/2013 No. of observation Main Standard deviate Varian Min./M Rate MAI Start date 31/12/2013 No. of observation	End date 13/06/2020 ons: 2353 ean: 26.5 ion: 9.9 nce: 97.1 Jax.: 5 / 89 nge: 84 DRID End date 13/06/2020 ons: 2323		LUXEM Start date 19/06/2015 No. of observation Ma Standard deviat Varian Min./M Ran PA Start date 31/12/2013 No. of observation	End date End date 13/06/2020 ons: 1583 ean: 19.5 ion: 6.8 nce: 46.2 fax.: 2 / 52 nge: 50 RIS End date 13/06/2020 ons: 2227
LON Start date 31/12/2013 No. of observation Main Standard deviate Varian Min./M Rate MAI Start date 31/12/2013 No. of observation Main	End date 13/06/2020 ons: 2353 ean: 26.5 ion: 9.9 nce: 97.1 Vax.: 5 / 89 nge: 84 DRID End date 13/06/2020 ons: 2323 ean: 24.3		LUXEM Start date 19/06/2015 No. of observation Ma Standard deviat Varian Min./M Ran PA Start date 31/12/2013 No. of observation Ma	End date 13/06/2020 ons: 1583 can: 19.5 ion: 6.8 nce: 46.2 ax.: 2 / 52 nge: 50 RIS End date 13/06/2020 ons: 2227 can: 39.3
LON Start date 31/12/2013 No. of observation Main Standard deviat Varia. Min./M Rat MAI Start date 31/12/2013 No. of observation Main Main Standard deviat	End date 13/06/2020 ons: 2353 ean: 26.5 ion: 9.9 nce: 97.1 Jax.: 5 / 89 nge: 84 DRID End date 13/06/2020 ons: 2323 ean: 24.3 ion: ion: 11.7		LUXEM Start date 19/06/2015 No. of observation Ma Standard deviat Varian Min./M Ran PA Start date 31/12/2013 No. of observation Ma Standard deviat	End date End date 13/06/2020 ons: 1583 can: 19.5 ion: 6.8 nce: 46.2 ax.: 2 / 52 nge: 50 RIS End date 13/06/2020 0ns: 2227 can: 39.3 ion: 14.5
LON Start date 31/12/2013 No. of observation Mainer Standard deviate Varial Min./Mainer MAI Start date 31/12/2013 No. of observation Mainer	End date 13/06/2020 ons: 2353 ean: 26.5 ion: 9.9 nce: 97.1 Vax.: 5 / 89 nge: 84 DRID End date 13/06/2020 0ns: 2323 ean: 24.3 11.7 nce: 138.0 138.0		LUXEM Start date 19/06/2015 No. of observation Ma Standard deviat Varian Min./M Ran PA Start date 31/12/2013 No. of observation Ma Standard deviat Varian	End date 13/06/2020 ons: 1583 can: 19.5 ion: 6.8 nce: 46.2 iax.: 2 / 52 nge: 50 RIS End date 13/06/2020 ons: 2227 can: 39.3 ion: 14.5 nce: 210.5
LON Start date 31/12/2013 No. of observation Main Standard deviate Varian Min./M Rate 31/12/2013 No. of observation Main Standard deviate Varian Min./M	End date 13/06/2020 ons: 2353 ean: 26.5 ion: 9.9 nce: 97.1 Jax.: 5 / 89 nge: 84 DRID End date 13/06/2020 ons: 2323 ean: 24.3 ion: 11.7 nce: 138.0 Jax.: 5 / 160		LUXEM Start date 19/06/2015 No. of observation Ma Standard deviat Varian Min./M Ran PA Start date 31/12/2013 No. of observation Ma Standard deviat Varian Min./M	End date 13/06/2020 ons: 1583 can: 19.5 ion: 6.8 nce: 46.2 iax.: $2/52$ ige: 50 RIS End date 13/06/2020 ons: 2227 can: 39.3 ion: 14.5 nce: 210.5 iax.: $6/122$

Series	No deterministic terms	An intercept	An intercept and a time trend
AMSTERDAM	0.42 (0.39, 0.46)	0.39 (0.35, 0.44)	0.39 (0.35, 0.44)
BERLIN	0.63 (0.58, 0.68)	0.61 (0.56, 0.67)	0.61 (0.56, 0.67)
BRUSSELS	0.52 (0.48, 0.56)	0.50 (0.46, 0.55)	0.50 (0.46, 0.55)
HELSINKI	0.57 (0.52, 0.61)	0.54 (0.50, 0.59)	0.54 (0.50, 0.59)
LONDON	0.54 (0.50, 0.59)	0.52 (0.47, 0.57)	0.52 (0.47, 0.57)
LUXEMBOURG	0.56 (0.51, 0.62)	0.54 (0.48, 0.61)	0.54 (0.48, 0.61)
MADRID	0.63 (0.58, 0.68)	0.62 (0.57, 0.67)	0.62 (0.57, 0.67)
PARIS	0.55 (0.51, 0.59)	0.53 (0.49, 0.58)	0.53 (0.49, 0.58)

Table 2: Estimates of d: White noise errors

We report the estimates of d and its 95% confidence band (in parenthesis). In bold, the selected specification for each series.

Series	d (95% band)	Intercept (t-value)	Time trend
AMSTERDAM	0.39 (0.35, 0.44)	26.0450 (8.77)	
BERLIN	0.61 (0.56, 0.67)	25.3864 (3.93)	
BRUSSELS	0.50 (0.46, 0.55)	20.9146 (4.17)	
HELSINKI	0.54 (0.50, 0.59)	20.4913 (4.57)	
LONDON	0.52 (0.47, 0.57)	23.0696 (5.23)	
LUXEMBOURG	0.54 (0.48, 0.61)	15.6394 (4.61)	
MADRID	0.62 (0.57, 0.67)	16.5249 (2.59)	
PARIS	0.53 (0.49, 0.58)	32.9226 (5.16)	

Table 3: Estimated coefficients in the selected model: White noise errors

The values in parenthesis in column 3 are the corresponding t-values.

Series	No deterministic terms	An intercept	An intercept and a time trend
AMSTERDAM	0.34 (0.29, 0.39)	0.26 (0.21, 0.30)	0.25 (0.20, 0.30)
BERLIN	0.38 (0.33, 0.43)	0.30 (0.25, 0.36)	0.29 (0.23, 0.36)
BRUSSELS	0.33 (0.28, 0.38)	0.26 (0.21, 0.32)	0.25 (0.21, 0.31)
HELSINKI	0.40 (0.35, 0.45)	0.33 (0.29, 0.39)	0.33 (0.29, 0.38)
LONDON	0.34 (0.30, 0.39)	0.26 (0.21, 0.31)	0.27 (0.22, 0.30)
LUXEMBOURG	0.32 (0.28, 0.38)	0.22 (0.16, 0.29)	0.21 (0.15, 0.28)
MADRID	0.34 (0.30, 0.39)	0.30 (0.25, 0.35)	0.30 (0.25, 0.35)
PARIS	0.41 (0.37, 0.46)	0.33 (0.28, 0.38)	0.33 (0.28, 0.38)

Table 4: Estimates of d: Autocorrelated errors

We report the estimates of d and its 95% confidence band (in parenthesis). In bold, the selected specification for each series.

Series	d (95% band)	Intercept (t-value)	Time trend (t-value)
AMSTERDAM	0.26 (0.21, 0.30)	25.6832 (21.18)	
BERLIN	0.29 (0.23, 0.36)	32.6088 (16.38)	-0.0046 (-2.92)
BRUSSELS	0.25 (0.21, 0.31)	26.3621 (15.22)	-0.0022 (1.97)
HELSINKI	0.33 (0.29, 0.39)	19.6888 (12.88)	
LONDON	0.26 (0.21, 0.31)	25.9193 (24.21)	
LUXEMBOURG	0.22 (0.16, 0.29)	19.2309 (34.57)	
MADRID	0.30 (0.25, 0.35)	23.5407 (16.78)	
PARIS	0.33 (0.28, 0.38)	42.2180 (13.79)	-0.0044 (-1.98)

 Table 5: Estimated coefficients in the selected model: Autocorrelated errors

The values in parenthesis in columns 3 and 4 are the corresponding t-values for the intercept and the time trend respectively.