

Schürger, Klaus

Working Paper

Laplace transforms and suprema of stochastic processes

Bonn Econ Discussion Papers, No. 10/2002

Provided in Cooperation with:

Bonn Graduate School of Economics (BGSE), University of Bonn

Suggested Citation: Schürger, Klaus (2002) : Laplace transforms and suprema of stochastic processes, Bonn Econ Discussion Papers, No. 10/2002, University of Bonn, Bonn Graduate School of Economics (BGSE), Bonn

This Version is available at:

<https://hdl.handle.net/10419/22839>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.

BONN ECON DISCUSSION PAPERS

Discussion Paper 10/2002

**Laplace transforms and suprema of stochastic
processes**

by

Klaus Schürger

May 2002



Bonn Graduate School of Economics
Department of Economics
University of Bonn
Adenauerallee 24 - 42
D-53113 Bonn

The Bonn Graduate School of Economics is
sponsored by the

Deutsche Post  World Net
MAIL EXPRESS LOGISTICS FINANCE

LAPLACE TRANSFORMS AND SUPREMA OF STOCHASTIC PROCESSES

KLAUS SCHÜRGER

Department of Statistics, University of Bonn

ABSTRACT. It is shown that moments of negative order as well as positive non-integral order of a nonnegative random variable X can be expressed by the Laplace transform of X . Applying these results to certain first passage times gives explicit formulae for moments of suprema of Bessel processes as well as strictly stable Lévy processes having no positive jumps.

KEY WORDS: Laplace transform, Bessel process, Lévy process.

0. INTRODUCTION

In the sequel (B_t) denotes a d -dimensional standard linear Brownian motion starting at $0 \in \mathbb{R}^d$ (denoted $\text{BM}(d)$). In Shiryaev (1999,p.251) a beautiful trick is used in order to show that if (B_t) is a $\text{BM}(1)$,

$$(0.0.1) \quad \mathbb{E} \left[\sup_{0 \leq s \leq 1} |B_s| \right] = \sqrt{\pi/2}.$$

In fact, the verification of (0.0.1) can be based on the stopping time

$$(0.0.2) \quad T(1) := \inf\{t \geq 0 \mid |B_t| = 1\}$$

and its Laplace transform

$$(0.0.3) \quad \varphi_1(t) = \mathbb{E}[\exp(-t T(1))] = \frac{1}{\cosh(\sqrt{2t})}, \quad t \geq 0.$$

The latter is easily obtained by applying the optional stopping theorem to the martingale

$$\cosh(sB_t) \exp(-s^2 t/2) \quad (t \geq 0) \quad \text{for fixed } s \geq 0;$$

see, e.g., Revuz/Yor (1991,p.68) or Rogers/Williams (1994,p.19). Although there is no explicit inversion of the Laplace transform in (0.0.3) in any particularly useful form, it turns out, however, that (0.0.3) contains enough information in order to yield (0.0.1). In fact, putting

$$(0.0.4) \quad M(t) = \sup_{0 \leq s \leq t} |B_s|, \quad t \geq 0,$$

we get by Brownian scaling, for any $t > 0$,

$$\begin{aligned} P\left(M(1) \leq t\right) &= P\left(\sup_{0 \leq s \leq 1} |B_{s/t^2}| \leq 1\right) = P\left(M(1/t^2) \leq 1\right) \\ &= P\left(T(1) \geq 1/t^2\right) = P\left((T(1))^{-1/2} \leq t\right), \end{aligned}$$

i.e.,

$$(0.0.5) \quad M(1) \text{ and } (T(1))^{-1/2} \text{ have the same distribution}$$

which implies

$$(0.0.6) \quad E[M(1)] = E\left[(T(1))^{-1/2}\right].$$

Next, using the density of a normal distribution with mean 0 and variance $s^2/2$ we get

$$(0.0.7) \quad s = \frac{2}{\sqrt{\pi}} \int_0^\infty \exp(-(t/s)^2) dt, \quad s > 0.$$

Hence if $X \geq 0$ is a random variable having Laplace transform φ_X we obtain from (0.0.7) by using the Fubini-Tonelli theorem,

$$(0.0.8) \quad E[X^{-1/2}] = \frac{2}{\sqrt{\pi}} \int_0^\infty \varphi_X(t^2) dt.$$

Applying (0.0.8) to $X = T(1)$ and taking into account (0.0.6) as well as (0.0.3) we arrive at

$$E[M(1)] = \frac{2}{\sqrt{\pi}} \int_0^\infty \frac{1}{\cosh(\sqrt{2}t)} dt.$$

Using the substitution $u = \exp(\sqrt{2}t)$ we end up with (0.0.1).

In the sequel we first extend (0.0.8) in two different ways (see Theorems 1.1 and 1.2 in the next section). Using the same pattern of proof as before allows us to obtain results similar to (0.0.1) for Bessel processes as well as for a certain class of Lévy processes.

1. CALCULATION OF MOMENTS VIA LAPLACE TRANSFORMS

We first derive an extension of (0.0.8). In order to achieve this it is natural to start with the identity

$$\Gamma(1/\tau) = \int_0^\infty u^{1/\tau-1} \exp(-u) du, \quad \tau > 0.$$

Using the substitution $u = (t/s)^\tau$ ($s > 0, \tau > 0$ being constants) we get

$$(1.0.1) \quad s = \frac{\tau}{\Gamma(1/\tau)} \int_0^\infty \exp\left(- (t/s)^\tau\right) dt, \quad s \geq 0, \quad \tau > 0.$$

1.1. Theorem. *Let $X \geq 0$ be a random variable having Laplace transform φ_X . Then*

$$(1.1.1) \quad E[X^{-r}] = \frac{1}{r \Gamma(r)} \int_0^\infty \varphi_X(t^{1/r}) dt, \quad r > 0.$$

Proof. Apply (1.0.1) to $s = X^{-r}$, $\tau = 1/r$ and use the Fubini-Tonelli theorem. \square

There are interesting connections between (1.1.1) and the "fractional calculus" (see Ross (1974) and Wolfe (1974)). It has been noticed in Wolfe (1974) that it is possible to calculate moments of positive non-integral order of a nonnegative random variable X by using the Laplace transform φ_X of X . In fact, we have the

1.2. Theorem. *Let $X \geq 0$ be a random variable having Laplace transform φ_X . Then,*

$$(1.2.1) \quad E[X^r] = \frac{r}{\Gamma(1-r)} \int_0^\infty \frac{1 - \varphi_X(t)}{t^{r+1}} dt, \quad 0 < r < 1.$$

More generally, we have for any integer $n \geq 0$ and $n < r < n + 1$

$$(1.2.2) \quad E[X^r] = \frac{r - n}{\Gamma(n + 1 - r)} \int_0^\infty \frac{(-1)^n (\varphi_X^{(n)}(0) - \varphi_X^{(n)}(t))}{t^{r+1-n}} dt.$$

Proof. In order to prove (1.2.1) we use the identity

$$(1.2.3) \quad \int_0^\infty \frac{1 - \exp(-st)}{t^{r+1}} dt = \frac{1}{r} \Gamma(1-r) s^r, \quad 0 < r < 1, \quad s \geq 0$$

which can be derived by using partial integration and the definition of the gamma function (see, e.g., Gradshteyn/Ryzhik (1965,p.333)). Applying (1.2.3) to $s = X$ and using the Fubini-Tonelli theorem we obtain (1.2.1). In a similar way we get (1.2.2), noting that

$$\varphi_X^{(n)}(t) = (-1)^n E[X^n \exp(-tX)], \quad t \geq 0, \quad n \geq 0.$$

\square

2. BESSEL PROCESSES

In the sequel (B_t) denotes a BM(d). Then $(|B_t|)$ is a realization of a d -dimensional Bessel process starting at $0 \in \mathbb{R}^d$ ($|B_t|$ denoting the Euclidean norm of B_t). Consider the stopping time

$$(2.0.1) \quad T(s) := \inf\{t \geq 0 \mid |B_t| = s\}, \quad s \geq 0.$$

The Laplace transform φ_s of $T(s)$ is given by

$$(2.0.2) \quad \varphi_s(t) = (s\sqrt{2t})^\lambda [2^\lambda \Gamma(\lambda + 1) I_\lambda(s\sqrt{2t})]^{-1}, \quad s > 0, \quad t > 0$$

(see, e.g., Gettoor/Sharpe (1979) and Gettoor(1979)). Here,

$$(2.0.3) \quad \lambda := (d - 2)/2 \geq -1/2,$$

and I_ν is denoting the modified Bessel function of order ν given by

$$(2.0.4) \quad I_\nu(t) = \sum_{m=0}^{\infty} \frac{(t/2)^{\nu+2m}}{m! \Gamma(\nu + m + 1)}, \quad t > 0, \quad \nu \geq -1/2.$$

In particular,

$$(2.0.5) \quad I_{-1/2}(t) = \sqrt{2/(\pi t)} \cosh(t), \quad t > 0.$$

(Note that, in the one-dimensional case, this yields the Laplace transform

$$(2.0.6) \quad \mathbb{E}[\exp(-t T(s))] = \frac{1}{\cosh(s\sqrt{2t})}, \quad s \geq 0, \quad t \geq 0.)$$

Putting

$$(2.0.7) \quad M(t) = \sup_{0 \leq s \leq t} |B_s|, \quad t \geq 0$$

we obtain by Brownian scaling

$$(2.0.8) \quad M(1) \text{ and } (T(1))^{-1/2} \text{ have the same distribution.}$$

By (1.1.1) this implies for any $r > 0$

$$E[(M(1))^r] = E[(T(1))^{-r/2}] = \frac{2}{r \Gamma(r/2)} \int_0^\infty \varphi_1(t^{2/r}) dt.$$

Using (2.0.2) and substituting $u = \sqrt{2}t^{1/r}$ yields, for any $r > 0$,

$$(2.0.9) \quad E[(M(1))^r] = \frac{4}{2^{(d+r)/2} \Gamma(d/2) \Gamma(r/2)} \int_0^\infty \frac{u^{d/2+r-2}}{I_\lambda(u)} du$$

($\lambda = (d - 2)/2$). In the case $d = 1$ we get from (2.0.9) (using (2.0.5))

$$E[(M(1))^r] = \frac{2}{2^{r/2} \Gamma(r/2)} \int_0^\infty \frac{u^{r-1}}{\cosh(u)} du, \quad r > 0.$$

The substitution $t = \exp(u)$, i.e. $u = \log t$, yields for any $r > 0$

$$(2.0.10) \quad \mathbb{E}[(M(1))^r] = \frac{4}{2^{r/2} \Gamma(r/2)} \int_1^\infty \frac{(\log t)^{r-1}}{t^2 + 1} dt \quad (d = 1)$$

In the special case $r = 2n + 1$ ($n \geq 0$) the integral in (2.0.10) equals

$$(2.0.11) \quad \int_1^\infty \frac{(\log t)^{2n}}{t^2 + 1} dt = \int_0^1 \frac{(\log t)^{2n}}{t^2 + 1} dt = \frac{\pi^{2n+1}}{2^{2n+2}} |E_{2n}|$$

(see Gradshteyn/Ryzhik (1965,p.549)), E_0, E_2, \dots denoting the Euler numbers determined by

$$\frac{1}{\cosh(t)} = \sum_{n=0}^\infty E_{2n} \frac{t^{2n}}{(2n)!}.$$

The first Euler numbers are $E_0 = 1, E_2 = -1, E_4 = 5, E_6 = -61, E_8 = 1385, E_{10} = -50521, E_{12} = 2702765, E_{14} = -199360981$. Combining (2.0.10) and (2.0.11) gives for $n = 0, 1, \dots$

$$(2.0.12) \quad \mathbb{E}[(M(1))^{2n+1}] = \sqrt{\frac{\pi}{2}} \left(\frac{\pi^2}{2}\right)^n \frac{n!}{(2n)!} |E_{2n}| \quad (d = 1)$$

In order to investigate the asymptotic behaviour of $\mathbb{E}[(M(1))^r]$ (as $r \rightarrow \infty$) we need

2.1. Lemma. *As $t \rightarrow \infty$*

$$(2.1.1) \quad \Gamma(t) \sim (t/e)^t \sqrt{2\pi/t}$$

and

$$(2.1.2) \quad I_\nu(t) \sim (2\pi t)^{-1/2} \exp(t)$$

independently of $\nu \geq -1/2$.

Proof. For a proof of (2.1.1) see Bender/Orszag (1978,p.275); a proof of (2.1.2) can be found in Courant/Hilbert (1966,p.526) (see also Bender/Orszag (1978,p.271)). \square

2.2. Proposition. *As $r \rightarrow \infty$*

$$(2.2.1) \quad \mathbb{E}[(M(1))^r] \sim \frac{4\sqrt{\pi}}{2^{d/2}} \frac{r^{(d+r-1)/2}}{\Gamma(d/2) e^{r/2}}.$$

In particular

$$(2.2.2) \quad \mathbb{E}[(M(1))^r] \sim 2\sqrt{2}(r/e)^{r/2} \quad (d = 1)$$

Proof. For the integral in (2.0.9) we get

$$\int_0^\infty \frac{u^{d/2+r-2}}{I_\lambda(u)} du \sim \frac{2\pi}{e^r} r^{d/2+r-1} \quad (r \rightarrow \infty)$$

by using some routine estimates and applying (2.1.2). Combining this with (2.1.1) gives (2.2.1). \square

2.3. Corollary. *For any $d \geq 1$ the distribution of $M(1)$ is uniquely determined by its moments*

$$\mu_n = E[(M(1))^n], \quad n = 0, 1, \dots$$

Proof. It suffices to show that

$$(2.3.1) \quad \sum_{n=0}^{\infty} \mu_{2n}^{-1/(2n)} = \infty$$

(see Feller (1966,p.224)). But, by (2.2.1),

$$\mu_{2n}^{1/(2n)} \sim \sqrt{2n/e} \quad (n \rightarrow \infty)$$

which implies (2.3.1). \square

2.4. Remark. Comparing (2.0.12) and (2.2.2) we obtain for the asymptotic behaviour of the Euler numbers E_{2n}

$$(2.4.1) \quad |E_{2n}| \sim \frac{8\sqrt{n}}{\sqrt{\pi}} \left(\frac{4n}{e\pi}\right)^{2n}.$$

3. STRICTLY STABLE LÉVY PROCESSES

Let $X = (X_t)$ be a one-dimensional Lévy process starting at 0 such that X_1 is not a.s. constant. Furthermore X is assumed to be a.s. càdlàg, i.e. almost all paths of X are right continuous and have finite left-hand limits at every point. In the sequel α denotes a real number such that

$$(3.0.1) \quad 1 < \alpha < 2.$$

The distribution μ of X_1 is assumed to be *strictly α -stable* with parameters (α, β, c) for constants $-1 \leq \beta \leq 1$ and $c > 0$, i.e. the characteristic function $\hat{\mu}$ of μ is of the form

$$(3.0.2) \quad \hat{\mu}(t) = \exp(-\Psi(t)), \quad t \in \mathbb{R}.$$

Here, Ψ is the *characteristic exponent* given by

$$(3.0.3) \quad \Psi(t) = c|t|^\alpha \left[1 - i\beta(\operatorname{sgn} t) \tan\left(\frac{\pi\alpha}{2}\right) \right], \quad t \in \mathbb{R}$$

where $\operatorname{sgn} t$ is equal to 1, 0, -1 when t is > 0 , $= 0$, < 0 , respectively (see Bertoin (1996,p.217) or Sato (1999,p.86)). It follows from the above assumptions that, for any $a > 0$,

$$(3.0.4) \quad (X_{at}) \text{ has the same law as } (a^{1/\alpha} X_t).$$

From now on we will additionally assume

$$(3.0.5) \quad \beta = -1.$$

This implies (see Bertoin (1996,p.217) or Sato (1999,p.346)) that the Lévy measure of X has support in $] -\infty, 0]$, and X has *no positive jumps*, i.e.

$$(3.0.6) \quad P\left(X_t \leq X_{t-} \text{ for all } t > 0\right) = 1.$$

Furthermore (see Sato (1999,p.350))

$$(3.0.7) \quad P\left(\limsup_{t \rightarrow \infty} X_t = \infty\right) = 1.$$

Let $R(s)$ denote the first passage time defined by

$$(3.0.8) \quad R(s) = \inf\{t > 0 | X_t > s\}, \quad s \geq 0.$$

Note that, by (3.0.7), $R(s) < \infty$ a.s., $s \geq 0$. For the following result see Sato (1999,pp.346,347).

3.1. Theorem. *We continue to assume (3.0.1) and (3.0.5). Then the following results hold.*

$$(a) \quad P\left(X_{R(s)} = s \text{ for all } s \geq 0\right) = 1.$$

(b) *The function*

$$(3.1.1) \quad \psi(t) := -\Psi(-it) = -\frac{c}{\cos(\frac{\pi\alpha}{2})} t^\alpha, \quad t \geq 0$$

(Ψ given by (3.0.3)) *is strictly increasing, continuous and satisfies $\psi(0) = 0$ and $\psi(t) \rightarrow \infty$ ($t \rightarrow \infty$).*

(c) *The Laplace transforms of the first passage times $R(s)$ ($s > 0$) are given by*

$$(3.1.2) \quad E[\exp(-s R(s))] = \exp(-s \psi^{-1}(t)), \quad s > 0, \quad t \geq 0$$

(ψ^{-1} denoting the inverse function of ψ).

It follows from (3.1.1) that ψ^{-1} is given by

$$(3.1.3) \quad \psi^{-1}(t) = \left(\frac{1}{c} \left| \cos\left(\frac{\pi\alpha}{2}\right) \right|\right)^{1/\alpha} t^{1/\alpha}, \quad t \geq 0.$$

Combining Theorems 3.1 and 1.1 gives

3.2. Proposition. *We continue to assume (3.0.1) and (3.0.5). Then the following results hold.*

(a) *For any $r > 0$,*

$$(3.2.1) \quad E\left[\left(\sup_{0 \leq s \leq 1} X_s\right)^r\right] = \frac{\alpha \Gamma(r)}{\Gamma(r/\alpha)} \left(\frac{c}{\left|\cos\left(\frac{\pi\alpha}{2}\right)\right|}\right)^{r/\alpha}.$$

(b) *We have, as $r \rightarrow \infty$,*

$$(3.2.2) \quad E\left[\left(\sup_{0 \leq s \leq 1} X_s\right)^r\right] \sim \sqrt{\alpha} \left(\frac{\alpha c r^{\alpha-1}}{\left|\cos\left(\frac{\pi\alpha}{2}\right)\right| e^{\alpha-1}}\right)^{r/\alpha}.$$

Proof. In order to prove (3.2.1) first note that it follows from (3.0.6) that

$$\widetilde{M}(t) := \sup_{0 \leq s \leq t} X_s \leq 1 \iff R(1) \geq t \quad \text{a.s.,} \quad t > 0,$$

which, by (3.0.4), implies that $\widetilde{M}(1)$ and $(R(1))^{-1/\alpha}$ have the same distribution. Hence, by Theorem 1.1 and (3.1.2),

$$\begin{aligned} \mathbb{E}[(\widetilde{M}(1))^r] &= \mathbb{E}[(R(1))^{-r/\alpha}] \\ &= \frac{\alpha}{r \Gamma(r/\alpha)} \int_0^\infty \exp\left(-\psi^{-1}(t^{\alpha/r})\right) dt, \quad r > 0. \end{aligned}$$

Taking into account (3.1.3) and putting

$$(3.2.3) \quad \widetilde{c} := \left(\frac{1}{c} \left| \cos\left(\frac{\pi\alpha}{2}\right) \right| \right)^{1/\alpha},$$

we get, using the substitution $u = \widetilde{c} t^{1/r}$,

$$\mathbb{E}[(\widetilde{M}(1))^r] = \frac{\alpha \Gamma(r)}{\Gamma(r/\alpha) \widetilde{c}^r},$$

i.e. (3.2.1). The assertion (3.2.2) is immediate from (3.2.1) and (2.1.1). \square

As a final result we mention

3.3. Proposition. *We continue to assume (3.0.1) and (3.0.5). Then, for any $s > 0$,*

$$(3.3.1) \quad \mathbb{E}[(R(s))^r] = \frac{\Gamma(1-\alpha r)}{\Gamma(1-r)} \left(\frac{|\cos(\frac{\pi\alpha}{2})|}{c} \right)^r s^{\alpha r}, \quad 0 \leq r < 1/\alpha$$

and

$$(3.3.2) \quad \mathbb{E}[(R(s))^r] = \infty, \quad r \geq 1/\alpha.$$

Proof. Put $\delta := s \widetilde{c}$ ($s > 0$; \widetilde{c} given by (3.2.3)). Taking into account (3.1.2), (1.2.1) and (3.1.3) we obtain, for any $0 < r < 1$,

$$\mathbb{E}[(R(s))^r] = \frac{r}{\Gamma(1-r)} \int_0^\infty \frac{1 - \exp(-\delta t^{1/\alpha})}{t^{r+1}} dt.$$

Using the substitution $u = \delta t^{1/\alpha}$ we arrive at

$$\mathbb{E}[(R(s))^r] = \frac{\alpha r \delta^{\alpha r}}{\Gamma(1-r)} \int_0^\infty \frac{1 - \exp(-u)}{u^{\alpha r+1}} du, \quad 0 < r < 1.$$

This clearly entails (3.3.2) and (using partial integration) also (3.3.1). \square

Acknowledgement

I wish to thank A.N. Shiryaev for some stimulating discussions concerning the approach mentioned in the introductory section.

References

- BENDER, C.M., ORSZAG, S.A. (1978) *Advanced Mathematical Methods for Scientists and Engineers*. McGraw-Hill. New York/Toronto.
- BERTOIN, J. (1996) *Lévy Processes*. Cambridge University Press. Cambridge.
- COURANT, R., HILBERT, D. (1966) *Methods of Mathematical Physics*, Vol. I. Interscience Publishers. New York.
- FELLER, W. (1966) *An Introduction to Probability Theory and Its Applications*, Vol. II. John Wiley. New York.
- GETOOR, R.K. (1979) The Brownian escape process. *Annals of Probability* 7, 864-867.
- GETOOR, R.K., SHARPE, M.J. (1979) Excursions of Brownian motion and Bessel processes. *Z. Wahrscheinlichkeitstheorie und verw. Gebiete* 47, 83-106.
- GRADSHTEYN, I.S., RYZHIK, I.M. (1965) *Tables of Integrals, Series and Products*. Academic Press. New York/London.
- REVUZ, D., YOR, M. (1991) *Continuous Martingales and Brownian Motion*. Springer-Verlag. Berlin.
- ROGERS, L.C.G., WILLIAMS, D. (1994) *Diffusions, Markov Processes and Martingales*. Vol. I: *Foundations*. 2nd Ed. John Wiley. New York.
- ROSS, B. (1975) A brief history and exposition of the fundamental theory of fractional calculus. In: *Fractional Calculus and Its Applications* (Ed.: B. Ross). *Lecture Notes in Math.* 457, 1-36.
- SATO, K. (1999) *Lévy Processes and Infinitely Divisible Distributions*. Cambridge University Press. Cambridge.

SHIRYAEV, A.N. (1999) *Essentials of Stochastic Finance*. World Scientific. Singapore

WOLFE, S.J. (1975) On moments of probability distribution functions. In: *Fractional Calculus and Its Applications* (Ed.: B. Ross). Lecture Notes in Math. 457, 306-316.