

Yan, Ji Gao

**Working Paper**

## Complete Convergence and Complete Moment Convergence for Maximal Weighted Sums of Extended Negatively Dependent Random Variables

IRTG 1792 Discussion Paper, No. 2018-040

**Provided in Cooperation with:**

Humboldt University Berlin, International Research Training Group 1792 "High Dimensional Nonstationary Time Series"

*Suggested Citation:* Yan, Ji Gao (2018) : Complete Convergence and Complete Moment Convergence for Maximal Weighted Sums of Extended Negatively Dependent Random Variables, IRTG 1792 Discussion Paper, No. 2018-040, Humboldt-Universität zu Berlin, International Research Training Group 1792 "High Dimensional Nonstationary Time Series", Berlin

This Version is available at:

<https://hdl.handle.net/10419/230751>

**Standard-Nutzungsbedingungen:**

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

**Terms of use:**

*Documents in EconStor may be saved and copied for your personal and scholarly purposes.*

*You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.*

*If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.*

IRTG 1792 Discussion Paper 2018-040



# Complete Convergence and Complete Moment Convergence for Maximal Weighted Sums of Extended Negatively Dependent Random Variables

Ji Gao YAN \*



\* Soochow University, PR China

This research was supported by the Deutsche Forschungsgemeinschaft through the International Research Training Group 1792 "High Dimensional Nonstationary Time Series".

<http://irtg1792.hu-berlin.de>  
ISSN 2568-5619

International Research Training Group 1792

## Complete Convergence and Complete Moment Convergence for Maximal Weighted Sums of Extended Negatively Dependent Random Variables

Ji Gao YAN

School of Mathematical Sciences, Soochow University, Suzhou 215006, P. R. China  
and

C.A.S.E.-Center for Applied Statistics and Economics, Humboldt-Universität zu Berlin,  
Spandauer Str. 1, 10178 Berlin, Germany

E-mail: [yanjigao@suda.edu.cn](mailto:yanjigao@suda.edu.cn)

**Abstract** In this paper, the complete convergence and complete moment convergence for maximal weighted sums of extended negatively dependent random variables are investigated. Some sufficient conditions for the convergence are provided. In addition, the Marcinkiewicz–Zygmund type strong law of large numbers for weighted sums of extended negatively dependent random variables is obtained. The results obtained in the article extend the corresponding ones for independent random variables and some dependent random variables.

**Keywords** Extended negatively dependent, complete convergence, complete moment convergence, maximal weighted sums, strong law of large numbers

**MR(2010) Subject Classification** 60F15

### 1 Introduction

In this paper, we consider a sequence  $\{X_n, n \geq 1\}$  of random variables defined on some probability space  $(\Omega, \mathcal{F}, P)$ . It is well known that the complete convergence plays an important role in establishing almost sure convergence of summation of random variables as well as weighted sums of random variables. The concept of the complete convergence was introduced by Hus and Robbins [11] as follows.

**Definition 1.1** A sequence of random variables  $\{U_n, n \geq 1\}$  is said to converge completely to a constant  $\theta$  if for any  $\varepsilon > 0$ ,

$$\sum_{n=1}^{\infty} P(|U_n - \theta| > \varepsilon) < \infty.$$

Chow [7] presented the following more general concept of the complete moment convergence.

**Definition 1.2** Let  $\{Z_n, n \geq 1\}$  be a sequence of random variables and  $a_n > 0, b_n > 0$ , and  $q > 0$ , if

$$\sum_{n=1}^{\infty} a_n E\{b_n^{-1}|Z_n| - \varepsilon\}_+^q < \infty \quad \text{for some or all } \varepsilon > 0,$$

then the above result was called the complete moment convergence.

As we know, the complete moment convergence implies complete convergence. Moreover, the complete moment convergence can more exactly describe the convergence rate of a sequence of random variables than the complete convergence. So, a study on complete moment convergence is of interest.

When  $\{X_n, n \geq 1\}$  is a sequence of independent and identically distributed random variables, Baum and Katz [3] proved the following remarkable result concerning the convergence rate of the tail probabilities  $P(|S_n| > \varepsilon n^r)$  for all  $\varepsilon > 0$ , where  $S_n = \sum_{i=1}^n X_i$  are the partial sums.

**Theorem 1.3** *Let  $r > 1/2$  and  $p > 1$ . Let  $\{X, X_n, n \geq 1\}$  be a sequence of independent and identical distributed random variables. Assume further that  $EX = 0$  if  $r \leq 1$ . Then the following statements are equivalent:*

- (i)  $E|X|^{p/r} < \infty$ ;
- (ii)  $\sum_{n=1}^{\infty} n^{p-2} P(|S_n| > \varepsilon n^r) < \infty$ , for all  $\varepsilon > 0$ .

In convenient, we call  $n^{p-2}$  as the weight function of the tail probabilities, and  $n^r$  as the boundary function in the tail probabilities. Many useful linear statistics are weighted sums of independent and identically distributed random variables, such as, least-squares estimators, nonparametric regression function estimators and jackknife estimates, and so on. However, in many stochastic model, the assumption that random variables are independent is not plausible. Increases in some random variables are often related to decreases in other random variables, so an assumption of dependence is more appropriate than an assumption of independence. One of the important dependence structure is the extended negatively dependent structure, which was introduced by Liu [12] as follows.

**Definition 1.4** *Random variables  $X_k, k = 1, \dots, n$  are said to be lower extended negatively dependent (LEND) if there is some  $M > 0$  such that, for all real numbers  $x_k, k = 1, \dots, n$ ,*

$$P\left\{\bigcap_{k=1}^n (X_k \leq x_k)\right\} \leq M \prod_{k=1}^n P\{X_k \leq x_k\}; \quad (1.1)$$

*they are said to be upper extended negatively dependent (UEND) if there is some  $M > 0$  such that, for all real numbers  $x_k, k = 1, \dots, n$ ,*

$$P\left\{\bigcap_{k=1}^n (X_k > x_k)\right\} \leq M \prod_{k=1}^n P\{X_k > x_k\}; \quad (1.2)$$

*and they are said to be extended negatively dependent (END) if they are both LEND and UEND. A sequence of infinitely many random variables  $\{X_k, k = 1, 2, \dots\}$  is said to be LEND/UEND/END if there is some  $M > 0$  such that for each positive integer  $n$ , the random variables  $X_1, X_2, \dots, X_n$  are LEND/UEND/END, respectively.*

The END structure covers all negative dependence structures and, more interestingly, it covers certain positive dependence structures. Some applications for END sequence have been found. See, for example, Liu [12] obtained the precise large deviations for dependent random variables with heavy tails. Liu [13] studied the sufficient and necessary conditions of moderate deviations for dependent random variables with heavy tails. Shen [18] gave the probability

inequalities. Chen [5], Yan [30] considered the SLLN for END random variables and applications to risk theory and renewal theory, and so on.

Recall that the sequence  $\{X_n, n \geq 1\}$  is stochastically dominated by a random variable  $X$  if

$$\sup_{n \geq 1} P\{|X_n| > t\} \leq CP\{|X| > t\}, \quad (1.3)$$

for some positive constant  $C$  and all  $t \geq 0$ . A real valued function  $l(x)$ , positive and measurable on  $[A, \infty)$  for some  $A > 0$ , is said to be slowly varying if

$$\lim_{x \rightarrow \infty} \frac{l(\lambda x)}{l(x)} = 1 \quad \text{for each } \lambda > 0.$$

Recently, Shen et al. [20] discussed the complete convergence of weighted sums of extended negatively dependent random variables and obtained the following result:

**Theorem 1.5** *Let  $1/2 < r \leq 1$  and  $\{X_n, n \geq 1\}$  be a sequence of mean zero END random variables, which is stochastically dominated by a random variable  $X$ . Let  $l(x) > 0$  be a slowly varying and monotone non decreasing function. Assume further that  $\{a_{ni}, 1 \leq i \leq n, n \geq 1\}$  is an array of real numbers such that*

$$\sum_{i=1}^n a_{ni}^2 = O(n).$$

If

$$E[|X|^{1/r} l(|X|^{1/r})] < \infty,$$

then for any  $\varepsilon > 0$ ,

$$\sum_{n=1}^{\infty} \frac{l(n)}{n} P\left(\left|\sum_{i=1}^n a_{ni} X_i\right| > \varepsilon n^r\right) < \infty.$$

The above result extended Theorem 1.3 in three aspects: independent and identically distributed random variables to END random variables without identical distribution; partial sums to weighted sums and changed the weight function of tail probability.

Motivated by the above works, we will further study the complete convergence for weighted sums of END random variables, the purpose of this article is threefold:

- (1) generalise the condition of weight  $\{a_{ni}, 1 \leq i \leq n, n \geq 1\}$  to  $\sum_{i=1}^n |a_{ni}|^\alpha = O(n)$  for some  $\alpha > 0$ ;
- (2) change the boundary function in the tail probabilities;
- (3) take into account the complete moment convergence for the maximal of weighted sums as well.

This work is organized as follows: some lemmas and main results on complete convergence and complete moment convergence for maximal weighted sums of END random variables are provided in Section 2. The proofs for the main results are presented in Section 3.

Before we present our main results, we note that  $C$  will be numerical constants whose value are without importance, and, in addition, may change between appearances.  $a_n = O(b_n)$  stands for  $a_n \leq Cb_n$  for all  $n \geq 1$  and  $I(A)$  is the indicator function on the set  $A$ . Denote  $X^+ = \max(X, 0)$  and  $X^- = \max(-X, 0)$ .

## 2 Some Lemmas and Main Results

To proceed the main results, we need some important lemmas.

**Lemma 2.1** ([12]) *Let  $\{X_n, n \geq 1\}$  be a sequence of END random variables.*

(i) *For each  $n \geq 1$ , if  $f_1, f_2, \dots, f_n$  are all nondecreasing (or nonincreasing) functions, then  $f_1(X_1), f_2(X_2), \dots, f_n(X_n)$  are also END random variables.*

(ii) *For each  $n \geq 1$ , there exists a constant  $M > 0$  such that*

$$E\left(\prod_{i=1}^n X_i^+\right) \leq M \prod_{i=1}^n EX_i^+.$$

**Lemma 2.2** ([24]) *Let  $p \geq 2$  and  $\{X_n, n \geq 1\}$  be a sequence of END random variables with  $EX_n = 0$  and  $E|X_n|^p < \infty$  for each  $n \geq 1$ . Then there exists a positive constant  $C_p$  depending only on  $p$  such that*

$$E\left|\sum_{i=1}^n X_i\right|^p \leq C_p \left[ \sum_{i=1}^n E|X_i|^p + \left(\sum_{i=1}^n E|X_i|^2\right)^{p/2} \right], \quad \text{for all } n \geq 1$$

and

$$E\left(\max_{1 \leq j \leq n} \left|\sum_{i=1}^j X_i\right|^p\right) \leq C_p (\log n)^p \left[ \sum_{i=1}^n E|X_i|^p + \left(\sum_{i=1}^n E|X_i|^2\right)^{p/2} \right], \quad \text{for all } n \geq 1.$$

**Lemma 2.3** ([1, 2]) *Let  $\{X_n, n \geq 1\}$  be a sequence of random variables, which is stochastically dominated by a random variable  $X$ . For any  $\alpha > 0$  and  $\beta > 0$ , the following two statements hold:*

$$\begin{aligned} E|X_n|^\alpha I(|X_n| \leq \beta) &\leq C_1 [E|X|^\alpha I(|X| \leq \beta) + \beta^\alpha P\{|X| > \beta\}]; \\ E|X_n|^\alpha I(|X_n| > \beta) &\leq C_2 E|X|^\alpha I(|X| > \beta), \end{aligned}$$

where  $C_1$  and  $C_2$  are positive constants. Consequently,  $E|X_n|^\alpha \leq CE|X|^\alpha$ .

**Lemma 2.4** *Let  $\{a_{ni}, 1 \leq i \leq n, n \geq 1\}$  be an array of real numbers such that*

$$\sum_{i=1}^n |a_{ni}|^\xi \leq n \quad \text{for some } \xi > 0. \quad (2.1)$$

Then

(i) *for every  $0 < \zeta < \xi$ ,*

$$\sum_{i=1}^n |a_{ni}|^\zeta \leq n; \quad (2.2)$$

(ii) *for all  $m \geq 1, \kappa > \xi$ ,*

$$\sum_{j=m}^{\infty} \#(I_{nj})(j+1)^{-\kappa/\xi} \leq C(m+1)^{1-\frac{\kappa}{\xi}}, \quad (2.3)$$

where  $I_{nj} = \{1 \leq i \leq n : n^{1/\xi}(j+1)^{-1/\xi} < |a_{ni}| \leq n^{1/\xi}j^{-1/\xi}\}$ , for each  $n \geq 1, j \geq 1$ . Obviously, it holds that for all  $n > m \geq 1, \kappa > \xi$ ,

$$\sum_{j=m}^{n-1} \#(I_{nj})(j+1)^{-\kappa/\xi} \leq C(m+1)^{1-\frac{\kappa}{\xi}}; \quad (2.4)$$

(iii) for arbitrary  $s \geq 1$ ,

$$\sum_{j=1}^s \sharp(I_{nj}) \leq s + 1. \quad (2.5)$$

We will use the following properties of slowly varying functions.

**Lemma 2.5** ([17]) *If  $\ell(x)$  is a function slowly varying at infinity, then for any  $r > 0$ ,*

$$C_1 n^{-r} \ell(n) \leq \sum_{i=n}^{\infty} i^{-1-r} \ell(i) \leq C_2 n^{-r} \ell(n),$$

and

$$C_3 n^r \ell(n) \leq \sum_{i=1}^n i^{-1+r} \ell(i) \leq C_4 n^r \ell(n),$$

where  $C_i, i = 1, 2, 3, 4$  are positive constants depending only on  $r$ .

**Lemma 2.6** *Let  $\{a_{ni}, 1 \leq i \leq n, n \geq 1\}$  be an array of real numbers satisfying*

$$\sum_{i=1}^n |a_{ni}|^\alpha = O(n) \quad \text{for some } \alpha > 0 \quad (2.6)$$

and  $X$  be a random variable. Let  $b_n = n^{1/\alpha} (\log n)^{3/\gamma}$  for some  $\gamma > 0$ .

(i) *If  $p > \max\{\alpha, \gamma(\beta + 1)/3\}$  for some  $\beta$ , then*

$$\begin{aligned} & \sum_{n=2}^{\infty} \frac{(\log n)^\beta}{n b_n^p} \sum_{i=1}^n E |a_{ni} X|^p I(|a_{ni} X| \leq b_n) \\ & \leq \begin{cases} CE|X|^\alpha, & \text{for } \alpha > \gamma(\beta + 1)/3, \\ CE|X|^\alpha \log(1 + |X|), & \text{for } \alpha = \gamma(\beta + 1)/3, \\ CE|X|^{\gamma(\beta+1)}, & \text{for } \alpha < \gamma(\beta + 1)/3. \end{cases} \end{aligned}$$

(ii) *If  $p = \alpha, \beta = 2$ , then*

$$\begin{aligned} & \sum_{n=2}^{\infty} \frac{(\log n)^2}{n b_n^\alpha} \sum_{i=1}^n E |a_{ni} X|^\alpha I(|a_{ni} X| > b_n) \\ & \leq \begin{cases} CE|X|^\alpha, & \text{for } \alpha > \gamma, \\ CE|X|^\alpha \log(1 + |X|), & \text{for } \alpha = \gamma, \\ CE|X|^\gamma, & \text{for } \alpha < \gamma. \end{cases} \end{aligned}$$

**Lemma 2.7** *Let  $0 < \alpha < 2$  and  $\gamma > 0$ . Suppose that the sequence  $\{X_{ni}, i \geq 1, n \geq 1\}$  is stochastically dominated by a random variable  $X$  with  $E|X|^\alpha < \infty$ . And  $\{a_{ni}, 1 \leq i \leq n, n \geq 1\}$  be an array of real numbers satisfying (2.6). Additionally, assume that  $EX_{ni} = 0$  for  $1 < \alpha < 2$  and  $b_n$  as in Lemma 2.6. Then*

$$\frac{1}{b_n} \max_{1 \leq j \leq n} \left| \sum_{i=1}^j E Y_{ni} \right| \leq C (\log n)^{-3\alpha/\gamma} E|X|^\alpha \rightarrow 0 \quad \text{as } n \rightarrow \infty, \quad (2.7)$$

where  $Y_{ni} = a_{ni} X_{ni} I(|a_{ni} X_{ni}| \leq b_n) + b_n I(a_{ni} X_{ni} > b_n) - b_n I(a_{ni} X_{ni} < -b_n)$  for each  $1 \leq i \leq n, n \geq 1$ .

Now, we present our main results as follows.

**Theorem 2.8** Let  $\{X_{ni}, i \geq 1, n \geq 1\}$  be an array of rowwise END random variables, which was stochastically dominated by a random variable  $X$ . For some  $0 < \alpha < 2, \gamma > 0, \{a_{ni}, 1 \leq i \leq n, n \geq 1\}$  be an array of real numbers satisfying (2.6) and  $b_n$  as in Lemma 2.6. Additionally, assume that  $EX_{ni} = 0$  for  $1 < \alpha < 2$ . If

$$\begin{cases} E|X|^\alpha < \infty, & \text{for } \alpha > \gamma, \\ E|X|^\alpha \log(1 + |X|) < \infty, & \text{for } \alpha = \gamma, \\ E|X|^\gamma < \infty, & \text{for } \alpha < \gamma, \end{cases} \quad (2.8)$$

then

$$\sum_{n=2}^{\infty} \frac{1}{n} P\left(\max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} X_{ni} \right| > \varepsilon b_n\right) < \infty \quad \text{for all } \varepsilon > 0. \quad (2.9)$$

**Theorem 2.9** Let  $\{X_{ni}, i \geq 1, n \geq 1\}$  be an array of rowwise END random variables, which was stochastically dominated by a random variable  $X$ . For some  $0 < \alpha < 2, \gamma > 0, \{a_{ni}, 1 \leq i \leq n, n \geq 1\}$  be an array of real numbers satisfying (2.6) and  $b_n$  as in Lemma 2.6. Additionally, assume that  $EX_{ni} = 0$  for  $1 < \alpha < 2$ . If (2.8) holds, then for  $0 < \tau < \alpha$ ,

$$\sum_{n=2}^{\infty} \frac{1}{n} E\left(\frac{1}{b_n} \max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} X_{ni} \right| - \varepsilon\right)_+^\tau < \infty \quad \text{for all } \varepsilon > 0. \quad (2.10)$$

**Remark 2.10** Under the conditions of Theorem 2.9, we can obtain that

$$\begin{aligned} \infty &> \sum_{n=2}^{\infty} \frac{1}{n} E\left(\frac{1}{b_n} \max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} X_{ni} \right| - \varepsilon\right)_+^\tau \\ &= \sum_{n=2}^{\infty} \frac{1}{n} \int_0^\infty P\left(\frac{1}{b_n} \max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} X_{ni} \right| - \varepsilon > t^{1/\tau}\right) dt \\ &\geq \int_0^\varepsilon \sum_{n=2}^{\infty} \frac{1}{n} P\left(\max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} X_{ni} \right| > b_n(\varepsilon + \varepsilon^{1/\tau})\right) dt \\ &\geq \varepsilon \sum_{n=2}^{\infty} \frac{1}{n} P\left(\max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} X_{ni} \right| > b_n(\varepsilon + \varepsilon^{1/\tau})\right) \quad \text{for all } \varepsilon > 0. \end{aligned} \quad (2.11)$$

Hence, from (2.11) we get that the complete moment convergence implies the complete convergence.

**Theorem 2.11** Conditions as those of Theorem 2.8. Let  $T_n = \sum_{i=1}^n a_{ni} X_{ni}$  for each  $n \geq 1$ , then

$$\lim_{n \rightarrow \infty} \frac{T_n}{b_n} = 0, \quad \text{a.s.} \quad (2.12)$$

### 3 Proofs of Main Results

*Proof of Lemma 2.4* (i) By Jensen's inequality and (2.1), we get

$$\sum_{i=1}^n |a_{ni}|^\zeta \leq \left(\sum_{i=1}^n |a_{ni}|^\xi\right)^{\frac{\zeta}{\xi}} \cdot n^{1-\frac{\zeta}{\xi}} \leq n.$$



(ii) By the denotes of  $I_{nj}$ , we see that for every  $n \geq 1$ ,  $\{I_{nj}, j \geq 1\}$  are pairwise disjoint and

$$\bigcup_{j \geq 1} I_{nj} = \{1 \leq i \leq n : 0 < |a_{ni}| \leq n^{1/\xi}\}$$

since  $\sum_{i=1}^n |a_{ni}|^\xi \leq n < \infty$ .

For all  $m \geq 1$ , by  $\kappa > \xi$  we have

$$\begin{aligned} n &\geq \sum_{i=1}^n |a_{ni}|^\xi = \sum_{j=1}^{\infty} \sum_{i \in I_{nj}} |a_{ni}|^\xi \\ &\geq \sum_{j=1}^{\infty} \#(I_{nj}) n(j+1)^{-1} \geq \sum_{j=m}^{\infty} \#(I_{nj}) n(j+1)^{-1} \\ &= \sum_{j=m}^{\infty} \#(I_{nj}) n(j+1)^{-\kappa/\xi} (j+1)^{\kappa/\xi-1} \\ &\geq \sum_{j=m}^{\infty} \#(I_{nj}) n(j+1)^{-\kappa/\xi} (m+1)^{\kappa/\xi-1}. \end{aligned}$$

Hence, (2.3) holds.

(iii) On the other hand, by (2.1), for arbitrary  $s \geq 1$ ,

$$n \geq \sum_{k=1}^n |a_{nk}|^\xi = \sum_{j=1}^{\infty} \sum_{k \in I_{nj}} |a_{nk}|^\xi \geq n \sum_{j=1}^s \frac{1}{j+1} \#(I_{nj}) \geq \frac{n}{s+1} \sum_{j=1}^s \#(I_{nj}),$$

which implies (2.5).

*Proof of Lemma 2.6* Without loss of generality, we may assume that

$$\sum_{i=1}^n |a_{ni}|^\alpha \leq n, \quad \text{for some } \alpha > 0.$$

Then, by Lemma 2.4,

$$\begin{aligned} &\sum_{n=2}^{\infty} \frac{(\log n)^\beta}{nb_n^p} \sum_{i=1}^n E|a_{ni}X|^p I(|a_{ni}X| \leq b_n) \\ &= \sum_{n=2}^{\infty} n^{-1-\frac{p}{\alpha}} (\log n)^{\beta-\frac{3p}{\gamma}} \sum_{i=1}^n |a_{ni}|^p E|X|^p I(|a_{ni}X| \leq n^{1/\alpha} (\log n)^{3/\gamma}) \\ &= \sum_{n=2}^{\infty} n^{-1-\frac{p}{\alpha}} (\log n)^{\beta-\frac{3p}{\gamma}} \sum_{j=1}^{\infty} \sum_{i \in I_{nj}} |a_{ni}|^p E|X|^p I(|a_{ni}X| \leq n^{1/\alpha} (\log n)^{3/\gamma}) \\ &\leq \sum_{n=2}^{\infty} n^{-1-\frac{p}{\alpha}} (\log n)^{\beta-\frac{3p}{\gamma}} \sum_{j=1}^{\infty} \#(I_{nj}) n^{p/\alpha} j^{-p/\alpha} E|X|^p I(|X| \leq (j+1)^{1/\alpha} (\log n)^{3/\gamma}) \\ &= \sum_{n=2}^{\infty} n^{-1} (\log n)^{\beta-\frac{3p}{\gamma}} \sum_{j=1}^{\infty} \#(I_{nj}) j^{-p/\alpha} E|X|^p I(|X| \leq (\log n)^{3/\gamma}) \\ &\quad + \sum_{n=2}^{\infty} n^{-1} (\log n)^{\beta-\frac{3p}{\gamma}} \sum_{j=1}^{\infty} \#(I_{nj}) j^{-p/\alpha} \\ &\quad \times \sum_{k=1}^j E|X|^p I(k^{1/\alpha} (\log n)^{3/\gamma} < |X| \leq (k+1)^{1/\alpha} (\log n)^{3/\gamma}) \end{aligned}$$

$$\triangleq I_1 + I_2.$$

First for  $I_1$ , if  $\alpha > \frac{\gamma(\beta+1)}{3}$ , by Lemma 2.4 and  $p > \alpha$ , we have

$$\begin{aligned} I_1 &\leq C \sum_{n=2}^{\infty} n^{-1} (\log n)^{\beta - \frac{3p}{\gamma}} E|X|^p I(|X| \leq (\log n)^{3/\gamma}) \\ &\leq C \sum_{n=2}^{\infty} n^{-1} (\log n)^{\beta - \frac{3p}{\gamma}} (\log n)^{3(p-\alpha)/\gamma} E|X|^\alpha I(|X| \leq (\log n)^{3/\gamma}) \\ &\leq CE|X|^\alpha \sum_{n=2}^{\infty} n^{-1} (\log n)^{\beta - \frac{3\alpha}{\gamma}} \\ &\leq CE|X|^\alpha. \end{aligned}$$

If  $\alpha \leq \frac{\gamma(\beta+1)}{3}$ , by Lemma 2.4 and  $p > \frac{\gamma(\beta+1)}{3}$ , we have

$$\begin{aligned} I_1 &\leq C \sum_{n=2}^{\infty} n^{-1} (\log n)^{\beta - \frac{3p}{\gamma}} E|X|^p I(|X| \leq (\log n)^{3/\gamma}) \\ &= C \sum_{n=2}^{\infty} n^{-1} (\log n)^{\beta - \frac{3p}{\gamma}} \sum_{m=2}^n E|X|^p I(\log(m-1) < |X|^{\gamma/3} \leq \log m) \\ &\leq C \sum_{m=2}^{\infty} E|X|^p I(\log(m-1) < |X|^{\gamma/3} \leq \log m) \sum_{n=m}^{\infty} n^{-1} (\log n)^{\beta - \frac{3p}{\gamma}} \\ &\leq C \sum_{m=2}^{\infty} (\log m)^{\beta - \frac{3p}{\gamma} + 1} E|X|^p I(\log(m-1) < |X|^{\gamma/3} \leq \log m) \\ &= C \sum_{m=2}^{\infty} (\log m)^{\beta - \frac{3p}{\gamma} + 1} E|X|^{\frac{\gamma(\beta+1)}{3}} E|X|^{p - \frac{\gamma(\beta+1)}{3}} I(\log(m-1) < |X|^{\gamma/3} \leq \log m) \\ &\leq C \sum_{m=2}^{\infty} (\log m)^{\beta - \frac{3p}{\gamma} + 1} (\log m)^{\frac{3}{\gamma}(p - \frac{\gamma(\beta+1)}{3})} E|X|^{\frac{\gamma(\beta+1)}{3}} I(\log(m-1) < |X|^{\gamma/3} \leq \log m) \\ &\leq CE|X|^{\frac{\gamma(\beta+1)}{3}}. \end{aligned}$$

Next for  $I_2$ , by Lemma 2.4 and  $p > \alpha$ , we have

$$\begin{aligned} I_2 &\leq C \sum_{n=2}^{\infty} n^{-1} (\log n)^{\beta - \frac{3p}{\gamma}} \sum_{k=1}^{\infty} E|X|^p I\left(k^{\frac{1}{\alpha}} < \frac{|X|}{(\log n)^{3/\gamma}} \leq (k+1)^{\frac{1}{\alpha}}\right) \sum_{j=k}^{\infty} \#(I_{nj}) j^{-p/\alpha} \\ &\leq C \sum_{n=2}^{\infty} n^{-1} (\log n)^{\beta - \frac{3p}{\gamma}} \sum_{k=1}^{\infty} (k+1)^{1 - \frac{p}{\alpha}} E|X|^p I\left(k^{\frac{1}{\alpha}} < \frac{|X|}{(\log n)^{3/\gamma}} \leq (k+1)^{\frac{1}{\alpha}}\right) \\ &= C \sum_{n=2}^{\infty} n^{-1} (\log n)^{\beta - \frac{3p}{\gamma}} \sum_{k=1}^{\infty} (k+1)^{1 - \frac{p}{\alpha}} E|X|^\alpha E|X|^{p-\alpha} I\left(k^{\frac{1}{\alpha}} < \frac{|X|}{(\log n)^{3/\gamma}} \leq (k+1)^{\frac{1}{\alpha}}\right) \\ &\leq C \sum_{n=2}^{\infty} n^{-1} (\log n)^{\beta - \frac{3\alpha}{\gamma}} \sum_{k=1}^{\infty} E|X|^\alpha I\left(k^{\frac{1}{\alpha}} < \frac{|X|}{(\log n)^{3/\gamma}} \leq (k+1)^{\frac{1}{\alpha}}\right) \\ &= C \sum_{n=2}^{\infty} n^{-1} (\log n)^{\beta - \frac{3\alpha}{\gamma}} E|X|^\alpha I(|X| > (\log n)^{3/\gamma}) \\ &= C \sum_{n=2}^{\infty} n^{-1} (\log n)^{\beta - \frac{3\alpha}{\gamma}} \sum_{m=n}^{\infty} E|X|^\alpha I(\log m < |X|^{\gamma/3} \leq \log(m+1)) \end{aligned}$$

$$= C \sum_{m=2}^{\infty} E|X|^{\alpha} I(\log m < |X|^{\gamma/3} \leq \log(m+1)) \sum_{n=2}^m n^{-1} (\log n)^{\beta - \frac{3\alpha}{\gamma}}.$$

Observing that

$$\sum_{n=2}^m n^{-1} (\log n)^{\beta - \frac{3\alpha}{\gamma}} \leq \begin{cases} C, & \text{for } \alpha > \frac{\gamma(\beta+1)}{3}, \\ C \log \log m, & \text{for } \alpha = \frac{\gamma(\beta+1)}{3}, \\ C(\log m)^{\beta - \frac{3\alpha}{\gamma} + 1}, & \text{for } \alpha < \frac{\gamma(\beta+1)}{3}, \end{cases}$$

we get

$$I_2 \leq \begin{cases} CE|X|^{\alpha}, & \text{for } \alpha > \frac{\gamma(\beta+1)}{3}, \\ CE|X|^{\alpha} \log(1+|X|), & \text{for } \alpha = \frac{\gamma(\beta+1)}{3}, \\ CE|X|^{\gamma(\beta+1)/3}, & \text{for } \alpha < \frac{\gamma(\beta+1)}{3}. \end{cases}$$

Therefore,

$$I = I_1 + I_2 \leq \begin{cases} CE|X|^{\alpha}, & \text{for } \alpha > \frac{\gamma(\beta+1)}{3}, \\ CE|X|^{\alpha} \log(1+|X|), & \text{for } \alpha = \frac{\gamma(\beta+1)}{3}, \\ CE|X|^{\gamma(\beta+1)/3}, & \text{for } \alpha < \frac{\gamma(\beta+1)}{3}. \end{cases}$$

The proof of Lemma 2.6 is completed.  $\square$

*Proof of Lemma 2.7* We will discuss (2.7) into two cases.

**Case I**  $0 < \alpha \leq 1$ .

By Lemma 2.3, Markov's inequality and  $E|X|^{\alpha} < \infty$  we have

$$\begin{aligned} \frac{1}{b_n} \max_{1 \leq j \leq n} \left| \sum_{i=1}^j EY_{ni} \right| &\leq \frac{1}{b_n} \sum_{i=1}^n |EY_{ni}| \\ &\leq \frac{C}{b_n} \sum_{i=1}^n E|a_{ni}X_{ni}| I(|a_{ni}X_{ni}| \leq b_n) + C \sum_{i=1}^n P(|a_{ni}X_{ni}| > b_n) \\ &\leq \frac{C}{b_n} \sum_{i=1}^n E|a_{ni}X| I(|a_{ni}X| \leq b_n) + C \sum_{i=1}^n P(|a_{ni}X| > b_n) \\ &\leq \frac{C}{b_n} \sum_{i=1}^n E|a_{ni}X|^{\alpha} |a_{ni}X|^{1-\alpha} I(|a_{ni}X| \leq b_n) + \frac{C}{b_n^{\alpha}} \sum_{i=1}^n |a_{ni}|^{\alpha} E|X|^{\alpha} \\ &\leq \frac{C}{b_n^{\alpha}} \sum_{i=1}^n |a_{ni}|^{\alpha} E|X|^{\alpha} \\ &\leq C(\log n)^{-3\alpha/\gamma} E|X|^{\alpha} \longrightarrow 0 \quad \text{as } n \rightarrow \infty. \end{aligned} \tag{3.1}$$

**Case II**  $1 < \alpha < 2$ .

For all  $1 \leq i \leq n, n \geq 1$ , denote

$$\begin{aligned} Z_{ni} &= a_{ni}X_{ni} - Y_{ni} \\ &= (a_{ni}X_{ni} + b_n)I(a_{ni}X_{ni} < -b_n) + (a_{ni}X_{ni} - b_n)I(a_{ni}X_{ni} > b_n). \end{aligned}$$

Then

$$0 < Z_{ni} = a_{ni}X_{ni} - b_n < a_{ni}X_{ni} \quad \text{for } a_{ni}X_{ni} > b_n,$$

and

$$a_{ni}X_{ni} < a_{ni}X_{ni} + b_n = Z_{ni} < 0 \quad \text{for } a_{ni}X_{ni} < -b_n.$$

Hence,

$$|Z_{ni}| < |a_{ni}X_{ni}|I(|a_{ni}X_{ni}| > b_n).$$

By  $EX_{ni} = 0$  for  $1 < \alpha < 2$ , Lemma 2.3 and  $E|X|^\alpha < \infty$  we get

$$\begin{aligned} \frac{1}{b_n} \max_{1 \leq j \leq n} \left| \sum_{i=1}^j EY_{ni} \right| &= \frac{1}{b_n} \max_{1 \leq j \leq n} \left| \sum_{i=1}^j EZ_{ni} \right| \\ &\leq \frac{1}{b_n} \sum_{i=1}^n E|Z_{ni}| \\ &\leq \frac{1}{b_n} \sum_{i=1}^n E|a_{ni}X_{ni}|I(|a_{ni}X_{ni}| > b_n) \\ &\leq \frac{C}{b_n} \sum_{i=1}^n E|a_{ni}X|I(|a_{ni}X| > b_n) \\ &\leq \frac{C}{b_n^\alpha} \sum_{i=1}^n |a_{ni}|^\alpha E|X|^\alpha I(|a_{ni}X| > b_n) \\ &\leq C(\log n)^{-3\alpha/\gamma} E|X|^\alpha \longrightarrow 0 \quad \text{as } n \rightarrow \infty. \end{aligned} \quad (3.2)$$

From (3.1) and (3.2), we see that (2.7) holds.  $\square$

*Proof of Theorem 2.8* By (2.6) and

$$a_{ni} = a_{ni}^+ - a_{ni}^-,$$

without loss of generality, we may assume that  $a_{ni} \geq 0$  and  $\sum_{i=1}^n a_{ni}^\alpha \leq n$ .

For fixed  $n \geq 1$ , let  $Z_{ni}$  as those in the proof of Lemma 2.7 and

$$\begin{aligned} A &= \bigcap_{i=1}^n \{Y_{ni} = a_{ni}X_{ni}\}, \\ B = \bar{A} &= \bigcup_{i=1}^n \{Y_{ni} \neq a_{ni}X_{ni}\} = \bigcup_{i=1}^n \{|a_{ni}X_{ni}| > b_n\}, \\ E_{ni} &= \left\{ \max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni}X_{ni} \right| > \varepsilon b_n \right\}. \end{aligned}$$

It is easy to check that for all  $\varepsilon > 0$ ,

$$\begin{aligned} E_{ni} &= E_{ni}A \cup E_{ni}B \\ &\subset \left\{ \max_{1 \leq j \leq n} \left| \sum_{i=1}^j Y_{ni} \right| > \varepsilon b_n \right\} \cup \left\{ \bigcup_{i=1}^n \{|a_{ni}X_{ni}| > b_n\} \right\}. \end{aligned}$$

Then, by Lemma 2.7, for  $n$  large enough, we have

$$P(E_{ni}) \leq P\left( \max_{1 \leq j \leq n} \left| \sum_{i=1}^j Y_{ni} \right| > \varepsilon b_n \right) + P\left( \bigcup_{i=1}^n \{|a_{ni}X_{ni}| > b_n\} \right)$$

$$\begin{aligned}
&\leq P\left(\max_{1 \leq j \leq n} \left| \sum_{i=1}^j (Y_{ni} - EY_{ni}) \right| > \varepsilon b_n - \max_{1 \leq j \leq n} \left| \sum_{i=1}^j EY_{ni} \right|\right) + \sum_{i=1}^n P(|a_{ni}X_{ni}| > b_n) \\
&\leq P\left(\max_{1 \leq j \leq n} \left| \sum_{i=1}^j (Y_{ni} - EY_{ni}) \right| > \frac{\varepsilon b_n}{2}\right) + \sum_{i=1}^n P(|a_{ni}X_{ni}| > b_n). \tag{3.3}
\end{aligned}$$

To prove (2.9), it suffices to show that

$$J_1 \triangleq \sum_{n=2}^{\infty} \frac{1}{n} P\left(\max_{1 \leq j \leq n} \left| \sum_{i=1}^j (Y_{ni} - EY_{ni}) \right| > \frac{\varepsilon b_n}{2}\right) < \infty, \tag{3.4}$$

$$J_2 \triangleq \sum_{n=2}^{\infty} \frac{1}{n} \sum_{i=1}^n P(|a_{ni}X_{ni}| > b_n) < \infty. \tag{3.5}$$

First for  $J_1$ , by Lemma 2.1, it is easy to see that  $\{Y_{ni} - EY_{ni}, 1 \leq i \leq n, n \geq 1\}$  is an array of rowwise END random variables. Hence, it follows from Markov's inequality, Lemmas 2.2 and 2.3 that

$$\begin{aligned}
J_1 &\leq C \sum_{n=2}^{\infty} \frac{1}{nb_n^2} E\left(\max_{1 \leq j \leq n} \left| \sum_{i=1}^j (Y_{ni} - EY_{ni}) \right|^2\right) \\
&\leq C \sum_{n=2}^{\infty} \frac{\log^2 n}{nb_n^2} \sum_{i=1}^n E(Y_{ni} - EY_{ni})^2 \\
&\leq C \sum_{n=2}^{\infty} \frac{\log^2 n}{nb_n^2} \sum_{i=1}^n E(Y_{ni})^2 \\
&\leq C \sum_{n=2}^{\infty} \frac{\log^2 n}{nb_n^2} \sum_{i=1}^n E|a_{ni}X_{ni}|^2 I(|a_{ni}X_{ni}| \leq b_n) + C \sum_{n=2}^{\infty} \frac{\log^2 n}{n} \sum_{i=1}^n P(|a_{ni}X_{ni}| > b_n) \\
&\leq C \sum_{n=2}^{\infty} \frac{\log^2 n}{nb_n^2} \sum_{i=1}^n E|a_{ni}X|^2 I(|a_{ni}X| \leq b_n) + C \sum_{n=2}^{\infty} \frac{\log^2 n}{n} \sum_{i=1}^n P(|a_{ni}X| > b_n) \\
&\leq C \sum_{n=2}^{\infty} \frac{\log^2 n}{nb_n^2} \sum_{i=1}^n E|a_{ni}X|^2 I(|a_{ni}X| \leq b_n) + C \sum_{n=2}^{\infty} \frac{\log^2 n}{nb_n^\alpha} \sum_{i=1}^n E|a_{ni}X|^\alpha I(|a_{ni}X| > b_n) \\
&\triangleq J_{11} + J_{12}. \tag{3.6}
\end{aligned}$$

From Lemma 2.6 (i) (for  $p = \beta = 2$ ), (ii) and (2.8) we obtain that  $J_{11} < \infty$ ,  $J_{12} < \infty$  and then  $J_1 < \infty$ .

Finally, we have

$$J_2 \leq \sum_{n=2}^{\infty} \frac{1}{nb_n^\alpha} \sum_{i=1}^n E|a_{ni}X_{ni}|^\alpha I(|a_{ni}X_{ni}| > b_n) \leq J_{12} < \infty.$$

The proof of Theorem 2.8 is completed.  $\square$

*Proof of Theorem 2.9* For all  $\varepsilon > 0$ , we have

$$\begin{aligned}
&\sum_{n=2}^{\infty} \frac{1}{n} E\left(\frac{1}{b_n} \max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni}X_{ni} \right| - \varepsilon\right)_+^\tau \\
&= \sum_{n=2}^{\infty} \frac{1}{n} \int_0^\infty P\left(\frac{1}{b_n} \max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni}X_{ni} \right| - \varepsilon > t^{\frac{1}{\tau}}\right) dt
\end{aligned}$$

$$\begin{aligned}
&= \sum_{n=2}^{\infty} \frac{1}{n} \int_0^1 P\left(\frac{1}{b_n} \max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} X_{ni} \right| > \varepsilon + t^{\frac{1}{\tau}}\right) dt \\
&\quad + \sum_{n=2}^{\infty} \frac{1}{n} \int_1^{\infty} P\left(\frac{1}{b_n} \max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} X_{ni} \right| > \varepsilon + t^{\frac{1}{\tau}}\right) dt \\
&\leq \sum_{n=2}^{\infty} \frac{1}{n} P\left(\max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} X_{ni} \right| > \varepsilon b_n\right) \\
&\quad + \sum_{n=2}^{\infty} \frac{1}{n} \int_1^{\infty} P\left(\max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} X_{ni} \right| > b_n t^{\frac{1}{\tau}}\right) dt \\
&\triangleq K_1 + K_2. \tag{3.7}
\end{aligned}$$

To prove (2.10), it suffices to show that  $K_1 < \infty$  and  $K_2 < \infty$ . By Theorem 2.8, we have  $K_1 < \infty$ . Next for  $K_2$ . For each  $1 \leq i \leq n, n \geq 1$  and all  $t \geq 1$ , let

$$Y'_{ni} = a_{ni} X_{ni} I(|a_{ni} X_{ni}| \leq b_n t^{1/\tau}) + b_n t^{1/\tau} I(a_{ni} X_{ni} > b_n t^{1/\tau}) - b_n t^{1/\tau} I(a_{ni} X_{ni} < -b_n t^{1/\tau});$$

$$Z'_{ni} = a_{ni} X_{ni} - Y'_{ni};$$

$$A' = \bigcap_{i=1}^n \{Y'_{ni} = a_{ni} X_{ni}\};$$

$$B' = \overline{A'} = \bigcup_{i=1}^n \{Y'_{ni} \neq a_{ni} X_{ni}\} = \bigcup_{i=1}^n \{|a_{ni} X_{ni}| > b_n t^{1/\tau}\};$$

$$E'_{ni} = \left\{ \max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} X_{ni} \right| > b_n t^{1/\tau} \right\}.$$

By Lemma 2.7, for all  $t \geq 1$ , and  $n$  large enough

$$\begin{aligned}
P(E'_{ni}) &\leq P\left(\max_{1 \leq j \leq n} \left| \sum_{i=1}^j Y'_{ni} \right| > b_n t^{1/\tau}\right) + P\left(\bigcup_{i=1}^n \{|a_{ni} X_{ni}| > b_n t^{1/\tau}\}\right) \\
&\leq P\left(\max_{1 \leq j \leq n} \left| \sum_{i=1}^j (Y'_{ni} - EY'_{ni}) \right| > b_n t^{1/\tau} - \max_{1 \leq j \leq n} \left| \sum_{i=1}^j EY'_{ni} \right|\right) \\
&\quad + \sum_{i=1}^n P(|a_{ni} X_{ni}| > b_n t^{1/\tau}) \\
&\leq P\left(\max_{1 \leq j \leq n} \left| \sum_{i=1}^j (Y'_{ni} - EY'_{ni}) \right| > \frac{1}{2} b_n t^{1/\tau}\right) + \sum_{i=1}^n P(|a_{ni} X_{ni}| > b_n t^{1/\tau}). \tag{3.8}
\end{aligned}$$

To prove  $K_2 < \infty$ , it suffices to show that

$$K_{21} \triangleq \sum_{n=2}^{\infty} \frac{1}{n} \int_1^{\infty} P\left(\max_{1 \leq j \leq n} \left| \sum_{i=1}^j (Y'_{ni} - EY'_{ni}) \right| > \frac{1}{2} b_n t^{1/\tau}\right) dt < \infty; \tag{3.9}$$

$$K_{22} \triangleq \sum_{n=2}^{\infty} \frac{1}{n} \int_1^{\infty} \sum_{i=1}^n P(|a_{ni} X_{ni}| > b_n t^{1/\tau}) dt < \infty. \tag{3.10}$$

By Lemma 2.1,  $\{Y'_{ni} - EY'_{ni}, 1 \leq i \leq n, n \geq 1\}$  is still an array of rowwise END random

variables. Hence, it follows from Markov's inequality, Lemmas 2.2 and 2.3 that

$$\begin{aligned}
K_{21} &\leq C \sum_{n=2}^{\infty} \frac{1}{n} \int_1^{\infty} \frac{1}{b_n^2 t^{2/\tau}} E \left( \max_{1 \leq j \leq n} \left| \sum_{i=1}^j (Y'_{ni} - EY'_{ni}) \right|^2 \right) dt \\
&\leq C \sum_{n=2}^{\infty} \frac{1}{n} \int_1^{\infty} \frac{\log^2 n}{b_n^2 t^{2/\tau}} \sum_{i=1}^n E (Y'_{ni} - EY'_{ni})^2 dt \\
&\leq C \sum_{n=2}^{\infty} \frac{\log^2 n}{n} \int_1^{\infty} \sum_{i=1}^n P(|a_{ni} X_{ni}| > b_n t^{1/\tau}) dt \\
&\quad + C \sum_{n=2}^{\infty} \frac{\log^2 n}{nb_n^2} \int_1^{\infty} t^{-2/\tau} \sum_{i=1}^n E |a_{ni} X_{ni}|^2 (|a_{ni} X_{ni}| \leq b_n t^{1/\tau}) dt \\
&\leq C \sum_{n=2}^{\infty} \frac{\log^2 n}{n} \int_1^{\infty} \sum_{i=1}^n P(|a_{ni} X| > b_n t^{1/\tau}) dt \\
&\quad + C \sum_{n=2}^{\infty} \frac{\log^2 n}{nb_n^2} \int_1^{\infty} t^{-2/\tau} \sum_{i=1}^n E |a_{ni} X|^2 (|a_{ni} X| \leq b_n) dt \\
&\quad + C \sum_{n=2}^{\infty} \frac{\log^2 n}{nb_n^2} \int_1^{\infty} t^{-2/\tau} \sum_{i=1}^n E |a_{ni} X|^2 (b_n < |a_{ni} X| \leq b_n t^{1/\tau}) dt \\
&\triangleq CK_{211} + CK_{212} + CK_{213}. \tag{3.11}
\end{aligned}$$

For  $0 < \tau < \alpha$  and (2.6), we get

$$\begin{aligned}
K_{211} &= C \sum_{n=2}^{\infty} \frac{\log^2 n}{n} \int_1^{\infty} \sum_{i=1}^n P \left( \frac{|a_{ni} X|^\tau}{b_n^\tau} > t \right) dt \\
&\leq C \sum_{n=2}^{\infty} \frac{\log^2 n}{nb_n^\tau} \sum_{i=1}^n E |a_{ni} X|^\tau I(|a_{ni} X| > b_n) \\
&\leq C \sum_{n=2}^{\infty} \frac{\log^2 n}{nb_n^\alpha} \sum_{i=1}^n E |a_{ni} X|^\alpha I(|a_{ni} X| > b_n) \\
&\leq CJ_{12} < \infty. \tag{3.12}
\end{aligned}$$

For  $0 < \tau < \alpha < 2$ , it follows from Lemma 2.6 and (2.8) that

$$K_{212} \leq C \sum_{n=2}^{\infty} \frac{\log^2 n}{nb_n^2} \sum_{i=1}^n E |a_{ni} X|^2 (|a_{ni} X| \leq b_n) < \infty. \tag{3.13}$$

Taking  $t = x^\tau$ , by Markov's inequality and Lemma 2.3 it follows that

$$\begin{aligned}
K_{213} &\leq C \sum_{n=2}^{\infty} \frac{\log^2 n}{nb_n^2} \int_1^{\infty} x^{\tau-3} \sum_{i=1}^n E |a_{ni} X|^2 (b_n < |a_{ni} X| \leq b_n x) dx \\
&= C \sum_{n=2}^{\infty} \frac{\log^2 n}{nb_n^2} \sum_{m=1}^{\infty} \int_m^{m+1} x^{\tau-3} \sum_{i=1}^n E |a_{ni} X|^2 (b_n < |a_{ni} X| \leq b_n x) dx \\
&\leq C \sum_{n=2}^{\infty} \frac{\log^2 n}{nb_n^2} \sum_{m=1}^{\infty} m^{\tau-3} \sum_{i=1}^n E |a_{ni} X|^2 (b_n < |a_{ni} X| \leq b_n(m+1)) \\
&\leq C \sum_{n=2}^{\infty} \frac{\log^2 n}{nb_n^2} \sum_{i=1}^n \sum_{m=1}^{\infty} \sum_{s=1}^m m^{\tau-3} E |a_{ni} X|^2 (b_n s < |a_{ni} X| \leq b_n(s+1))
\end{aligned}$$

$$\begin{aligned}
&= C \sum_{n=2}^{\infty} \frac{\log^2 n}{nb_n^2} \sum_{i=1}^n \sum_{s=1}^{\infty} E|a_{ni}X|^2 (b_n s < |a_{ni}X| \leq b_n(s+1)) \sum_{m=s}^{\infty} m^{\tau-3} \\
&= C \sum_{n=2}^{\infty} \frac{\log^2 n}{nb_n^2} \sum_{i=1}^n \sum_{s=1}^{\infty} s^{\tau-2} E|a_{ni}X|^2 (b_n s < |a_{ni}X| \leq b_n(s+1)) \\
&= C \sum_{n=2}^{\infty} \frac{\log^2 n}{nb_n^2} \sum_{i=1}^n \sum_{s=1}^{\infty} s^{\tau-2} E|a_{ni}X|^{\tau} |a_{ni}X|^{2-\tau} (b_n s < |a_{ni}X| \leq b_n(s+1)) \\
&\leq C \sum_{n=2}^{\infty} \frac{\log^2 n}{nb_n^{\tau}} \sum_{i=1}^n E|a_{ni}X|^{\tau} I(|a_{ni}X| > b_n) \\
&\leq C \sum_{n=2}^{\infty} \frac{\log^2 n}{nb_n^{\alpha}} \sum_{i=1}^n E|a_{ni}X|^{\alpha} I(|a_{ni}X| > b_n) \\
&\leq CJ_{12} < \infty.
\end{aligned} \tag{3.14}$$

By (3.11)–(3.14), we get (3.9). On the other hand, it is obviously to see that

$$K_{22} \leq CK_{211} < \infty,$$

which completes the proof of Theorem 2.9.  $\square$

*Proof of Theorem 2.11* For all  $\varepsilon > 0$ , from (2.9) we know that

$$\begin{aligned}
\infty &> \sum_{n=2}^{\infty} n^{-1} P\left(\max_{1 \leq j \leq n} |T_j| > \varepsilon b_n\right) \\
&\geq \sum_{i=0}^{\infty} \sum_{n=2^i}^{2^{i+1}-1} (2^{i+1}-1)^{-1} P\left(\max_{1 \leq j \leq n} |T_j| > \varepsilon b_n\right) \\
&\geq C \sum_{i=0}^{\infty} \sum_{n=2^i}^{2^{i+1}-1} (2^{i+1}-1)^{-1} P\left(\max_{1 \leq j \leq 2^i} |T_j| > \varepsilon b_{2^{i+1}}\right) \\
&\geq C \sum_{i=1}^{\infty} P\left(\max_{1 \leq j \leq 2^i} |T_j| > \varepsilon b_{2^{i+1}}\right).
\end{aligned} \tag{3.15}$$

It follows from Borel–Cantelli lemma that

$$P\left(\max_{1 \leq j \leq 2^i} |T_j| > \varepsilon b_{2^{i+1}}, i.o.\right) = 0. \tag{3.16}$$

Hence

$$\frac{\max_{1 \leq j \leq 2^i} |T_j|}{b_{2^{i+1}}} \longrightarrow 0, \quad \text{a.s. as } i \rightarrow \infty. \tag{3.17}$$

For all positive integers  $n$ , there exists a non-negative integer  $i$ , such that  $2^{i-1} \leq n < 2^i$ . Thus

$$\begin{aligned}
\max_{2^{i-1} \leq n \leq 2^i} \frac{|T_n|}{b_n} &\leq \frac{\max_{1 \leq n \leq 2^i} |T_j|}{b_{2^{i-1}}} \\
&= \frac{\max_{1 \leq n \leq 2^i} |T_j|}{b_{2^{i+1}}} \cdot 2^{2/\alpha} \cdot \left(\frac{i+1}{i-1}\right)^{3/\gamma} \rightarrow 0, \quad \text{a.s. as } i \rightarrow \infty.
\end{aligned} \tag{3.18}$$

We have

$$\lim_{n \rightarrow \infty} \frac{|T_n|}{b_n} = 0, \quad \text{a.s.} \tag{3.19}$$



The proof of Theorem 2.11 is completed.  $\square$

**Acknowledgements** The author is most grateful to the Editor-in-Chief, Associate Editor and anonymous referee for careful reading of the manuscript and valuable suggestions which helped in improving an earlier version of this paper.

## References

- [1] Adler, A., Rosalsky, A.: Some general strong laws for weighted sums of stochastically dominated random variables. *Stoch. Anal. Appl.*, **5**, 1–16 (1987)
- [2] Adler, A., Rosalsky, A., Taylor, R. L.: Strong laws of large numbers for weighted sums of random elements in normed linear spaces. *Int. J. Math. Math. Sci.*, **12**, 507–530 (1989)
- [3] Baum, L. E., Katz, M.: Convergence rates in the law of large numbers. *Trans. Amer. Math. Soc.*, **120**(1), 108–123 (1965)
- [4] Birkel, T.: A note on the strong law of large numbers for positively dependent random variables. *Stat. Prob. Letters*, **7**, 17–20 (1989)
- [5] Chen, Y., Chen, A., Ng, K.: The strong law of large numbers for extended negatively dependent random variables. *J. Appl. Prob.*, **47**(4), 908–922 (2010)
- [6] Cheng, F. X., Yan, J. G., Yang, L. J.: Extended Glivenko Cantelli theorem in nonparametric regression. *Comm. Statist. Theory Methods*, **43**(17), 3720–3725 (2014)
- [7] Chow, Y. S.: On the rate of moment convergence of sample sums and extremes. *Bull. Institute Math.*, **16**(3), 177–201 (1988)
- [8] Etemadi, N.: An elementary proof of the strong law of large numbers. *Z. Wahrsch. Verw. Gebiete.*, **55**, 119–122 (1981)
- [9] Etemadi, N.: On the strong law of large numbers for nonnegative random variables. *J. Multivariate Anal.*, **13**, 187–193 (1983)
- [10] Heyde, C. C.: A supplement to the strong law of large numbers. *J. Appl. Prob.*, **12**, 173–175 (1975)
- [11] Hsu, P. L., Robbins, H.: Complete convergence and the law of large numbers. *Proc. Natl. Acad. Sci. USA*, **33**, 25–31 (1947)
- [12] Liu, L.: Precise large deviations for dependent random variables with heavy tails. *Stat. Prob. Letters*, **79**, 1290–1298 (2009)
- [13] Liu, L.: Necessary and sufficient conditions for moderate deviations of dependent random variables with heavy tails. *Sci. China Ser. A*, **53**(6), 1421–1434 (2010)
- [14] Qiu, D. H., Chen, P. Y.: Limiting behavior of weighted sums of NOD random variables. *J. Math. Res. Exposition*, **31**, 1081–1091 (2011)
- [15] Qiu, D. H., Chen, P. Y.: Complete and complete moment convergence for weighted sums of widely orthant dependent random variables. *Acta Math. Sin., Engl. Ser.*, **30**(9), 1539–1548 (2014)
- [16] Qiu, D. H., Hu, T. C.: Strong limit theorems for weighted sums of widely orthant dependent random variables. *J. Math. Res. Appl.*, **34**(1), 105–113 (2014)
- [17] Seneta, E.: Regularly varying function. Lecture Notes in Math., Vol. 508. Springer, Berlin, 1976
- [18] Shen, A.: Probability inequalities for END sequence and their applications. *J. Inequal. Appl.*, **98**, 1–12 (2011)
- [19] Shen, A., Wu, R.: Strong and weak convergence for asymptotically almost negatively associated random variables. *Discret Dyn. Nat. Soc.*, **7**, Article ID: 235012 (2013)
- [20] Shen, A., Xue, M., Wang, W.: Complete convergence for weighted sums of extended negatively dependent random variables. *Comm. Statist. Theory Methods*, **46**(3), 1433–1444 (2017)
- [21] Stout, W.: Almost Sure Convergence, Academic Press, New York, 1974
- [22] Wang, X. J., Deng, X., Zheng, L. L., et al.: Complete convergence for arrays of rowwise negatively superadditive-dependent random variables and its applications. *Statistics: J. Theor. Appl. Stat.*, **48**(4), 834–850 (2014)
- [23] Wang, X. J., Hu, S. H.: The consistency of the nearest neighbor estimator of the density function based on WOD samples. *J. Math. Anal. Appl.*, **429**, 497–512 (2015)

- [24] Wang, X. J., Li, X. Q., Hu, S. H., et al.: On complete convergence for an extended negatively dependent sequence. *Comm. Statist. Theory Methods*, **43**, 2923–2937 (2014)
- [25] Wang, X. J., Xu, C., Hu, T. C., et al.: On complete convergence for widely Orthant-dependent random variables and its applications in nonparametric regression models. *TEST*, **23**, 607–629 (2014)
- [26] Wang, X. J., Zheng, L. L., Xu, C., et al.: Complete consistency for the estimator of nonparameter regression models based on extended negatively dependent errors. *Statistics: J. Theor. Appl. Stat.*, **49**, 396–407 (2015)
- [27] Wu, Y. F., Song, M. Z., Wang, C. H.: Complete moment convergence and mean convergence for arrays of rowwise extended negatively dependent random variables. *The Scientific World Journal*, Artical ID 478612, 7 pages, doi: [10.1155/2014/478612](https://doi.org/10.1155/2014/478612) (2014)
- [28] Yan, J. G., Su, C.: Precise asymptotics of U-statistics. *Acta Math. Sin., Chin. Ser.*, **50**, 517–526 (2007)
- [29] Yan, J. G.: The uniform convergence and precise asymptotics of generalized stochastic order statistics. *J. Math. Anal. Appl.*, **343**, 644–653 (2008)
- [30] Yan, J. G.: Strong stability of a type of Jamison weighted sums for END random variables. *J. Korean Math. Soc.*, **54**(3), 897–907 (2017)
- [31] Yan, J. G.: Almost sure convergence for weighted sums of WNOD random variables and its applications in nonparametric regression models. *Comm. Statist. Theory Methods*, 2017
- [32] Yang, W. Z., Liu, T. T., Wang, X. J., et al.: On the Bahadur representation of sample quantiles for widely orthant dependent sequences. *Filomat*, **28**(7), 1333–1343 (2014)
- [33] Yang, W. Z., Wang, X. J., Wang, X. H., et al.: The consistency for estimator of nonparametric regression model based on NOD errors. *J. Inequal. Appl.*, 13 pages, Article ID 140 (2012)



Please check it.

Any updates for [27,31]?



# IRTG 1792 Discussion Paper Series 2018

For a complete list of Discussion Papers published, please visit [irtg1792.hu-berlin.de](http://irtg1792.hu-berlin.de).

- 001 "Data Driven Value-at-Risk Forecasting using a SVR-GARCH-KDE Hybrid" by Marius Lux, Wolfgang Karl Härdle and Stefan Lessmann, January 2018.
- 002 "Nonparametric Variable Selection and Its Application to Additive Models" by Zheng-Hui Feng, Lu Lin, Ruo-Qing Zhu and Li-Xing Zhu, January 2018.
- 003 "Systemic Risk in Global Volatility Spillover Networks: Evidence from Option-implied Volatility Indices " by Zihui Yang and Yinggang Zhou, January 2018.
- 004 "Pricing Cryptocurrency options: the case of CRIX and Bitcoin" by Cathy YH Chen, Wolfgang Karl Härdle, Ai Jun Hou and Weining Wang, January 2018.
- 005 "Testing for bubbles in cryptocurrencies with time-varying volatility" by Christian M. Hafner, January 2018.
- 006 "A Note on Cryptocurrencies and Currency Competition" by Anna Almosova, January 2018.
- 007 "Knowing me, knowing you: inventor mobility and the formation of technology-oriented alliances" by Stefan Wagner and Martin C. Goossen, February 2018.
- 008 "A Monetary Model of Blockchain" by Anna Almosova, February 2018.
- 009 "Deregulated day-ahead electricity markets in Southeast Europe: Price forecasting and comparative structural analysis" by Antanina Hryshchuk, Stefan Lessmann, February 2018.
- 010 "How Sensitive are Tail-related Risk Measures in a Contamination Neighbourhood?" by Wolfgang Karl Härdle, Chengxiu Ling, February 2018.
- 011 "How to Measure a Performance of a Collaborative Research Centre" by Alona Zharova, Janine Tellingner-Rice, Wolfgang Karl Härdle, February 2018.
- 012 "Targeting customers for profit: An ensemble learning framework to support marketing decision making" by Stefan Lessmann, Kristof Coussement, Koen W. De Bock, Johannes Haupt, February 2018.
- 013 "Improving Crime Count Forecasts Using Twitter and Taxi Data" by Lara Vomfell, Wolfgang Karl Härdle, Stefan Lessmann, February 2018.
- 014 "Price Discovery on Bitcoin Markets" by Paolo Pagnottoni, Dirk G. Baur, Thomas Dimpfl, March 2018.
- 015 "Bitcoin is not the New Gold - A Comparison of Volatility, Correlation, and Portfolio Performance" by Tony Klein, Hien Pham Thu, Thomas Walther, March 2018.
- 016 "Time-varying Limit Order Book Networks" by Wolfgang Karl Härdle, Shi Chen, Chong Liang, Melanie Schienle, April 2018.
- 017 "Regularization Approach for Network Modeling of German EnergyMarket" by Shi Chen, Wolfgang Karl Härdle, Brenda López Cabrera, May 2018.
- 018 "Adaptive Nonparametric Clustering" by Kirill Efimov, Larisa Adamyan, Vladimir Spokoiny, May 2018.
- 019 "Lasso, knockoff and Gaussian covariates: a comparison" by Laurie Davies, May 2018.

**IRTG 1792, Spandauer Straße 1, D-10178 Berlin**  
**<http://irtg1792.hu-berlin.de>**

This research was supported by the Deutsche  
Forschungsgemeinschaft through the IRTG 1792.



# IRTG 1792 Discussion Paper Series 2018

For a complete list of Discussion Papers published, please visit [irtg1792.hu-berlin.de](http://irtg1792.hu-berlin.de).

- 020 "A Regime Shift Model with Nonparametric Switching Mechanism" by Haiqiang Chen, Yingxing Li, Ming Lin and Yanli Zhu, May 2018.
- 021 "LASSO-Driven Inference in Time and Space" by Victor Chernozhukov, Wolfgang K. Härdle, Chen Huang, Weining Wang, June 2018.
- 022 " Learning from Errors: The case of monetary and fiscal policy regimes" by Andreas Tryphonides, June 2018.
- 023 "Textual Sentiment, Option Characteristics, and Stock Return Predictability" by Cathy Yi-Hsuan Chen, Matthias R. Fengler, Wolfgang Karl Härdle, Yanchu Liu, June 2018.
- 024 "Bootstrap Confidence Sets For Spectral Projectors Of Sample Covariance" by A. Naumov, V. Spokoiny, V. Ulyanov, June 2018.
- 025 "Construction of Non-asymptotic Confidence Sets in 2 -Wasserstein Space" by Johannes Ebert, Vladimir Spokoiny, Alexandra Suvorikova, June 2018.
- 026 "Large ball probabilities, Gaussian comparison and anti-concentration" by Friedrich Götze, Alexey Naumov, Vladimir Spokoiny, Vladimir Ulyanov, June 2018.
- 027 "Bayesian inference for spectral projectors of covariance matrix" by Igor Silin, Vladimir Spokoiny, June 2018.
- 028 "Toolbox: Gaussian comparison on Euclidian balls" by Andzhey Koziuk, Vladimir Spokoiny, June 2018.
- 029 "Pointwise adaptation via stagewise aggregation of local estimates for multiclass classification" by Nikita Puchkin, Vladimir Spokoiny, June 2018.
- 030 "Gaussian Process Forecast with multidimensional distributional entries" by Francois Bachoc, Alexandra Suvorikova, Jean-Michel Loubes, Vladimir Spokoiny, June 2018.
- 031 "Instrumental variables regression" by Andzhey Koziuk, Vladimir Spokoiny, June 2018.
- 032 "Understanding Latent Group Structure of Cryptocurrencies Market: A Dynamic Network Perspective" by Li Guo, Yubo Tao and Wolfgang Karl Härdle, July 2018.
- 033 "Optimal contracts under competition when uncertainty from adverse selection and moral hazard are present" by Natalie Packham, August 2018.
- 034 "A factor-model approach for correlation scenarios and correlation stress-testing" by Natalie Packham and Fabian Woebbecking, August 2018.
- 035 "Correlation Under Stress In Normal Variance Mixture Models" by Michael Kalkbrener and Natalie Packham, August 2018.
- 036 "Model risk of contingent claims" by Nils Detering and Natalie Packham, August 2018.
- 037 "Default probabilities and default correlations under stress" by Natalie Packham, Michael Kalkbrener and Ludger Overbeck, August 2018.
- 038 "Tail-Risk Protection Trading Strategies" by Natalie Packham, Jochen Papenbrock, Peter Schwendner and Fabian Woebbecking, August 2018.

**IRTG 1792, Spandauer Straße 1, D-10178 Berlin**  
**<http://irtg1792.hu-berlin.de>**

This research was supported by the Deutsche  
Forschungsgemeinschaft through the IRTG 1792.



# IRTG 1792 Discussion Paper Series 2018

For a complete list of Discussion Papers published, please visit [irtg1792.hu-berlin.de](http://irtg1792.hu-berlin.de).

- 039 "Penalized Adaptive Forecasting with Large Information Sets and Structural Changes" by Lenka Zbonakova, Xinjue Li and Wolfgang Karl Härdle, August 2018.
- 040 "Complete Convergence and Complete Moment Convergence for Maximal Weighted Sums of Extended Negatively Dependent Random Variables" by Ji Gao YAN, August 2018.

**IRTG 1792, Spandauer Straße 1, D-10178 Berlin**  
**<http://irtg1792.hu-berlin.de>**

This research was supported by the Deutsche  
Forschungsgemeinschaft through the IRTG 1792.