

Kuczmazewska, Anna; Yan, Ji Gao

**Working Paper**

## On complete convergence in Marcinkiewicz-Zygmund type SLLN for random variables

IRTG 1792 Discussion Paper, No. 2018-041

**Provided in Cooperation with:**

Humboldt University Berlin, International Research Training Group 1792 "High Dimensional Nonstationary Time Series"

*Suggested Citation:* Kuczmazewska, Anna; Yan, Ji Gao (2018) : On complete convergence in Marcinkiewicz-Zygmund type SLLN for random variables, IRTG 1792 Discussion Paper, No. 2018-041, Humboldt-Universität zu Berlin, International Research Training Group 1792 "High Dimensional Nonstationary Time Series", Berlin

This Version is available at:

<https://hdl.handle.net/10419/230752>

**Standard-Nutzungsbedingungen:**

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.


Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

**Terms of use:**

*Documents in EconStor may be saved and copied for your personal and scholarly purposes.*

*You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.*

*If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.*



# On complete convergence in Marcinkiewicz-Zygmund type SLLN for random variables

Anna Kuczmaszewska <sup>\*</sup>  
Ji Gao YAN <sup>\*2</sup>



<sup>\*</sup> Lublin University of Technology, Poland  
<sup>\*2</sup> Soochow University, PR China

This research was supported by the Deutsche  
Forschungsgemeinschaft through the  
International Research Training Group 1792  
"High Dimensional Nonstationary Time Series".

<http://irtg1792.hu-berlin.de>  
ISSN 2568-5619

# On complete convergence in Marcinkiewicz-Zygmund type SLLN for random variables<sup>1</sup>

Anna Kuczmazewska

Department of Applied Mathematics, Lublin University of Technology,  
Nadbystrzycka 38 D, 20-618 Lublin, Poland

Jigao Yan<sup>2</sup>

School of Mathematical Sciences, Soochow University, Suzhou 215006, China  
C.A.S.E. - Center for Applied Statistics and Economics, Humboldt-Universität zu Berlin,  
Spandauer Str. 1, 10178 Berlin, Germany

**Abstract:** We consider a generalization of Baum-Katz theorem for random variables satisfying some cover conditions. Consequently, we get the result for many dependent structure, such as END,  $\varrho^*$ -mixing,  $\varrho^-$ -mixing and  $\varphi$ -mixing, etc.

**MSC (2010):** 60F15.

**Keywords:** Complete convergence; Marcinkiewicz-Zygmund type SLLN; Extended negatively dependent; Mixing dependency; Weakly mean bounded.

## 1 Introduction

In this paper, we consider a sequence  $\{X_n, n \geq 1\}$  of random variables defined on some probability space  $(\Omega, \mathcal{F}, P)$ . Hsu and Robbins [10] introduced the following concept of complete convergence. A sequence  $\{X_n, n \geq 1\}$  is said to converge completely to a constant  $C$  if

$$\sum_{n=1}^{\infty} P(|X_n - C| > \varepsilon) < \infty,$$

for all  $\varepsilon > 0$ . For independent and identically distributed (i.i.d., in short) random variables  $\{X, X_n, n \geq 1\}$ , let  $S_n = \sum_{k=1}^n X_k, n \geq 1$  be the partial sums, Hsu and Robbins [10] proved that  $S_n/n$  converge completely to  $EX$ , provided  $DX < \infty$ . Erdős [8] proved the converse theorem. This Hsu-Robbins-Erdős's theorem was generalized in different ways. Katz [13], Baum and Katz [1], and Chow [7] formed the following generalization of Marcinkiewicz-Zygmund type.

**Theorem 1.1** *Let  $\{X, X_n, n \geq 1\}$  be a sequence of i.i.d. random variables and let  $\alpha p \geq 1, \alpha > 1/2$ . Then the following statements are equivalent:*

- (i)  $E|X|^p < \infty$  and  $EX = 0$  if  $p \geq 1$ ;
  - (ii)  $\sum_{n=1}^{\infty} n^{\alpha p - 2} P(|S_n| > \varepsilon n^{\alpha}) < \infty$  for all  $\varepsilon > 0$ ;
  - (iii)  $\sum_{n=1}^{\infty} n^{\alpha p - 2} P(\max_{1 \leq i \leq n} |S_i| > \varepsilon n^{\alpha}) < \infty$  for all  $\varepsilon > 0$ ;
- If  $\alpha p > 1, \alpha > 1/2$  the above are also equivalent to*
- (iv)  $\sum_{n=1}^{\infty} n^{\alpha p - 2} P(\sup_{i \geq n} i^{-\alpha} |S_i| > \varepsilon) < \infty$  for all  $\varepsilon > 0$ .

---

<sup>1</sup>Supported by the National Natural Science Foundation of China (No. 11571250)

<sup>2</sup>**Correspondence:** yanjigao@suda.edu.cn (Jigao Yan), Tel: (86)512-65112637, Fax: (86)512-65222691

In many stochastic models, the assumption that random variables are independent is not plausible. Increases in some random variables are often related to decreases in other random variables, so an assumption of dependence is more appropriate than an assumption of independence. One of the important dependence structure is the extended negatively dependent structure, which was introduced by Liu [18] as follows.

**Definition 1.1** *Random variables  $X_k, k = 1, \dots, n$  are said to be lower extended negatively dependent (LEND) if there is some  $M > 0$  such that, for all real numbers  $x_k, k = 1, \dots, n$ ,*

$$P \left\{ \bigcap_{k=1}^n (X_k \leq x_k) \right\} \leq M \prod_{k=1}^n P\{X_k \leq x_k\}; \quad (1.1)$$

*they are said to be upper extended negatively dependent (UEND) if there is some  $M > 0$  such that, for all real numbers  $x_k, k = 1, \dots, n$ ,*

$$P \left\{ \bigcap_{k=1}^n (X_k > x_k) \right\} \leq M \prod_{k=1}^n P\{X_k > x_k\}; \quad (1.2)$$

*and they are said to be extended negatively dependent (END) if they are both LEND and UEND. A sequence of infinitely many random variables  $\{X_k, k = 1, 2, \dots\}$  is said to be LEND/UEND/END if there is some  $M > 0$  such that, for each positive integer  $n$ , the random variables  $X_1, X_2, \dots, X_n$  are LEND/UEND/END, respectively.*

In the case  $M = 1$ , the formula of END random variables reduces to the notion of negatively orthant dependent (NOD, in short) random variables which was introduced by Joag-Dev and Proschan [12]. They also pointed out that negatively associated (NA, in short) random variables are NOD random variables and then END.

As pointed out in Liu [18], the END structure covers many negative dependence structures and, more interestingly, it covers certain positive dependence structures. Hence, studying the limiting behavior of END random variables is of great significance. There are more and more literatures appeared. See, for example, Liu [18] obtained the precise large deviations for dependent random variables with heavy tails. Shen [25] presented some probability inequalities and gave some applications. Wu et al [33] considered some complete moment convergence and mean convergence theorems for the partial sums of END random variables. Qiu et al [24], Wang et al [31, 32] and Shen et al [27] investigated some results on complete convergence of END random variables. Chen et al [6], Yan [36, 37] considered the SLLN for END random variables, and so on.

Another group of dependencies is formed by mixing type structures defined by special sequences of mixing coefficients. Some of them, however defined in a way that is significantly different from the negative dependence structures, have many similar properties which allow us to use in consideration methods and tools similar to those used in END case. In further consideration we will deal with three types of mixing dependencies:  $\varrho^*$ -mixing,  $\varphi$ -mixing and  $\varrho^-$ -mixing.

**Definition 1.2** A sequence of random variable  $\{X_n, n \geq 1\}$  is said to be a  $\varrho^*$ -mixing sequence if there exists  $k \in \mathbb{N}$  such that

$$\varrho^*(k) = \sup_{S, T} \left( \sup_{X \in L^2(\mathcal{F}_S), Y \in L^2(\mathcal{F}_T)} \frac{\text{cov}(X, Y)}{\sqrt{\text{Var}(X) \cdot \text{Var}(Y)}} \right) < 1,$$

where  $S, T$  are the finite subsets of positive integers such that  $\text{dist}(S, T) \geq k$  and  $\mathcal{F}_W$  is the  $\sigma$ -field generated by the random variable  $\{X_i, i \in W \subset \mathbb{N}\}$ .

Bradley [2] and Miller [19] studied various limit properties of random fields under the assumption  $\varrho^*(k) \rightarrow 0, k \rightarrow \infty$ . We refer to the results obtained under the condition  $\varrho^*(k) < 1$  for some  $k \in \mathbb{N}$  which is important in estimating the moments of partial and maxima of partial sums, see Bryc and Smolenski [3] and Peligrad [20]. Peligrad [20], Utev and Peligrad [29] investigated the properties of the maximum of partial sums and used them to obtain an invariance principle, Peligrad and Gut [21] presented Rosenthal-type maximal inequality and rate convergence for the Marcinkiewicz-Zygmund type SLLN, Cai [4] obtained SLLN and complete convergence for random variables with different distributions.

**Definition 1.3** A sequence of random variables  $\{X_n, n \geq 1\}$  is called to be  $\varphi$ -mixing (or uniformly strong mixing) if

$$\varphi(n) = \sup_{k \geq 1, A \in \mathcal{F}_1^k, P(A) > 0, B \in \mathcal{F}_{k+n}^\infty} |P(B|A) - P(B)| \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

where  $\mathcal{F}_n^m$  is the  $\sigma$ -field generated by random variables  $X_n, X_{n+1}, \dots, X_m$ .

A concept of  $\varphi$ -mixing dependence was introduced independently by Rozanov and Volkonski [23] and Ibragimov [11]. A number of limit theorems for  $\varphi$ -mixing random variables have been established by many authors. We refer to Wang *at al* [30] (Rosenthal type maximal inequality, Hájek-Rényi type inequality, SLLN), Tuyen [28] (SLLN), and Chen *at al* [5] (complete convergence and Marcinkiewicz-Zygmund type SLLN of moving averages processes) and Kuczmaszewska [16] (complete convergence for NA,  $\varrho^*$ -mixing and  $\varphi$ -mixing sequences satisfying Petrov's condition).

**Definition 1.4** A sequence  $\{X_n, n \geq 1\}$  is called  $\varrho^-$ -mixing, if

$$\varrho^-(n) = \sup\{\varrho^-(S, T) : S, T \subset \mathbb{N}, \text{dist}(S, T) \geq n\} \rightarrow 0, \text{ as } n \rightarrow \infty,$$

where

$$\varrho^-(S, T) = 0 \vee \sup\{\text{corr}(f(X_i, i \in S), g(X_j, j \in T))\},$$

and the supremum is taken over all coordinatewise increasing real functions  $f$  on  $R^S$  and  $g$  on  $R^T$ .

Some results concerning the complete moment convergence, the complete convergence and strong law of large numbers of Marcinkiewicz-Zygmund type for moving average process generated by  $\varrho^-$ -mixing sequences one can find in Zhang [38]. We also refer to Wang and Lu [34].

Most of results concerning limit theorems are formulated for identically distributed random variables. Pruss [22] introduced the following concept of regular cover which allowed to consider sequences without identical distribution.

**Definition 1.5** *Let  $X_1, X_2, \dots, X_n$  be random variables, and  $X$  be a random variable possibly defined on a different probability space. Then,  $X_1, X_2, \dots, X_n$  are said to be a regular cover of (the distribution of)  $X$  provided we have*

$$E(G(X)) = \frac{1}{n} \sum_{k=1}^n E(G(X_k)), \quad (1.3)$$

for any measurable function  $G$  for which both sides make sense.

In this paper, we are interested in generalizations of the Baum-Katz result. Under some cover condition weaker than (1.3), Kuczmaszewska [14] extended the result to the case of negatively associated (NA, in short) sequence. They got the following result

**Theorem 1.2** (Kuczmaszewska [14]). *Let  $\{X_n, n \geq 1\}$  be a sequence of NA random variables and  $X$  be a random variable possibly defined on a different space satisfying the condition*

$$\frac{1}{n} \sum_{k=1}^n P(|X_k| > x) = c \cdot P(|X| > x) \quad (1.4)$$

for all  $x > 0$ , all  $n \geq 1$  and some positive constant  $c$ . Let  $\alpha p > 1$  and  $\alpha > 1/2$ . Moreover, additionally assume that for  $p \geq 1$   $EX_n = 0$  for all  $n \geq 1$ . Then the following statements are equivalent:

- (i)  $E|X|^p < \infty$ ;
- (ii)  $\sum_{n=1}^{\infty} n^{\alpha p - 2} P(\max_{1 \leq i \leq n} |S_i| > \varepsilon n^\alpha) < \infty$  for all  $\varepsilon > 0$ .

Though condition (1.4) is weak in some sense, it remains a strong condition, we even get the same result for arbitrary random variables with some rough conditions. Gut [9] introduced the following concept of weakly mean dominated

**Definition 1.6** *We say that the array  $\{X_{nk}, 1 \leq k \leq n, n \geq 1\}$  is weakly mean dominated (WMD, in short) by the random variable  $X$  if, for some  $\gamma > 0$ ,*

$$\frac{1}{n} \sum_{k=1}^n P(|X_{nk}| > x) \leq \gamma P(|X| > x), \quad (1.5)$$

for all  $x > 0$  and all  $n$ .

A. Kuczmaszewska and Z. A. Lagodowski [15] introduced another structure which can also be used to prove results for non-identically distributed random variables.

**Definition 1.7** Random variables  $\{X_k, k \geq 1\}$  are weakly mean bounded (WMB, in short) by random variable  $X$  (possibly defined on a different probability space) iff there exist some constant  $\gamma_1, \gamma_2 > 0$  such that for all  $x > 0$  and  $n \geq 1$

$$\gamma_1 \cdot P(|X| > x) \leq \frac{1}{n} \sum_{k=1}^n P(|X_k| > x) \leq \gamma_2 \cdot P(|X| > x). \quad (1.6)$$

Obviously, if a sequence  $\{X_k, k \geq 1\}$  and a random variable  $X$  satisfy WMB condition, they must satisfy WMD ones. The aim of this paper is to consider the analogous generalization of the Baum-Katz theorem for a sequence of random variables satisfying WMD or WMB sense and some usual conditions (Marcinkiewicz-Zygmund type inequality and Rosenthal type inequality). The main results are provided in Section 2. Some lemmas and the proofs of the main results are presented in Section 3.

As usual, we note that  $C$  will be numerical constants whose value are without importance, and, in addition, may change between appearances.  $I(A)$  is the indicator function on the set  $A$ . Denote  $X^+ = \max(X, 0)$  and  $X^- = \max(-X, 0)$ .

## 2 Main Results

Before presenting our main results, we first give the following assumptions.

**Hypothesis.** Let  $\{X_n, n \geq 1\}$  be an arbitrary sequence of random variables satisfying for every  $\theta \geq 2$  and  $n \geq 1$

$$E \left( \left| \sum_{i=1}^n f_i(X_i) \right| \right)^\theta \leq C \left[ \sum_{i=1}^n E|f_i(X_i)|^\theta + \left( \sum_{i=1}^n E|f_i(X_i)|^2 \right)^{\theta/2} \right] \quad (2.1)$$

and

$$E \left( \max_{1 \leq k \leq n} \left| \sum_{i=1}^k f_i(X_i) \right| \right)^\theta \leq C \log^\theta n \left[ \sum_{i=1}^n E|f_i(X_i)|^\theta + \left( \sum_{i=1}^n E|f_i(X_i)|^2 \right)^{\theta/2} \right], \quad (2.2)$$

whenever  $f_1, f_2, \dots, f_n$  are all nondecreasing (or non-increasing) functions,  $E f_i(X_i) = 0$  and  $E|f_i(X_i)|^\theta < \infty$ , for all  $1 \leq i \leq n$ .

**Remark 2.1** A lot of dependent structures, for example such as  $\rho^*$ -mixing,  $\varphi$ -mixing, NA, ND, END, etc., satisfy (2.1) and (2.2) in Hypothesis.

**Theorem 2.1** Suppose  $\alpha p > 1$ ,  $\alpha > 1/2$ . Let  $\{X_n, n \geq 1\}$  be an arbitrary sequence of random variables with  $EX_n = 0$  for all  $n \geq 1$  if  $p > 1$  and  $X$  be a random variable possibly defined on a different probability space satisfying (1.5) for all  $x > 0$ , all  $n \geq 1$  and some positive constant  $\gamma$ . Assume that  $\{X_n, n \geq 1\}$  satisfies the conditions of Hypothesis. Then  $E|X|^p < \infty$  implies that for all  $\varepsilon > 0$

$$\sum_{n=1}^{\infty} n^{\alpha p - 2} P \left( \max_{1 \leq k \leq n} |S_k| > \varepsilon n^\alpha \right) < \infty, \quad (2.3)$$

and

$$\sum_{n=1}^{\infty} n^{\alpha p-2} P(\sup_{i \geq n} i^{-\alpha} |S_i| > \varepsilon) < \infty, \quad (2.4)$$

where  $S_k = \sum_{i=1}^k X_i$ ,  $1 \leq k \leq n$ .

The next theorem presents the necessary condition for (2.3) under assumption that random variables  $\{X_n, n \geq 1\}$  and  $X$  satisfy WMB condition (1.6).

**Example 2.1** We give an example of (1.6). Suppose that  $P(X_k = 1 - \frac{1}{k}) = P(X_k = 2 - \frac{1}{k}) = 1/2$  for  $k = 1, 2, \dots$ . Then

$$\frac{1}{n} \sum_{k=1}^n P(|X_k| > x) \rightarrow \begin{cases} 1, & 0 \leq x < 1, \\ 1/2, & 1 \leq x < 2, \\ 0, & x \geq 2. \end{cases}$$

that is, if the mean dominating random variable,  $X$ , is such that  $P(X = 1) = P(X = 2) = 1/2$ , then (1.6) is satisfied.

**Theorem 2.2** Suppose  $\alpha p > 1$ ,  $\alpha > 1/2$ . Let  $\{X_n, n \geq 1\}$  be a sequence of random variables satisfying (2.1) for  $\theta = 2$  and  $X$  be a random variable possibly defined on a different probability space satisfying (1.6) for all  $x > 0$ , all  $n \geq 1$  and some positive constants  $\gamma_1$  and  $\gamma_2$ . Then (2.3) implies  $E|X|^p < \infty$ .

As a consequence of Theorem 2.1 and Theorem 2.2 by Lemma 3.2 and Lemma 3.3 we get the following result.

**Corollary 2.1** Suppose  $\alpha p > 1$ ,  $\alpha > 1/2$ . Let  $\{X_n, n \geq 1\}$  be a sequence of END,  $\varrho^*$ -mixing,  $\varrho^-$ -mixing or  $\varphi$ -mixing random variables and  $X$  be a random variable possibly defined on a different probability space satisfying (1.6) for all  $x > 0$ , all  $n \geq 1$  and some positive constants  $\gamma_1$  and  $\gamma_2$ . Moreover, we assume  $EX_n = 0$  for all  $n \geq 1$  if  $p > 1$  and  $\sum_{n=1}^{\infty} \varphi^{\frac{1}{2}}(n) < \infty$  in case of  $\varphi$ -mixing sequence. Then the following statements are equivalent:

- (i)  $E|X|^p < \infty$ ;
- (ii)  $\sum_{n=1}^{\infty} n^{\alpha p-2} P(\max_{1 \leq i \leq n} |S_i| > \varepsilon n^{\alpha}) < \infty$  for all  $\varepsilon > 0$ .

**Remark 2.2** Since END and  $\varrho^-$ -mixing random variables include NA random variables, our result also holds for NA case.

### 3 Some Lemmas and Proofs

To prove the main results of the paper, we need the following important lemmas.

**Lemma 3.1** (cf. Liu [18]) Let  $\{X_n, n \geq 1\}$  be a sequence of END random variables. For each  $n \geq 1$ , if  $f_1, f_2, \dots, f_n$  are all nondecreasing ( or nonincreasing ) functions, then random variables  $f_1(X_1), f_2(X_2), \dots, f_n(X_n)$  are also END.



**Lemma 3.2** (cf. Wang et al.[31]) Let  $p \geq 2$  and  $\{X_n, n \geq 1\}$  be a sequence of END random variables with  $EX_n = 0$  and  $E|X_n|^p < \infty$  for each  $n \geq 1$ . Then there exists a positive constant  $C_p$  depending only on  $p$  such that

$$E \left| \sum_{i=1}^n X_i \right|^p \leq C_p \left[ \sum_{i=1}^n E|X_i|^p + \left( \sum_{i=1}^n E|X_i|^2 \right)^{p/2} \right], \quad \text{for all } n \geq 1.$$

and

$$E \left( \max_{1 \leq j \leq n} \left| \sum_{i=1}^j X_i \right|^p \right) \leq C_p (\log n)^p \left[ \sum_{i=1}^n E|X_i|^p + \left( \sum_{i=1}^n E|X_i|^2 \right)^{p/2} \right], \quad \text{for all } n \geq 1.$$

**Lemma 3.3** (cf. Utev and Peligrad[29], Wang and Lu[34], Wang et al.[30]) Let  $p \geq 2$  and  $\{X_n, n \geq 1\}$  be a sequence of  $\varrho^*$ -mixing,  $\varrho^-$ -mixing or  $\varphi$ -mixing random variables with  $EX_n = 0$  and  $E|X_n|^p < \infty$  for each  $n \geq 1$ . Moreover, if  $X_n, n \geq 1$  are  $\varphi$ -mixing we assume that  $\sum_{n=1}^{\infty} \varphi^{\frac{1}{2}}(n) < \infty$ . Then there exists a positive constant  $C_p$  depending only on  $p$  such that

$$E \left( \max_{1 \leq j \leq n} \left| \sum_{i=1}^j X_i \right|^p \right) \leq C_p \left[ \sum_{i=1}^n E|X_i|^p + \left( \sum_{i=1}^n E|X_i|^2 \right)^{p/2} \right], \quad \text{for all } n \geq 1. \quad (3.1)$$

**Lemma 3.4** (cf. Gut [9]) Let  $\{X_n, n \geq 1\}$  be a sequence of random variables satisfying a weak dominating condition with mean dominating random variable  $X$ , i.e. for some  $c > 0$

$$\frac{1}{n} \sum_{i=1}^n P(|X_i| > x) \leq cP(|X| > x).$$

Let  $r > 0$  and for some  $A > 0$

$$\begin{aligned} X'_i &= X_i I(|X_i| \leq A), \quad X''_i = X_i I(|X_i| > A), \\ X_i^* &= X_i I(|X_i| \leq A) - AI(X_i < -A) + AI(X_i > A), \end{aligned}$$

and

$$\begin{aligned} X' &= XI(|X| \leq A), \quad X'' = XI(|X| > A), \\ X^* &= XI(|X| \leq A) - AI(X < -A) + AI(X > A). \end{aligned}$$

Then

- (i) if  $E|X|^r < \infty$ , then  $\frac{1}{n} \sum_{i=1}^n E|X_i|^r \leq CE|X|^r$ ;
- (ii)  $\frac{1}{n} \sum_{i=1}^n E|X'_i|^r \leq C(E|X'|^r + A^r P(|X| > A))$  for any  $A > 0$ ;
- (iii)  $\frac{1}{n} \sum_{i=1}^n E|X''_i|^r \leq CE|X''|^r$  for any  $A > 0$ ;
- (iv)  $\frac{1}{n} \sum_{i=1}^n E|X_i^*|^r \leq CE|X^*|^r$  for any  $A > 0$ .

Now, we present the proofs of the main results step by step.

**Proof of Theorem 2.1.** We first take

$$0 < p' < p, \quad \frac{1}{\alpha p} < q < 1$$

such that

$$\alpha(p - p') > \alpha(p - p')q > 1, \quad \text{and} \quad p - p' > 1 \text{ if } p > 1.$$

For all  $1 \leq i \leq n, n \geq 1$ , denote that

$$\begin{aligned} X_{ni}^{(1)} &= -n^{\alpha q} I(X_i < -n^{\alpha q}) + X_i I(|X_i| \leq n^{\alpha q}) + n^{\alpha q} I(X_i > n^{\alpha q}); \\ X_{ni}^{(2)} &= (X_i - n^{\alpha q}) I(X_i > n^{\alpha q}); \\ X_{ni}^{(3)} &= -(X_i + n^{\alpha q}) I(X_i < -n^{\alpha q}). \end{aligned}$$

Then, for every  $1 \leq i \leq n, n \geq 1$

$$X_i = X_{ni}^{(1)} + X_{ni}^{(2)} - X_{ni}^{(3)} \quad \text{and} \quad X_{ni}^{(2)} \geq 0, X_{ni}^{(3)} \geq 0.$$

Thus,

$$\begin{aligned} & \sum_{n=1}^{\infty} n^{\alpha p-2} P \left( \max_{1 \leq k \leq n} |S_k| > \varepsilon n^{\alpha} \right) \\ & \leq \sum_{n=1}^{\infty} n^{\alpha p-2} P \left( \max_{1 \leq k \leq n} \left| \sum_{i=1}^k X_{ni}^{(1)} \right| > \varepsilon n^{\alpha}/3 \right) + \sum_{n=1}^{\infty} n^{\alpha p-2} P \left( \max_{1 \leq k \leq n} \left| \sum_{i=1}^k X_{ni}^{(2)} \right| > \varepsilon n^{\alpha}/3 \right) \\ & \quad + \sum_{n=1}^{\infty} n^{\alpha p-2} P \left( \max_{1 \leq k \leq n} \left| \sum_{i=1}^k X_{ni}^{(3)} \right| > \varepsilon n^{\alpha}/3 \right) \\ & = \sum_{n=1}^{\infty} n^{\alpha p-2} P \left( \max_{1 \leq k \leq n} \left| \sum_{i=1}^k X_{ni}^{(1)} \right| > \varepsilon n^{\alpha}/3 \right) + \sum_{n=1}^{\infty} n^{\alpha p-2} P \left( \sum_{i=1}^n X_{ni}^{(2)} > \varepsilon n^{\alpha}/3 \right) \\ & \quad + \sum_{n=1}^{\infty} n^{\alpha p-2} P \left( \sum_{i=1}^n X_{ni}^{(3)} > \varepsilon n^{\alpha}/3 \right) \\ & \triangleq I_1 + I_2 + I_3. \quad (\text{in say}) \end{aligned}$$

To prove (2.3), it suffices to show that  $I_k < \infty, k = 1, 2, 3$ .

For  $I_1$ , we first prove that

$$n^{-\alpha} \max_{1 \leq k \leq n} \left| \sum_{i=1}^k E X_{ni}^{(1)} \right| \rightarrow 0, \quad n \rightarrow \infty. \quad (3.2)$$

We will do it in three cases.

**Case I.** Let  $\alpha \leq 1$ . Then  $\alpha p > 1$  implies  $p > 1$  and, according to the assumption,

$EX_n = 0, n \geq 1$ . It follows from Lemma 3.4 that

$$\begin{aligned}
& n^{-\alpha} \max_{1 \leq k \leq n} \left| \sum_{i=1}^k EX_{ni}^{(1)} \right| \\
& \leq n^{-\alpha} \max_{1 \leq k \leq n} \left| \sum_{i=1}^k [EX_i I(|X_i| \leq n^{\alpha q}) + n^{\alpha q} P(|X_i| > n^{\alpha q})] \right| \\
& = n^{-\alpha} \max_{1 \leq k \leq n} \left| \sum_{i=1}^k [EX_i I(|X_i| > n^{\alpha q}) + n^{\alpha q} P(|X_i| > n^{\alpha q})] \right| \\
& \leq 2n^{-\alpha} \sum_{i=1}^n E|X_i| I(|X_i| > n^{\alpha q}) \leq 2n^{1-\alpha} E|X| I(|X| > n^{\alpha q}) \\
& \leq Cn^{1-\alpha} \cdot n^{\alpha q(1-(p-p'))} E|X|^{p-p'} \leq Cn^{-[\alpha q(p-p')-1]-\alpha(1-q)} \rightarrow 0, \quad n \rightarrow \infty.
\end{aligned}$$

**Case II.** Let  $\alpha > 1$  and  $p > 1$ . We have

$$\begin{aligned}
& n^{-\alpha} \max_{1 \leq k \leq n} \left| \sum_{i=1}^k EX_{ni}^{(1)} \right| \leq n^{-\alpha} \sum_{i=1}^n [E|X_i| I(|X_i| \leq n^{\alpha q}) + n^{\alpha q} P(|X_i| > n^{\alpha q})] \\
& \leq Cn^{1-\alpha} [E|X| I(|X| \leq n^{\alpha q}) + n^{\alpha q} P(|X| > n^{\alpha q})] \leq Cn^{1-\alpha} \rightarrow 0, \quad n \rightarrow \infty.
\end{aligned}$$

**Case III.** Let  $\alpha > 1$  and  $p \leq 1$ . We have

$$\begin{aligned}
& n^{-\alpha} \max_{1 \leq k \leq n} \left| \sum_{i=1}^k EX_{ni}^{(1)} \right| \leq n^{-\alpha} \sum_{i=1}^n [E|X_i| I(|X_i| \leq n^{\alpha q}) + n^{\alpha q} P(|X_i| > n^{\alpha q})] \\
& \leq Cn^{1-\alpha} [E|X| I(|X| \leq n^{\alpha q}) + n^{\alpha q} P(|X| > n^{\alpha q})] \\
& \leq Cn^{1-\alpha} \cdot n^{\alpha q(1-(p-p'))} E|X|^{p-p'} \leq Cn^{-[\alpha q(p-p')-1]-\alpha(1-q)} \rightarrow 0, \quad n \rightarrow \infty.
\end{aligned}$$

By (3.2), to prove  $I_1 < \infty$ , we prove only that

$$I_1^* \triangleq \sum_{n=1}^{\infty} n^{\alpha p-2} P \left( \max_{1 \leq k \leq n} \left| \sum_{i=1}^k (X_{ni}^{(1)} - EX_{ni}^{(1)}) \right| > \varepsilon n^{\alpha}/6 \right) < \infty. \quad (3.3)$$

From Hypothesis, for each  $n \geq 1$ ,  $\{X_{ni}^{(1)} - EX_{ni}^{(1)}, 1 \leq i \leq n\}$  remain satisfy the inequalities in Hypothesis. By  $\alpha q(p-p') > 1$  and  $0 < q < 1$ , we have for  $p \leq 2$

$$\alpha - \frac{1}{2} - \alpha q \left( 1 - \frac{p-p'}{2} \right) > \alpha - \frac{1}{2} - \alpha \left( 1 - \frac{p-p'}{2} \right) = \frac{\alpha(p-p')-1}{2} > 0.$$

By taking

$$\tau > \max \left\{ 2, p, \frac{p-(p-p')q}{1-q}, \frac{\alpha p-1}{\alpha-\frac{1}{2}}, \frac{\alpha p-1}{\alpha-\frac{1}{2}-\alpha q \left( 1 - \frac{p-p'}{2} \right)} \right\},$$

Chebyshev's inequality and Hypothesis we get

$$\begin{aligned}
I_1^* &\leq C \sum_{n=1}^{\infty} n^{\alpha p-2-\alpha\tau} E \left( \max_{1 \leq k \leq n} \left| \sum_{i=1}^k (X_{ni}^{(1)} - EX_{ni}^{(1)}) \right| \right)^{\tau} \\
&\leq C \sum_{n=1}^{\infty} n^{\alpha p-2-\alpha\tau} \log^{\tau} n \sum_{i=1}^n E \left| X_{ni}^{(1)} \right|^{\tau} + C \sum_{n=1}^{\infty} n^{\alpha p-2-\alpha\tau} \log^{\tau} n \left( \sum_{i=1}^n E \left| X_{ni}^{(1)} \right|^2 \right)^{\tau/2} \\
&\triangleq I_{11}^* + I_{12}^*. \text{ (in say)}
\end{aligned}$$

Again by Hypothesis

$$\begin{aligned}
I_{11}^* &\leq C \sum_{n=1}^{\infty} n^{\alpha p-2-\alpha\tau} \log^{\tau} n \sum_{i=1}^n [E|X_i|^{\tau} I(|X_i| \leq n^{\alpha q}) + n^{\alpha q\tau} P(|X_i| > n^{\alpha q})] \\
&\leq C \sum_{n=1}^{\infty} n^{\alpha p-1-\alpha\tau} \log^{\tau} n \cdot n^{\alpha q(\tau-(p-p'))} E|X|^{p-p'} \\
&\leq C \sum_{n=1}^{\infty} n^{-\alpha(1-q)\left(\tau-\frac{p-q(p-p')}{1-q}\right)-1} \log^{\tau} n < \infty,
\end{aligned}$$

and

$$I_{12}^* \leq C \sum_{n=1}^{\infty} n^{\alpha p-2-\alpha\tau} \log^{\tau} n \left\{ \sum_{i=1}^n [E|X_i|^2 I(|X_i| \leq n^{\alpha q}) + n^{2\alpha q} P(|X_i| > n^{\alpha q})] \right\}^{\tau/2}.$$

Now, we prove  $I_{12}^* < \infty$  in two cases:  $p > 2$  and  $0 < p \leq 2$ .

Let  $p > 2$ .

$$I_{12}^* \leq C \sum_{n=1}^{\infty} n^{\alpha p-2-\alpha\tau} \log^{\tau} n \cdot n^{\tau/2} (EX^2)^{\tau/2} \leq C \sum_{n=1}^{\infty} n^{-(\alpha-\frac{1}{2})\left(\tau-\frac{\alpha p-1}{\alpha-\frac{1}{2}}\right)-1} \log^{\tau} n < \infty.$$

For  $0 < p \leq 2$  we have

$$\begin{aligned}
I_{12}^* &\leq C \sum_{n=1}^{\infty} n^{\alpha p-2-\alpha\tau} \log^{\tau} n \cdot \left( n^{\alpha q(2-(p-p'))} \cdot n \right)^{\tau/2} \left( E|X|^{p-p'} \right)^{\tau/2} \\
&\leq C \sum_{n=1}^{\infty} n^{\alpha p-2-\tau\left[\alpha-\frac{1}{2}-\alpha q\left(1-\frac{p-p'}{2}\right)\right]} \log^{\tau} n < \infty.
\end{aligned}$$

This ends the proof of  $I_1 < \infty$ . Next for  $I_2 < \infty$  and each  $1 \leq i \leq n, n \geq 1$ , let

$$Y_{ni}^{(2)} = (X_i - n^{\alpha q})I(n^{\alpha q} < X_i \leq n^{\alpha q} + n^{\alpha}) + n^{\alpha}I(X_i > n^{\alpha q} + n^{\alpha}).$$

Then  $Y_{ni}^{(2)} \geq 0$  and

$$X_{ni}^{(2)} = Y_{ni}^{(2)} + (X_i - n^{\alpha q} - n^{\alpha})I(X_i > n^{\alpha q} + n^{\alpha}).$$

Thus,

$$\begin{aligned}
I_2 &\leq \sum_{n=1}^{\infty} n^{\alpha p-2} P\left(\sum_{i=1}^n Y_{ni}^{(2)} > \frac{\varepsilon n^{\alpha}}{6}\right) \\
&\quad + \sum_{n=1}^{\infty} n^{\alpha p-2} P\left(\sum_{i=1}^n (X_i - n^{\alpha q} - n^{\alpha}) I(X_i > n^{\alpha q} + n^{\alpha}) > \frac{\varepsilon n^{\alpha}}{6}\right) \\
&\leq \sum_{n=1}^{\infty} n^{\alpha p-2} P\left(\sum_{i=1}^n Y_{ni}^{(2)} > \frac{\varepsilon n^{\alpha}}{6}\right) + \sum_{n=1}^{\infty} n^{\alpha p-2} \sum_{i=1}^n P(X_i > n^{\alpha q} + n^{\alpha}) \\
&\triangleq I_{21} + I_{22}. \text{ (in say)}
\end{aligned}$$

By Lemma 3.4,

$$\begin{aligned}
I_{22} &\leq \sum_{n=1}^{\infty} n^{\alpha p-2} \sum_{i=1}^n P(|X_i| > n^{\alpha}) \leq C \sum_{n=1}^{\infty} n^{\alpha p-1} P(|X| > n^{\alpha}) \\
&= C \sum_{n=1}^{\infty} n^{\alpha p-1} \sum_{i=n}^{\infty} P(i^{\alpha} < |X| \leq (i+1)^{\alpha}) \\
&= C \sum_{i=1}^{\infty} P(i^{\alpha} < |X| \leq (i+1)^{\alpha}) \sum_{n=1}^i n^{\alpha p-1} \\
&\leq C \sum_{i=1}^{\infty} i^{\alpha p} P(i^{\alpha} < |X| \leq (i+1)^{\alpha}) \leq CE|X|^p < \infty.
\end{aligned} \tag{3.4}$$

Next we prove only that  $I_{21} < \infty$ . We first show that

$$n^{-\alpha} \sum_{i=1}^n EY_{ni}^{(2)} \rightarrow 0, \quad n \rightarrow \infty. \tag{3.5}$$

If  $p > 1$ , then by Lemma 3.4

$$\begin{aligned}
0 &\leq n^{-\alpha} \sum_{i=1}^n EY_{ni}^{(2)} \leq n^{-\alpha} \sum_{i=1}^n EX_i I(X_i > n^{\alpha q}) \\
&\leq n^{1-\alpha} EXI(X > n^{\alpha q}) \leq n^{1-\alpha} \cdot n^{\alpha q(1-(p-p'))} E|X|^{p-p'} \\
&\leq Cn^{-[\alpha q(p-p')-1]-\alpha(1-q)} \rightarrow 0, \quad n \rightarrow \infty.
\end{aligned}$$

If  $0 < p \leq 1$ , then by Lemma 3.4

$$\begin{aligned}
0 &\leq n^{-\alpha} \sum_{i=1}^n EY_{ni}^{(2)} \leq n^{-\alpha} \sum_{i=1}^n [EX_i I(|X_i| \leq 2n^{\alpha}) + n^{\alpha} P(|X_i| > 2n^{\alpha})] \\
&\leq Cn^{1-\alpha} [EXI(|X| \leq 2n^{\alpha}) + 2n^{\alpha} P(|X| > 2n^{\alpha}) + n^{\alpha} P(|X| > 2n^{\alpha})] \\
&\leq Cn^{1-\alpha} \left[ n^{\alpha(1-(p-p'))} + n^{\alpha-\alpha q(p-p')} \right] \\
&\leq Cn^{-[\alpha q(p-p')-1]} \rightarrow 0, \quad n \rightarrow \infty.
\end{aligned}$$

By (3.5), to prove  $I_{21} < \infty$ , it is sufficient to show that

$$I_{21}^* = \sum_{n=1}^{\infty} n^{\alpha p-2} P \left( \left| \sum_{i=1}^n \left( Y_{ni}^{(2)} - EY_{ni}^{(2)} \right) \right| > \frac{\varepsilon n^{\alpha}}{12} \right) < \infty.$$

We will prove it in two cases:  $0 < p < 2$  and  $p \geq 2$ .

For  $0 < p < 2$ , by Chebyshev's inequality, Hypothesis, Lemma 3.4 and (3.4) we have

$$\begin{aligned} I_{21}^* &\leq C \sum_{n=1}^{\infty} n^{\alpha p-2-2\alpha} E \left| \sum_{i=1}^n \left( Y_{ni}^{(2)} - EY_{ni}^{(2)} \right) \right|^2 \leq C \sum_{n=1}^{\infty} n^{\alpha p-2-2\alpha} \sum_{i=1}^n E \left| Y_{ni}^{(2)} \right|^2 \\ &\leq C \sum_{n=1}^{\infty} n^{\alpha p-2-2\alpha} \sum_{i=1}^n [E|X_i|^2 I(n^{\alpha q} < X_i \leq n^{\alpha q} + n^{\alpha}) + n^{2\alpha} P(|X_i| > n^{\alpha q} + n^{\alpha})] \\ &\leq C \sum_{n=1}^{\infty} n^{\alpha p-2-2\alpha} \sum_{i=1}^n E|X_i|^2 I(|X_i| \leq 2n^{\alpha}) + C \sum_{n=1}^{\infty} n^{\alpha p-2} \sum_{i=1}^n P(|X_i| > n^{\alpha}) \\ &\leq C \sum_{n=1}^{\infty} n^{\alpha p-1-2\alpha} E|X|^2 I(|X| \leq 2n^{\alpha}) + C \sum_{n=1}^{\infty} n^{\alpha p-1} P(|X| > n^{\alpha}) \\ &\leq C \sum_{n=1}^{\infty} n^{\alpha p-1-2\alpha} \sum_{i=1}^n E|X|^2 I(2(i-1)^{\alpha} < |X| \leq 2i^{\alpha}) + CE|X|^p \\ &= C \sum_{i=1}^{\infty} E|X|^2 I(2(i-1)^{\alpha} < |X| \leq 2i^{\alpha}) \sum_{n=i}^{\infty} n^{\alpha p-1-2\alpha} + CE|X|^p \\ &\leq C \sum_{i=1}^{\infty} i^{\alpha p-2\alpha} E|X|^2 I(2(i-1)^{\alpha} < |X| \leq 2i^{\alpha}) + CE|X|^p \\ &\leq CE|X|^p < \infty. \end{aligned}$$

Let  $p \geq 2$ . By taking

$$\kappa > \max \left\{ p, \frac{\alpha p - 1}{\alpha - \frac{1}{2}}, \frac{2(\alpha p - 1)}{\alpha(p - p') - 1} \right\}$$

Chebyshev's inequality and Hypothesis we have

$$\begin{aligned} I_{21}^* &\leq C \sum_{n=1}^{\infty} n^{\alpha p-2-\alpha\kappa} E \left| \sum_{i=1}^n \left( Y_{ni}^{(2)} - EY_{ni}^{(2)} \right) \right|^{\kappa} \\ &\leq C \sum_{n=1}^{\infty} n^{\alpha p-2-\alpha\kappa} \sum_{i=1}^n E \left| Y_{ni}^{(2)} \right|^{\kappa} + C \sum_{n=1}^{\infty} n^{\alpha p-2-\alpha\kappa} \left( \sum_{i=1}^n E \left| Y_{ni}^{(2)} \right|^2 \right)^{\kappa/2} \\ &\triangleq I_{211}^* + I_{212}^* \text{ (in say)} \end{aligned}$$

It is easy to see that

$$I_{212}^* \leq C \sum_{n=1}^{\infty} n^{\alpha p-2-\alpha\kappa} \left( \sum_{i=1}^n EX_i^2 \right)^{\kappa/2} \leq C \sum_{n=1}^{\infty} n^{\alpha p-2-\alpha\kappa+\frac{\kappa}{2}} < \infty.$$

On the other hand, by Lemma 3.4 and (3.4) we get

$$\begin{aligned}
I_{211}^* &\leq C \sum_{n=1}^{\infty} n^{\alpha p-2-\alpha\kappa} \sum_{i=1}^n [E|X_i|^\kappa I(n^{\alpha q} < X_i \leq n^\alpha + n^{\alpha q}) + n^{\alpha\kappa} P(|X_i| > n^\alpha + n^{\alpha q})] \\
&\leq C \sum_{n=1}^{\infty} n^{\alpha p-2-\alpha\kappa} \sum_{i=1}^n [E|X_i|^\kappa I(|X_i| \leq 2n^\alpha) + n^{\alpha\kappa} P(|X_i| > n^\alpha)] \\
&\leq C \sum_{n=1}^{\infty} n^{\alpha p-1-\alpha\kappa} [E|X|^\kappa I(|X| \leq 2n^\alpha) + n^{\alpha\kappa} P(|X| > n^\alpha)] \\
&\leq C \sum_{n=1}^{\infty} n^{\alpha p-1-\alpha\kappa} E|X|^\kappa I(|X| \leq 2n^\alpha) + CE|X|^p \\
&= C \sum_{n=1}^{\infty} n^{\alpha p-1-\alpha\kappa} \sum_{i=1}^n E|X|^\kappa I(2(i-1)^\alpha < |X| \leq 2i^\alpha) + CE|X|^p \\
&= C \sum_{i=1}^{\infty} E|X|^\kappa I(2(i-1)^\alpha < |X| \leq 2i^\alpha) \sum_{n=i}^{\infty} n^{\alpha p-1-\alpha\kappa} + CE|X|^p \\
&\leq C \sum_{i=1}^{\infty} n^{\alpha p-\alpha\kappa} E|X|^\kappa I(2(i-1)^\alpha < |X| \leq 2i^\alpha) + CE|X|^p \\
&\leq CE|X|^p < \infty.
\end{aligned}$$

To prove the second thesis of Theorem 2.1 it is enough to show that (2.3) implies (2.4). For  $0 < \varepsilon < 1$  and  $\alpha p > 1$  we have

$$\begin{aligned}
&\sum_{n=1}^{\infty} n^{\alpha p-2} P\left(\sup_{k \geq n} k^{-\alpha} |S_k| > \varepsilon\right) \\
&= \sum_{j=1}^{\infty} \sum_{n=2^{j-1}}^{2^j-1} n^{\alpha p-2} P\left(\sup_{k \geq n} k^{-\alpha} |S_k| > \varepsilon\right) \leq C \sum_{j=1}^{\infty} 2^{j(\alpha p-1)} P\left(\sup_{k \geq 2^{j-1}} k^{-\alpha} |S_k| > \varepsilon\right) \\
&\leq C \sum_{j=1}^{\infty} 2^{j(\alpha p-1)} \sum_{i=j}^{\infty} P\left(\max_{2^{i-1} \leq k < 2^i} k^{-\alpha} |S_k| > \varepsilon\right) \\
&\leq C \sum_{i=1}^{\infty} P\left(\max_{2^{i-1} \leq k < 2^i} k^{-\alpha} |S_k| > \varepsilon\right) \sum_{j=1}^i 2^{j(\alpha p-1)} \leq C \sum_{i=1}^{\infty} 2^{i(\alpha p-1)} P\left(\max_{2^{i-1} \leq k < 2^i} k^{-\alpha} |S_k| > \varepsilon\right) \\
&\leq C \sum_{i=1}^{\infty} 2^{i(\alpha p-1)} P\left(\max_{2^{i-1} \leq k < 2^i} |S_k| > \varepsilon 2^{(i-1)\alpha}\right) \leq C \sum_{i=1}^{\infty} 2^{i(\alpha p-1)} P\left(\max_{1 \leq k \leq 2^i} |S_k| > \varepsilon 2^{-\alpha} 2^{i\alpha}\right) \\
&\leq C \sum_{i=1}^{\infty} n^{\alpha p-2} P\left(\max_{1 \leq k \leq n} |S_k| > \varepsilon 2^{-2\alpha} n^\alpha\right) < \infty.
\end{aligned}$$

This ends the proof of Theorem 2.1. □

**Proof of Theorem 2.2.** In order to prove the result, it suffices to show that

$$\sum_{n=1}^{\infty} n^{\alpha p-1} P(|X| > n^{\alpha}) < \infty. \quad (3.6)$$

By (2.3), we first note that for  $0 < \varepsilon < 1$

$$\begin{aligned} & \sum_{n=1}^{\infty} n^{\alpha p-2} P\left(\max_{1 \leq i \leq n} |X_i| > n^{\alpha}\right) \leq \sum_{n=1}^{\infty} n^{\alpha p-2} P\left(\max_{1 \leq i \leq n} |X_i| > \varepsilon n^{\alpha}\right) \\ &= \sum_{n=1}^{\infty} n^{\alpha p-2} P\left(\max_{1 \leq i \leq n} |S_i - S_{i-1}| > \varepsilon n^{\alpha}\right) \quad (S_0 = 0) \\ &\leq 2 \sum_{n=1}^{\infty} n^{\alpha p-2} P\left(\max_{1 \leq i \leq n} |S_i| > \varepsilon n^{\alpha}/2\right) < \infty. \end{aligned} \quad (3.7)$$

Therefore,

$$P\left(\max_{1 \leq i \leq n} |X_i| > n^{\alpha}\right) \rightarrow 0, \quad \text{as } n \rightarrow \infty. \quad (3.8)$$

Moreover, by Lemma ?? and (1.6)

$$\begin{aligned} \gamma_1 \cdot n P(|X| > n^{\alpha}) &\leq \sum_{k=1}^n P(|X_k| > n^{\alpha}) \\ &= \sum_{k=1}^n P\left(|X_k| > n^{\alpha}, \max_{1 \leq i < k} |X_i| > n^{\alpha}\right) + \sum_{k=1}^n P\left(|X_k| > n^{\alpha}, \max_{1 \leq i < k} |X_i| \leq n^{\alpha}\right) \\ &= \sum_{k=1}^n P\left(X_k^+ > n^{\alpha}, \max_{1 \leq i < k} |X_i| > n^{\alpha}\right) + \sum_{k=1}^n P\left(X_k^- > n^{\alpha}, \max_{1 \leq i < k} |X_i| > n^{\alpha}\right) \\ &\quad + P\left(\bigcup_{k=1}^n \left\{|X_k| > n^{\alpha}, \max_{1 \leq i < k} |X_i| \leq n^{\alpha}\right\}\right) \\ &= \sum_{k=1}^n P\left(X_k^+ > n^{\alpha}, \max_{1 \leq i < k} |X_i| > n^{\alpha}\right) + \sum_{k=1}^n P\left(X_k^- > n^{\alpha}, \max_{1 \leq i < k} |X_i| > n^{\alpha}\right) \\ &\quad + P\left(\max_{1 \leq i \leq n} |X_i| > n^{\alpha}\right), \end{aligned} \quad (3.9)$$

since the sets  $\{|X_k| > n^{\alpha}, \max_{1 \leq i < k} |X_i| \leq n^{\alpha}\}, 1 \leq k \leq n$  are disjoint.

By Hypothesis,  $\{X_n^+, n \geq 1\}$  still satisfy the inequality in Hypothesis. It follows from Cauchy-Schwarz inequality and Hypothesis that

$$\begin{aligned} & \sum_{k=1}^n P\left(X_k^+ > n^{\alpha}, \max_{1 \leq i < k} |X_i| > n^{\alpha}\right) = E\left[\sum_{k=1}^n I\left(X_k^+ > n^{\alpha}, \max_{1 \leq i < k} |X_i| > n^{\alpha}\right)\right] \\ &= E\left[\sum_{k=1}^n I\left(X_k^+ > n^{\alpha}\right) I\left(\max_{1 \leq i < k} |X_i| > n^{\alpha}\right)\right] \leq E\left[\sum_{k=1}^n I\left(X_k^+ > n^{\alpha}\right) I\left(\max_{1 \leq i \leq n} |X_i| > n^{\alpha}\right)\right] \end{aligned}$$



$$\begin{aligned}
&= E \left[ \sum_{k=1}^n \{I(X_k^+ > n^\alpha) - P(X_k^+ > n^\alpha)\} I\left(\max_{1 \leq i \leq n} |X_i| > n^\alpha\right) \right] \\
&\quad + E \left[ \sum_{k=1}^n P(X_k^+ > n^\alpha) I\left(\max_{1 \leq i \leq n} |X_i| > n^\alpha\right) \right] \\
&\leq \sqrt{E \left( \sum_{k=1}^n \{I(X_k^+ > n^\alpha) - P(X_k^+ > n^\alpha)\} \right)^2 P\left(\max_{1 \leq i \leq n} |X_i| > n^\alpha\right)} \\
&\quad + P\left(\max_{1 \leq i \leq n} |X_i| > n^\alpha\right) \sum_{k=1}^n P(X_k^+ > n^\alpha) \\
&\leq \sqrt{C_2 \sum_{k=1}^n E \{I(X_k^+ > n^\alpha) - P(X_k^+ > n^\alpha)\}^2 P\left(\max_{1 \leq i \leq n} |X_i| > n^\alpha\right)} \\
&\quad + P\left(\max_{1 \leq i \leq n} |X_i| > n^\alpha\right) \sum_{k=1}^n P(X_k^+ > n^\alpha) \\
&\leq \sqrt{C_2 \sum_{k=1}^n P(X_k^+ > n^\alpha) P\left(\max_{1 \leq i \leq n} |X_i| > n^\alpha\right)} + P\left(\max_{1 \leq i \leq n} |X_i| > n^\alpha\right) \sum_{k=1}^n P(X_k^+ > n^\alpha) \\
&\leq \sqrt{C_2 \gamma_2 n P(|X| > n^\alpha) P\left(\max_{1 \leq i \leq n} |X_i| > n^\alpha\right)} + \gamma_2 P\left(\max_{1 \leq i \leq n} |X_i| > n^\alpha\right) \cdot n P(|X| > n^\alpha) \\
&\leq \frac{\gamma_1 n}{4} P(|X| > n^\alpha) + \frac{\gamma_2 C_2}{\gamma_1} P\left(\max_{1 \leq i \leq n} |X_i| > n^\alpha\right) \\
&\quad + \gamma_2 P\left(\max_{1 \leq i \leq n} |X_i| > n^\alpha\right) \cdot n P(|X| > n^\alpha),
\end{aligned} \tag{3.10}$$

where we used the following inequality

$$\sqrt{ab} \leq \frac{a\gamma_1}{4\gamma_2 C_2} + \frac{\gamma_2 C_2}{\gamma_1} b, \quad a \geq 0, b \geq 0.$$

Similarly, we can have

$$\begin{aligned}
&\sum_{k=1}^n P\left(X_k^- > n^\alpha, \max_{1 \leq i < k} |X_i| > n^\alpha\right) \\
&\leq \frac{\gamma_1 n}{4} P(|X| > n^\alpha) + \frac{\gamma_2 C_2}{\gamma_1} P\left(\max_{1 \leq i \leq n} |X_i| > n^\alpha\right) \\
&\quad + \gamma_2 P\left(\max_{1 \leq i \leq n} |X_i| > n^\alpha\right) \cdot n P(|X| > n^\alpha),
\end{aligned} \tag{3.11}$$

Now we see that (3.9), (3.10) and (3.11) lead to

$$\begin{aligned} \frac{\gamma_1 n}{2} P(|X| > n^\alpha) &\leq \frac{2\gamma_2 C_2 + \gamma_1}{\gamma_1} P\left(\max_{1 \leq i \leq n} |X_i| > n^\alpha\right) \\ &\quad + 2\gamma_2 n P\left(\max_{1 \leq i \leq n} |X_i| > n^\alpha\right) P(|X| > n^\alpha) \end{aligned}$$

By (3.8), for sufficiently large  $n$  we have

$$P\left(\max_{1 \leq i \leq n} |X_i| > n^\alpha\right) < \frac{\gamma_1}{8\gamma_2},$$

and consequently

$$nP(|X| > n^\alpha) \leq \frac{4(2\gamma_2 C_2 + \gamma_1)}{\gamma_1} P\left(\max_{1 \leq i \leq n} |X_i| > n^\alpha\right). \quad (3.12)$$

Relations (3.7) and (3.12) give (3.6) and in conclusion we get the desired condition  $E|X|^p < \infty$ . This ends the proof of Theorem 2.2.  $\square$

## References

- [1] Baum, L.E., Katz, M. Convergence rates in the law of large numbers. *Trans. Amer. Math. Soc.* 120, 1965, 108-123.
- [2] Bradley, R. C. On the spectral density and asymptotic normality of weakly dependent random fields. *J. of Theor. Probab.* vol. 5, no. 2, 1992, 355-73.
- [3] Bryc, W., Smolenski, W. Moment conditions for almost sure convergence of weakly correlated random variables. *Proc. of Amer. Math. Soc.* vol. 119, no. 2, 1993, 629-35.
- [4] Cai, G.H. Strong law of large numbers for  $\varrho^*$ -mixing sequences with different distributions *Disc. Dynamics in Nat. Soc.* vol. 2006, Article ID 27648, 7 pages, 2006
- [5] Chen, P., Hu, T., C., Volodin A. Limit behaviour of moving average processes under  $\phi$ -mixing assumption. *Statist. Probab. Lett.* 79, 2009, 105-111.
- [6] Chen, Y. Chen, A., Ng K., The strong law of large numbers for extended negatively dependent random variables, *J. Appl. Probab.* 47, 4, 2010, 908-922.
- [7] Chow, Y.S. Delayed sums and Borel summability of independent, identically distributed random variables. *Bull. Inst. Math. Acad. Sinica* 1, 1973, 207-220.
- [8] Erdős, P. On a theorem of Hsu and Robbins. *Ann. Math. Stat.* 20, 1949, 286-291.
- [9] Gut, A. Complete convergence for arrays. *Periodica Math. Hungar.* 25, 1992, 51-75.
- [10] Hsu, P.L., Robbins, H. Complete convergence and the law of large numbers. *Proc. Nat. Acad. Sci. USA* 33, 1947, 25-31.

- [11] Ibragimov, I. A. (1962) Some limit theorems for stationary processes. *Theory Probab. Appl.* 7 1962, 349-382.
- [12] Joag-Dev, K., Proschan, F. Negative association of random variables with applications. *The Ann. Stat.* 11, 1983, 286-295.
- [13] Katz, M. The probability in the tail of a distribution. *Ann. Math. Stat.* 34, 1963, 312-318.
- [14] Kuczmaszewska, A. On complete convergence in Marcinkiewicz-Zygmund type SLLN for negatively associated random variables. *Acta. Math. Hungar.* 128(1-2), 2010, 116-130.
- [15] Kuczmaszewska, A., Lagodowski, Z. A. Convergence rates in the SLLN for some classes of dependent random fields, *J. Math. Anal. Appl.* 380, 2011, 571-584.
- [16] Kuczmaszewska, Convergence rate in the Petrov SLLN for dependent random variables. *Acta. Math. Hungar.* 148, 1, 2016, 56-72.
- [17] Lagodowski, Z. A. An approach to complete convergence theorems for dependent random fields via application of Fuk-Nagaev inequality. *J. Math. Anal. Appl.* 437, 2016, 380-395.
- [18] Liu, L. Precise large deviations for dependent random variables with heavy tails. *Stat. Probab. Lett.* 79, 2009, 1290C1298. doi:10.1016/j.spl.2009.02.001.
- [19] Miller, C. Three theorems on  $\varrho^*$ -mixing random fields. *J. Theor. Probab.* 7, 1994, 867-882.
- [20] Peligrad, M. Maximum of partial sums and an invariance principle for a class of weak dependent random variables. *Proc. Amer. Math. Soc.* vol. 126, no. 4, 1998, 1181-1189.
- [21] M. Peligrad, M., Gut, A. Almost-sure results for a class of dependent random variables. *J. Theor. Probab.* vol. 12, no. 1, 1999, 87-104.
- [22] Pruss, A.R. Randomly sampled Riemann sums and complete convergence in the law of large numbers for a case without identical distribution. *Proc. Amer. Math. Soc.* 124, 1996, 919-929.
- [23] Rozanov, Y.A., Volkonski, V.A. Some limit theorems for random function. *Theory Probab.* 4, 1959, 186-207.
- [24] Qiu, D.H., Chen, P.Y., Antonini, R.G., Volodin, A. On the complete convergence for arrays of rowwise extended negatively dependent random variables. *J. Korean Math. Soc.* 50, 2, 2013, 379-392.
- [25] Shen, A. Probability inequalities for END sequence and their applications, *J. Inequal. Appl.* 98, 2011, 1-12.

- [26] Shen, A. Complete convergence for weighted sums of END random variables and its applications to nonparametric regression models. *J. Nonparametric Stat.* 28, 4, 2016, 702-715. -
- [27] Shen, A., Xue, M., Wang, W. Complete convergence for weighted sums of extended negatively dependent random variables. *Communications in Stat.- Theory and Methods* 46, 3, 2017, 1433-1444.
- [28] Tuyen, D. Q. A strong law for mixing random variables, *Periodica Math. Hung.* 38, 1999, 131-136.
- [29] Utev, S., Peligrad, M., Maximal inequalities and an invariance principle for a class of weakly dependent random variables. *J. Theor. Probab.* vol. 16, no. 1, 2003, 101-115.
- [30] Wang, X., Hu, S., Yang, W., Shen, Y., On complete convergence for weighted sums of  $\varphi$ -mixing random variables, *J. Ineq. Appl.* 2010, Article ID 372390, 13 pages doi:10.1155/2010/372390.
- [31] Wang, X.J., Li, X.Q., Hu, S.H., Wang, X.H. On complete convergence for an extended negatively dependent sequence. *Communications in Statistics- Theory and Methods* 43, 2014, 2923-2937.
- [32] Wang, X.J., Zheng, L.L., Xu, C., Hu, S.H. Complete consistency for the estimator of nonparameter regression models based on extended negatively dependent errors. *Statistics: A Journal of Theoretical and Applied Stat.* 49, 2015, 396-407.
- [33] Wu, Y.F., Song, M.Z., Wang, C.H. Complete moment convergence and mean convergence for arrays of rowwise extended negatively dependent random variables. *The Scientific World Journal*, Artical ID 478612, 7 pages. 2014, doi: 10.1155/2014/478612.
- [34] Wang, J.F., Lu, F. B., Inequalities of maximum of partial sums and weak convergence for a class of weak dependent random variables. *Acta, Math. Sin.* 22, 2006, 693-700.
- [35] Yan, J.G. The Uniform Convergence and Precise Asymptotics of Generalized Stochastic Order Statistics. *J. Math. Anal. Appl.* 343, 2008, 644-653.
- [36] Yan, J.G. Strong Stability of a type of Jamison Weighted Sums for END Random Variables. *J. Korean Math. Soc.* 54, 3, 2017, 897-907.
- [37] Yan, J.G. Almost sure convergence for weighted sums of WNOD random variables and its applications in nonparametric regression models. *Communications in Stat.- Theory and Methods*. 2017.
- [38] Zhang, Y., Complete moment convergence for moving average process generated by  $\rho^-$ -mixing random variables. *J. Ineq. Appl.* 2015, 2015:245, DOI 10.1186/s13660-015-0766-5


# IRTG 1792 Discussion Paper Series 2018

For a complete list of Discussion Papers published, please visit  
[irtg1792.hu-berlin.de](http://irtg1792.hu-berlin.de).

- 001 "Data Driven Value-at-Risk Forecasting using a SVR-GARCH-KDE Hybrid" by Marius Lux, Wolfgang Karl Härdle and Stefan Lessmann, January 2018.
- 002 "Nonparametric Variable Selection and Its Application to Additive Models" by Zheng-Hui Feng, Lu Lin, Ruo-Qing Zhu and Li-Xing Zhu, January 2018.
- 003 "Systemic Risk in Global Volatility Spillover Networks: Evidence from Option-implied Volatility Indices " by Zihui Yang and Yinggang Zhou, January 2018.
- 004 "Pricing Cryptocurrency options: the case of CRIX and Bitcoin" by Cathy YH Chen, Wolfgang Karl Härdle, Ai Jun Hou and Weining Wang, January 2018.
- 005 "Testing for bubbles in cryptocurrencies with time-varying volatility" by Christian M. Hafner, January 2018.
- 006 "A Note on Cryptocurrencies and Currency Competition" by Anna Almosova, January 2018.
- 007 "Knowing me, knowing you: inventor mobility and the formation of technology-oriented alliances" by Stefan Wagner and Martin C. Goossen, February 2018.
- 008 "A Monetary Model of Blockchain" by Anna Almosova, February 2018.
- 009 "Deregulated day-ahead electricity markets in Southeast Europe: Price forecasting and comparative structural analysis" by Antanina Hryshchuk, Stefan Lessmann, February 2018.
- 010 "How Sensitive are Tail-related Risk Measures in a Contamination Neighbourhood?" by Wolfgang Karl Härdle, Chengxiu Ling, February 2018.
- 011 "How to Measure a Performance of a Collaborative Research Centre" by Alona Zharova, Janine Tellingner-Rice, Wolfgang Karl Härdle, February 2018.
- 012 "Targeting customers for profit: An ensemble learning framework to support marketing decision making" by Stefan Lessmann, Kristof Coussement, Koen W. De Bock, Johannes Haupt, February 2018.
- 013 "Improving Crime Count Forecasts Using Twitter and Taxi Data" by Lara Vomfell, Wolfgang Karl Härdle, Stefan Lessmann, February 2018.
- 014 "Price Discovery on Bitcoin Markets" by Paolo Pagnottoni, Dirk G. Baur, Thomas Dimpfl, March 2018.
- 015 "Bitcoin is not the New Gold - A Comparison of Volatility, Correlation, and Portfolio Performance" by Tony Klein, Hien Pham Thu, Thomas Walther, March 2018.
- 016 "Time-varying Limit Order Book Networks" by Wolfgang Karl Härdle, Shi Chen, Chong Liang, Melanie Schienle, April 2018.
- 017 "Regularization Approach for Network Modeling of German EnergyMarket" by Shi Chen, Wolfgang Karl Härdle, Brenda López Cabrera, May 2018.
- 018 "Adaptive Nonparametric Clustering" by Kirill Efimov, Larisa Adamyan, Vladimir Spokoiny, May 2018.
- 019 "Lasso, knockoff and Gaussian covariates: a comparison" by Laurie Davies, May 2018.

**IRTG 1792, Spandauer Straße 1, D-10178 Berlin**  
**<http://irtg1792.hu-berlin.de>**

This research was supported by the Deutsche  
Forschungsgemeinschaft through the IRTG 1792.



# IRTG 1792 Discussion Paper Series 2018

For a complete list of Discussion Papers published, please visit  
[irtg1792.hu-berlin.de](http://irtg1792.hu-berlin.de).

- 020 "A Regime Shift Model with Nonparametric Switching Mechanism" by Haiqiang Chen, Yingxing Li, Ming Lin and Yanli Zhu, May 2018.
- 021 "LASSO-Driven Inference in Time and Space" by Victor Chernozhukov, Wolfgang K. Härdle, Chen Huang, Weining Wang, June 2018.
- 022 "Learning from Errors: The case of monetary and fiscal policy regimes" by Andreas Tryphonides, June 2018.
- 023 "Textual Sentiment, Option Characteristics, and Stock Return Predictability" by Cathy Yi-Hsuan Chen, Matthias R. Fengler, Wolfgang Karl Härdle, Yanchu Liu, June 2018.
- 024 "Bootstrap Confidence Sets For Spectral Projectors Of Sample Covariance" by A. Naumov, V. Spokoiny, V. Ulyanov, June 2018.
- 025 "Construction of Non-asymptotic Confidence Sets in 2 -Wasserstein Space" by Johannes Ebert, Vladimir Spokoiny, Alexandra Suvorikova, June 2018.
- 026 "Large ball probabilities, Gaussian comparison and anti-concentration" by Friedrich Götze, Alexey Naumov, Vladimir Spokoiny, Vladimir Ulyanov, June 2018.
- 027 "Bayesian inference for spectral projectors of covariance matrix" by Igor Silin, Vladimir Spokoiny, June 2018.
- 028 "Toolbox: Gaussian comparison on Euclidian balls" by Andzhey Koziuk, Vladimir Spokoiny, June 2018.
- 029 "Pointwise adaptation via stagewise aggregation of local estimates for multiclass classification" by Nikita Puchkin, Vladimir Spokoiny, June 2018.
- 030 "Gaussian Process Forecast with multidimensional distributional entries" by Francois Bachoc, Alexandra Suvorikova, Jean-Michel Loubes, Vladimir Spokoiny, June 2018.
- 031 "Instrumental variables regression" by Andzhey Koziuk, Vladimir Spokoiny, June 2018.
- 032 "Understanding Latent Group Structure of Cryptocurrencies Market: A Dynamic Network Perspective" by Li Guo, Yubo Tao and Wolfgang Karl Härdle, July 2018.
- 033 "Optimal contracts under competition when uncertainty from adverse selection and moral hazard are present" by Natalie Packham, August 2018.
- 034 "A factor-model approach for correlation scenarios and correlation stress-testing" by Natalie Packham and Fabian Woebbecking, August 2018.
- 035 "Correlation Under Stress In Normal Variance Mixture Models" by Michael Kalkbrener and Natalie Packham, August 2018.
- 036 "Model risk of contingent claims" by Nils Detering and Natalie Packham, August 2018.
- 037 "Default probabilities and default correlations under stress" by Natalie Packham, Michael Kalkbrener and Ludger Overbeck, August 2018.
- 038 "Tail-Risk Protection Trading Strategies" by Natalie Packham, Jochen Papenbrock, Peter Schwendner and Fabian Woebbecking, August 2018.

**IRTG 1792, Spandauer Straße 1, D-10178 Berlin**  
**<http://irtg1792.hu-berlin.de>**

This research was supported by the Deutsche  
Forschungsgemeinschaft through the IRTG 1792.



# **IRTG 1792 Discussion Paper Series 2018**

For a complete list of Discussion Papers published, please visit [irtg1792.hu-berlin.de](http://irtg1792.hu-berlin.de).

- 039 "Penalized Adaptive Forecasting with Large Information Sets and Structural Changes" by Lenka Zbonakova, Xinjue Li and Wolfgang Karl Härdle, August 2018.
- 040 "Complete Convergence and Complete Moment Convergence for Maximal Weighted Sums of Extended Negatively Dependent Random Variables" by Ji Gao YAN, August 2018.
- 041 "On complete convergence in Marcinkiewicz-Zygmund type SLLN for random variables" by Anna Kuczmaszewska and Ji Gao YAN, August 2018.

**IRTG 1792, Spandauer Straße 1, D-10178 Berlin**  
**<http://irtg1792.hu-berlin.de>**

This research was supported by the Deutsche  
Forschungsgemeinschaft through the IRTG 1792.