

Muratov, Oleg

Working Paper

All-pay auctions with reserve price and bid cap

Discussion Papers, No. 21-06

Provided in Cooperation with:

Department of Economics, University of Bern

Suggested Citation: Muratov, Oleg (2021) : All-pay auctions with reserve price and bid cap, Discussion Papers, No. 21-06, University of Bern, Department of Economics, Bern

This Version is available at:

<https://hdl.handle.net/10419/242857>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.



<https://creativecommons.org/licenses/by/4.0/>



^b
**UNIVERSITÄT
BERN**

Faculty of Business, Economics
and Social Sciences

Department of Economics

All-Pay Auctions with Reserve Price and Bid Cap

Oleg Muratov

21-06

April, 2021

DISCUSSION PAPERS

Schanzeneckstrasse 1
CH-3012 Bern, Switzerland
<http://www.vwi.unibe.ch>

All-Pay Auctions with Reserve Price and Bid Cap*

Oleg Muratov

April 23, 2021

Abstract

I characterize equilibria in an all-pay auction with a reserve price and a bid cap. I consider the cases of two and three players. I show that equilibrium bidding is characterized by atoms at 0, the reserve price, and the bid cap, as well as continuous bidding above the bid cap. If the valuations are high enough, the range for continuous bidding shrinks completely. I show that for some parameter ranges there exist multiple equilibria. Under three players, I show that there exist equilibria with the following features: the player with the non-top valuation can have a positive rent; the players with completely different valuations can actively compete for the single prize; the player with positive payoff can have different payoff in different equilibria.

1 Introduction

Contests, and, in particular, all-pay auctions, are used to model competitive environments where part-taking is costly, a limited number of prizes is available, and

*e-mail: oleg.muratov@vwi.unibe.ch; This article includes work from Chapter 2 of my PhD thesis, Muratov (2019). I am indebted to my thesis advisors, Ron Siegel, Kalyan Chatterjee, and Rohit Lamba for their guidance and support. I am grateful to Igor Letina for the detailed discussion and feedback. I would also like to thank the participants of the 2020 Contests: Theory and Evidence conference. I acknowledge and appreciate the financial support from Swiss National Science Foundation (Grant number 100018_185202).

the payoffs of players often involve an all-or-nothing component. For instance, all-pay auctions have been used to model and analyze lobbying and campaign spendings (Hillman and Riley (1989), Baye et al. (1993), Che and Gale (1998)) and patent and R&D races (Moldovanu and Sela (2003), Che and Gale (2003)).

Theoretical results in the standard complete information all-pay auction with the single prize and multiple players are fully characterized in Baye et al. (1996). They show that in equilibrium only the player with the strongly highest prize valuation has a positive payoff. Baye et al. (1996) also show that more than 2 players can actively bid only in non-generic cases of either (a) 3 and more players having the same highest-ranked valuation of the prize, or (b) 2 and more players having the same second highest-ranked valuation of the prize.

Clark and Riis (1998) generalize the analysis to include multiple prizes. Siegel (2009) generalizes the all-pay auctions to all-pay *contests*, allowing for general asymmetries between the players and general cost functions. One of the results of Siegel (2009) is that under regularity conditions, there are pre-determined sets of “winners” and “losers” in the contest, such that “winners” get their positive payoffs, same across equilibria, and “losers” get zero.

The previous literature has not considered all-pay auctions with a combination of a reserve price and a bid cap, or a combination of a bid cap and more than two players. However, such environments are natural to occur. Consider, for instance, sport tournaments. In the NBA there is a salary cap for the team.¹ NBA also has a minimum team salary, so it constitutes an example of a contest with the reserve and the cap.² Similarly, there are salary caps and franchise fees in the NFL.³ In the R&D-race, if a patent awarded to the single researcher, a contest organizer has incentives to limit the total expenditures. This way, there would be less money non-productively spent on duplicating projects. Together with sunk cost, typical for the R&D, such environment would be similar to the contest with

¹See, for instance, Salary Cap Overview.

²Fee for expansion teams.

³ The business of owning an NFL team.

the reserve and the cap. Studying all-pay auctions with the bid cap is important from the theoretical point of view, because such mechanism was shown to be the optimal one when the agents' performance is evaluated by a lenient reviewer, in Letina et al. (2020). Understanding, how equilibrium and the efforts profile change with the addition non-zero fixed cost of effort is an important robustness check.

The bid caps and the reserve price in the all-pay auctions have been considered separately. The two-player all-pay auction with a cap is analyzed in Che and Gale (1998). Szech (2015) shows the importance of the tie-breaking rule in the all-pay auction with the cap scenario. The role of the reserve price in the all-pay auction is studied in Siegel (2014). In this paper I study single prize all-pay auctions with 2 and 3 bidders in the presence of both the reserve price and the bid cap. Studying this environment allows to derive new insights. Thus, I show that there are settings, in which the same player can be either a "winner" or a "looser", depending on what equilibrium is played (see Corollary 1); settings, where a "winner" can have different payoffs across equilibria (see Proposition 4.iii); settings, where three players with completely different payoffs can bid actively in equilibrium (see Proposition 4.iv); and also settings, where the player with a non-top valuation is a "winner" in an equilibrium (as illustrated in examples for Propositions 4.iii and 4.iv). Thus, my results complement the results of Baye et al. (1996) and Siegel (2009).

I fully characterize equilibria in all-pay auctions with the reserve and the bid cap. I show that with 2 players, the type of equilibrium behavior depends on the sizes of valuation for the prize. There are three equilibria regimes possible. The first regime happens if the second-highest valuation is between the reserve and the cap: in equilibrium there are atoms at 0 and the reserve, as well as continuously distributed mass above the reserve. The support is expanding with the increase of the second-highest prize. The second regime occurs if both valuations are above the reserve. For this case, there appears an atom at the cap, and there is continuous bidding in the midrange, between the reserve and the cap. The

tie-breaking rule becomes important in a non-trivial way. The player with the tie-breaking adjusted advantage has economic rent, and the range for continuous bidding shrinks with respect to the valuation of the player with the tie-breaking adjusted disadvantage. The third regime occurs when both valuations are high, and the range for continuous bidding has shrunk completely: in equilibrium, there is only atomic bidding, with atoms at 0, the reserve, and the cap. In case of three players, I show that equilibria have a similar structure. However, characterization is more complex, and there is no longer such natural ordering between types of equilibria and a second- or a third order valuation.

There exist an application of my results to the study of class of games of information design. In Muratov (2021), I show that there is a mapping between an all-pay auction with the reserve and the cap, on the one hand, and the competing senders information design game, as in Boleslavsky and Cotton (2018). Hence, understanding equilibria in the setting of the current paper is useful for characterizing equilibria in related information design games.

2 Model: Two Bidders Case

Consider an All-Pay Auction with one prize and two bidders, indexed 1 and 2. Suppose, the two bidders value the prize at V_1 and V_2 , and have to submit their bids, denoted b_1 and b_2 . The player, who bids the most, wins the prize. Each player has to pay his actual bid, regardless of whether he wins or loses.

Suppose, that in addition to standard assumptions, there is a reserve price, r , and also a bid cap, κ , such that $0 < r < \kappa$. It is without loss to normalize $\kappa = 1$. A player, who bids below the reserve price can never win the item, even if his bid turns out to be the largest; any bid larger than 1, will be disregarded. Finally, in case of a tie, let bidder 1 obtain the prize with probability ρ_1 , and bidder 2 - with probability ρ_2 , such that $\rho_1 + \rho_2 = 1$, $\rho_1 \in (0, 1)$.

Let the bidders be risk-neutral expected utility maximizers. Denote their

utility functions as $u_1()$ and $u_2()$, respectively. For a pair of bids (b_1, b_2) the bidders' payoffs are

$$u_1(b_1, b_2) = \begin{cases} V_1 - b_1, & \text{if } b_1 > b_2, b_1 \in [r, 1] \\ \rho \times V_1 - b_1, & \text{if } b_1 = b_2, b_1 \in [r, 1] \\ -b_1, & \text{else.} \end{cases}$$

$$u_2(b_1, b_2) = \begin{cases} V_2 - b_2, & \text{if } b_2 > b_1, b_2 \in [r, 1] \\ (1 - \rho) \times V_2 - b_2, & \text{if } b_2 = b_1, b_2 \in [r, 1] \\ -b_2, & \text{else.} \end{cases}$$

Let us first state the result on how in general the equilibrium bidding looks. This result is valid for the two and the three bidders case, which is considered in detail in section 3.

Lemma 1. *The bids are distributed in the subset of $[0, 1]$. The subset has the following properties:*

- (i) *There is no mass of bids in $(0, r)$*
- (ii) *There can be atoms at $\{0\}$, $\{r\}$, and $\{1\}$. There can be at most one player placing an atom at $\{r\}$. There can be no other atoms*
- (iii) *If there is mass of bids in a set $B \subseteq (r, 1)$, then B is an interval: $B = (r, \bar{b})$, $r < \bar{b} \leq 1$. At least two players bid everywhere in (r, \bar{b}) . If a third player also bids in the interior of B , he distributes it continuously in an interval (\underline{b}, \bar{b}) , where $\underline{b} \in (r, \bar{b})$, but is not uniquely determined*

The proof of this lemma is in the appendix A.1.

Three qualitatively different types bidding behavior emerge in equilibria, depending on how high the valuations are. When one of the valuations is low, like in the region A, there can be no atoms besides the ones at 0 and r , with the remaining mass distributed continuously in a subset of $(r, 1)$; when one of the valuations is

middle, like in the region B, there are atoms at 0, r , and 1, as well as continuously distributed mass in the subset of $(r, 1)$; when both valuations are high, like in the regions C and D, there are atoms only. We will now list the results for a strict subset of $(V_1, V_2) \in [r, +\infty)^2$, namely the union of the regions A, A', B, B', C, and D, as depicted on figure 1. The remaining cases are symmetric to the ones that will be described, and the results for them can be stated by permuting the valuations, V_i , and the tie-breaking probabilities.⁴




Figure 1: Equilibrium Regions

Proposition 1.i. *If $V_2 < 1$ and $V_2 \leq V_1$ (region A), in the unique equilibrium, the players' bids are distributed according to the CDFs:*

$$G_1(b) = \begin{cases} 0, & \text{if } b < r \\ \frac{b}{V_2}, & \text{if } b \in [r, V_2) \\ 1, & \text{if } b \geq V_2; \end{cases} \quad G_2(b) = \begin{cases} \frac{r+V_1-V_2}{V_1}, & \text{if } b < r \\ \frac{b+V_1-V_2}{V_1}, & \text{if } b \in [r, V_2) \\ 1, & \text{if } b \geq V_2. \end{cases}$$

Player 1 has a positive equilibrium payoff $U^ = V_1 - V_2$; Player 2 has a zero equilibrium payoff.*

Proposition 1.ii. *If $V_2 = V_1 \leq 1$ (region A'), there are multiple equilibria. In*

⁴Also note that there are trivial equilibria for a subset of parameters: if both valuations are below the reserve price, both agents will bid 0 with certainty; if just valuation is above the reserve price, the player with that valuation will bid r with certainty; and if both valuations are high enough, i.e. if $(V_1, V_2) > \left(\frac{1}{\rho_1}, \frac{1}{\rho_2}\right)$, both players will bid 1 with certainty.

every equilibrium the bidding behavior can be described by the CDFs:

$$G_1(b) = \begin{cases} \frac{r-t_1}{V_2}, & \text{if } b < r \\ \frac{b}{V_2}, & \text{if } b \in [r, V_2) \\ 1, & \text{if } b \geq V_2; \end{cases} \quad G_2(b) = \begin{cases} \frac{r-t_2}{V_2}, & \text{if } b < r \\ \frac{b}{V_2}, & \text{if } b \in [r, V_2) \\ 1, & \text{if } b \geq V_2; \end{cases}$$

where $t_i \in [0, r]$, $t_1 \times t_2 = 0$; so that t_i is a free parameter, that stands for the size of the atom that player i might have at r , with at most one player having an atom there. Both players have equilibrium payoffs of zero.

Overall, if player 1 has a strictly higher valuation, the equilibrium is unique and only player 1 has a non-arbitrarily sized atom at r ; if the valuations are equal, there are multiple equilibria, with multiplicity being driven by two sources: the identity of the player, who has an atom at r (either player can have an atom there, but not both); and the size of the atom at r (any probability no greater than $\frac{r}{V_2}$).

The characterization above is exhaustive, as shown in the Appendix B.1.

Beside the possibility of multiplicity in case of equal valuations, the equilibrium for this case has some standard properties of All-Pay Auctions equilibria, similar to the ones shown in Baye et al. (1996), Siegel (2009), and Siegel (2014): players bid up to the lower valuation; the player with the (strictly) higher valuation has the (strictly) positive payoff, $V_1 - V_2$; in the interior of $\{0\} \cup \{r\} \cup (r, 1]$ the bidding is continuous. With the knowledge of this facts, it is straightforward to establish the values of CDFs in (b, V_2) :

$$V_i G_j(b) - b = \max\{V_i - V_j, 0\}, \forall b \in (r, V_2).$$

The major difference with the standard setting of, for example, Baye et al. (1996), is that in equilibrium there is no mass in $(0, r)$. Atoms at 0 and/or r compensate for that gap, taking also into account that the player with the strictly positive payoff would never have 0 in his best response set. The appendix proves that the behavior described here is exhaustive for all equilibria. Figure 2 shows the plots

of CDFs and comparative statics with respect to the increase in the V_2 .




Figure 2: Distributions of bids and comparative statics with respect to V_2

Note that in regions A and A', the upper bound of the support of bids is increasing in V_2 . At some point, after V_2 becomes greater than 1, there is nowhere left for the support to expand, since there is a bid cap of 1. This is where the equilibrium bidding behavior switches to regions B and B', for which the following results hold:

Proposition 2.i. *If $1 < V_2 \leq \frac{1-\rho_1 r}{\rho_2}$, $\rho_2^2 V_2 - \rho_2 < \rho_1^2 V_1 - \rho_1$ (region B), in the unique equilibrium, the players' bids are distributed according to the CDFs:*

$$G_1(b) = \begin{cases} 0, & \text{if } b < r \\ \frac{b}{V_2}, & \text{if } b \in \left[r, \frac{1}{\rho_1} - \frac{\rho_2}{\rho_1} V_2 \right) \\ \frac{1-\rho_2 V_2}{\rho_1 V_2}, & \text{if } b \in \left[\frac{1}{\rho_1} - \frac{\rho_2}{\rho_1} V_2, 1 \right) \\ 1, & \text{if } b \geq 1; \end{cases} \quad G_2(b) = \begin{cases} \frac{r+U^*}{V_1}, & \text{if } b < r \\ \frac{b+U^*}{V_1}, & \text{if } b \in \left[r, \frac{1}{\rho_1} - \frac{\rho_2}{\rho_1} V_2 \right) \\ \frac{1-\rho_2 V_2 + \rho_1 U^*}{\rho_1 V_1}, & \text{if } b \in \left[\frac{1}{\rho_1} - \frac{\rho_2}{\rho_1} V_2, 1 \right) \\ 1, & \text{if } b \geq 1, \end{cases}$$

where $U^* = \frac{\rho_2 - \rho_1 + \rho_1^2 V_1 - \rho_2^2 V_2}{\rho_1^2} > 0$ is the equilibrium payoff of player 1. Player 2 has a zero payoff.

Proposition 2.ii. *If $1 < V_2 \leq \frac{1-\rho_1 r}{\rho_2}$, $\rho_2^2 V_2 - \rho_2 = \rho_1^2 V_1 - \rho_1$ (region B'), there are multiple equilibria. In every equilibrium the bidding behavior can be described by*

the CDFs:

$$G_1(b) = \begin{cases} \frac{r-t_1}{V_2}, & \text{if } b < r \\ \frac{b}{V_2}, & \text{if } b \in \left[r, \frac{1}{\rho_1} - \frac{\rho_2}{\rho_1} V_2 \right) \\ \frac{1-\rho_2 V_2}{\rho_1 V_2}, & \text{if } b \in \left[\frac{1}{\rho_1} - \frac{\rho_2}{\rho_1} V_2, 1 \right) \\ 1, & \text{if } b \geq 1; \end{cases} \quad G_2(b) = \begin{cases} \frac{r-t_2}{V_1}, & \text{if } b < r \\ \frac{b}{V_1}, & \text{if } b \in \left[r, \frac{1}{\rho_1} - \frac{\rho_2}{\rho_1} V_2 \right) \\ \frac{1-\rho_2 V_2}{\rho_1 V_1}, & \text{if } b \in \left[\frac{1}{\rho_1} - \frac{\rho_2}{\rho_1} V_2, 1 \right) \\ 1, & \text{if } b \geq 1, \end{cases}$$

where, similarly to Proposition 1.ii, $t_i \in [0, r]$, $t_1 \times t_2 = 0$. Both players have equilibrium payoffs of zero.

It is shown in the appendix B.2 that the above characterization is exhaustive.

There are several changes in equilibrium bidding in regions B and B', in comparison to A and A': first, there appear atoms at the bid of 1 for both players; second, the range for continuous bidding, $\left[r, \frac{1}{\rho_1} - \frac{\rho_2}{\rho_1} V_2 \right)$ now shrinks in V_2 ; and third, player 1's payoff is no longer equal to $V_1 - V_2$, unless $\rho_1 = \frac{1}{2}$. So, favoritism is effective under the bid cap and high enough valuations: if $\rho_1 > \rho_2$, player can have payoff that is higher than standard payoff of $V_1 - V_2$, and vice versa. Player 2's payoff, nevertheless, remains equal to 0.

What remains the same as in the regions A and A', is the density at the continuous part of bidding: in order to make the opponent indifferent between bidding slightly more and slightly less, the density must be the inverse of their valuation, as is standard. The upper bound of the continuous part of the support, \bar{b} , the sizes of atoms, and the equilibrium payoff of player 1 are determined by the indifference of players between bidding \bar{b} and bidding 1:

$$\begin{aligned} V_1 \times G_2(\bar{b}) - \bar{b} &= V_1(G_2(\bar{b}) + \rho_1(1 - G_2(\bar{b}))) - 1 = U^* \\ V_2 \times G_1(\bar{b}) - \bar{b} &= V_2(G_1(\bar{b}) + \rho_2(1 - G_1(\bar{b}))) - 1 = 0. \end{aligned}$$

From the conditions above, together with the fact that $G'_i(b) = \frac{1}{V_j}$, $\forall b \in (r, \bar{b})$, it follows that $\bar{b} = \frac{1-\rho_2 V_2}{\rho_1}$, and $U^* = \frac{\rho_2 - \rho_1 + \rho_1^2 V_1 - \rho_2^2 V_2}{\rho_1^2}$, as stated in the propositions

2.i, 2.ii.

Figure 3 shows the CDFs for a typical pair of valuations in region B and the tie-breaking rule $\rho_1 = \rho_2 = 1/2$, as well as the comparative statics with respect to the increase in V_2 .




Figure 3: Distributions of bids

Finally, when V_2 is high enough, and the upper bound of continuous bidding, \bar{b} , reaches the bid cap, any continuous bidding disappears. In regions C and D, in equilibria, the bidding distributions are atomic only.

Proposition 3.i. *If $(V_1, V_2) \in \left[\frac{1-r\rho_2}{\rho_1}, \frac{1}{\rho_1}\right] \times \left[\frac{1-r\rho_1}{\rho_2}, \frac{1}{\rho_2}\right]$, (region C), there is an equilibrium in which the players bid according to*

$$b_1 = \begin{cases} 0, & \text{with probability } \frac{1-\rho_2 V_2}{\rho_1 V_2} \\ 1, & \text{with probability } \frac{V_2-1}{\rho_1 V_2} \end{cases} \quad b_2 = \begin{cases} 0, & \text{with probability } \frac{1-\rho_1 V_1}{\rho_2 V_1} \\ 1, & \text{with probability } \frac{V_1-1}{\rho_2 V_1} \end{cases};$$

and both players have zero payoffs.

Proposition 3.ii. *If $(V_1, V_2) \in \left\{\frac{1-\rho_2 r}{\rho_1}\right\} \times \left[\frac{1-r\rho_1}{\rho_2}, \frac{1}{\rho_2}\right]$, there is a class of equilibria in which the players bid according to*

$$b_1 = \begin{cases} 0, & \text{with probability } (1-t) \times \frac{1-\rho_2 V_2}{\rho_1 V_2} \\ r, & \text{with probability } t \times \frac{1-\rho_2 V_2}{\rho_1 V_2} \\ 1, & \text{with probability } \frac{V_2-1}{\rho_1 V_2} \end{cases} \quad b_2 = \begin{cases} 0, & \text{with probability } \frac{1-\rho_1 V_1}{\rho_2 V_1} \\ 1, & \text{with probability } \frac{V_1-1}{\rho_2 V_1} \end{cases};$$

where $t \in [0, 1]$ is a free parameter, and both players have zero payoffs.

Proposition 3.iii. *If $(V_1, V_2) \in \left[\frac{1-\rho_2 r}{\rho_1}, \infty\right) \times \left[\frac{1-\rho_1 r}{\rho_2}, \frac{1}{\rho_2}\right]$, (region D), there is an equilibrium, in which the players bid according to*

$$b_1 = \begin{cases} r, & \text{with probability } \frac{1-\rho_2 V_2}{\rho_1 V_2} \\ 1, & \text{with probability } \frac{V_2-1}{\rho_1 V_2} \end{cases} \quad b_2 = \begin{cases} 0, & \text{with probability } \frac{V_1 \rho_1 - (1-r)}{V_1 \rho_1} \\ 1, & \text{with probability } \frac{1-r}{\rho_1 V_1} \end{cases};$$

and player 1 has a positive equilibrium payoff $u_1^* = V_1 - \frac{1-\rho_2 r}{\rho_1}$

The above characterization in propositions 3.i-3.iii is exhaustive, because under such high valuations and the bidding space limited by 1 from above, there can be no mass in the interior of $(r, 1)$. Thus, the bidding behavior is completely characterized by sizes of atoms at 0, r , and 1.

Let us also consider the setting of 3.iii, but let us switch the role of the players. It follows then that for the region $(V_1, V_2) \in \left[\frac{1-\rho_2 r}{\rho_1}, \frac{1}{\rho_2}\right] \times \left[\frac{1-\rho_1 r}{\rho_2}, \infty\right)$ there is an equilibrium with player one mixing between bidding 0 and 1, bringing him a zero payoff, and player two mixing between bidding r and 1, bringing him a positive payoff of $V_2 - \frac{1-\rho_1 r}{\rho_2}$. This observation, combined with the results of propositions 3.i-3.iii allows to state the following:

Corollary 1. *In the non-empty region $(V_1, V_2) \in \left[\frac{1-\rho_2 r}{\rho_1}, \frac{1}{\rho_1}\right] \times \left[\frac{1-\rho_1 r}{\rho_2}, \frac{1}{\rho_2}\right]$, there are three qualitatively different equilibria. In this region, each player can be a "winner" (have a positive payoff) or a "loser" (have a zero payoff), depending on what equilibrium is played. Also in one equilibrium both players are "losers", despite $V_1 \neq V_2$ in that region, generically.*

Augmenting the above characterization with the regions, derived by switching the roles of two players; and also by the region $(V_1, V_2) > \left(\frac{1}{\rho_1}, \frac{1}{\rho_2}\right)$, where the equilibrium is to bid 1 with certainty, completes the characterization of equilibria.

To see how the equilibrium conditions were derived for region D, for example, use the following conditions:

$$V_1 \times \mathbb{P}\{b_2 = 0\} - r = V_1 \times (\mathbb{P}\{b_2 = 0\} + \rho_1(1 - \mathbb{P}\{b_2 = 0\})) - 1 \geq 0,$$

$$V_2 \times (\mathbb{P}\{b_1 = r\} + \rho_2(1 - \mathbb{P}\{b_1 = r\})) - 1 = 0,$$

$$V_2 \times \mathbb{P}\{b_1 = r\} - r \leq 0.$$

Solving the above system of equalities and inequalities allows for the characterization of bidding probabilities, equilibrium payoffs, and the region in the proposition 3.iii. Similar logic applies to propositions 3.i and 3.ii.

2.1 Characterizing expenditures

Knowing the equilibrium distributions of bids allows to compute various expected characteristics of bids, such as, the individual expenditures of each player, the sum of expenditures, the variance of expenditures, and so on. Depending on the exact economic context, and the interpretation of the bids (productive effort in preparation for an exam, a sports team budget, R&D spendings) the All-Pay Auction organizer might have different objectives - she or he might want to maximize the total average spendings/efforts; make sure that individual levels are above a certain threshold; compress the discrepancy between individual spendings; or achieve some other goals. Here we provide the expressions for individual average spendings and their sum, as examples of the target variables, and the ways they behave in our environment.

In a working paper, Muratov (2021), I show that there is a mapping between all-pay auctions and a class of information design games, in which competing entrepreneurs-senders try to persuade a single receiver to invest into their project, rather than the opponent's, as the one studied in Boleslavsky and Cotton (2018). Average expenses in the all-pay auctions environment correspond to prior expected qualities in the information design environment. Thus, the expressions for the expenditures in terms of prize valuations are essential for characterizing this

mapping.

Denote bidder one's expenditures, and bidder two's expenditures as α_1 , and α_2 , respectively. For the valuations in the region A, $V_2 < 1, V_2 \leq V_1$, the expenditures are

$$(\alpha_1; \alpha_2) = \left(\frac{V_2}{2} + \frac{r^2}{2V_2}; \frac{V_2^2 - r^2}{2V_1} \right).$$

In the region A', $V_1 = V_2 \leq 1$, the expenditures are

$$(\alpha_1; \alpha_2) = \left(\frac{V_2}{2} - \frac{r(r - 2t_1)}{2V_2}; \frac{V_2}{2} - \frac{r(r - 2t_2)}{2V_2} \right),$$

where $t_i \in [0, r]$, $t_1 \times t_2 = 0$, representing the multiplicity of equilibria in this region.

In the region B, $1 < V_2 \leq \frac{1 - \rho_1 r}{\rho_2}$, $\rho_2^2 V_2 - \rho_2 < \rho_1^2 V_1 - \rho_1$, the expenditures are

$$(\alpha_1; \alpha_2) = \left(\frac{\rho_1^2 (r^2 + V_2^2) + (\rho_2 - \rho_1)(V_2 - 1)^2}{2\rho_1^2 V_2}; \frac{\rho_2^2 V_2^2 + \rho_1 (2 - \rho_1 r^2) - 1}{2\rho_1^2 V_1} \right).$$

Note that if the tie-breaking rule is fair, $\rho_1 = 0.5$, the expenditures in this region coincide with the expenditures in region A. In B', the expenditures are

$$\alpha_1 = \frac{V_2(\rho_2^2 V_2 - 2(\rho_2 - \rho_1)) + 1 - \rho_1(2 + r(r - 2t_1)\rho_1)}{2V_2\rho_1^2},$$

$$\alpha_2 = \frac{V_2^2\rho_2^2 + \rho_1(2 - r(r - 2t_2)\rho_1) - 1}{2V_1\rho_1^2},$$

$t_i \in [0, r]$, $t_1 \times t_2 = 0$.

Finally, in the regions C, C', and D, the expenditures are:

-

$$(\alpha_1; \alpha) = \left(\frac{V_2 - 1}{\rho_1 V_2}; \frac{V_1 - 1}{\rho_2 V_1} \right),$$

$$\text{in C, } (V_1, V_2) \in \left[\frac{1-r\rho_2}{\rho_1}, \frac{1}{\rho_1} \right] \times \left[\frac{1-r\rho_1}{\rho_2}, \frac{1}{\rho_2} \right];$$

-

$$(\alpha_1; \alpha_2) = \left(\frac{V_2 - 1 + r \times t(1 - \rho_2 V_2)}{\rho_1 V_2}; \frac{1 - r}{1 - \rho_2 r} \right),$$

$$t \in [0, 1], \text{ in C', } (V_1, V_2) \in \left\{ \frac{1-\rho_2 r}{\rho_1} \right\} \times \left[\frac{1-r\rho_1}{\rho_2}, \frac{1}{\rho_2} \right];$$

-

$$(\alpha_1, \alpha_2) = \left(\frac{V_2(1 - r\rho_2) + 1 - r}{V_2\rho_1}, \frac{1 - r}{\rho_1 V_1} \right),$$

$$\text{in D, } (V_1, V_2) \in \left[\frac{1-(1-\rho)r}{\rho}, \infty \right) \times \left[\frac{1-\rho r}{1-\rho}, \frac{1}{1-\rho} \right].$$

Applying the expenditures results, on figure 4, I plot the sum of the bidders' expenditures as a function of V_2 , fixing various values of V_1 : low ($r < V < 1$), average-low ($1 < V_1 < \frac{1-\rho_2 r}{\rho_1}$), average-high ($\frac{1-\rho_2 r}{\rho_1} < V_1 < \frac{1}{\rho_1}$), and high ($V_1 > \frac{1}{\rho_1}$).⁵ Together with the expenditures in the all-pay auction with the reserve price r and the bid cap of 1, the graphs also contain the expenditures in the all-pay auction without any restrictions; with just the reserve price r ; and just the bid cap of 1. It can be seen that there is no uniform ranking of the four formats in terms of the total expenditures; but the all-pay auction *with* the reserve price *and* the bid cap can provide the largest total expenditures for some ranges of valuations.

3 Three Bidders Case

Studying the case of two bidders in the all-pay auction in the presence of the reserve price and the bid cap has brought new insights, if we compare the results to the ones in the settings of Baye et al. (1996) and Che and Gale (1998). Thus, for a range of parameters, $(V_1, V_2) \in C \cap D$, in equilibrium bidding CDFs are

⁵For parameters with multiple equilibria, the equilibria with maximal expenditures were chosen. Also note that discontinuities in the expenditures are due to multiplicity of equilibria on the boundaries of regions.




Figure 4: Revenue comparison for different values of V_1

characterized by atoms only, but not purely at the top. In the same range of parameters there are three types equilibria. In one, both players bid 0 and 1 only, in another, player 1 bids r instead of 0 and has a positive payoff, and in the third type it is player 2 who bids r instead of 0 and has a positive payoff. Thus, in this region, it is possible that the player with initial disadvantage, i.e. lower valuation of the prize, in equilibrium gets a positive payoff, unlike in the setting of Baye et al. (1996), and that phenomenon is not driven by the tie-breaking rule.

Studying this environment with the third bidder allows to perform a robustness check of the results, derived in the two players case; and it also brings new insights, not possible under the two bidders. Thus, multiple equilibria are possible even for ranges of parameters that result in equilibria with some continuous bidding. The nature of this multiplicity is of a different kind than the one in Baye et al. (1996) with three or more players: in addition to the degree of freedom in when one of the three players joins the continuous bidding, there is multiplicity of the positive equilibrium rent, and the equilibrium support, which all affects the shapes of CDFs. Another new finding is that it is possible for all three players to bid actively for a range of completely asymmetric valuations $V_1 \neq V_2 \neq V_3$, with two players' bidding involving continuous support, and the third player bidding at the top and the bottom.

Throughout this section only the symmetric tie-breaking rule will be considered, i.e. 1/2-1/2 chance of getting the item if the two bidders tie, and 1/3-each chance of getting the item if the three bidders tie.

Let us first identify the restrictions on the valuations of player 3, V_3 , such that equilibria from section 2 hold; i.e. player 3 stays inactive and effectively only players 1 and 2 are the bidders.

Proposition 4.i. *There are non-empty regions of parameters, such that only players 1 and 2 are active bidders. Depending on how high V_1 and V_2 are, the results are:*

- *For the parameters $V_1 \geq V_2 \geq V_3$, $V_2 \in [r, 1)$, there are equilibria as in the*

environments of propositions 1.i and 1.ii. Moreover, if $V_2 > V_3$, these are all the equilibria

- For the parameters $V_1 \geq V_2 \in [1, 2-r)$, $V_3 \leq \frac{3V_1V_2}{4+3V_1-5V_2+V_2^2}$, there are equilibria as in the environments of propositions 2.i and 2.ii, under $\rho_1 = \rho_2 = 1/2$.
- For the parameters $(V_1, V_2) \in [2-r, 2]^2$, $V_3 \leq \frac{3V_1V_2}{4+V_1(V_2-1)-V_2}$, there are equilibria as in the environments of propositions 3.i and 3.ii.
- For the parameters $(V_1, V_2) \in [2-r, \infty) \times [2-r, 2]$, $V_3 \leq \frac{3V_1V_2}{3V_1+V_2+r(4-V_2)-4}$, there is equilibrium as in the environment of proposition 3.iii.

If $V_3 < 1$, these are all the equilibria.

The values for V_3 in the regions of the equilibria regimes in the above proposition follow from postulating the player 1's and 2's behavior to be as in the results of the section 2, and making it unprofitable for the player 3 to bid anywhere above 0.

Note that when the valuation of the third player is below the bid cap, $V_3 < 1$, the analysis is essentially the same as with the two bidders: the third bidder is always inactive.

The case of $V_1 \geq V_2 = V_3 = v$ with $v \in [r, 1]$ leads to the multiplicity of equilibria: two bidders, 1 and 2 are always active in (r, v) , while the third player joins the bidding in (\tilde{b}, v) , $\tilde{b} \in [r, v]$.⁶ The indeterminacy of \tilde{b} is what creates the multiplicity of equilibria.⁷

Proposition 4.ii. *Suppose $V_1 \geq V_2 = V_3 = v$, and $v \in [r, 1)$. Then, for each $\tilde{b} \in [r, v]$ there is an equilibrium where players bid according to the CDFs:*

$$- G_1(b) = \begin{cases} 0, & \text{if } b < r \\ \frac{b}{v} \times \sqrt{\frac{V_1}{V_1+b-v}}, & \text{if } b \in [r, \tilde{b}) \\ \frac{b}{v} \times \sqrt{\frac{V_1}{V_1+b-v}}, & \text{if } b \in [\tilde{b}, v] \end{cases}, \quad - G_2(b) = \begin{cases} \frac{r+V_1-v}{\sqrt{V_1(\tilde{b}+V_1-v)}}, & \text{if } b \in [0, r) \\ \frac{b+V_1-v}{\sqrt{V_1(\tilde{b}+V_1-v)}}, & \text{if } b \in [r, \tilde{b}) \\ \frac{\sqrt{b+V_1-v}}{\sqrt{V_1}}, & \text{if } b \in [\tilde{b}, v] \end{cases}$$

⁶Players 2 and 3 can switch their roles.

⁷Such multiplicity of equilibria is of the same nature as the one in Baye et al. (1996).

$$- G_3(b) = \begin{cases} \frac{\sqrt{\tilde{b}+V_1-v}}{\sqrt{V_1}}, & \text{if } b \in [0, \tilde{b}) \\ \frac{\sqrt{b+V_1-v}}{\sqrt{V_1}}, & \text{if } b \in [\tilde{b}, v] \end{cases}.$$

If $V_1 = V_2$, player 1 can have an atom at 0, with the total mass of 0 and r being $\frac{r}{v} \times \sqrt{\frac{V_1}{V_1+\tilde{b}-v}}$. These CDFs describe all equilibria for such set of parameters.

Overall, in this case the higher value bidder has an atom at the reserve price, while the other two bidders have atoms at the 0. When two players are active in the interior, their bids are uniformly distributed, and when three players are active the bids are distributed in such way, that the maximum of any two bids is distributed uniformly.

The proof for this case is located in C.2.

For the case of the second- and third-order valuations still being equal to each other and taking medium values, the distributions of bids are similar to the case of results on Proposition 4.ii, but now there appears an atom at the bid cap, 1. There is no explicit way to express the equilibrium objects, namely the bidder one's equilibrium payoff and the upper bound of the continuous part of the support. They follow from the system of non-linear equations.

Proposition 4.iii. *For the set of valuations $V_1 \gtrless V_2 = V_3 = v$, $V_1 \geq 1$, $v \in [1, 3]$, there is a non-empty subset, such that equilibrium CDFs are*

$$\begin{aligned} G_1(b) &= \frac{b}{v} \times \sqrt{\frac{V_1}{\tilde{b}+U}} \mathbb{I}_{\{b \in [r, \tilde{b})\}} + \frac{b}{v} \times \sqrt{\frac{V_1}{b+U}} \mathbb{I}_{\{b \in [\tilde{b}, \hat{b})\}} + \frac{\hat{b}}{v} \times \sqrt{\frac{V_1}{\hat{b}+U}} \mathbb{I}_{\{b \in [\hat{b}, 1)\}} + \mathbb{I}_{\{b \geq 1\}} \\ G_2(b) &= \frac{r+U}{\sqrt{V_1(\tilde{b}+U)}} \mathbb{I}_{\{b \in [0, r)\}} + \frac{b+U}{\sqrt{V_1(\tilde{b}+U)}} \mathbb{I}_{\{b \in [r, \tilde{b})\}} + G_3(b) \mathbb{I}_{\{b \geq \tilde{b}\}} \\ G_3(b) &= \frac{\sqrt{\tilde{b}+U}}{\sqrt{V_1}} \mathbb{I}_{\{b \in [0, \tilde{b})\}} + \frac{\sqrt{b+U}}{\sqrt{V_1}} \mathbb{I}_{\{b \in [\tilde{b}, \hat{b})\}} + \frac{\sqrt{\hat{b}+U}}{\sqrt{V_1}} \mathbb{I}_{\{b \in [\hat{b}, 1)\}} + \mathbb{I}_{\{b \geq 1\}} \end{aligned}$$

where U , player 1's positive equilibrium rent, and \hat{b} follow from the system of

equations:

$$\begin{cases} \frac{1}{3} + \frac{1}{3}\sqrt{\frac{U+\hat{b}}{V_1}} + \frac{1}{3}\frac{U+\hat{b}}{V_1} & = \frac{1+U}{V_1} \\ \frac{1}{3}\sqrt{\frac{U+\hat{b}}{V_1}} + \frac{1}{6}\frac{\hat{b}}{v} + \frac{1}{6}\frac{U+\hat{b}}{V_1} + \frac{1}{3}\frac{\hat{b}}{v}\sqrt{\frac{U+\hat{b}}{V_1}} & = \frac{1}{v}\sqrt{\frac{U+\hat{b}}{V_1}}. \end{cases} \quad (\text{A})$$

Moreover, for a non-empty subset of valuations, two solutions, (U, \hat{b}) , to the above system, constitute equilibrium objects. If $U = 0$, player 1 can have an atom at 0, with the total mass of 0 and r being $\frac{r}{v}\sqrt{\frac{V_1}{\hat{b}}}$.

In the appendix C.3 I derive the algebraic expressions for the regions of valuations, such that the type of equilibrium from the Proposition 4.iii holds. Here I demonstrate the results with several numerical examples.

Let $r = 0.03$, $V_1 = 2.56$, $V_2 = V_3 = 1.5$. Then, the solution to system A that satisfies the equilibrium requirements $U \geq 0$, $\hat{b} \geq r$ is, approximately,

$$U = 1.07954, \hat{b} = 0.5541.$$

Note that since one of the players 2 or 3 can join the bidding at any $\tilde{b} \in [r, \hat{b}]$, which causes the multiplicity of equilibria. We can plot equilibrium CDFs for this type of equilibria for some choice of \tilde{b} .




Figure 5: Distributions of bids in the environment of 4.iii

Consider now different valuations of players 2 and 3 $V_2 = V_3 = 2.57$, fixing

$r = 0.03$ and $V_1 = 2.56$, as before. The solutions to system A that satisfy the equilibrium conditions are, approximately:

$$\begin{cases} U^1 &= 0.103775, \hat{b}^1 = 0.04028 \\ U^2 &= 0.009232, \hat{b}^2 = 0.05451. \end{cases}$$

Notice that this example demonstrates that under some parameters a slightly disadvantaged player 1 can have a positive equilibrium rent, and moreover, some parameters can lead to two classes of equilibria.⁸

Comparative statics of equilibrium U and \hat{b} with respect to v is not trivial due to non-linearity of the condition of system A and the fact that there can be two pairs (U, \hat{b}) that satisfy the equilibrium condition. On the figure 6, we plot the graphs of U and \hat{b} with respect to the value of v , given $V_1 = 2.5$, $r = 0.01$; and compare them with the behavior of U and \hat{b} for the case of players 1 and 2 being active (i.e. the case of $V_3 = 0$). Note the multiplicity of equilibria for v high enough.




Figure 6: **Comparative statics of U and \hat{b} in 4.iii**

Proposition 4.iv. *There exists a non-empty subset of $(V_1, V_2, V_3) \in (1, +\infty)$ such that there is an equilibrium, where player 1 (2) has an atom at r (0), bids continuously in (r, \hat{b}) , $\hat{b} > r$, and has an atom at 1, while player 3 only bids 0 and 1 with non-zero probabilities. Player 1 has a non-negative equilibrium payoff U . The equilibrium objects (U, \hat{b}, P_3) , where P_3 is the probability of player 3 bidding*

⁸There are multiple equilibria within each class, because player 3 can join the continuous bidding anywhere in (r, \hat{b}) , and that also affects the behavior of players 1 and 2. Each class is represented by a single solution (U, \hat{b}) to the system A.

0, follow from the system

$$\begin{cases} \frac{1}{3}V_1 + \frac{1}{6}P_3V_1 + \frac{1}{6}\frac{\hat{b}+U}{P_3} + \frac{1}{3}(\hat{b}+U) & = 1+U \\ \frac{1}{3}V_2 + \frac{1}{6}P_3V_2 + \frac{1}{6}\frac{\hat{b}}{P_3} + \frac{1}{3}\hat{b} & = 1 \\ \frac{1}{3}V_3 + \frac{1}{6}\frac{\hat{b}+U}{V_1P_3} + \frac{1}{6}\frac{\hat{b}}{V_2P_3} + \frac{1}{3}\frac{(\hat{b}+U)\hat{b}}{V_1V_2P_3^2} & = 1 \end{cases} \quad (\text{B})$$

If $U = 0$, player 1 can have an atom at 0, arbitrarily splitting the mass between 0 and r .

In the appendix C.4 I describe how to establish the region of parameters, in which this type of equilibrium holds. Here I provide examples of parameters for this type of equilibria to demonstrate that such region is non-empty.

For example, for the parameters $V_1 = 2.7$, $V_2 = 2$, $V_3 = 2.66$, $r = 0.07$, the solution to system B, that satisfies equilibrium conditions, is (approximately) $U = 0.729137$, $\hat{b} = 0.165146$, $P_3 = 0.72$. Note that for this solution, the values of CDFs at \hat{b} are less than one, $G_1(\hat{b}) = \frac{\hat{b}}{V_2P_3} = 0.114652$, $G_2(\hat{b}) = \frac{\hat{b}+U}{V_1P_3} = 0.4599$; and that player 3 does not have incentives to bid anywhere in in (r, \hat{b}) :

$$\begin{aligned} V_3G_1(b)G_2(b) - b &= V_3 \times \frac{b}{V_2P_3} \times \frac{b+U}{V_1P_3} - b \\ &= b \left(V_3 \frac{1}{V_2P_3} \times \frac{b+U}{V_1P_3} - 1 \right) \leq b \left(V_3 \frac{1}{V_2P_3} \times \frac{\hat{b}+U}{V_1P_3} - 1 \right) = b \times (-0.150715). \end{aligned}$$

Another thing to notice about this example is that player 1's payoff is larger than $V_1 - V_2$ and $V_1 - V_3$. So, the standard payoff result, that under some assumptions in equilibrium the top player's payoff is $V_1 - \max_{i \neq 1} V_i$ does not hold in this environment for this type of equilibrium.

Consider another example of parameters and the correspondent solutions. For $V_1 = 2.62$, $V_2 = 2.621$, $V_3 = 2.875$, $r = 0.01$, there is a couple of solutions that satisfy equilibrium conditions: $U^I = 0.12$, $\hat{b}^I = 0.017578$, $P_3^I = 0.249$; and $U^{II} = 0.00075$, $\hat{b}^{II} = 0.0425$, $P_3^{II} = 0.1454$. Beside the multiplicity of equilibria for these parameters, it is also noteworthy that player 1, the one with positive

rent, has the lowest valuation in this example.

Let us now describe two types of equilibria, where all players are active and bids' distributions are atomic only. The first of such two types is the one, where every players' equilibrium payoff is zero, and they all bid 0 and 1:

Proposition 5.i. *There exists a non-empty subset of $(V_1, V_2, V_3) \in (1, 4)^3$ such that there is an equilibrium, where players bid 0 and 1, only. Player i bids 0 with a probability P_i , as in*

$$\begin{cases} P_1 &= \frac{\sqrt{3V_1(V_2-4)(V_3-4)}}{2\sqrt{(4-V_1)V_2V_3}} - \frac{1}{2} \\ P_2 &= \frac{\sqrt{3V_2(V_1-4)(V_3-4)}}{2\sqrt{(4-V_2)V_1V_3}} - \frac{1}{2} \\ P_3 &= \frac{\sqrt{3V_3(V_1-4)(V_2-4)}}{2\sqrt{(4-V_3)V_1V_2}} - \frac{1}{2}. \end{cases} \quad (C)$$

The region is defined by the conditions $(\cap_i \{0 \leq P_i \leq 1\}) \cap (\cap_i \{V_i \prod_{j \neq i} P_j \leq r\})$.

The probabilities in the above proposition follow from solving the system

$$\begin{aligned} &V_i \left(\mathbb{P}\{\text{bid}_j = 0\} \mathbb{P}\{\text{bid}_k = 0\} + \frac{1}{2}(1 - \mathbb{P}\{\text{bid}_j = 0\}) \mathbb{P}\{\text{bid}_k = 0\} + \right. \\ &\left. + \frac{1}{2}(1 - \mathbb{P}\{\text{bid}_j = 0\}) \mathbb{P}\{\text{bid}_k = 0\} + \frac{1}{3}(1 - \mathbb{P}\{\text{bid}_j = 0\})(1 - \mathbb{P}\{\text{bid}_k = 0\}) \right) - 1 = 0, \\ &(i, j, k) \in \{1, 2, 3\}, i \neq k \neq j, \end{aligned}$$

with respect to $\mathbb{P}\{\text{bid}_i = 0\} \doteq P_i$. There are two solutions to that system, but only one allows for joint possibility of $0 \leq P_i \leq 1$. The restrictions, defining the region, follow from feasibility of probabilities and absence of profitable deviation to bidding $b \in (0, 1)$.

Another type of equilibrium with atomic distributions is the one, where one player has a positive equilibrium payoff and bids r instead of 0:

Proposition 5.ii. *There exists a non-empty subset of $(V_1, V_2, V_3) \in (1, 4)^3$ such that there is an equilibrium, where player 1 bids r and 1, with probabilities P_1 and $1 - P_1$, and has a payoff $U \geq r$; and players 2,3 bid 0 and 1, only. Player $i \in \{2, 3\}$*

bids 0 with a probability P_i , as in

$$\begin{cases} P_1 &= \frac{\sqrt{3V_1(V_2-4)(V_3-4)}}{2\sqrt{(4-V_1)V_2V_3}} - \frac{1}{2} \\ P_2 &= \frac{\sqrt{3V_2(V_1-4)(V_3-4)}}{2\sqrt{(4-V_2)V_1V_3}} - \frac{1}{2} \\ P_3 &= \frac{\sqrt{3V_3(V_1-4)(V_2-4)}}{2\sqrt{(4-V_3)V_1V_2}} - \frac{1}{2}. \end{cases} \quad (\text{F})$$

Player 1's payoff is

$$U = 3 - r - \frac{\sqrt{3(4-V_1)V_1(V_2(V_3-2) - 2V_3)}}{2\sqrt{(4-V_2)V_2(4-V_3)V_3}} - \frac{V_1}{2}.$$

The region is defined by the conditions $(\cap_i \{0 \leq P_i \leq 1\}) \cap (\cap_{i \in \{2,3\}} \{V_i \prod_{j \neq i} P_j \leq r\}) \cap \{U \geq 0\}$. If $U = 0$, player 1 is indifferent between r and 0, and can split the mass between these two bids.

Note that after we were to consider all permutations of valuations in all of the above propositions of the section 3, we would have characterized all types of equilibria possible under three players, except for those where all three players are inactive; and those that result in players bidding 1 with certainty.

4 Conclusion

In this paper the equilibria of the all-pay auctions with the reserve price and the bid cap were characterized. It was shown that equilibrium bidding is characterized by atoms at 0, the reserve price, and the cap, as well as continuous bidding between the reserve and the cap. It was shown that multiple equilibria are possible for some parameter ranges. In case of three players, the following interesting features can emerge in equilibria: the player with a slight disadvantage can have the positive payoff; three players with completely different valuations can all actively compete for the single prize; the “winning” player can have different payoffs across different equilibria.

A Appendix

A.1 General Characterization of Equilibria

Let us show that all the equilibria in All-Pay Auctions with a reserve price and a bid cap are follow the characterization in lemma 1. We do so in the series of lemma below.

Lemma A.1. *There is no mass of bids above 1*

Proof. Suppose by way of contradiction that there is at least one player, who places a bid greater than 1 with a non-zero probability. Recall that 1 is the bid cap. Thus, players who place bids above 1 have the same chances of obtaining the item as if they were bidding 1 exactly, but they have to pay more because of higher bids. Thus, bidding above 1 is strictly dominated. \square

Lemma A.2. *There is no mass of bids in $(0, r)$*

Proof. Suppose by way of contradiction that there is at least one player i , for whom $\mathbb{P}\{\text{bid}_i \in (0, r)\} > 0$. Since r is the reserve price, if the actual bid of player i realizes as $b \in (0, r)$, player i does not obtain the object, but has to pay a positive amount of money. Thus, a strategy of player i that involves bidding in the subset of $(0, r)$ with some non-zero probability is dominated by a similar strategy that replaces the bids in $(0, r)$ with the zero-bid, otherwise remaining the same. \square

Lemma A.3. *There are no atoms in $(r, 1)$.*

Proof. Suppose, by way of contradiction, that there is at least one player, i , who in equilibrium has an atom of size $P > 0$ at a bid $b_i \in (r, 1)$. Any other player would not have an atom at b_i , since bidding slightly above increases the costs continuously, but causes a discontinuous upward jump in the probability of winning the object. If nobody else besides i bids anywhere in the region of b_i , i would be better off lowering the b_i bid. If there is a player $k \neq i$ who bids with a non-zero probability in the interval $(b_i - \varepsilon, b_i)$, we can choose a small $\varepsilon' \in (0, \varepsilon)$, such that

replacing any bid from $(b_i - \varepsilon', b_i)$ to $b_i + \varepsilon$ increases the expenditures by at most $2\varepsilon'$ but increases the probability of winning the object by at least P . We are free to choose ε' so that the net change of k 's payoff is positive. Thus, k would have a gap in bids below b_i . This again would cause i to decrease b_i \square

Lemma A.4. *There can be no more than one atom at r*

Proof. Suppose, by way of contradiction, that two or more players have atoms at r . Then, any of them would be better off by bidding slightly higher than r . \square

Let us introduce the following notation $\bar{b} \doteq \sup\{\cup_i BR_i \cap (r, 1)\}$, $\underline{b} \doteq \inf\{\cup_i BR_i \cap (r, 1)\}$. Then, the following result holds

Lemma A.5. *The set $\cup_i\{BR_i \cap (\underline{b}, \bar{b})\}$ is closed relative to (\underline{b}, \bar{b}) .*

Proof. If the set $\cup_i\{BR_i \cap (\underline{b}, \bar{b})\}$ is empty, the result holds trivially. If, on the contrary, $\cup_i\{BR_i \cap (\underline{b}, \bar{b})\}$ is non-empty, it means there are at least two players who bid in the interior of (\underline{b}, \bar{b}) (it cannot be that only one player bids there, since for him it doing so would increase the expenditures, but not increase the probability of winning). Consider any player $k \in \{1, 2, 3\}$, for whom $BR_k \cap (\underline{b}, \bar{b}) \neq \emptyset$. For all bids b in $BR_k \cap (\underline{b}, \bar{b}) \neq \emptyset$ we can write down $V_k \times \prod_{j \neq k} G_j(b) - b = U_k$, where G_j is a respective CDF of player j 's bids, and U_k is k 's equilibrium payoff. U_k is a point in \mathbb{R} , hence, a closed set. For bids b strictly greater than r , $G_j(b)$ is a continuous function for all $j \in \{1, 2, 3\}$, due to the absence of atoms. Thus, $V_k \times \prod_{j \neq k} G_j(b) - b$ is also a continuous function. For a continuous set, the pre-image of a closed set is a closed set. This establishes the claim of the lemma above. \square

Lemma A.6. *The set $\cup_i\{BR_i \cap (\underline{b}, \bar{b})\}$ is an interval.*

Proof. Let $\cup_i\{BR_i \cap (\underline{b}, \bar{b})\}$ be non-empty. Suppose, by way of contradiction, that there is a point b^* in (\underline{b}, \bar{b}) that does not belong to a Best Response Set of any of the players. By $\cup_i\{BR_i \cap (\underline{b}, \bar{b})\}$ being closed relative to (\underline{b}, \bar{b}) , there is an open interval, that contains b^* , any point of which is not in any of the Best Response Sets. Pick

the largest such interval and denote it $(\underline{b}^*, \overline{b}^*)$. \overline{b}^* must be in the Best Response Set of some player. However, since nobody bids in $(\underline{b}^*, \overline{b}^*)$, the probability of winning the item from bidding \overline{b}^* is the same as from bidding \underline{b}^* , while \overline{b}^* is more costly. This makes \overline{b}^* less profitable than \underline{b}^* , leading to a contradiction. \square

Lemma A.7. *If $\cup_i \{BR_i \cap (\underline{b}, \overline{b})\} = (\underline{b}, \overline{b})$ is non-empty, all players have atoms in the set $\{0, r\}$. Moreover, there are at least two players having an atom at 0.*

Proof. Suppose, by way of contradiction, that there is no mass in $\{0, r\}$. That would mean, by absence of atoms in $(r, 1)$, that probability of winning the item from bidding \underline{b} is zero, while the expenditures are strictly positive. That, in turn results in a negative payoff for any player from bidding \underline{b} , thus contradicting to \underline{b} being in a Best Response set. The fact that at least two players have atoms at 0 therefore follows from the Lemma A.4. \square

Lemma A.8. *For players i and k , $i \neq k$, who have atoms at 0 $G_i(b) \times V_k = G_k(b) \times V_i$, for $b \in (\underline{b}, \overline{b})$, being a point of increase of both G_i and G_k*

Proof. Players, who have atoms at 0, have an equilibrium payoff of 0. A point of an increase of a CDF of a player must yield him an equilibrium payoff. So, for the two players, i and k , it holds that

$$V_i G_j(b) G_k(b) - b = 0$$

$$V_k G_j(b) G_i(b) - b = 0,$$

from which the assertion of the lemma follows, given that $G_j(b) > 0$. \square

Lemma A.9. *For a player, who has an atom at zero, his CDF is increasing on an interval $(a, \tilde{b}) \subseteq (a, \overline{b})$, $a \geq \underline{b}$, this CDF is increasing on (a, \overline{b}) .*

Proof. Denote by i the player who has an increasing CDF on (a, \tilde{b}) and an atom at zero. Suppose, by way of contradiction, that the CDF of player i is constant on $(\tilde{b}, c) \subseteq (\tilde{b}, \hat{b})$, $c > \tilde{b}$. By the Lemma A.6, there is always a pair of players bidding

in (\tilde{b}, c) , and by the Lemma A.7, at least one of them has an atom at zero, and, hence, a zero equilibrium payoff. Call that player k . From the Lemma A.8, it holds that $G_i(\tilde{b})V_k = G_k(\tilde{b})V_i > 0$. Let $\varepsilon > 0$ be such that at $\tilde{b} + \varepsilon$ player k is still bidding. Then, it must be that for any $x \in (\tilde{b}, \tilde{b} + \varepsilon)$, the payoff of i from bidding such x would be weakly lower than the payoff of k :

$$V_i \prod_{j \neq i} G_j(x) - x \leq V_k \prod_{j \neq k} G_j(x) - x \Rightarrow V_i G_k(x) \leq V_k G_i(x),$$

which, together with the Lemma A.8 contradicts to G_i being constant on $(\tilde{b}, \tilde{b} + \varepsilon)$ and G_k being strictly increasing there. \square

Lemma A.10. *If (r, \bar{b}) is non-empty, at least one player, who has an atom at zero, randomizes continuously on (r, \bar{b}) .*

Proof. A different way to write the statement of the Lemma above is $\underline{b} = r$. Otherwise, bidding anywhere in (r, \underline{b}) would be a profitable deviation, since it would yield the same probability of obtaining the item as from bidding \underline{b} , but would incur smaller expenditures. \square

Lemma A.11. *If $\bar{b} > r$, and if a player has an atom at r , this player randomizes continuously on (r, \bar{b}) .*

Proof. Denote the player, who has an atom at r by i . The other two players, j and k , must have atoms at zero and, hence, a zero equilibrium payoff. Denote by $u_i^* \geq 0$ the equilibrium payoff of player i . Suppose, by way of contradiction, that player i rejoins bidding at some $\tilde{b} \in (r, \bar{b})$. This implies that bidding in (r, \tilde{b}) gives him a weakly lower payoff, than u_i^* . Denote by P the mass that i places at r . From players j and k having a zero equilibrium payoff, it follows that in the interval (r, \tilde{b}) , the CDFs of j and k are given by $G_j(b) = \frac{b}{P \times V_k}$, and $G_k(b) = \frac{b}{P \times V_j}$. Above \tilde{b} , in order for equilibrium payoffs to hold, the CDFs of all three players must be given by $G_i(b) = \frac{b\sqrt{V_i}}{\sqrt{V_j V_k (b + u_i^*)}}$, $G_j(b) = \sqrt{\frac{(b + u_i^*)V_j}{V_k V_i}}$, $G_k(b) = \sqrt{\frac{(b + u_i^*)V_k}{V_j V_i}}$. Since there can be no atoms in $(r, 1)$, smooth-pasting must hold: CDFs must be continuous at \tilde{b} .

That allows to pin down the size of i 's atom at r : $P = \frac{\tilde{b}\sqrt{V_1}}{\sqrt{V_j V_k (\tilde{b} + u_i^*)}}$. That, in turn, allows to determine i 's payoff from an arbitrary bid b in $[r, \tilde{b}]$:

$$V_i G_j(b) G_k(b) - b = V_i \frac{b^2}{P^2 V_k V_j} - b = V_i \frac{b^2 (\tilde{b} + u_i^*)}{\tilde{b}^2 V_i} - b = \frac{b^2 (\tilde{b} + u_i^*)}{\tilde{b}^2} - b.$$

If we plug in $b = r$ into the expression above, we get that the payoff from bidding r is equal to u_i^* , i 's equilibrium payoff, only if $\tilde{b} = r$. Otherwise, the payoff from bidding r is lower, and player i would not be willing to put an atom at r , contradicting the condition of the lemma. \square

B Appendix

B.1 Completeness of characterization in 1.i and 1.ii.

For the case $V_1 > V_2$, recall the general characterization lemma 1, and also apply the following steps:

- There is no mass strictly above V_2 , as player 2 won't ever bid strictly above his valuation.
- 0 is never in a Best Response set of player 1, since he can secure a positive payoff by bidding slightly above V_2 .
- player 2 has 0 in the Best Response set.
- A strategy profile according to which player 1 bids r with a unit probability is not an equilibrium. If it was an equilibrium, player 2 would always bid slightly above r outbidding player 1. However, player 1 can always make sure he has a positive payoff.
- $\mathbb{P}\{bid_1 > r\} > 0$. Follows from the point above.
- $\mathbb{P}\{bid_2 \geq r\} > 0$. Otherwise, player 1 wouldn't bid above r .

- In equilibrium for both players the subset $BR \cap (r, V_2)$ is closed relative to (r, V_2) . As follows from lemma 1.

- $BR_j \cap (r, V_2)$ is an interval for both players, which also follows from lemma 1.

Denote $\inf(BR_i \cap (r, V_2)) = \underline{b}$, and $\sup(BR_i \cap (r, V_2)) = \bar{b}$.

- $\bar{b} = V_2$.

Suppose that $\bar{b} < V_2$. Then player 2 can place a bid somewhere between \bar{b} and V_2 , win with probability 1 and have a positive payoff. This contradicts to 0 being in his best response set, which was established above.

- For players 1 and 2 distributions of their best responses on (\underline{b}, V_2) have densities and are equal to $\frac{1}{V_2}$ and $\frac{1}{V_1}$, respectively.

For a bid $x \in (\underline{b}, V_2)$ of player j it must hold that $V_j \times \mathbb{P}_{\{bid_k < x\}} - x = u$, where u is player-specific equilibrium payoff. So, for $x, x' : \underline{b} \leq x < x' \leq V_2$ it holds that $V_j \times \mathbb{P}_{\{bid_k < x\}} - x = V_j \times \mathbb{P}_{\{bid_k < x'\}} - x' \Rightarrow \mathbb{P}_{\{bid_k < x'\}} - \mathbb{P}_{\{bid_k < x\}} = (x' - x)/V_j \Rightarrow \lim_{x' \rightarrow x+0} \frac{\mathbb{P}_{\{bid_k < x'\}} - \mathbb{P}_{\{bid_k < x\}}}{x' - x} = 1/V_j$, which establishes existence and expression for the right derivative, with similar logic applying to the left derivative.

- Player 1 has an atom at r .

Otherwise, player 1 would not be able to reach a unit value of her/his CDF by V_2 .

- $\underline{b} = r$, as follows from lemma 1.

- Player 2 has an atom at 0.

In order for him to reach a unit value of CDF by V_2 .

The sizes of atoms for both players are uniquely pinned down.

For $V_1 = V_2$ the following steps apply:

- There is no mass in $(0, r)$
- There is no mass above $V_1 = V_2 = v$.
- There are no atoms above r .
- There can be no two atoms at r .
- $\min_i \inf(BR_i) = 0$
- In equilibrium for both players the subset $BR \cap (r, V)$ is closed relative to (r, v) .
- $BR \cap (r, v)$ is non-empty. Otherwise, the player with 0 in his best response would be able to have a positive payoff.
- $BR_i \cap (r, v)$ is an interval for both players.
- $BR_1 \cap (r, v) = BR_2 \cap (r, v)$.
- $\sup(BR_1 \cap (r, v)) = \sup(BR_2 \cap (r, V)) = v$. Otherwise, the player with 0 in his best response would be able to have a positive payoff.
- PDFs of distributions of bids of both players in the interval $BR \cap (r, V_2)$ are equal to $\frac{1}{v}$.
- $\inf(BR \cap (r, V)) = r$.
- The fact that CDFs have to reach value of 1 at v , and the value of densities in (r, V) pin down that $\mathbb{P}[bid_1 \in \{0, r\}] = \mathbb{P}[bid_2 \in \{0, r\}] = \frac{r}{v}$. Absence of two atoms at r means also that $\mathbb{P}\{bid_1 = r\} \times \mathbb{P}\{bid_2 = r\} = 0$.

B.2 Completeness of characterization in propositions 2.i and 2.ii

Start with the case of $\rho_2^2 V_2 - \rho_2 < \rho_1^2 V_1 - \rho_1$:

- There can be no equilibria such that at least one of the players bids 1 with a unit probability.

If player 1 always bids 1 the only way for player 2 to win the prize is to also bid 1, which will result in player 2 getting the prize with probability ρ_2 . This in turn will lead to player 2's payoff of $\rho_2 \times V_2 - 1 \leq \rho_2(1 - \rho_1) - 1 < 0$. Thus player 2 will bid 0 in response to player 1's bid of 1, hence player 1 has incentives to lower his bid. If player 2 always bids 1, depending on player 1's valuation, he would always bid 0 or 1. In both cases player 2 would be better off lowering his bid (by a small amount, or down to 0, respectively).

- There can be no two atoms at r .
- $\min_{i \in \{1,2\}}(\inf(BR_i)) = 0$.
- A profile such that both players bid only a 0 and a 1 with some probabilities is not an equilibrium. Suppose, by contradiction, there is such an equilibrium profile. It would result in $\mathbb{P}\{\text{bid}_i = 0\} = \frac{1 - \rho_k V_k}{V_k \rho_i}$, which results in profitable deviation to bidding r for any of players.
- For the similar reason, a profile such that one player bids r and 1 only, and the other bids 0 and 1 only, is not an equilibrium. Thus, in equilibrium, the probability of a bid in $(r, 1)$ is greater than zero.
- $BR_i \cap (r, 1)$ is an interval for both players.
- In an equilibrium $BR_1 \cap (r, 1) = BR_2 \cap (r, 1)$, and $\inf\{BR_i \cap (r, 1)\} = r$.
- Densities for players 1 and 2 in interval $(r, \sup(BR \cap 1))$ are equal to $\frac{1}{V_2}$ and $\frac{1}{V_1}$, respectively.
- Either $\sup(BR_1 \cap (r, 1)) = \sup(BR_2 \cap (r, 1)) = 1$ or $\mathbb{P}\{\text{bid}_j = 1\} > 0$ for both players. If it wasn't so, the player with 0 in his best response would be able to achieve a positive payoff by bidding 1 with a higher frequency.

- $\sup(BR_1 \cap (r, 1)) = \sup(BR_2 \cap (r, 1))$ is equal either to $1/\rho_2 - (\rho_1/\rho_2)V_1$ or to $1/\rho_1 - (\rho_2/\rho_1)V_2$. Let player j have 0 in his best response. Then \bar{b} is determined by him being indifferent between bidding 1 and 0, and \bar{b} and 0:

$$V_j \mathbb{P}\{bid_k < \bar{b}\} - \bar{b} = V_j (\rho_j(1 - \mathbb{P}\{bid_k < \bar{b}\}) + \mathbb{P}\{bid_k < \bar{b}\}) - 1 = 0.$$

- Only one player has 0 in his best response. Both of them cannot be indifferent between 0 and $1/\rho_2 - (\rho_1/\rho_2)V_1$, or 0 and $1/\rho_1 - (\rho_2/\rho_1)V_2$.
- In equilibrium it is player 2 who has 0 in his best response. Had it been player 1 who has 0 in a best response, \bar{b} would be $1/\rho_2 - (\rho_1/\rho_2)V_1$, from 1's indifference. Then, condition on 1's density and 2's indifference between $1/\rho_2 - (\rho_1/\rho_2)V_1$ and 1 would lead to such values of 1's cdf in $(r, 1/\rho_2 - (\rho_1/\rho_2)V_1)$, that would give a second player a negative payoff from bidding there. He would be better off not including $(r, 1/\rho_2 - (\rho_1/\rho_2)V_1)$ into his BR as he can always secure a payoff of 0.
- $(BR_1 \cap (r, 1)) = (BR_2 \cap (r, 1)) = (r, 1/\rho_1 - (\rho_2/\rho_1)V_2)$.
- Values of 1's CDF at $(r, 1/\rho_1 - (\rho_2/\rho_1)V_2) \cup 1$ are pinned down by 2's indifference between 0 and $(r, 1/\rho_1 - (\rho_2/\rho_1)V_2) \cup 1$, and by 1's PDF; hence, 1's atom at r is uniquely defined.
- Values of 2's CDF at $(r, 1/\rho_1 - (\rho_2/\rho_1)V_2) \cup 1$ are pinned down by 1's indifference across $[r, 1/\rho_1 - (\rho_2/\rho_1)V_2) \cup 1$, and by 2's PDF; hence, 2's atom at 0 is uniquely defined.

The case $\rho_2^2 V_2 - \rho_2 = \rho_1^2 V_1 - \rho_1$ is similar, but now the upper bound of interior bidding, $\sup(BR \cap (r, 1)) = 1/\rho_1 - (\rho_2/\rho_1)V_2$ and $1/\rho_1 - (\rho_2/\rho_1)V_2 = 1/\rho_2 - (\rho_1/\rho_2)V_1$. This makes both players indifferent between bidding at 0 and at 1, resulting in a zero payoff and multiplicity of equilibria due to unidentified value of atom at r ; and the identity of the player who would bid r .

C Proofs for the Three Bidders Case

C.1 Proof of completeness of 4.i, for $V_3 < 1$

We know from the Proposition 1 that all the equilibria are characterized by atoms at $\{0\}$, $\{r\}$, and $\{1\}$, and also by continuously distributed mass in (r, \bar{b}) . Since $V_3 < 1$, player three will never have mass at 1. His equilibrium strategy, therefore, should be characterized by atoms at $\{0\}$, $\{r\}$, and mass in (\underline{b}, \bar{b}) . Moreover, if $\mathbb{P}\{\text{bid}_3 = r\} > 0$ and $\mathbb{P}\{\text{bid}_3 \in (\underline{b}, \bar{b})\} > 0$, then it must be that $\underline{b} = r$. Let us consider the following possibilities: (i) player 3 has mass at $\{0\} \cup (\underline{b}, \bar{b})$, $\underline{b} \geq r$; (ii) player 3 has mass at $\{r\} \cup (r, \bar{b})$ and, possibly, mass at $\{0\}$; (iii) player 3 has mass at $\{r\}$ and, possibly, mass at $\{0\}$.

(i) Since player 3 is willing to bid 0, his equilibrium payoff must also be zero.

So, for bids $b \in (\underline{b}, \bar{b})$, we can write down:

$$V_3 G_1(b) G_2(b) - b = 0.$$

If $\underline{b} > r$, both of the other two players must be active in (r, \bar{b}) . So, for $b \in (\underline{b}, \bar{b})$ the following system must hold:

$$V_1 G_2(b) G_3(b) - b = u_1^*$$

$$V_2 G_1(b) G_3(b) - b = u_2^*$$

$$V_3 G_1(b) G_2(b) - b = 0,$$

where $u_i^* \geq 0$ are the equilibrium payoffs with $u_1^* \times u_2^* = 0$. If it is the case that $u_2^* = 0$, then it must be that in the region (\underline{b}, \bar{b}) , $G_2(b) = \sqrt{\frac{(b+u_1^*)V_2}{V_1 \times V_3}}$, $G_3(b) = \sqrt{\frac{(b+u_1^*)V_3}{V_1 \times V_2}}$. G_3 must reach the value of 1 at \bar{b} , so we get that $\bar{b} = \frac{V_1 \times V_2}{V_3} - u_1^*$. But evaluating G_2 at $b = \bar{b}$ gives $G_2(\bar{b}) = \frac{V_2}{V_3} > 1$, a contradiction. For the case $u_1^* = 0$ the symmetric argument holds.

If $\underline{b} = r$, and only one of the other two players is active in (r, \bar{b}) , let player

1 be active in (r, \bar{b}) . Let the total mass that player 2 has at $\{0\}$ and/or $\{r\}$ be P_2 . For bids in (r, \bar{b}) it must hold that

$$\begin{aligned} V_1 P_2 G_3(b) - b &= u_1^* \\ V_3 P_2 G_1(b) - b &= 0 \\ &\Rightarrow \\ G_3(b) &= \frac{b + u_1^*}{P_2 V_1}, G_1(b) = \frac{b}{V_3 P_2}. \end{aligned}$$

G_3 must reach the value of 1 at \bar{b} , so $\bar{b} = P_2 V_1 - u_1^*$. Check the value of G_1 at \bar{b} : $G_1(\bar{b}) = \frac{V_1}{V_3} - \frac{u_1^*}{V_3 P_2}$. Since $V_1 > V_3$, it is necessary that player 1 has a strictly positive payoff, $u_1^* > 0$. Moreover, in order for the probability masses of bids of players 1 and 2 to be completely spent and equal to one, players 1 and 2 must have atoms at $b = 1$. Checking that their incentives to bid at 1 are satisfied, we have

$$\begin{aligned} V_1 \left(\frac{1}{2} + \frac{1}{2} P_2 \right) - 1 &= u_1^* \\ V_2 \left(\frac{1}{2} + \frac{1}{2} G_1(\bar{b}) \right) - 1 &= 0. \end{aligned}$$

Using these conditions to solve for u_1^* , P_2 and \bar{b} , we have

$$\begin{aligned} P_2 &= \frac{(V_1 - 2)V_2}{V_1 V_2 - 2(2 - V_2)V_3}, & u_1^* &= \frac{(V_1 - 2)(V_1 V_2 - (2 - V_2)V_3)}{V_1 V_2 - 2(2 - V_2)V_3} \\ G_1(\bar{b}) &= \frac{2 - V_2}{V_2}, & \bar{b} &= \frac{(V_1 - 2)(2 - V_2)V_3}{V_1 V_2 - 2(2 - V_2)V_3}. \end{aligned}$$

For the objects to correspond to an equilibrium, it is necessary that $V_2 < 2$. Let us check that player two is weakly worse off from bidding in (r, \bar{b}) : his payoff from bidding $b = \bar{b}$ is:

$$2 - V_2 - \frac{(V_1 - 2)(2 - V_2)V_3}{V_1 V_2 - 2(2 - V_2)V_3} \leq 0.$$

If $V_1 > 2$ and $V_1V_2 - 2(2 - V_2)V_3 > 0$, the above condition implies $V_1V_2 - 2(2 - V_2)V_3 < V_1V_3 - 2V_3$, which is not true, given the condition $V_2 > 1 > V_3$. If it is the case that $V_1 < 2$ and $V_1V_2 - 2(2 - V_2)V_3 < 0$, it can be seen that these two conditions are incompatible with \bar{b} being weakly lower than V_3 (the latter being necessary for player 3 to be willing to bid up to \bar{b}).

(ii) The other two players must have atoms at zero, and, therefore, zero payoffs. \bar{b} cannot be greater than V_3 . If there are no atoms at 1, either player one or player two could be better off by bidding slightly below their valuation and winning the item with certainty. If there are atoms at 1, arguments similar to part (i) of this proof apply.

(iii) In order for player 3 to be willing to bid r , both of the other two players must have mass at 0 and zero payoffs. It is not an equilibrium for any of them to bid 0 with probability one, since that would lead to possibility of profitable deviations. Player 3 must have a non-negative payoff from bidding r , so $\mathbb{P}\{\text{bid}_1 = 0\}\mathbb{P}\{\text{bid}_2 = 0\}V_3 = r + u_3^*$, $u_3^* \geq 0$. If there is bidding in $(r, \bar{b}) \neq \emptyset$, 1's and 2's indifference with the fact that 3's mass is spent completely on r , by assumption, give

$$G_1(b) = \frac{b}{V_2},$$

$$G_2(b) = \frac{b}{V_1}.$$

Together with absence of atoms in (r, \bar{b}) this means that $\mathbb{P}\{\text{bid}_1 = 0\} = \frac{r}{V_2}$, and $\mathbb{P}\{\text{bid}_2 = 0\} = \frac{r}{V_1}$. Then, 3's payoff from bidding r is $\frac{r^2}{V_1V_2}V_3 - r$, which is less than zero, given $r < 1$, $V_1 > V_3$, $V_2 > V_3$ and $\min_{i \in \{1,2,3\}}\{V_i\} > r$. This contradicts 3's willingness to bid r .

If there is no bidding in (r, \bar{b}) , players 1 and 2 must also have atoms at 1. From 1's and 2's indifference between bidding 0 and 1, we have that $\mathbb{P}\{\text{bid}_1 = 0\} = \frac{2-V_2}{V_2}$, $\mathbb{P}\{\text{bid}_2 = 0\} = \frac{2-V_1}{V_1}$. For players 1 and 2 not to be

willing to bid slightly above r , it must be that $\frac{2-V_2}{V_2}V_1 \leq r$ and $\frac{2-V_1}{V_1}V_2 \leq r$. Then, player 3's payoff from bidding r is $\frac{2-V_2}{V_2}\frac{2-V_1}{V_1}V_3 - r \leq \frac{r}{V_1}\frac{r}{V_2}V_3 - r < 0$, which contradicts player 3's willingness to bid r .

C.2 Proof of Proposition 4.ii

It is straightforward to check that the CDFs described in the Proposition statement do constitute equilibria. Let us, therefore, show that those equilibria are all equilibria possible for the range of parameters stated in the Proposition. So far we know that all equilibria can be described by atoms at $\{0\}$, $\{r\}$, and $\{1\}$, and continuously distributed mass of bids in (r, \bar{b}) . Let us follow the steps:

- (i) There cannot be any mass above $V_2 = V_3 = v$, continuous, or atoms, as for players 2 and 3 it is more expensive to bid above v , than what they can get from the item if they win it. So, $\bar{b} \leq v$, in all equilibria.
- (ii) Player 1 always has a positive equilibrium payoff: he can always bid slightly above v , which makes his payoff bounded below by $V_1 - v > 0$. This implies that player 1 cannot have an atom at 0, since it implies a zero equilibrium payoff.
- (iii) In equilibrium it is never the case that all the bids concentrate on $\{0\}$ and $\{r\}$. If it was the case, player 2, or player 3 would have a profitable deviation of bidding just slightly above r and winning the item with certainty. Overall, it implies that there is some continuously distributed mass of bids in $(r, \bar{b}) \neq \emptyset$.
- (iv) $\bar{b} = v$, otherwise either player 2, or player 3 would have a profitable deviation.
- (v) Since player 1 has a positive payoff, he must be bidding somewhere in (\underline{b}, v) , $\underline{b} \geq r$. Every point in (r, v) is a point of increase at least for one player. In order to have at least one other player bidding in (r, v) , player 1 must have

an atom at r . This, in turn, implies that for player 1 every point in (r, v) is a point of increase.

(vi) Player 1's equilibrium payoff is equal to $V_1 - v$. This follows from the fact that the CDFs of bids reach the value of 1 by $b = v$, and at least one of player 2's and player 3's CDF reach value of 1 exactly at $b = v$. Since $b = v$ is in the best response set, the payoff of player 1 from bidding v is his equilibrium payoff and is equal to

$$V_1 \times \underbrace{G_2(v) \times G_3(v)}_1 - v.$$

To summarize, what we know so far is that player 1 has an atom at r and a continuously distributed mass in (r, v) and a positive payoff of $V_1 - v$. Players 2 and 3 have atoms at 0. At least one of them has a continuously distributed mass in (r, v) , while the other could re-join bidding in (\tilde{b}, v) , $\tilde{b} \in [r, v]$, or just bid 0 with probability 1. Without loss, we let player 2 bid continuously (r, v) . We, therefore, have the following equilibrium conditions:

$$\begin{aligned} V_1 G_2(b) G_3(0) - b &= V_1 - v \\ V_2 G_1(b) G_3(0) - b &= 0 \\ V_3 G_1(b) G_2(b) - b &\leq 0, \text{ for } b \in [r, \tilde{b}), \\ V_1 G_2(b) G_3(b) - b &= V_1 - v \\ V_2 G_1(b) G_3(b) - b &= 0 \\ V_3 G_1(b) G_2(b) - b &= 0, \text{ for } b \in [\tilde{b}, v]. \end{aligned}$$

Taking into account that there is no mass in $(0, r)$, player 1 has an atom at r , players 2 and 3 have atoms at 0, and solving for G_1, G_2, G_3 following from the conditions above, we get the equilibrium CDFs, stated in the proposition, with the \tilde{b} being the ‘‘degree of freedom’’ of this type of equilibria, and causing their multiplicity.

C.3 Proof of Proposition 4.iii and region characterization

Let us first show how the system A was derived.

In the type of equilibrium, such that player 1 has a positive rent, three players bid continuously somewhere in $(\tilde{b}, \hat{b}) \subseteq [r, 1]$, and also have atoms at 1, the following must hold:

$$\begin{aligned}
V_1 G_2(b_1) G_3(b_1) - b_1 &= U > 0, \forall b_1 \in (\tilde{b}, \hat{b}] \\
V_1 \left(G_2(\hat{b}) G_3(\hat{b}) + \frac{1}{2}(1 - G_2(\hat{b})) G_3(\hat{b}) + \frac{1}{2} G_2(\hat{b})(1 - G_3(\hat{b})) + \right. \\
&\quad \left. + \frac{1}{3}(1 - G_2(\hat{b}))(1 - G_3(\hat{b})) \right) - 1 = U \\
V_2 G_1(b_2) G_3(b_2) - b_2 &= 0, \forall b_2 \in (\tilde{b}, \hat{b}] \\
V_2 \left(G_1(\hat{b}) G_3(\hat{b}) + \frac{1}{2}(1 - G_1(\hat{b})) G_3(\hat{b}) + \frac{1}{2} G_1(\hat{b})(1 - G_3(\hat{b})) + \right. \\
&\quad \left. + \frac{1}{3}(1 - G_1(\hat{b}))(1 - G_3(\hat{b})) \right) - 1 = 0 \\
V_3 G_1(b_3) G_2(b_3) - b_3 &= 0, \forall b_3 \in (\tilde{b}, \hat{b}] \\
V_3 \left(G_1(\hat{b}) G_2(\hat{b}) + \frac{1}{2}(1 - G_1(\hat{b})) G_2(\hat{b}) + \frac{1}{2} G_1(\hat{b})(1 - G_2(\hat{b})) + \right. \\
&\quad \left. + \frac{1}{3}(1 - G_1(\hat{b}))(1 - G_2(\hat{b})) \right) - 1 = 0,
\end{aligned}$$

which follows from the indifference of players across the increase points for continuous bidding, and also the indifference between bidding 1 and \hat{b} , the upper-bound of the continuous bidding of all three players. Since $V_2 = V_3 = v$, it must be that the indifference conditions for players 2 and 3 are identical, and it also means that $G_2(b) = G_3(b)$ for $b \in (\tilde{b}, 1]$. So, after some transformations, we can state that $G_2(b) = G_3(b) = \frac{\sqrt{b+U}}{\sqrt{V_1}}$, $G_1(b) = \frac{b}{v} \times \frac{\sqrt{V_1}}{\sqrt{b+U}}$, for $b \in (\tilde{b}, \hat{b}]$. Since player 2's and 3's indifference conditions are identical, we have two equations instead of three, the condition of players' willingness to bid 1 transforms into

$$\begin{aligned}
V_1 \left(\frac{1}{3} + \frac{1}{3} \frac{\sqrt{U + \hat{b}}}{\sqrt{V_1}} + \frac{1}{3} \frac{U + \hat{b}}{V_1} \right) - 1 &= U \\
v \left(\frac{1}{3} + \frac{1}{6} \frac{\sqrt{U + \hat{b}}}{\sqrt{V_1}} + \frac{\hat{b}}{6v} \times \frac{\sqrt{V_1}}{\hat{b} + U} + \frac{1}{3} \frac{\hat{b}}{v} \right) - 1 &= 0,
\end{aligned}$$

from which it is straightforward to derive system A.

Denoting $t \doteq \sqrt{\frac{U + \hat{b}}{V_1}}$ and solving for t in the first and the second equation of system A, we have

$$\begin{cases} t = \frac{3 + 2U - \hat{b}}{V_1} - 1 \\ t = \frac{\frac{\hat{b}}{6v} + \frac{U + \hat{b}}{6V_1}}{\frac{1}{v} - \frac{\hat{b}}{3v} - \frac{1}{3}}. \end{cases}$$

Since we are interested in finding $U \in \mathbb{R}_+$ and $\hat{b} \in \mathbb{R}_+$, it must be that $t > 0$. So, from the second equation for t it follows that $3 > \hat{b} + v$ must hold. Combining the last inequality with $\hat{b} \geq 0$ it must be that the region for this type of equilibria must be included into $v < 3$.

Setting the two expressions for t equal to each other, we have that

$$\frac{3 + 2U - \hat{b}}{V_1} - 1 = \frac{\frac{\hat{b}}{6v} + \frac{U + \hat{b}}{6V_1}}{\frac{1}{v} - \frac{\hat{b}}{3v} - \frac{1}{3}}.$$

Expressing U in terms of \hat{b} , V_1 , and v ,

$$U = \frac{12\hat{b} - v\hat{b} - V_1\hat{b} - 2\hat{b}^2 + 6V_1 + 6v - 2V_1v - 18}{12 - 4\hat{b} - 5v}.$$

Plugging this expression for U back into the first equation of the system (A), above, we have an equation for \hat{b}

$$\sqrt{\frac{(6 - \hat{b} - 2v)V_1 - 6(1 - \hat{b})(3 - \hat{b} - v)}{(12 - 4\hat{b} - 5v)V_1}} = \frac{2\hat{b}V_1 - v(3 - \hat{b} - V_1)}{(12 - 4\hat{b} - 5v)V_1}. \quad ((B))$$

Raising both sides of the equation to the power of 2, we have a cubic equation in

\hat{b} ,

$$A\hat{b}^3 + B\hat{b}^2 + C\hat{b} + D = 0,$$

where coefficients are

$$A := -24V_1$$

$$B := 9v^2 - 42(v - 4)V_1$$

$$C := -3(12V_1(10 - V_1) - V_1v(68 - 3V_1) + v^2(6 + 8V_1))$$

$$D := 9v^2 + 216V_1 - 162vV_1 + 24v^2V_1 - 72V_1^2 + 54vV_1^2 - 9v^2V_1^2.$$

At least one root of the cubic equations is always a real number. The other two roots are real numbers whenever the cubic discriminant is real. We can check that the root which is always real is greater than 1. Notice that as $\hat{b} \rightarrow +\infty$ in the expression $A\hat{b}^3 + B\hat{b}^2 + C\hat{b} + D$ the term $A\hat{b}^3$ becomes dominant. Since $A = -24V_1$ and V_1 is always positive, we get that $A\hat{b}^3 + B\hat{b}^2 + C\hat{b} + D \rightarrow -\infty$ as $\hat{b} \rightarrow +\infty$. Now, if we plug in the value of $\hat{b} = 1$, we get that $A\hat{b}^3 + B\hat{b}^2 + C\hat{b} + D = A + B + C + D = 9(4 - v)(v - 1)V_1^2$, which is always non-negative given that $v \in (1, 3)$. Hence, the expression $A\hat{b}^3 + B\hat{b}^2 + C\hat{b} + D$ changes the sign, as \hat{b} increases from 1 to infinity. Thus, the root, which is always real, is greater than 1.

So, it is necessary for the kind of equilibrium we are looking for to exist that the cubic discriminant is positive:

$$\mathcal{D} = 18 \times A \times B \times C \times D - 4 \times B^3 \times D + B^2 \times C^2 - 4 \times A \times C^3 - 27 \times A^2 \times D^2 > 0,$$

which can be also simplified to

$$\mathcal{D}^* = -(12v^3 + 49vV_1(8 + 3V_1) - 4v^2(3 + 34V_1) - 4V_1(64 + 3V_1(13 + 6V_1))) > 0. \quad (\mathcal{D}^*)$$




Figure 7: Region for positive value of cubic discriminant

The graph (7) depicts the region of v and V_1 for which the discriminant is positive taking into account also that v must be less than 3. Denote the set of the points $(V_1, v), v \in [1, 3]$ for which the cubic discriminant \mathcal{D} is equal to zero as $\tilde{\Delta}$.

In principle, if there are multiple solutions (U, \hat{b}) to the system (A) that all meet the condition $U \geq 0, 0 \leq r \leq \hat{b} \leq 1$, that would mean that there are multiple equilibria candidates of the type we are looking for. From an observation above, there could be at most two real values of \hat{b} that lie below 1. To each of those two real \hat{b} corresponds one real U , as follows from the equation (B). Since we are interested in such pairs (U, \hat{b}) , that $U \geq 0$, finding pairs (V_1, v) such that $U = 0$ is informative for establishing the region, for which the current type of equilibria holds. Setting $U = 0$ in the system (A), we get that it can hold for $V_1 = v$ and also for $V_1 = 3$ when $\hat{b} = 0$. So, at least one U among the solutions

to the system (A) is positive for $V_1 \geq v$. To see that at least one U from the solution to (A) is positive, note that the expressions there are continuous in all parameters and variables, except for the cases of $v = 0$ or $V_1 = 0$. Then, by setting, for example, $V_1 = 1.75$, $v = 1.5$ numerically solving the system (A) yields approximately $(U, \hat{b}) = (0.263643, 0.569679)$ and $(U, \hat{b}) = (-0.299109, 0.350856)$.

However, it can also be that the lower of U among the two solutions to the system (A) is equal to 0 at $V_1 = v$. If this is the case, this means that there is a subset of $(V_1, v | V_1 < v)$ where there are two solutions to (A) such that $U > 0$, and, hence, two equilibrium candidates.

To find this subset, notice that the two elements U from the solutions coincide when the two respective \hat{b} coincide. The two solutions \hat{b} coincide when the cubic discriminant defined in the condition (\mathcal{D}^*) is equal to zero. Hence in order to pin down the pair of parameters (V_1, v) for which both solutions U are equal to zero, we must intersect the straight line $V_1 = v$ with the $\tilde{\Delta}$, i.e. set of points that make the value of the cubic discriminant \mathcal{D} equal to zero. Plugging $V_1 = v$ into the expression for \mathcal{D}^* we get $\mathcal{D}^*|_{V_1=v} = -(16 - 7V_1)^2 V_1$. This expression is equal to zero for $V_1 = \frac{16}{7}$ and $V_1 = 0$, but the latter is not in the region we are currently interested in. So, for pairs (V_1, v) such that $\{v < 3, V_1 \geq \frac{16}{7}, V_1 \leq v\}$, there are two solutions (U, \hat{b}) to the system (A) such that the element U is positive.

Overall, in terms of non-negativity of U , one of the necessary conditions for a pair of (U, \hat{b}) to be an equilibrium object is that $V_1 \geq v$ for $v \in [1, \frac{16}{7})$, and $\mathcal{D} \geq 0$ for $v \in [\frac{16}{7}, 3]$.

Another restriction for the solution (U, \hat{b}) to correspond to an equilibrium is that $\hat{b} \geq r$, i.e. the upper-bound for the continuous part of bids distribution is above the reserve price. To establish the subset of (V_1, v) for which $\hat{b} \geq r$, first set $\hat{b} = r$ in the system (A) and solve for the U and V_1 . This way we get V_1 expressed

in terms of v and r , for which it holds that $\hat{b} = r$. There are two such V_1 's:

$$V_{1;I,II} = \frac{1}{3(4-v)(2-r-v)} \left(36 - 60r + 4(7-r)r^2 - 27v + (34-7r)rv + 4(1-r)v^2 \right. \\ \left. \pm (1-r)(12-4r-5v)\sqrt{r^2 - r(6-v) + (3-v)^2} \right).$$

The values of $V_{1;I}$ and $V_{1;II}$ coincide for three values of v : $v = \frac{4}{5}(3-r)$, $v = \frac{1}{2} \left(6 - r - \sqrt{3r(4-r)} \right)$, and $v = \frac{1}{2} \left(6 - r + \sqrt{3r(4-r)} \right)$. The last value is greater than 3 for $r > 0$, so it is not among the values of (V_1, v) we are interested in. The values of $(4/5)(3-r)$ and $\frac{1}{2}(6-r-\sqrt{3r(4-r)})$ coincide for $r = \frac{1}{7}$. For $r > \frac{1}{7}$, setting $v = \frac{4}{5}(3-r)$ makes $\sqrt{r^2 - r(6-v) + (3-v)^2}$ not a real number. Thus, for $r \in [0, \frac{1}{7})$, $V_{1;I}$ and $V_{1;II}$ intersect twice, while for $r \in [\frac{1}{7}, 1]$ they intersect once, at $v = \frac{1}{2} \left(6 - r - \sqrt{3r(4-r)} \right)$. Notice that for this value of v the value of $V_{1;I}$ and $V_{1;II}$ is also $\frac{1}{2} \left(6 - r - \sqrt{3r(4-r)} \right)$. This means that the two curves, $V_{1;I}$ and $V_{1;II}$ intersect the second time exactly at the 45-degree line.

Consider the curve $V_{1;I}$. For values of $v < 2-r$, the correspondent values of $V_{1;I}$ are negative, hence, are out of the region of our current interest. There is a discontinuity at $v = 2-r$, and $V_{1;I}$ becomes positive for $v \in \left(2-r, \frac{1}{2} \left(6 - r - \sqrt{3r(4-r)} \right) \right]$. The right limit $\lim_{b \rightarrow 2-r+0} V_{1;I}$ is plus-infinity. This curve crosses the 45-degree line at $v = \frac{8}{3}(1-r)$.

Consider now the curve $V_{1;II}$. Notice that for $v \rightarrow 2-r$ the right limit and the left limit of $V_{1;II}$ coincide, and, using the L'Hospital's rule, are equal to $\frac{3(2-r)^2}{2(2+r)}$.

Overall, the curves $V_{1;I}$ and $V_{1;II}$ intersect twice for $r < \frac{1}{7}$ and once for $r \geq \frac{1}{7}$, at the border of their domain. For $r < \frac{1}{7}$ at $v = 2-r$ $V_{1;I}$ intersects $V_{1;II}$ from above and then reaches $V_{1;II}$. Thus, for $r < \frac{1}{7}$ the union of points lying on the curves $V_{1;I}$ and $V_{1;II}$ geometrically represent a loop.

Recall that curves $V_{1;I}$ and $V_{1;II}$ represent the set of points (V_1, v) for which the element \hat{b} of one of the solutions (U, \hat{b}) to the system (A) is equal to r . Recall also that the set $\tilde{\Delta}$ represents the set of points for which the two solutions (U, \hat{b}) coincide. So, when the elements \hat{b} of the two solutions coincide *and* are equal to r ,

the points (V_1, v) from the graph of $V_{1;I} \cup V_{1;II}$ and from the graph of $\tilde{\Delta}$ coincide. In other words, the two sets of curves intersect. Notice that they cannot intersect once. This is because given the shape of $V_{1;I} \cup V_{1;II}$, it would have to get to the south-east of $\tilde{\Delta}$, in order for the two curves to intersect twice. However, to the south-east of $\tilde{\Delta}$ there are no solutions such that \hat{b} is less than 1, so there it cannot be that $\hat{b} = r$. Thus, the graphs of $V_{1;I} \cup V_{1;II}$ and of $\{(V_1, v) : \mathcal{D} = 0\}$ intersect only once and the graphs are also tangent there. In other words, $(V_{1;I} \cup V_{1;II}) \cap \tilde{\Delta}$ is a singleton. Denote that point $(V_1^{\tilde{\Delta}}, v^{\tilde{\Delta}})$.

Consider the case that the graph of $V_{1;I} \cup V_{1;II}$ is a loop, i.e. $r < \frac{1}{7}$. We can distinguish four regions of (V_1, v) (all regions also imply that the cubic discriminant is non-negative):

$$(M) \{(V_1, v) | V_1 \leq \min\{V_{1;I}, V_{1;II}\}, v \leq v^{\tilde{\Delta}}\}$$

$$(N) \{(V_1, v) | V_1 \in (\min\{V_{1;I}, V_{1;II}\}, \max\{V_{1;I}, V_{1;II}\}), v < 2 - r\}$$

$$(N') \{(V_1, v) | V_1 \in (\min\{V_{1;I}, V_{1;II}\}, \max\{V_{1;I}, V_{1;II}\}), 2 - r \leq v \leq v^{\tilde{\Delta}}\}$$

$$(K) (A \cup B \cup B')^C$$

Recall that the set of points $V_{1;I} \cup V_{1;II}$ represent the parameters for which the system (A) has a solution such that $\hat{b} = r$. So, as we cross the graph of $V_{1;I} \cup V_{1;II}$ go from the interior of one of the regions above to the interior of the other region, \hat{b} from one of the solutions would cross r , and when we go from the interior of region (M) to the interior of region (K) two solutions would cross r . Let us look at some typical elements of the sets (M)-(K) to understand the ranking of \hat{b} relative to r from the solutions from each of the regions. Set $r = 0.03$. Check that the points $(2.62, 2.2)$, $(2.9, 2.2)$, $(2.62, 2.63)$, $(2.9, 2.63)$ lie in the interiors of regions (M), (N), (N'), and (K), respectively. The correspondent pairs of solutions are:

$$- (U, \hat{b}) = (0.526087, 0.125546) \text{ and } (U, \hat{b}) = (-0.070028, 0.0798147) \text{ for the point } (V_1, v) = (2.62, 2.2);$$

- $(U, \hat{b}) = (0.809835, 0.0975269)$ and $(U, \hat{b}) = (-0.0201097, 0.020638)$ for the point $(V_1, v) = (2.9, 2.2)$;
- $(U, \hat{b}) = (0.133118, 0.0176779)$ and $(U, \hat{b}) = (0.00617516, 0.0409543)$ for the point $(V_1, v) = (2.62, 2.63)$;
- $(U, \hat{b}) = (0.471844, -0.0553398)$ and $(U, \hat{b}) = (-0.0125482, 0.0138343)$ for the point $(V_1, v) = (2.9, 2.63)$;

Overall, the \hat{b} from the solution that corresponds to a higher U crosses r from above as we go from either of regions (M) and (N) to either of regions (N') and (K); while the \hat{b} from the solution that corresponds to a low U crosses r from above as we go from either of regions (M) and (N') to either of regions (N) and (K);

Consider now the case that $r \geq \frac{1}{7}$, so that $V_{1,I}$ and $V_{1,II}$ intersect only once at $v = \frac{1}{2} \left(6 - r - \sqrt{3r(4-r)} \right)$. For this case we can distinguish only three regions:

$$(\tilde{M}) \{(V_1, v) | V_1 \leq \min\{V_{1,I}; V_{1,II}\}, v \leq v^{\tilde{A}}\}$$

$$(\tilde{N}) \{(V_1, v) | V_1 \in (\min\{V_{1,I}; V_{1,II}\}, \max\{V_{1,I}; V_{1,II}\})\}$$

$$(\tilde{K}) (A \cup B)^C$$

In this case consider typical elements of those sets. For $r = 0.5$ the points $(V_1, v) = (1.2, 1.1)$, $(V_1, v) = (1.575, 1.1)$, $(V_1, v) = (1.575, 1.65)$ lie in (\tilde{M}) , (\tilde{N}) , and (\tilde{K}) , respectively. The correspondent pairs of solutions are:

- $(U, \hat{b}) = (0.100275, 0.903202)$ and $(U, \hat{b}) = (-0.490029, 0.550519)$ for the point $(V_1, v) = (1.2, 1.1)$;
- $(U, \hat{b}) = (0.475719, 0.902897)$ and $(U, \hat{b}) = (-0.384299, 0.419833)$ for the point $(V_1, v) = (1.575, 1.1)$;
- $(U, \hat{b}) = (-0.0960184, 0.467774)$ and $(U, \hat{b}) = -0.2728940.411679)$ for the point $(V_1, v) = (1.575, 1.65)$;

Thus, the \hat{b} from the solution that corresponds to a higher U crosses r from above as we go from either of regions (\tilde{M}) and (\tilde{N}) to the region (\tilde{K}) ; and the \hat{b} from the solution that corresponds to a low U crosses r from above as we go from the region (\tilde{M}) to either of regions (\tilde{N}) and (\tilde{K}) ;

Note also that in the case $r \geq \frac{1}{7}$ the curve $V_{1,II}$ intersects the 45-degree line twice and the larger value of v at the intersection is $\frac{1}{2} \left(6 - r - \sqrt{3r(4-r)} \right)$ (which is also a point at which $V_{1,II}$ and $V_{1,I}$ coincide). For $r \geq \frac{1}{7}$ the value of $\frac{1}{2} \left(6 - r - \sqrt{3r(4-r)} \right)$ is less than or equal to $\frac{16}{7}$. The point of tangency of $V_{1,I} \cup V_{1,II}$ with $\tilde{\Delta}$ (the points for which cubic discriminant is zero), is below the higher intersection of $V_{1,II}$ with a 45 degree line, which itself is below the point of tangency of a 45 degree line and the set of points $\tilde{\Delta}$. This means that for $r \geq \frac{1}{7}$ one of the necessary conditions for a pair (U, \hat{b}) to be an equilibrium object is that $V_1 \geq v$, since for this region only the solutions with higher U are such that U is non-negative.

Overall, recall that for a pair U, \hat{b} to be an equilibrium object, it needs to be a solution to the system (A), U must be non-negative, and \hat{b} needs to be weakly greater than r . For $r \leq \frac{1}{7}$ the equilibrium (equilibria) exists (exist) in the following regions:

(Q) $V_1 \in [v, +\infty)$ for $v \in [1, 2-r]$, $V_1 \in [v, V_{1,I}]$ for $v \in (2-r, \frac{16}{7})$. There is one equilibrium pair U, \hat{b} , which corresponds to the solution to the system (A) with a higher U

(Q') $V_1 \geq v$, $v \in [\frac{16}{7}, \frac{8}{3}(1-r))$. Similar to the region described above, there is one equilibrium pair (U, \hat{b}) , which corresponds to the solution to the system (A) with a higher U .

(R) $V_1 < \min\{v, V_{1,I}\}$, $v \in [\frac{16}{7}, \frac{8}{3}(1-r))$, $\mathcal{D} \geq 0$. There are two pairs, which solutions to the system (A) and which are also equilibrium objects. This is because the lower U , which is a solution, is now non-negative. And elements \hat{b} which are parts of solution are now not less than r . So, in this sub-region

there are two equilibria possible. Note also that player 1 has a positive rent here, $U \geq 0$, even though he has lower valuation V_1 than the other two players, who have v .

(S) $V_1 \in [V_{1,I}, v]$, $v \in \left[\frac{8}{3}(1-r), \frac{1}{2} \left(6-r-\sqrt{3r(4-r)} \right) \right]$. There is one pair, which solutions to the system (A) and which is an equilibrium object. Note, however, that in this pair U is lower, than in the other pair. However, since in the pair with the higher U the element \hat{b} is lower than r , that pair cannot be an equilibrium.

As for $r > \frac{1}{7}$, the equilibrium exists only in the region $V_1 \in [v, +\infty)$ for $v \in [1, 2-r]$, $V_1 \in [v, V_{1,I}]$ for $v \in (2-r, \frac{16}{7})$. An equilibrium pair (U, \hat{b}) is the solution to the system (A) with the higher value of U .

C.4 Proposition 4.iv region characterization

Without loss, let player 1 be the one with the non-negative payoff $u_1 \geq 0$. Denote the size of player 3's atom at 0 by P_3 . The behavior of CDFs of player 1 and player 2 in the continuous part (bidding in (r, \bar{b})) follows from

$$\begin{aligned} V_1 G_2(b) P_3 - b &= U \\ V_2 G_1(b) P_3 - b &= 0 \\ \Rightarrow \\ G_1(b) &= \frac{b}{V_2 \times P_3}, G_2(b) = \frac{b+U}{V_1 \times P_3}. \end{aligned}$$

The equilibrium objects, that need to be determined, are, therefore, P_3 , i.e. with what probability is player 3 inactive; \hat{b} , the upper-bound of the support of continuous bidding; and U , player 1's equilibrium payoff. Determining these equilibrium objects and their expressions through V_1 , V_2 , and V_3 will allow to determine the subset of parameters, for which the current type of equilibrium holds. These objects can be inferred from the indifference of players 1 and 2 between bidding \hat{b}

and 1; and indifference of player 3 between bidding 0 and 1. Writing down these conditions:

$$V_1 \times G_2(\hat{b}) \times P_3 - \hat{b} =$$

$$V_1 \left(G_2(\hat{b})P_3 + \frac{1}{2}(1 - G_2(\hat{b}))P_3 + \frac{1}{2}G_2(\hat{b})(1 - P_3) + \frac{1}{3}(1 - G_2(\hat{b}))(1 - P_3) \right) - 1 = U,$$

$$V_2 \times G_1(\hat{b}) \times P_3 - \hat{b} =$$

$$V_2 \left(G_1(\hat{b})P_3 + \frac{1}{2}(1 - G_1(\hat{b}))P_3 + \frac{1}{2}G_1(\hat{b})(1 - P_3) + \frac{1}{3}(1 - G_1(\hat{b}))(1 - P_3) \right) - 1 = 0,$$

$$V_3 \left(G_1(\hat{b})G_2(\hat{b}) + \frac{1}{2}(1 - G_1(\hat{b}))G_2(\hat{b}) + \frac{1}{2}G_1(\hat{b})(1 - G_2(\hat{b})) + \frac{1}{3}(1 - G_1(\hat{b}))(1 - G_2(\hat{b})) \right) - 1 = 0.$$

Using the fact that $G_1(\hat{b}) = \frac{\hat{b}}{V_2 \times P_3}$ and $G_2(\hat{b}) = \frac{\hat{b} + U}{V_1 \times P_3}$, we can rewrite the above conditions as a system of three equations:

$$\frac{1}{3}V_1 + \frac{1}{6}P_3V_1 + \frac{1}{6}\frac{\hat{b} + U}{P_3} + \frac{1}{3}(\hat{b} + U) = 1 + U \quad (1)$$

$$\frac{1}{3}V_2 + \frac{1}{6}P_3V_2 + \frac{1}{6}\frac{\hat{b}}{P_3} + \frac{1}{3}\hat{b} = 1 \quad (2)$$

$$\frac{1}{3}V_3 + \frac{1}{6}\frac{\hat{b} + U}{V_1P_3} + \frac{1}{6}\frac{\hat{b}}{V_2P_3} + \frac{1}{3}\frac{(\hat{b} + U)\hat{b}}{V_1V_2P_3^2} = 1. \quad (3)$$

Let us denote U and \hat{b} that follow from the equations (1) and (2) as U^* and \hat{b}^* , respectively. Then,

$$U^* = \frac{P_3(2 + P_3)(V_1 - V_2)}{4P_3 - 1}, \hat{b}^* = \frac{P_3(6 - (2 + P_3)V_2)}{1 + 2P_3}.$$

Plugging the expression for \hat{b}^* into the equation (3), solving for U , and denoting that solution as \tilde{U} , we get

$$\tilde{U} = \frac{P_3((2 + P_3)V_2 - 6)}{1 + 2P_3} + \frac{P_3V_1(6 + V_2(2V_3(1 + 2P_3) - 8 - 13P_3))}{3(V_2 - 4)}.$$

From now on, let us consider the two sub-cases: (a) $P_3 > \frac{1}{4}$; and (b) $P_3 < \frac{1}{4}$:

(a), $P_3 > \frac{1}{4}$ In this case U^* can only be non-negative if $V_1 \geq V_2$. So, this is one of the necessary conditions. Another necessary condition is that $V_2 \leq 3$, which follows from the fact that \hat{b}^* must be non-negative. Let us establish the remaining necessary conditions in terms of V_1 , V_2 , V_3 , and r from the facts that: U^* and \tilde{U} must cross and they must be non-negative; $P_3 \in (\frac{1}{4}, 1]$; $\hat{b} \in [r, 1]$; $G_i(\hat{b}) \leq 1$, $i = 1, 2$; $V_3 G_1(\hat{b}) G_2(\hat{b}) \leq \hat{b}$.

Let us consider U_1^* and \tilde{U} as functions of P_3 , $U^*(P_3)$, and $\tilde{U}(P_3)$. We are interested in their intersection (if it exists) at some $P_3^* \in (\frac{1}{4}, 1]$. Since $P_3^* = 0$ is not the intersection we are interested in, we can consider the intersection(s) of the functions $v^*(P_3) = \frac{\tilde{U}(P_3)}{P_3}$, and $\tilde{v}(P_3) = \frac{U^*(P_3)}{P_3}$. Note that under the restriction that $V_2 \leq 3$, \tilde{v} is strictly decreasing in V_3 . If V_3 is small enough so that $\tilde{v}(1) \geq v^*(1)$ (i.e. if $V_3 \leq \frac{4V_1V_2+5V_2-4-3V_1-V_2^2}{V_1V_2}$), then $\tilde{v}(\frac{1}{4}) > 0$ and $\tilde{v}'(1) > 0$.⁹ For $P_3 > 0$, \tilde{v} can change monotonicity at most once, thus, for $P_3 \in (\frac{1}{4}, 1]$ the two curves cross only once: $v^*(P_3)$ is decreasing in P_3 with $\lim_{P_3 \downarrow \frac{1}{4}} v^*(P_3) = +\infty$; \tilde{v} is increasing in P_3 if the two curves cross at $P_3 \in (\frac{1}{4}, 1]$. So, another necessary condition is that $V_3 \leq \frac{4V_1V_2+5V_2-4-3V_1-V_2^2}{V_1V_2}$. Note also, that since \tilde{v} is increasing in V_2 and v^* is decreasing in V_2 given our restrictions, the point of intersection of the two curves is decreasing in V_2 .

Note that \hat{b}^* is always weakly less than 1 under our restrictions. \hat{b}^* needs also to be weakly greater than r when v^* and \tilde{v} cross. \hat{b}^* achieves maximum with respect to P_3 at $P_3 = \frac{\sqrt{3}\sqrt{V_2(4-V_2)}}{2V_2} - \frac{1}{2} \in [0, 1]$ for $V_2 \in [1, 3]$. Let us therefore solve for such values of V_3 , at which at the intersection of \tilde{v} and v^* the value of \hat{b}^* is equal to r , and for values of V_3 between the pairs identified above, it will be satisfied that $\hat{b}^* > r$. Treating, for now, V_3 as endogenous parameter and solving

⁹This is because $\tilde{v}'(1)$ and $\tilde{v}(\frac{1}{4})$ are decreasing in V_3 , given our restriction that $V_2 \leq 3$; for $\tilde{v}'(1)$ to be weakly positive, it needs to hold that $V_3 \leq \frac{16+V_2(-8+13V_1+V_2)}{4V_1V_2}$, and for $\tilde{v}(\frac{1}{4})$ to be weakly positive, it needs to hold that $V_3 \leq \frac{13}{4} + \frac{(4-V_2)^2}{V_1V_2}$; both of the latter two inequalities are implied by $V_3 \leq \frac{4V_1V_2+5V_2-4-3V_1-V_2^2}{V_1V_2}$.

the system

$$\begin{aligned}\tilde{v}(P_3; V_3) &= v^*(P_3; V_3) \\ \hat{b}^*(P_3) &= r\end{aligned}\tag{C}$$

with respect to P_3 and V_3 gives us two pairs of roots:

$$\begin{aligned}P_3^I &= \frac{3 - r - V_2 - \psi}{V_2} \\ V_3^I &= \frac{1}{V_1 V_2 (8 - 8r - 3V_2)} \times \\ &\quad (16r^3 - 144 + 36V_1 + V_2(90 - 13V_2) - V_1 V_2 (6V_2 - 2) + \\ &\quad 2r^2(11V_2 - 56) - r(V_1(12 + 23V_2) - 240 + V_2(112 - 13V_2)) \\ &\quad - (48 - 12V_1 - 14V_2 + 3V_1 V_2 + 16r^2 - r(64 - 14V_2))\psi) \\ P_3^{II} &= \frac{3 - r - V_2 + \psi}{V_2} \\ V_3^{II} &= \frac{1}{V_1 V_2 (8 - 8r - 3V_2)} \times \\ &\quad (16r^3 - 144 + 36V_1 + V_2(90 - 13V_2) - V_1 V_2 (6V_2 - 2) + \\ &\quad 2r^2(11V_2 - 56) - r(V_1(12 + 23V_2) - 240 + V_2(112 - 13V_2)) \\ &\quad + (48 - 12V_1 - 14V_2 + 3V_1 V_2 + 16r^2 - r(64 - 14V_2))\psi),\end{aligned}\tag{1}$$

where $\psi = \sqrt{(3 - r)^2 - (6 - r)V_2 + V_2^2}$. Note that $P_3^I \leq P_3^{II}$. Thus, the necessary conditions that follow from the requirement $\hat{b} \geq r$ is that $V_3 \geq V_3^I$ whenever $P_3^I \in [0.25, 1]$ and $V_3 \leq V_3^{II}$ whenever $P_3^{II} \in [\frac{1}{4}, 1]$.

From the restriction $G_1(\hat{b}) \leq 1 \iff \frac{\hat{b}}{V_2 P_3} \leq 1$ follows the necessary condition $V_3 \geq \frac{9V_2 + 14V_1 V_2 - 17V_1 - 9V_2^2}{V_1(5V_2 - 8)}$ whenever $\frac{2 - V_2}{V_2} \in [\frac{1}{4}, 1]$. This is implied by the following observations: $G_1(\hat{b}^*) = \frac{\hat{b}^*}{V_2 P_3} = \frac{6 - (2 + P_3)V_2}{(1 + 2P_3)V_2}$ is decreasing in P_3 ; the P_3 corresponding to the intersection of \tilde{v} and v^* is increasing in V_3 ; so V_3 must be high enough so that when at the cross of \tilde{v} and v^* , $G_1(\hat{b})$ is weakly less than 1; $G_1(\hat{b})$ is exactly equal to 1 at the intersection of \tilde{v} and v^* when V_3 is equal to $\frac{9V_2 + 14V_1 V_2 - 17V_1 - 9V_2^2}{V_1(5V_2 - 8)}$, and this occurs at $P_3 = \frac{2 - V_2}{V_2}$, which establishes the considered condition.

By the similar logic, from the restriction $G_2(\hat{b}) \leq 1$ follows the restriction $V_3 \geq \frac{32V_2 - 9V_1 - 3(8 - \sqrt{9V_1^2 - 16(4 - V_2)(V_2 - 1)})}{8V_2}$, whenever $\frac{V_1 - 4V_2 + 8 - \sqrt{9V_1^2 - 16(4 - V_2)(V_2 - 1)}}{4(V_1 + V_2)} \in [\frac{1}{4}, 1]$ (with the latter being implied by $V_1 \geq V_2$).

Finally, it must be non-profitable for player 3 to bid anywhere in $[r, \hat{b}]$, which is formally written down as

$$\begin{aligned} V_3 G_1(b) G_2(b) - b &\leq 0, \forall b \in [r, \hat{b}] \iff \\ V_3 \times \frac{b(b+U)}{V_1 V_2 P_3^2} &\leq b \forall b \in [r, \hat{b}] \iff \\ V_3 \times \frac{\hat{b}+U}{V_1 V_2 P_3^2} &\leq 1. \end{aligned}$$

Using the expressions for \tilde{U} and \hat{b}^* , $V_3 \times \frac{\hat{b}+U}{V_1 V_2 P_3^2}$ can be rewritten as $\frac{V_3(6+V_2(2V_3(1+2P_3)-8-13P_3))}{3(V_2-4)V_2P_3} \doteq \phi(P_3)$. Note that if $\phi(P_3) = 1$ for some $P_3^* \in [\frac{1}{4}, 1]$, then $\phi(P_3)' \geq 0$.¹⁰ Using the fact that the point of intersection of v^* and \tilde{v} , P_3^\dagger , is increasing in V_1 ,¹¹, we can establish the upper bound on V_1 , such that if V_1 is below that bound, $V_3 G_1(\hat{b}) G_2(\hat{b}) \leq \hat{b}$. That bound follows from the system

$$\begin{aligned} v^*(P_3) &= \tilde{v}(P_3) \\ V_3 G_1(\hat{b}^*) G_2(\hat{b}^*) &= \hat{b}^*, \end{aligned}$$

and is

$$\begin{aligned} \overline{V_1} &= \frac{1}{(V_2 - V_3)^2(36 + V_2^2(V_3 - 1) + V_3(4V_3 - 13) - V_2(5 + V_3(43 + V_3(4V_3 - 25))))} \times \\ &\times ((V_2(3V_2 - 12 + (13 - 4V_3)V_3)(V_3((23 - 8V_3)V_3 - 24) + \\ &V_2^2(3 + 4(V_3 - 4)V_3) - 2V_2(6 + V_3(2V_3(16 + (V_3 - 7)V_3) - 35))))). \end{aligned}$$

¹⁰To see this, note that the root of $\phi(P_3) = 1$ is $P_3^* = \frac{2(3-V_2(4-V_3))V_3}{V_2((13-4V_3)V_3+3V_2-12)}$; while the derivative of $\phi(P_3)$ is $\phi(P_3)' = \frac{2(3-V_2(4-V_3))V_3}{3P_3^2(4-V_2)V_2}$. The minimum of that derivative, given the constraints $P_3^* \in [\frac{1}{4}, 1]$ is 0; thus $\phi(P_3)$ is weakly increasing in the parameters region of our interest.

¹¹As can be seen by taking the implicit derivative P_3 with respect to V_1 that follows from $F(P_3) = v^* - \tilde{v}$.

Combining all the necessary conditions established above, we have the necessary and sufficient condition for the type of equilibrium with players 1 and 2 bidding continuously in (r, \hat{b}) , and having atoms at 0, r , and 1; while player 3 only has atoms at 0 and 1, with the size of atom at 0 being greater than $\frac{1}{4}$:

$$V_1 \geq V_2$$

$$V_2 \leq 3$$

$$V_3 \leq \frac{4V_1V_2 + 5V_2 - 4 - 3V_1 - V_2^2}{V_1V_2}$$

$$V_3 \geq \frac{1}{V_1V_2(8 - 8r - 3V_2)} \times$$

$$(16r^3 - 144 + 36V_1 + V_2(90 - 13V_2) - V_1V_2(6V_2 - 2) +$$

$$2r^2(11V_2 - 56) - r(V_1(12 + 23V_2) - 240 + V_2(112 - 13V_2))$$

$$- (48 - 12V_1 - 14V_2 + 3V_1V_2 + 16r^2 - r(64 - 14V_2))\psi),$$

$$\text{when } \frac{3 - r - V_2 - \psi}{V_2} \in \left[\frac{1}{4}, 1\right]$$

$$V_3 \leq \frac{1}{V_1V_2(8 - 8r - 3V_2)} \times$$

$$(16r^3 - 144 + 36V_1 + V_2(90 - 13V_2) - V_1V_2(6V_2 - 2) +$$

$$2r^2(11V_2 - 56) - r(V_1(12 + 23V_2) - 240 + V_2(112 - 13V_2))$$

$$+ (48 - 12V_1 - 14V_2 + 3V_1V_2 + 16r^2 - r(64 - 14V_2))\psi),$$

$$\text{when } \frac{3 - r - V_2 + \psi}{V_2} \in \left[\frac{1}{4}, 1\right]$$

$$V_3 \geq \frac{9V_2 + 14V_1V_2 - 17V_1 - 9V_2^2}{V_1(5V_2 - 8)}$$

$$V_3 \geq \frac{32V_2 - 9V_1 - 3 \left(8 - \sqrt{9V_1^2 - 16(4 - V_2)(V_2 - 1)}\right)}{8V_2}$$

$$V_1 \leq \frac{1}{(V_2 - V_3)^2(36 + V_2^2(V_3 - 1) + V_3(4V_3 - 13) - V_2(5 + V_3(43 + V_3(4V_3 - 25))))} \times$$

$$\times ((V_2(3V_2 - 12 + (13 - 4V_3)V_3)(V_3((23 - 8V_3)V_3 - 24) +$$

$$V_2^2(3 + 4(V_3 - 4)V_3) - 2V_2(6 + V_3(2V_3(16 + (V_3 - 7)V_3) - 35))))),$$

where $\psi = \sqrt{(3 - r)^2 - (6 - r)V_2 + V_2^2}$.

(b), $P_3 < \frac{1}{4}$. In this sub-case, the exact algebraic expression for the region is not presented. However, the procedure, that can used to derive the respective region is described. Recall the expressions for U^* , and \tilde{U} :

$$U^* = \frac{P_3(2 + P_3)(V_1 - V_2)}{4P_3 - 1},$$

$$\tilde{U} = \frac{P_3((2 + P_3)V_2 - 6)}{1 + 2P_3} + \frac{P_3V_1(6 + V_2(2V_3(1 + 2P_3) - 8 - 13P_3))}{3(V_2 - 4)}.$$

The root of $U^* = \tilde{U}$, $P_3 = 0$ is not a root we are interested in, as well as the cases $P_3 = \frac{1}{4}$, or $P_3 = -\frac{1}{2}$. In lieu of $U^* = \tilde{U}$, we can therefore work with the cubic equation in P_3 :

$$3(2 + P_3)(V_1 - V_2)(1 + 2P_3)(V_2 - 4) = 3((2 + P_3)V_2 - 6)(4P_3 - 1)(V_2 - 4) + V_1(6 + V_2(2V_3(1 + 2P_3) - 8 - 13P_3))(4P_3 - 1)(1 + 2P_3).$$

The root that is real and always exists is negative. Therefore for there to exist a root $P_3 \in (0, 1/4)$, it is necessary that a cubic discriminant is positive. Using the notation,

$$A = 4V_1V_2(13 - 4V_3)$$

$$B = -3(3(-4 + V_2)V_2 + 4V_1(3 + V_2(-4 + V_3)))$$

$$C = 9(4 + V_1 - 2V_2)(-4 + V_2)$$

$$D = -9(-4 + V_1 + V + 2) + V_1V_2(-1 + V_3),$$

we have that a necessary condition for the equilibrium with $P_3 \in (0, 1/4)$ to exist is

$$18 \times A \times B \times C \times D - 4 \times B^3 \times D + B^2 \times C^2 - 4 \times A \times C^3 - 27 \times A^2 D^2 \geq 0.$$

From the fact that we need $\hat{b} \geq r \geq 0$, and using the expression for \hat{b}^* ,

$$\hat{b}^* = \frac{P_3(6 - (2 + P_3)V_2)}{1 + 2P_3},$$

it follows that it is necessary that $V_2 \leq 3$.

Under the restrictions that $1 \leq V_1 \leq V_2 \leq V_3$, the function $\tilde{v}(P_3) = \frac{\tilde{U}(P_3)}{P_3}$ is increasing in P_3 on $P_3 \in (0, 1/4)$. If it was decreasing in P_3 , or if it achieved a maximum for some $P_3 = P^* \in (0, 1/4)$, the function \tilde{U} would be negative on $P_3 \in (0, 1/4)$, contradicting the equilibrium condition of non-negative players' payoffs. We, therefore, must require that $\tilde{U}(1/4) \geq 0$, which can be written down as

$$-16 + 6V_2 - \frac{(V_1(8 + V_2(-15 + 4V_3)))}{4 - V_2} \geq 0.$$

Besides the restriction that $\hat{b} \geq 0$, we need to impose more requirement in order to insure $\hat{b} \geq r$. Similarly to as we did before, in case $P_3 > 1/4$, we need to consider roots (P_3, V_3) of the system C. However, now we need to account for the fact that there are up to two intersections of U^* and \tilde{U} possible. The P_3 that follows from higher intersection of U^* and \tilde{U} is decreasing in V_3 and the P_3 that follows from the lower intersection of U^* and \tilde{U} is increasing in V_3 . There are also two intersections of $\hat{b}^* = r$ with respect to P_3 , and those values of P_3 that lie between the points of intersection are the ones for which the condition $\hat{b}^* \geq r$ is satisfied. From the expressions of the roots of the system, (P_3^I, V_3^I) , and (P_3^{II}, V_3^{II}) it is possible to say that P_3^{II} corresponds to the higher intersection of $\tilde{b}^* = r$ and P_3^I - to the lower. But it is not possible to say whether either of V_3^I or V_3^{II} corresponds to the lower or higher intersection of \tilde{U} and U^* . However, an implicit derivative of P_3 that follows from $\tilde{U} = U^*$ with respect to V_3 can be evaluated at $(P_3, V_3) = (P_3^I, V_3^I)$ and $(P_3, V_3) = (P_3^{II}, V_3^{II})$. The parameters with $\frac{dP_3}{dV_3} > 0$ would correspond to the lower intersection of \tilde{U} and U^* , and vice versa.

Using the implicit function derivative,

$$\begin{aligned}
\frac{dP_3}{dV_3} &= -\frac{\frac{\partial(\tilde{U}-U^*)}{\partial V_3}}{\frac{\partial(\tilde{U}-U^*)}{\partial P_3}} \\
&= \frac{2(2P_3+1)V_1V_2}{3(V_2-4)\left(\frac{12(P_3(P_3(7V_2-16)+V_2+8)+V_2-1)}{(8P_3^2+2P_3-1)^2} + \frac{2V_1(V_2(-4P_3(2P_3-1)(4V_3-13)-2V_3-7)+54)}{3(1-4P_3)^2(V_2-4)}\right)} \\
&\doteq \chi(P_3; V_3).
\end{aligned}$$

Then, the restrictions that follow from $\hat{b} \geq r$ are:

- for the region of parameters (V_1, V_2, V_3, r) such that $P_3^I > 1/4$ and $P_3^{II} > 1/4$, there are no subsets such that $\hat{b} \geq r$;
- for the region of parameters (V_1, V_2, V_3, r) such that $P_3^I < 1/4$ and $P_3^{II} > 1/4$, the subset of parameters such that $\chi(P_3^I, V_3^I) < 0$ (i.e. the parameters for which (P_3^I, V_3^I) corresponds to the higher intersection of U^* and \tilde{U}), we need to exclude $V_3 > V_3^I$; remainder of the region $P_3^I < 1/4$ and $P_3^{II} > 1/4$ can be included;
- for the region of parameters (V_1, V_2, V_3, r) such that $P_3^I < 1/4$ and $P_3^{II} < 1/4$,
 - the subset of parameters for which $\chi(P_3^I; V_3^I) < 0$ and $\chi(P_3^{II}; V_3^{II}) < 0$ should be included entirely;
 - the subset $\chi(P_3^I; V_3^I) < 0$ and $\chi(P_3^{II}; V_3^{II}) > 0$ is empty, since $P_3^I < P_3^{II}$, but $\chi < 0$ means higher intersection of \tilde{U} and U^* ;
 - from the subset for which $\chi(P_3^I; V_3^I) > 0$ and $\chi(P_3^{II}; V_3^{II}) < 0$ exclude $V_3 < \min\{V_3^I, V_3^{II}\}$;
 - from the subset for which $\chi(P_3^I; V_3^I) > 0$ and $\chi(P_3^{II}; V_3^{II}) > 0$ exclude $V_3 \in ([V_3^I, V_3^{II}])^C$.

The remaining restrictions follow from making $G_1(\hat{b}) \leq 1$, $G_2(\hat{b}) \leq 1$, and making sure it is non-profitable for player 3 to bid anywhere in $b \in (0, 1)$.

In equilibrium of this type, $G_1(\hat{b}) = \hat{b}/(V_2 P_3)$. Replacing \hat{b} with the expression for \hat{b}^* , and treating it like a function of P_3 , we have that

$$G_1(P_3) = \frac{6 - (2 + P_3)V_2}{V_2(1 + 2P_3)}.$$

This expression is decreasing in P_3 under the restriction $V_2 \leq 3$ that we have previously established. $G_1(P_3) = 1$ at $P_3 = \frac{2}{V_2} - 1$. Since we need to have $G_1 \leq 1$ for some $P_3 \leq 1/4$, we need to impose $V_2 \geq \frac{8}{5}$. In order to insure that $G_1 \leq 1$ for the intersection of \tilde{U} and U^* , let us solve the system

$$\begin{aligned} G_1(P_3) &= 1 \\ \tilde{U} &= U^* \end{aligned}$$

with respect to P_3 and V_3 . We get that the root of that system is $P_3^\dagger = \frac{2}{V_2} - 1$ and $V_3^\dagger = \frac{V_1(17-14V_2)+9(V_2-1)V_2}{V_1(8-5V_2)}$. If P_3^\dagger is below 0 (happens for $V_2 \geq 2$), no additional restrictions are needed. If $P_3^\dagger \in (0, 1/4)$, we need to insure that equilibrium $P_3 \geq P_3^\dagger$, since $G_1(P_3)$ is decreasing in P_3 for at least one of the two non-trivial roots of $\tilde{U} = U^*$. $G_1(P_3)$ does not depend on V_3 . As we saw, equilibrium P_3 can increase or decrease in V_3 , depending on whether it corresponds to the higher or the lower intersection of \tilde{U} and U^* . Check how many roots of $U^* = \tilde{U}$ there are in $P_3 \in (0, 1/4)$. If $2(V_2 - V_1) \leq \frac{2V_1(V_2(V_3-4)+3)}{3(V_2-4)} + 2V_2 - 6$, there is one root, that is the one which is decreasing in V_3 . For this case we need to restrict $V_3 \leq V_3^\dagger$. If $2(V_2 - V_1) > \frac{2V_1(V_2(V_3-4)+3)}{3(V_2-4)} + 2V_2 - 6$, there are two roots. For this case we need to understand the sign of $\frac{dP_3}{dV_3}$. Again using the function $\chi(P_3, V_3)$, we need to have $V_3 \leq V_3^\dagger$ if $\chi(P_3^\dagger, V_3^\dagger) < 0$.

Consider the restriction $G_2(\hat{b}) \leq 1$. In equilibrium $G_2(\hat{b}) = \frac{U+\hat{b}}{V_1 P_3}$. Using the expressions for \tilde{U} and \hat{b}^* , G_2 can be rewritten, as a function of P_3 :

$$G_2(P_3) = \frac{V_2(4P_3V_3 - 13P_3 + 2V_3 - 8) + 6}{3(V_2 - 4)},$$

which is linear in P_3 , and, besides, does not depend on V_1 . Given that we are already restricted to $V_2 \leq 3$, G_2 is increasing in P_3 if $V_3 \leq 13/4$ and vice versa. Solving the system

$$\begin{aligned} U^* &= \tilde{U} \\ G_2(P_3) &= 1 \end{aligned}$$

with respect to P_3 and V_1 , we get the root

$$\begin{aligned} P_3^\dagger &= \frac{2V_2V_3 - 11V_2 + 18}{13V_2 - 4V_2V_3} \\ V_1^\dagger &= \frac{V_2(V_2(-4(V_3 - 8)V_3 - 55) - 24V_3 + 51)}{9(V_2(V_3 - 4) + 3)}. \end{aligned}$$

So, if ($V_3 \leq 13/4$ and $P_3^\dagger \geq 1/4$), or if ($V_3 \geq 13/4$ and $P_3^\dagger \leq 0$), there are no additional restrictions. We need to exclude all the parameters such that ($V_3 \leq 13/4$ and $P_3^\dagger \leq 0$), or ($V_3 \geq 13/4$ and $P_3^\dagger \geq 1/4$). For $P_3^\dagger \in [0, 1/4]$, we need to also consider the number of roots of $\tilde{U} = U^*$ and, potentially, the sign of $\frac{dP_3}{dV_1}$ of the P_3 that follows from $\tilde{U} = U^*$, derivative being evaluated at $(P_3^\dagger, V_1^\dagger)$. If there is one non-trivial root of $\tilde{U} = U^*$ in $P_3 \in (0, 1/4)$, it is the root that is increasing in V_1 . Then V_1 must be $\leq V_1^\dagger$, if $V_3 \leq 13/4$, and V_1 must be $\geq V_1^\dagger$ if $V_3 \geq 13/4$. For the case of two non-trivial roots of $U^* = \tilde{U}$ in $(0, 1/4)$, determine the sign of $\frac{dP_3}{dV_1}$ at $(P_3, V_1) = (P_3^\dagger, V_1^\dagger)$. If $\frac{dP_3}{dV_1}|_{(P_3^\dagger, V_1^\dagger)} \leq 0$ AND $V_3 \leq 13/4$, V_1 must be weakly below V_1^\dagger , while if $\frac{dP_3}{dV_1}|_{(P_3^\dagger, V_1^\dagger)} \geq 0$ AND $V_3 \geq 13/4$, V_1 must be weakly above V_1^\dagger . The sign of $\frac{dP_3}{dV_1}$ is determined from the equation $\tilde{U} = U^*$ using the implicit function derivative:

$$\begin{aligned} \frac{dP_3}{dV_1} &= -\frac{\frac{\partial(U^* - \tilde{U})}{\partial V_1}}{\frac{\partial(U^* - \tilde{U})}{\partial P_3}} = \\ &= -\frac{2(2P_3 + 1)(P_3V_2(4V_3 - 13) + V_2(-V_3) + V_2 + 9)}{3(4P_3 - 1)(V_2 - 4) \left(\frac{12(P_3(P_3(7V_2 - 16) + V_2 + 8) + V_2 - 1)}{(8P_3^2 + 2P_3 - 1)^2} + \frac{2V_1(V_2(-4P_3(2P_3 - 1)(4V_3 - 13) - 2V_3 - 7) + 54)}{3(1 - 4P_3)^2(V_2 - 4)} \right)} \\ &\doteq \delta(P_3; V_1). \end{aligned}$$

The last remaining restriction is from non-profitability of player 3's deviation to bidding in the interior of $(0, 1)$, which happens if and only if $V_3 G_1(\hat{b}) G_2(\hat{b}) \leq \hat{b}$.

The latter can be rewritten as $V_3 \frac{\hat{b}+U}{V_1 V_2 P_3^2} \leq 1$, or

$$\tau(P_3) \doteq \frac{V_3(V_2(4P_3V_3 - 13P_3 + 2V_3 - 8) + 6)}{3P_3(V_2 - 4)V_2} \leq 1.$$

$\tau(P_3)$ is increasing whenever $3 - V_2(4 - V_3) \geq 0$ and vice versa.

Let us solve for the roots (P_3, V_1) of the system

$$\frac{V_3(V_2(4P_3V_3 - 13P_3 + 2V_3 - 8) + 6)}{3P_3(V_2 - 4)V_2} = 1$$

$$U^* = \tilde{U},$$

and get the root

$$P_3^{\S} = \frac{2V_3(V_2(V_3 - 4) + 3)}{V_2(3V_2 + V_3(13 - 4V_3) - 12)}$$

$$V_1^{\S} = (V_2(3V_2 + V_3(13 - 4V_3) - 12) (V_2^2(4(V_3 - 4)V_3 + 3) - 2V_2(V_3(2V_3((V_3 - 7)V_3 + 16) - 35) + 6) + V_3((23 - 8V_3)V_3 - 24))) / ((V_2 - V_3)^2 (V_2^2(V_3 - 1) - V_2(V_3(V_3(4V_3 - 25) + 43) + 5) + V_3(4V_3 - 13) + 36)).$$

As before, we need to check for the number of roots of $U^* = \tilde{U}$, and the sign of $\frac{dP_3}{dV_1} |_{(P_3^{\S}, V_1^{\S})}$. If $(P_3^{\S} \geq 1/4$ and $3 + V_2(V_3 - 4) \geq 0)$, there are no new restrictions needed, all parameters such that $(P_3^{\S} \geq 1/4$ and $3 + V_2(V_3 - 4) \leq 0)$ should be excluded. If there is one non-trivial root of $\tilde{U} = U^*$ in $(0, 1/4)$, this is the one that is increasing in V_1 . So if $(P_3^{\S} \leq 1/4$ and $3 + V_2(V_3 - 4) \geq 0)$, we must exclude $V_1 > V_1^{\S}$; while if $(P_3^{\S} \leq 1/4$ and $3 + V_2(V_3 - 4) \leq 0)$, we must exclude $V_1 < V_1^{\S}$. If there are two roots of $\tilde{U} = U^*$ in $(0, 1/4)$, determine the sign of $\frac{dP_3}{dV_1}$ at $(P_3, V_1) = (P_3^{\S}, V_1^{\S})$. If $\frac{dP_3}{dV_1} |_{(P_3^{\S}, V_1^{\S})} \geq 0$ AND $3 + V_2(V_3 - 4) \leq 0$, V_1 must be weakly above V_1^{\S} , while if $\frac{dP_3}{dV_1} |_{(P_3^{\S}, V_1^{\S})} \leq 0$ AND $3 + V_2(V_3 - 4) \geq 0$, V_1 must be

weakly below V_1^{\S} . The sign of the derivative $\frac{dP_3}{dV_1}$ again can be determined using the function $\delta(P_3; V_1)$.

References

- Baye, Michael R, Dan Kovenock, and Casper G De Vries**, “Rigging the lobbying process: an application of the all-pay auction,” *The american economic review*, 1993, *83* (1), 289–294.
- , – , and – , “The all-pay auction with complete information,” *Economic Theory*, 1996, *8* (2), 291–305.
- Boleslavsky, Raphael and Christopher Cotton**, “Limited capacity in project selection: Competition through evidence production,” *Economic Theory*, 2018, *65* (2), 385–421.
- Che, Yeon-Koo and Ian L Gale**, “Caps on political lobbying,” *The american economic review*, 1998, *88* (3), 643–651.
- and – , “Optimal design of research contests,” *American Economic Review*, 2003, *93* (3), 646–671.
- Clark, Derek J and Christian Riis**, “Competition over more than one prize,” *The American Economic Review*, 1998, *88* (1), 276–289.
- Hillman, Arye L and John G Riley**, “Politically contestable rents and transfers,” *Economics & Politics*, 1989, *1* (1), 17–39.
- Letina, Igor, Shuo Liu, and Nick Netzer**, “Delegating performance evaluation,” *Theoretical Economics*, 2020, *15* (2), 477–509.
- Moldovanu, Benny and Aner Sela**, “Patent licensing to Bertrand competitors,” *International Journal of Industrial Organization*, 2003, *21* (1), 1–13.
- Muratov, Oleg**, “Essays on Information Design in Economic Theory,” 2019.

– , “Mapping an Information Design Game into an All-Pay Auction,” *Mimeo*, 2021.

Siegel, Ron, “All-Pay Contests,” *Econometrica*, 2009, *77*, 71–92.

– , “Asymmetric all-pay auctions with interdependent valuations,” *Journal of Economic Theory*, 2014, *153*, 684–702.

Szech, Nora, “Tie-breaks and bid-caps in all-pay auctions,” *Games and Economic Behavior*, 2015, *92*, 138–149.