

Magnus, Jan R.

**Working Paper**

## Gauss on least-squares and maximum-likelihood estimation

Tinbergen Institute Discussion Paper, No. TI 2021-105/III

**Provided in Cooperation with:**

Tinbergen Institute, Amsterdam and Rotterdam

*Suggested Citation:* Magnus, Jan R. (2021) : Gauss on least-squares and maximum-likelihood estimation, Tinbergen Institute Discussion Paper, No. TI 2021-105/III, Tinbergen Institute, Amsterdam and Rotterdam

This Version is available at:

<https://hdl.handle.net/10419/248787>

**Standard-Nutzungsbedingungen:**

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

**Terms of use:**

*Documents in EconStor may be saved and copied for your personal and scholarly purposes.*

*You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.*

*If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.*

TI 2021-105/III  
Tinbergen Institute Discussion Paper

# Gauss on Least-Squares and Maximum-Likelihood Estimation

*Jan Magnus<sup>1</sup>*

<sup>1</sup> Vrije Universiteit Amsterdam

Tinbergen Institute is the graduate school and research institute in economics of Erasmus University Rotterdam, the University of Amsterdam and Vrije Universiteit Amsterdam.

Contact: [discussionpapers@tinbergen.nl](mailto:discussionpapers@tinbergen.nl)

More TI discussion papers can be downloaded at <https://www.tinbergen.nl>

Tinbergen Institute has two locations:

Tinbergen Institute Amsterdam  
Gustav Mahlerplein 117  
1082 MS Amsterdam  
The Netherlands  
Tel.: +31(0)20 598 4580

Tinbergen Institute Rotterdam  
Burg. Oudlaan 50  
3062 PA Rotterdam  
The Netherlands  
Tel.: +31(0)10 408 8900

# Gauss on least-squares and maximum-likelihood estimation<sup>1</sup>

Jan R. Magnus

Department of Econometrics and Data Science,  
Vrije Universiteit Amsterdam, Amsterdam, The Netherlands; and  
Tinbergen Institute, Amsterdam, The Netherlands

November 26, 2021

*Abstract:* Gauss' 1809 discussion of least squares, which can be viewed as the beginning of mathematical statistics, is reviewed. The general consensus seems to be that Gauss' arguments are at fault, but we show that his reasoning is in fact correct, given his self-imposed restrictions, and persuasive without these restrictions.

*Key words:* Gauss, least squares, maximum likelihood, history

*JEL Classification:* B16, C10

An old question in probability theory is the following: suppose we throw with two fair dice, how many times do we need to throw so that the probability of at least one double-6 is at least  $1/2$ . In the first half of the seventeenth century, the Chevalier de Méré, a well-known gambler, thought that he needed 24 throws. This problem has become famous because it intrigued Pascal and Fermat, and the solution is contained in a letter of Pascal to Fermat dated July 29, 1654.<sup>2</sup> With some caution we can take 1654 as the birth year of probability theory. It took a while to understand the basic rules of probability and sixty years later, in a letter to the Swiss philosopher and mathematician Louis Bourguet dated March 22, 1714, Leibniz still maintained that it was equally likely to throw twelve with two dice than to throw eleven, because “l'un et l'autre ne ce peut faire que d'une seule manière” (one or the other can be done in only one way).

Mathematical statistics is much younger and, with similar caution, we select as its beginning the publication of Gauss' famous 1809 monograph.<sup>3</sup> Legendre (1805) had published his method of least squares four years earlier,

---

<sup>1</sup>I am grateful to Franco Peracchi and Steven Tijms for useful comments.

<sup>2</sup>The correct answer is 25 and is obtained by showing that the equation  $1 - (35/36)^n = 1/2$  has the solution  $n \approx 24.6$ .

<sup>3</sup>Others would choose Laplace (1774) as the beginning of statistics, which is equally reasonable.

but he developed his method as an approximation tool and no randomness is assumed. Gauss (1809), in contrast, works in the context of random variables and distributions; see e.g. Pearson (1978), Stigler (1986), and Gorroochurn (2016) for historical details.

Some satisfaction seems to be derived in finding mistakes in the writings of great minds, and Leibniz' error is quoted frequently. Rather than laughing at Leibniz' mistake, we should realize just how difficult the beginnings of probability theory were, and that things that we now consider easy are not easy because we are so clever but because they have sunk into common knowledge.

Similarly, most Gauss commentators have found his 1809 treatment of least squares at fault. For example, Stigler (1986, pp. 141–143) considers it a “logical aberration . . . essentially both circular and non sequitur” and Gorroochurn (2016, p. 163) writes that “his reasoning contains an inherent circularity because the normal distribution emerges as a consequence of the postulate of the arithmetic mean, which is in fact a consequence of the normality assumption!” The purpose of this note is to demonstrate that it is not Gauss who is at fault but his commentators.

In modern notation, Gauss starts with the linear model

$$y = X\beta + u, \tag{1}$$

where he assumes that the errors  $u_i$  are independent and identically distributed (iid) with mean zero and common variance  $\sigma^2$ , which we set equal to one without loss of generality. Since the  $u_i$  are iid, they have a common density function, say  $\phi(u_i)$ , and the logarithm of the joint density becomes  $\sum_{i=1}^n \log \phi(u_i)$ . Gauss wishes to estimate  $\beta$  by maximizing the joint density. In other words, he wants to derive the maximum-likelihood estimator for  $\beta$ .

Gauss is aware of the fact that *if* he assumes normality of the errors, then the joint density will be of the form

$$\sum_{i=1}^n \log \phi(u_i) = a - b \sum_{i=1}^n u_i^2, \tag{2}$$

so that (under normality) maximizing the likelihood is the same as minimizing the sum of squared deviations. Gauss makes life unnecessarily difficult for himself by working in a Bayesian framework, assuming a flat bounded prior for each of the  $\beta_j$ , so that the posterior also has bounded support. But in essence, Gauss showed (for the first time) that in the standard linear model under normality the maximum-likelihood estimator is equal to the least-squares formula.

This is an important result in itself and Gauss could have stopped there. But he did *not* want to assume at the outset that the errors are normally distributed. Instead he wants to show that normality of the errors is not only sufficient but also necessary for the maximum-likelihood estimator to be equal to the least-squares formula. In this attempt he fails, not because his argument is wrong (as most Gauss scholars seem to believe), but because his (correct) argument is not general, which he fully realizes.

Let us reexamine his argument. Gauss (1809, book II, section III, §177) proves the following result (in modern notation).

**Proposition (Gauss, 1809):** *Let  $y_1, y_2, \dots, y_n$  ( $n \geq 3$ ) be a sequence of independent and identically distributed observations from an absolutely-continuous distribution with  $E(y_i) = \mu$  and  $\text{var}(y_i) = 1$ . Assume that the  $n$  realizations of  $y_i$  take only two values with frequencies  $n_1$  and  $n_2$ , respectively ( $n_1 \geq 1$ ,  $n_2 \geq 1$ ,  $n_1 \neq n_2$ ). Then, the average  $\bar{y}$  is the maximum-likelihood estimator of  $\mu$  if and only if the  $y_i$  are normally distributed.*

Before we prove the proposition, some comment is in order on Gauss' assumption that the  $n$  realizations of  $y_i$  take only two values. This seems to contradict the fact that the  $y_i$  follow an absolutely-continuous distribution. Of course, there is a difference between observations (random variables) from an absolutely-continuous distribution and observations (the realized values). Some statistical concepts have two terms (estimator, estimate; predictor, prediction) to emphasize this difference, but most (like observation) don't. The random variables follow an absolutely-continuous distribution, but the realizations take on specific values, and Gauss assumes that they take one or the other of two values. This is a rather heroic assumption, but it is not inconsistent or wrong. Gauss himself simply says *supponendo itaque* (by supposing therefore) as if this were a logical continuation of his argument, and provides no further comment.

To prove the proposition, Gauss argues as follows. Let  $u_i = y_i - \mu$ . Since the  $u_i$  are iid, they have a common density function, say  $\phi(u_i)$ . First assume that  $\phi$  is the standard-normal density. Then the loglikelihood  $L(\mu)$  can be written as in (2). This is maximized if and only if the sum of squares is minimized, which occurs when  $\sum_i (y_i - \mu) = 0$ , that is when  $\hat{\mu} = \bar{y}$ . Note that the additional assumption on the realizations of  $y_i$  is not required.

Now assume that  $\hat{\mu} = \bar{y}$ . Gauss needs to show that this implies that  $\phi$  is the standard-normal density. As assumed, the  $y_i$  can only take two distinct values, say  $z_1$  ( $n_1$  times) and  $z_2$  ( $n_2$  times), where  $n = n_1 + n_2$ . Then, letting

$$d = z_1 - z_2, \quad r = n_1/n, \quad (3)$$

he obtains

$$y_i - \bar{y} = \begin{cases} d(1-r) & \text{if } y_i = z_1, \\ -dr & \text{if } y_i = z_2. \end{cases} \quad (4)$$

Setting  $L'(\mu) = 0$  then gives

$$L'(\mu) = \sum_{i=1}^n \frac{\phi'(u_i)}{\phi(u_i)} = \frac{n_1 \phi'[d(1-r)]}{\phi[d(1-r)]} + \frac{n_2 \phi'(-dr)}{\phi(-dr)} = 0, \quad (5)$$

which can be rewritten as

$$f[d(1-r)] = f[-dr], \quad f(x) = \frac{\phi'(x)}{x\phi(x)}. \quad (6)$$

For each given value of  $r$  ( $0 < r < 1$ ,  $r \neq 1/2$ ), this has to hold for every value of  $d$ , and it is easy to see (*unde facile colligitur* in Gauss' words) that this implies that  $f$  is a constant. (We have to exclude  $r = 1/2$  because this would only imply that  $f$  is symmetric around zero.) Hence, we must solve the equation  $\phi'(x) = -kx\phi(x)$ , where  $k$  is a constant. The solution to this differential equation was known to be

$$\phi(x) = A \exp(-kx^2/2) \quad (7)$$

for some constant  $A$ . Since  $\phi$  represents a distribution it must integrate to one which implies that the constant  $A$  takes the value  $A = \sqrt{k/(2\pi)}$ , as proved a few decades earlier by Laplace (1774) in a *theoremata elegans*, a fact gracefully acknowledged by Gauss.<sup>4</sup> In our case,  $\sigma = 1$  and hence  $k = 1$ . Hence  $\phi$  is the standard-normal distribution, and the proof is complete.

The presented proof follows Gauss' argument closely except that he sets  $n_1 = 1$  and  $n_2 = n - 1$  (and tacitly assumes that  $n \geq 3$ ). The proposition tells us how far Gauss came into proving the necessity of the normality assumption. The answer is: not very far, because his conditions are rather restrictive. Two centuries later we can get a little further. In particular, Kagan et al. (1973, Theorem 7.4.1), building on an earlier result in Kagan et al. (1965), established that, in general, linear estimators of location parameters are admissible if and only if the random variables are normally distributed; and they applied the approach through admissibility to the linear model in Kagan et al. (1973, Section 7.7).

To link the linear model  $y = X\beta + u$  to the proposition, Gauss thus makes three simplifying assumptions:

---

<sup>4</sup>Gauss often uses epithets for his colleagues, typically *clarissimus*. Laplace is *illustrissimus* (Gauss abbreviates *ill.*). Only Newton is *summus*.

1. The design matrix  $X$  has only one column, namely the vector of ones, so that we only have a constant term in the model, there is only one  $\beta$  to estimate, and  $u_i = y_i - \beta$ .
2. The realizations of  $y_i$  only take two distinct values, say  $z_1$  ( $n_1$  times) and  $z_2$  ( $n_2$  times), where  $n = n_1 + n_2$  and  $n_1 \neq n_2$ .
3. The optimum is attained at  $\hat{\beta} = \bar{y}$ .

The third assumption is a perfectly reasonable assumption as we are considering iid random variables  $y_i$  with common mean  $\beta$  and common variance. So, unless we expect Gauss to discuss shrinkage estimators, what alternative is there to estimate  $\beta$ ? Under these three assumptions, Gauss shows that the  $y_i$  must be normally distributed.

After establishing the proposition, Gauss argues that it is thus *reasonable* to assume normality. This is a qualitative statement which can be challenged, but it is not incorrect. Gauss was primarily interested in the justification of least squares, not in pushing the normal distribution, and he fully realized that his qualitative jump from the special to the general case was not mathematically solid. Not completely happy with his restrictive assumptions, Gauss (1823) considered the same model again. This time he asked a different question, namely: which linear unbiased estimator has the smallest variance? This resulted in what we now call the Gauss–Markov theorem and it does not rely on normality of the errors.

## References

- Gauss, C. F. (1809). *Theoria Motus Corporum Coelestium*. Perthes et Besser, Hamburg. (Translated as “Theory of Motion of the Heavenly Bodies Moving About the Sun in Conic Sections” by C. H. Davis (1857). Little, Brown, Boston. Reprinted in 1963, Dover, New York).
- Gauss C. F. (1823). *Theoria Combinationis Observationum Erroribus Minimis Obnoxia*. Dieterich, Göttingen. (Translated as “Theory of the Combination of Observations Least Subject to Errors” by G. W. Stewart (1995). SIAM, Philadelphia).
- Gorroochurn, P. (2016). *Classic Topics on the History of Modern Mathematical Statistics*. John Wiley, Hoboken, NJ.
- Kagan, A. M., Yu. V. Linnik, and C. R. Rao (1965). On a characterization of the normal law based on a property of the sample average. *Sankhya, Series A*, 27, 405–406.



- Kagan, A. M., Yu. V. Linnik, and C. R. Rao (1973). *Characterization Problems in Mathematical Statistics* (translated from the Russian by B. Ramachandran). John Wiley, New York.
- Laplace, P. S. (1774). Mémoire sur la Probabilité des Causes par les Évènemens. Oeuvres Complètes, vol. 8, 27–65. (Translated as “Memoir on the Probability of Causes of Events” by S. Stigler (1986), *Statistical Science*, 1, 364–378.)
- Legendre A. M. (1805). *Nouvelles Méthodes Pour la Détermination des Orbites des Comètes*. Courcier, Paris.
- Pearson, K. (1978). *The History of Statistics in the 17th and 18th Centuries Against the Changing Background of Intellectual, Scientific and Religious Thought. Lectures by Karl Pearson given at University College London during academic sessions 1921–1933* (Edited by E. S. Pearson). Charles Griffin, London.
- Stigler, S. M. (1986). *The History of Statistics: The Measurement of Uncertainty Before 1900*. The Belknap Press of Harvard University Press, Cambridge, MA.