Ghomrasni, Raouf

Working Paper

On local times of ranked continuous semimartingales: application to portfolio generating functions

SFB 649 Discussion Paper, No. 2005,043

Provided in Cooperation with:
Collaborative Research Center 649: Economic Risk, Humboldt University Berlin

Suggested Citation: Ghomrasni, Raouf (2005) : On local times of ranked continuous semimartingales: application to portfolio generating functions, SFB 649 Discussion Paper, No. 2005,043, Humboldt University of Berlin, Collaborative Research Center 649 - Economic Risk, Berlin

This Version is available at:
https://hdl.handle.net/10419/25062

SFB 649 Discussion Paper 2005-043

On Local Times of Ranked Continuous Semimartingales; Application to Portfolio Generating Functions

Raouf Ghomrasni*

* Institute of Mathematics, Technische Universität Berlin, Germany

This research was supported by the Deutsche Forschungsgemeinschaft through the SFB 649 "Economic Risk".

http://sfb649.wiwi.hu-berlin.de
ISSN 1860-5664

SFB 649, Humboldt-Universität zu Berlin
Spandauer Straße 1, D-10178 Berlin

On Local Times of Ranked Continuous Semimartingales; Application to Portfolio Generating Functions

Raouf Ghomrasni*

June 13th, 2005

We derive the decomposition of the ranked continuous semimartingales i.e. order-statistics processes. We apply it to portfolios generated by functions of the ranked market weights. Thus we generalize recent results of Fernholz.

Key words and phrases: Portfolio-generating function, continuous semimartingale, local time, ranked processes.
MSC2000: 60H05, 60J65.

1 Introduction

The distribution of capital, i.e., the family of ranked market weights, starting with the largest weight and going to the smallest, is of central importance in stochastic portfolio theory, as are functionally generated portfolios. Usually we identify stocks by their *names*, i.e., their subscripts, X_1, X_2, X_3, etc. However, with regards to the distribution of capital, it is advantageous to identify the stocks by their *ranks* rather than their names. Portfolios generated by functions of the ranked market weights first appeared in Fernholz ([2]). Fernholz derivations require that the processes $X_1, \cdots, X_n$ be *pathwise mutually nondegenerate*. Here, we shall extend Fernholz's results in its full generality. The relative return of a functionally generated portfolio satisfies a stochastic differential equation wich depends on the *local times* associated with the changes in rank among the stocks.

Section 2 of the paper contains the decomposition of the ranked continuous semimartingales i.e. order-statistics processes. As an interesting byproduct, we obtain an extension of Ouknine's formula [5, 6, 7]. In section 3 we use these decompositions to portfolio generated by functions of the ranked market weights. Thus we generalize recent results of Fernholz [2, 3].

*Fakultät II, Institut für Mathematik, Sekr. MA 7-5, Technische Universität Berlin, Straße des 17. Juni 136 , D-10623 Berlin, Germany, e-mail: ghomrasni@math.tu-berlin.de. Financial support of the Deutsche Forschungsgemeinschaft through the SFB 649 "Economic Risk" is gratefully acknowledged.

2 Decomposition of Ranked Continuous Semimartingales

Definition 2.1 *Let $X_1, \cdots, X_n$ be continuous semimartingales. For $1 \leq k \leq n$, the k-th rank process of $X_1, \cdots, X_n$ is defined by*

$$X^{(k)} = \max_{i_1 < \cdots < i_k} \min(X_{i_1}, \cdots, X_{i_k})$$

where $1 \leq i_1$ and $i_k \leq n$.

Note that, according to Definition 2.1, for $t \in I\!R^+$,

$$\max_{1 \leq i \leq n} X_i(t) = X^{(1)}(t) \geq X^{(2)}(t) \geq \cdots \geq X^{(n)}(t) = \min_{1 \leq i \leq n} X_i(t)$$

so that at any given time, the values of the rank processes represent the values of the original processes arranged in descending order (i.e. the (reverse) order statistics).

The following proposition shows that the rank processes derived from continuous semimartingales can be expressed in terms of the original processes, adjusted by local times.

Proposition 2.1 *Let $X_1, \cdots, X_n$ be continuous semimartingales. For $k \in \{1, 2, \cdots, n\}$, Let $u(k) = (u_t(k),\, t \geq 0) : \Omega \times [0, \infty[\rightarrow \{1, 2, \cdots, n\}$ be any predictable process with the property:*

$$X_t^{(k)} = X_{u_t(k)}(t)$$

Then the k-th rank processes $X^{(k)}, k = 1, \cdots, n$, are continuous semimartingales and we have:

$$\begin{aligned} X^{(k)}(t) = X^{(k)}(0) &+ \sum_{i=1}^{n} \int_0^t 1_{\{u_s(k)=i\}}\, dX_i(s) \;+\; \frac{1}{2} \sum_{i=1}^{n} \int_0^t 1_{\{u_s(k)=i\}}\, d_s\, L_s^0((X^{(k)} - X_i)^+) \\ &- \frac{1}{2} \sum_{i=1}^{n} \int_0^t 1_{\{u_s(k)=i\}}\, d_s\, L_s^0((X^{(k)} - X_i)^-). \end{aligned}$$

where $L_t^0(X)$ is the local time of the continuous semimartingale X at 0.

Proof: We adapt here the proof given by Chitashvili and Mania ([1]) for the decomposition of the maximum of semimartingales (i.e. $k = 1$), we have,

$$X_t^{(k)} - X_0^{(k)} = \sum_{i=1}^{n} \int_0^t 1_{\{u_s(k)=i\}}\, dX_s^i \;+\; \sum_{i=1}^{n} \int_0^t 1_{\{u_s(k)=i\}}\, d\,(X_s^{(k)} - X_s^i)$$

Where we used the property $\sum_{i=1}^{n} 1_{\{u_s(k)=i\}} = 1$ it follows,

$$X_t^{(k)} - X_0^{(k)} = \sum_{i=1}^{n} \int_0^t 1_{\{u_s(k)=i\}}\, dX_s^i + \sum_{i=1}^{n} \int_0^t 1_{\{u_s(k)=i\}}\, d\,(X_s^{(k)} - X_s^i)^+ - \sum_{i=1}^{n} \int_0^t 1_{\{u_s(k)=i\}}\, d\,(X_s^{(k)} - X_s^i)^-$$

We note the fact:

$$\{u_s(k) = i\} \subset \{X_s^{(k)} = X_i(s)\}$$

Therefore, using the following formula

$$\frac{1}{2}\, L_t^0(X) = \int_0^t 1_{\{(X_s=0)\}}\, d\, X_s$$

which is valid for non-negative continuous semimartingales X.

□

We can give a more explicit decomposition as follows:

Corollary 2.1 *Let $X_1, \cdots, X_n$ be continuous semimartingales. Then the k-th rank processes $X^{(k)}, k = 1, \cdots, n$, are continuous semimartingales and we have:*

$$X^{(k)}(t) = X^{(k)}(0) + \sum_{i=1}^{n} \int_0^t 1_{\{u_s(k)=i\}}\, dX_i(s) + \frac{1}{2} \sum_{i=k+1}^{n} L_t^0(X^{(k)} - X^{(i)}) - \frac{1}{2} \sum_{i=1}^{k-1} L_t^0(X^{(i)} - X^{(k)}).$$

where $L_t^0(X)$ is the local time of the continuous semimartingale X at 0.

Proof: We fix $i = 1, \cdots, n$ and we deal with $\int_0^t 1_{\{u_s(k)=i\}}\, d_s\, L_s^0((X^{(k)} - X_i)^+)$

$$\begin{aligned}\int_0^t 1_{\{u_s(k)=i\}}\, d_s\, L_s^0((X^{(k)} - X_i)^+) &= \sum_{j=1}^{n} \int_0^t 1_{\{u_s(k)=i\} \cap \{X_i = X^{(j)}\}}\, d_s\, L_s^0((X^{(k)} - X_i)^+) \\ &= \sum_{j=1}^{n} \int_0^t 1_{\{u_s(k)=i\} \cap \{X_i = X^{(j)}\}}\, d_s\, L_s^0((X^{(k)} - X^{(j)})^+)\end{aligned}$$

We, conclude by noting:

$$\sum_{i=1}^{n} \int_0^t 1_{\{u_s(k)=i\} \cap \{X_i = X^{(j)}\}}\, d_s\, L_s^0((X^{(k)} - X^{(j)})^+) = L_t^0((X^{(k)} - X^{(j)})^+)$$

Indeed,

$$\begin{aligned}&\int_0^t 1_{\{u_s(k)=i\} \cap \{X_i = X^{(j)}\}}\, d_s\, L_s^0((X^{(k)} - X^{(j)})^+) \\ &= \int_0^t 1_{\{u_s(k)=i\} \cap \{X_i = X^{(j)}\}} \times 1_{\{X^{(k)} - X^{(j)} = 0\}}\, d_s\, L_s^0((X^{(k)} - X^{(j)})^+) \\ &= \int_0^t 1_{\{u_s(k)=i\} \cap \{X_i = X^{(k)}\}} \times 1_{\{X^{(k)} - X^{(j)} = 0\}}\, d_s\, L_s^0((X^{(k)} - X^{(j)})^+) \\ &= \int_0^t 1_{\{u_s(k)=i\}} d_s\, L_s^0((X^{(k)} - X^{(j)})^+)\end{aligned}$$

□

In the particular case considered by Fernholz (see [2]) of rank processes derived from pathwise mutually nondegenerate absolutely continuous semimartingales

Corollary 2.2 (**Fernholz**)
Let $X_1, \cdots, X_n$ be pathwise mutually nondegenerate absolutely continuous semimartingales, and for $t \in [0, T]$, let p_t be the random permutation of $\{1., n\}$ such that for $k = 1, \cdots, n$,

$$X_{p_t(k)(t)} = X^{(k)}(t), \qquad \text{and} \qquad p_t(k) < p_t(k+1) \quad \text{if} \quad X^{(k)}(t) = X^{(k+1)}(t).$$

Then the k-th rank processes $X^{(k)}, k = 1, \cdots, n$, are continuous semimartingales and we have:

$$X^{(k)}(t) = X^{(k)}(0) + \sum_{i=1}^{n} \int_0^t 1_{\{p_s(k)=i\}}\, dX_i(s) + \frac{1}{2} L_t^0(X^{(k)} - X^{(k+1)}) - \frac{1}{2}\, L_t^0(X^{(k-1)} - X^{(k)}).$$

where $L_t^0(X)$ is the local time of the continuous semimartingale X at 0.

Now, we state the solution to Problem 4.1.13 stated in Fernholz's book:

Corollary 2.3 *Let $X_1, \cdots, X_n$ be continuous semimartingales. Then the k-th rank processes $X^{(k)}, k = 1, \cdots, n$, are continuous semimartingales and we have:*

$$\begin{aligned} X^{(k)}(t) =& X^{(k)}(0) + \sum_{i=1}^{n} \int_0^t 1_{\{p_s(k)=i\}}\, dX_i(s) + \frac{1}{2} L_t^0(X^{(k)} - X^{(k+2)}) + \frac{1}{2} L_t^0(X^{(k)} - X^{(k+1)}) \\ & - \frac{1}{2}\, L_t^0(X^{(k-2)} - X^{(k)}) - \frac{1}{2}\, L_t^0(X^{(k-1)} - X^{(k)}). \end{aligned}$$

where $L_t^0(X)$ is the local time of the continuous semimartingale X at 0.

Theorem 2.1 *Let $X_1, \cdots, X_n$ be continuous semimartingales. Then we have:*

$$\sum_{i=1}^{n} L_t^0(X^{(i)}) = \sum_{i=1}^{n} L_t^0(X_i)$$

where $L_t^0(X)$ is the local time of the continuous semimartingale X at 0.

Proof:

We recall first that $L_t^0(Z) = L_t^0(Z^+)$ for every semimartingale Z. Hence, it is enough to consider the case where $X_1, \cdots, X_n$ are non-negatives continuous semimartingales. We need to check the following equality holds:

$$\sum_{i=1}^{n} 1_{\{X^{(i)}=0\}}\, dX^{(i)} = \sum_{i=1}^{n} 1_{\{X_i=0\}}\, dX_i$$

From Corollary 2.1, we have

$$d\, X^{(k)} = \sum_{i=1}^{n} 1_{\{u_t(k)=i\}}\, dX_i + \sum_{i=k+1}^{n} 1_{\{X^{(k)}-X^{(i)}=0\}} d\,(X^{(k)} - X^{(i)}) - \sum_{i=1}^{k-1} 1_{\{X^{(i)}-X^{(k)}=0\}} d\,(X^{(i)} - X^{(k)}).$$

It follows:

$$\begin{aligned} 1_{\{X^{(k)}=0\}}\, d\, X^{(k)} =& \sum_{i=1}^{n} 1_{\{u_t(k)=i\}}\, 1_{\{X^{(k)}=0\}}\, 1_{\{X_i=X^{(k)}\}} dX_i + \sum_{i=k+1}^{n} 1_{\{X^{(k)}-X^{(i)}=0\}} 1_{\{X^{(k)}=0\}}\, d\,(X^{(k)} - X^{(i)}) \\ & - \sum_{i=1}^{k-1} 1_{\{X^{(i)}-X^{(k)}=0\}} 1_{\{X^{(k)}=0\}}\, d\,(X^{(i)} - X^{(k)}). \\ =& \sum_{i=1}^{n} 1_{\{u_t(k)=i\}}\, 1_{\{X_i=0\}} dX_i + \sum_{i=k+1}^{n} 1_{\{X^{(i)}=0\}} 1_{\{X^{(k)}=0\}}\, d\,(X^{(k)} - X^{(i)}) \\ & - \sum_{i=1}^{k-1} 1_{\{X^{(i)}=0\}} 1_{\{X^{(k)}=0\}}\, d\,(X^{(i)} - X^{(k)}). \end{aligned}$$

Now we may choose our $u_t(\cdot)$ as a bijection on $\{1, \cdots, n\}$, we denote its inverse by $v_t(\cdot)$

$$1_{\{X^{(k)}=0\}}\, d\, X^{(k)} = \sum_{i=1}^{n} 1_{\{v_t(i)=k\}}\, 1_{\{X_i=0\}} dX_i \; + \sum_{i=k+1}^{n} 1_{\{X^{(i)}=0\}} 1_{\{X^{(k)}=0\}}\, d\,(X^{(k)} - X^{(i)}) \\ - \sum_{i=1}^{k-1} 1_{\{X^{(i)}=0\}} 1_{\{X^{(k)}=0\}}\, d\,(X^{(i)} - X^{(k)}).$$

By summation over the index $k = 1, \cdots, n$ we obtain our result.

□

In particular,

Corollary 2.4 Ouknine's formula
Let X and Y be semimartingales. It is shown that

$$L_t^0(X \vee Y) + L_t^0(X \wedge Y) = L_t^0(X) + L_t^0(Y)$$

where $L_t^0(X)$ $(t \geq 0)$ denotes the local time at 0 of X.

3 Portfolio Generating Functions

Theorem 3.1 *Let $\mathcal{M}$ be a market of stocks $X_1, \cdots, X_n$. Let $\mathbf{S}$ be a function defined on a neighborhood U of Δ^n. Suppose that there exists a positive C^2 function S defined on U . Then $\mathbf{S}$ generates the portfolio π such that for $k = 1, \cdots, n$,*

$$\pi_{p_t(k)}(t) \;=\; \big(D_k \log S(\mu_{(\cdot)}(t)) + 1 - \sum_{j=1}^{n} \mu_{(j)}(t)\, D_j \log S(\mu_{(\cdot)}(t))\big)\mu_{(k)}(t),$$

for all $t \in [0, T]$, a.s., with a drift process Θ that satisfies

$$d\Theta(t) = \frac{-1}{2\mathbf{S}(\mu(t))} \sum_{i,j=1}^{n} D_{ij} S(\mu_{(\cdot)}(t))\, \mu_{(i)}(t) \mu_{(j)}(t) \tau_{(ij)}(t) dt \\ - \frac{1}{2} \sum_{k=1}^{n} \pi_{p_t(k)}(t) \sum_{i=k+1}^{n} dL_t^0((\log(\mu_{(k)}) - \log(\mu_{(i)}))) \\ + \frac{1}{2} \sum_{k=1}^{n} \pi_{p_t(k)}(t) \sum_{i=1}^{k-1} dL_t^0((\log(\mu_{(i)}) - \log(\mu_{(k)})))$$

Hence,

$$
\begin{aligned}
d\,\log(Z_\pi(t)/Z_\mu(t)) &= \sum_{i=1}^{n} D_i \log \mathbf{S}(\mu_{(\cdot)}(t))\, d\mu_{(i)}(t) \\
&\quad - \frac{1}{2}\sum_{i,j=1}^{n} D_i \log \mathbf{S}(\mu_{(\cdot)}(t)) D_j \log \mathbf{S}(\mu_{(\cdot)}(t)) \mu_{(i)}(t)\mu_{(j)}(t)\tau_{(ij)}(t)dt \\
&\quad - \frac{1}{2}\sum_{k=1}^{n} \pi_{p_t(k)}(t) \sum_{i=1}^{n} 1_{\{u_t(k)=i\}} dL_t^0((\log(\mu_{(k)}) - \log(\mu_i))^+) \\
&\quad + \frac{1}{2}\sum_{k=1}^{n} \pi_{p_t(k)}(t) \sum_{i=1}^{n} 1_{\{u_t(k)=i\}} dL_t^0((\log(\mu_{(k)}) - \log(\mu_i))^-)
\end{aligned}
$$

Remark 3.1 *The theorem above extend Fernholz's theorem where the assumption for $X_1, \cdots, X_n$ of pathwise mutually nondegenerate was needed.*

Proof :
Itô's formula implies that:

$$
\begin{aligned}
d\log \mathbf{S}(\mu_{(\cdot)}(t)) &= \sum_{i=1}^{n} D_i \log \mathbf{S}(\mu_{(\cdot)}(t)) d\,\mu_{(i)}(t) \\
&\quad + \frac{1}{2\,\mathbf{S}(\mu_{(\cdot)}(t))} \sum_{i,j=1}^{n} D_{ij}\mathbf{S}(\mu_{(\cdot)}(t))\, d\, < \mu_i, \mu_j >_t \\
&\quad - \frac{1}{2}\sum_{i,j=1}^{n} D_i \log \mathbf{S}(\mu_{(\cdot)}(t)) D_j \log \mathbf{S}(\mu_{(\cdot)}(t))\, d\, < \mu_i, \mu_j >_t
\end{aligned}
$$

Now let us consider the relative return process $\log(Z_\pi(t)/Z(t))$. We have

$$
\begin{aligned}
d\,\log(Z_\pi(t)/Z_\mu(t)) &= \sum_{i=1}^{n} \pi_i(t) d\log\mu_i(t) + \gamma_\pi^\star\, dt \\
&= \sum_{k=1}^{n} \pi_{p_t(k)}(t)\, d\,\log\mu_{(k)}(t) + \frac{1}{2}\sum_{k=1}^{n} \pi_{p_t(k)}(t) \sum_{i=1}^{n} 1_{\{u_t(k)=i\}} dL_t^0((\log(\mu_{(k)}) - \log(\mu_i))^+) \\
&\quad - \frac{1}{2}\sum_{k=1}^{n} \pi_{p_t(k)}(t) \sum_{i=1}^{n} 1_{\{u_t(k)=i\}} dL_t^0((\log(\mu_{(k)}) - \log(\mu_i))^-) + \gamma_\pi^\star\, dt \\
&= \sum_{k=1}^{n} \frac{\pi_{p_t(k)}(t)}{\mu_{(k)}(t)}\, d\,\mu_{(k)}(t) + \frac{1}{2}\sum_{k=1}^{n} \pi_{p_t(k)}(t) \sum_{i=1}^{n} 1_{\{u_t(k)=i\}} dL_t^0((\log(\mu_{(k)}) - \log(\mu_i))^+) \\
&\quad - \frac{1}{2}\sum_{k=1}^{n} \pi_{p_t(k)}(t) \sum_{i=1}^{n} 1_{\{u_t(k)=i\}} dL_t^0((\log(\mu_{(k)}) - \log(\mu_i))^-) + \gamma_\pi^\star\, dt \\
&\quad - \frac{1}{2}\sum_{k=1}^{n} \pi_{p_t(k)}(t) d < \log\mu_k(\cdot) >_t
\end{aligned}
$$

Let assume that the weights $\pi_i, i = 1, \cdots, n$ satisfy:

$$
\frac{\pi_{p_t(k)}(t)}{\mu_{(k)}(t)} = D_k \log \mathbf{S}(\mu_{(\cdot)}(t)) + \varphi(t)
$$

In this case:

$$\begin{aligned}\sum_{i=1}^{n}\frac{\pi_{p_t(i)}(t)}{\mu_{(i)}(t)}\,d\mu_{(i)}(t) &= \sum_{i=1}^{n} D_i\log \mathbf{S}(\mu_{(\cdot)}(t))\,d\mu_{(i)}(t) + \varphi(t)\sum_{i=1}^{n} d\mu_{(i)}(t)\\ &= \sum_{i=1}^{n} D_i\log \mathbf{S}(\mu_{(\cdot)}(t))\,d\mu_{(i)}(t)\end{aligned}$$

Now we consider the last summation $\gamma_\pi^\star\,dt - \frac{1}{2}\sum_{k=1}^{n}\pi_{p_t(k)}(t)d < \log\mu_k(\cdot) >_t$. It follows

$$\gamma_\pi^\star\,dt - \frac{1}{2}\sum_{k=1}^{n}\pi_{p_t(k)}(t)d < \log\mu_k(\cdot) >_t = \sum_{i,j=1}^{n} D_i\log \mathbf{S}(\mu_{(\cdot)}(t))D_j\log \mathbf{S}(\mu_{(\cdot)}(t))\mu_{(i)}(t)\mu_{(j)}(t)\tau_{(ij)}(t)$$

Finally,

$$\begin{aligned}d\,\log(Z_\pi(t)/Z_\mu(t)) &= \sum_{i=1}^{n} D_i\log \mathbf{S}(\mu_{(\cdot)}(t))\,d\mu_{(i)}(t)\\ &\quad - \frac{1}{2}\sum_{i,j=1}^{n} D_i\log \mathbf{S}(\mu_{(\cdot)}(t))D_j\log \mathbf{S}(\mu_{(\cdot)}(t))\mu_{(i)}(t)\mu_{(j)}(t)\tau_{(ij)}(t)dt\\ &\quad + \frac{1}{2}\sum_{k=1}^{n}\pi_{p_t(k)}(t)\sum_{i=1}^{n} 1_{\{u_t(k)=i\}}dL_t^0((\log(\mu_{(k)}) - \log(\mu_i))^+)\\ &\quad - \frac{1}{2}\sum_{k=1}^{n}\pi_{p_t(k)}(t)\sum_{i=1}^{n} 1_{\{u_t(k)=i\}}dL_t^0((\log(\mu_{(k)}) - \log(\mu_i))^-)\end{aligned}$$

□

Corollary 3.1 *In the particular case studied by Fernholz, where we assume the stocks $X_1, \cdots, X_n$ are pathwise mutually nondegerate, we have:*

$$\begin{aligned}d\,\log(Z_\pi(t)/Z_\mu(t)) &= \sum_{i=1}^{n} D_i\log \mathbf{S}(\mu_{(\cdot)}(t))\,d\mu_{(i)}(t)\\ &\quad - \frac{1}{2}\sum_{i,j=1}^{n} D_i\log \mathbf{S}(\mu_{(\cdot)}(t))D_j\log \mathbf{S}(\mu_{(\cdot)}(t))\mu_{(i)}(t)\mu_{(j)}(t)\tau_{(ij)}(t)dt\\ &\quad + \frac{1}{2}\sum_{k=1}^{n-1}(\pi_{p_t(k+1)}(t) - \pi_{p_t(k)}(t))dL_t^0((\log(\mu_{(k)}) - \log(\mu_{(k+1)}))\end{aligned}$$

3.1 Examples of Rank-Dependent Portfolios

As considered by Fernholz, we have more generally:

Example. (The biggest stock) Let $\mathbf{S} = x_{(1)}$.

$$d\,\log(Z_\pi(t)/Z_\mu(t)) = \frac{1}{(\mu_{(1)}(t))}\,d\mu_{(1)}(t) - \frac{1}{2}\tau_{(11)}(t)\,dt \\ - \frac{1}{2}\pi_{p_t(1)}(t)\sum_{i=1}^{n} 1_{\{u_t(1)=i\}} dL_t^0((\log(\mu_{(1)}) - \log(\mu_i)))$$

Hence,

$$d\,\log(Z_\pi(t)/Z_\mu(t)) = d\,\log\mu_{(1)}(t) - \frac{1}{2}\pi_{p_t(1)}(t)\sum_{i=1}^{n} 1_{\{u_t(1)=i\}} dL_t^0((\log(\mu_{(1)}) - \log(\mu_i)))$$

Equivalently,

$$d\,\log(Z_\pi(t)/Z_\mu(t)) = d\,\log\mu_{(1)}(t) - \frac{1}{2}\pi_{p_t(1)}(t)\sum_{i=1}^{n} dL_t^0((\log(\mu_{(1)}) - \log(\mu_{(i)})))$$

Since the local time component of the drift process is decreasing, the long-term relative performance of π will suffer if there are many changes of leadership in the market.

Example. (The size effect)
The *size effect* is the observed tendency of small stocks to have higher long-term returns than large stocks. Let $1 < m < n$ and suppose

$$\mathbf{S}_L(x) = x_{(1)} + \cdots + x_{(m)}.$$

Then the drift $d\Theta$ is given by:

$$d\Theta_t := -\frac{1}{2}\sum_{k=1}^{m}\frac{\mu_{(k)}}{S_L}\sum_{i=m+1}^{n} d\,L_t^0(\log\mu_{(k)} - \log\mu_{(i)})$$

If we specify, the Fernholz's case, we obtain:

$$d\Theta_t := -\frac{1}{2}\frac{\mu_{(m)}}{S_L}\,d\,L_t^0(\log\mu_{(m)} - \log\mu_{(m+1)})$$

Similarly,

$$\mathbf{S}_S(x) = x_{(m+1)} + \cdots + x_{(n)}.$$

Then the drift $d\Theta$ is given by:

$$d\Theta_t := +\frac{1}{2}\sum_{k=m+1}^{n}\frac{\mu_{(k)}}{S_L}\sum_{i=1}^{m-1} d\,L_t^0(\log\mu_{(i)} - \log\mu_{(k)})$$

Since the drift process is monotonically increasing, it is likely that the return on small-stock index will eventually be greater than that of the large-stock index. Hence, the higher long-term return of small-stock index is due to the increasing drift process, and the relative level of small-stock risk is irrelevant.

References

[1] R. J. Chitashvili *and* M. G. Mania, *Decomposition of the maximum of semimartingales and generalized Itô's formula*. New trends in probability and statistics, Vol. **1** (Bakuriani, 1990), 301–350.

[2] R. Fernholz, *Equity portfolios generated by functions of ranked market weights.* Finance Stoch. **5** (2001) 469–486.

[3] R. Fernholz, *Stochastic portfolio theory.* Springer-Verlag, New York, 2002.

[4] R. Fernholz *and* I. Karatzas, *The implied liquidity premium for equities* Preprint (http://www.intechjanus.com/research/research.html).

[5] Y. Ouknine, *Temps local du produit et du sup de deux semimartingales.* Séminaire de Probabilités, XXIV, 1988/89 Lecture Notes in Math. **1426** (1990), 477–479.

[6] Y. Ouknine, *Généralisation d'un lemme de S. Nakao et applications.* Stochastics **23** (1988), 149–157.

[7] J. A. Yan, *A formula for local times of semimartingales.* Northeast. Math. J. **1** (1985), 138–140.

SFB 649 Discussion Paper Series

For a complete list of Discussion Papers published by the SFB 649, please visit http://sfb649.wiwi.hu-berlin.de.

001 "Nonparametric Risk Management with Generalized Hyperbolic Distributions" by Ying Chen, Wolfgang Härdle and Seok-Oh Jeong, January 2005.
002 "Selecting Comparables for the Valuation of the European Firms" by Ingolf Dittmann and Christian Weiner, February 2005.
003 "Competitive Risk Sharing Contracts with One-sided Commitment" by Dirk Krueger and Harald Uhlig, February 2005.
004 "Value-at-Risk Calculations with Time Varying Copulae" by Enzo Giacomini and Wolfgang Härdle, February 2005.
005 "An Optimal Stopping Problem in a Diffusion-type Model with Delay" by Pavel V. Gapeev and Markus Reiß, February 2005.
006 "Conditional and Dynamic Convex Risk Measures" by Kai Detlefsen and Giacomo Scandolo, February 2005.
007 "Implied Trinomial Trees" by Pavel Čížek and Karel Komorád, February 2005.
008 "Stable Distributions" by Szymon Borak, Wolfgang Härdle and Rafal Weron, February 2005.
009 "Predicting Bankruptcy with Support Vector Machines" by Wolfgang Härdle, Rouslan A. Moro and Dorothea Schäfer, February 2005.
010 "Working with the XQC" by Wolfgang Härdle and Heiko Lehmann, February 2005.
011 "FFT Based Option Pricing" by Szymon Borak, Kai Detlefsen and Wolfgang Härdle, February 2005.
012 "Common Functional Implied Volatility Analysis" by Michal Benko and Wolfgang Härdle, February 2005.
013 "Nonparametric Productivity Analysis" by Wolfgang Härdle and Seok-Oh Jeong, March 2005.
014 "Are Eastern European Countries Catching Up? Time Series Evidence for Czech Republic, Hungary, and Poland" by Ralf Brüggemann and Carsten Trenkler, March 2005.
015 "Robust Estimation of Dimension Reduction Space" by Pavel Čížek and Wolfgang Härdle, March 2005.
016 "Common Functional Component Modelling" by Alois Kneip and Michal Benko, March 2005.
017 "A Two State Model for Noise-induced Resonance in Bistable Systems with Delay" by Markus Fischer and Peter Imkeller, March 2005.

SFB 649, Spandauer Straße 1, D-10178 Berlin
http://sfb649.wiwi.hu-berlin.de

This research was supported by the Deutsche Forschungsgemeinschaft through the SFB 649 "Economic Risk".



018 "Yxilon – a Modular Open-source Statistical Programming Language" by Sigbert Klinke, Uwe Ziegenhagen and Yuval Guri, March 2005.
019 "Arbitrage-free Smoothing of the Implied Volatility Surface" by Matthias R. Fengler, March 2005.
020 "A Dynamic Semiparametric Factor Model for Implied Volatility String Dynamics" by Matthias R. Fengler, Wolfgang Härdle and Enno Mammen, March 2005.
021 "Dynamics of State Price Densities" by Wolfgang Härdle and Zdeněk Hlávka, March 2005.
022 "DSFM fitting of Implied Volatility Surfaces" by Szymon Borak, Matthias R. Fengler and Wolfgang Härdle, March 2005.
023 "Towards a Monthly Business Cycle Chronology for the Euro Area" by Emanuel Mönch and Harald Uhlig, April 2005.
024 "Modeling the FIBOR/EURIBOR Swap Term Structure: An Empirical Approach" by Oliver Blaskowitz, Helmut Herwartz and Gonzalo de Cadenas Santiago, April 2005.
025 "Duality Theory for Optimal Investments under Model Uncertainty" by Alexander Schied and Ching-Tang Wu, April 2005.
026 "Projection Pursuit For Exploratory Supervised Classification" by Eun-Kyung Lee, Dianne Cook, Sigbert Klinke and Thomas Lumley, May 2005.
027 "Money Demand and Macroeconomic Stability Revisited" by Andreas Schabert and Christian Stoltenberg, May 2005.
028 "A Market Basket Analysis Conducted with a Multivariate Logit Model" by Yasemin Boztuğ and Lutz Hildebrandt, May 2005.
029 "Utility Duality under Additional Information: Conditional Measures versus Filtration Enlargements" by Stefan Ankirchner, May 2005.
030 "The Shannon Information of Filtrations and the Additional Logarithmic Utility of Insiders" by Stefan Ankirchner, Steffen Dereich and Peter Imkeller, May 2005.
031 "Does Temporary Agency Work Provide a Stepping Stone to Regular Employment?" by Michael Kvasnicka, May 2005.
032 "Working Time as an Investment? – The Effects of Unpaid Overtime on Wages, Promotions and Layoffs" by Silke Anger, June 2005.
033 "Notes on an Endogenous Growth Model with two Capital Stocks II: The Stochastic Case" by Dirk Bethmann, June 2005.
034 "Skill Mismatch in Equilibrium Unemployment" by Ronald Bachmann, June 2005.

This research was supported by the Deutsche Forschungsgemeinschaft through the SFB 649 "Economic Risk".



035 "Uncovered Interest Rate Parity and the Expectations Hypothesis of the Term Structure: Empirical Results for the U.S. and Europe" by Ralf Brüggemann and Helmut Lütkepohl, April 2005.
036 "Getting Used to Risks: Reference Dependence and Risk Inclusion" by Astrid Matthey, May 2005.
037 "New Evidence on the Puzzles. Results from Agnostic Identification on Monetary Policy and Exchange Rates." by Almuth Scholl and Harald Uhlig, July 2005.
038 "Discretisation of Stochastic Control Problems for Continuous Time Dynamics with Delay" by Markus Fischer and Markus Reiss, August 2005.
039 "What are the Effects of Fiscal Policy Shocks?" by Andrew Mountford and Harald Uhlig, July 2005.
040 "Optimal Sticky Prices under Rational Inattention" by Bartosz Maćkowiak and Mirko Wiederholt, July 2005.
041 "Fixed-Prize Tournaments versus First-Price Auctions in Innovation Contests" by Anja Schöttner, August 2005.
042 "Bank finance versus bond finance: what explains the differences between US and Europe?" by Fiorella De Fiore and Harald Uhlig, August 2005.
043 "On Local Times of Ranked Continuous Semimartingales; Application to Portfolio Generating Functions" by Raouf Ghomrasni, June 2005.

This research was supported by the Deutsche Forschungsgemeinschaft through the SFB 649 "Economic Risk".