

Onuchic, Paula

**Article**  
**Informed intermediaries**

Theoretical Economics

**Provided in Cooperation with:**

The Econometric Society

*Suggested Citation:* Onuchic, Paula (2022) : Informed intermediaries, Theoretical Economics, ISSN 1555-7561, The Econometric Society, New Haven, CT, Vol. 17, Iss. 1, pp. 57-87,  
<https://doi.org/10.3982/TE4072>

This Version is available at:

<https://hdl.handle.net/10419/253516>

**Standard-Nutzungsbedingungen:**

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

**Terms of use:**

*Documents in EconStor may be saved and copied for your personal and scholarly purposes.*

*You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.*

*If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.*



<https://creativecommons.org/licenses/by-nc/4.0/>

## Supplement to “Informed intermediaries” (*Theoretical Economics*, Vol. 17, No. 1, January 2022, 57–87)

PAULA ONUCHIC  
Department of Economics, New York University

### APPENDIX: PROOF OF PROPOSITION 2 (CONTINUED)

#### S.1 Guesses

Surplus sharing:

$$\begin{aligned}\beta(I, v_s, U, v_b) &= \beta^I \geq \frac{1}{2}, & \beta(U, v_s, I, v_b) &= 1 - \beta^I \leq \frac{1}{2}, \\ \beta(I, v_s, I, v_b) &= \frac{1}{2}.\end{aligned}$$

Efficient trading if and only if an informed agent is involved:

$$\mathcal{I}(i_s, v_s, i_b, v_b) = 1 \Leftrightarrow (i_s, i_b) \neq (U, U) \quad \text{and} \quad V_{v_s 0}^{i_s} - V_{v_s 1}^{i_s} + V_{v_b 1}^{i_b} - V_{v_b 0}^{i_b} > 0.$$

#### S.2 Stationary distribution

Given the guesses for  $\mathcal{I}$ , the inflow equal to outflow equations for the stationary distribution become

$$\mu_{L1}^U(\eta + \lambda(\mu_{H0}^I + \mu_{L0}^I)) = \eta\mu_{L0}^U, \tag{S.1}$$

$$\mu_{H0}^U(\eta + \lambda(\mu_{L1}^I + \mu_{H1}^I)) = \eta\mu_{H1}^U, \tag{S.2}$$

$$\mu_{L1}^I(\eta + \lambda(\mu_{H0}^U + \mu_{H0}^I)) = (\eta + \lambda\mu_{L1}^U)\mu_{L0}^I, \tag{S.3}$$

$$\mu_{H0}^I(\eta + \lambda(\mu_{L1}^U + \mu_{L1}^I)) = (\eta + \lambda\mu_{H0}^U)\mu_{H1}^I. \tag{S.4}$$

Combine (S.1) and (S.2), and using  $\mu_{L1}^U + \mu_{L0}^U = \mu_{H0}^U + \mu_{H1}^U = \frac{1-\phi}{2}$  (since half of the uninformed agents have high valuation and half have low valuation), I get

$$\mu_{L1}^U(2\eta + \lambda(\mu_{H0}^I + \mu_{L0}^I)) = \mu_{H0}^U(2\eta + \lambda(\mu_{L1}^I + \mu_{H1}^I)) = \frac{\eta(1-\phi)}{2}. \tag{S.5}$$

Similarly use (S.3) and (S.4) and  $\mu_{L1}^I + \mu_{L0}^I = \mu_{H0}^I + \mu_{H1}^I$  to get

$$(\eta + \lambda\mu_{L1}^U)(\mu_{H0}^I + \mu_{L0}^I) = (\eta + \lambda\mu_{H0}^U)(\mu_{H1}^I + \mu_{L1}^I). \tag{S.6}$$

---

Paula Onuchic: [p.onuchic@nyu.edu](mailto:p.onuchic@nyu.edu)

Use (S.5) and (S.6) to get  $\mu_{L1}^U + \mu_{L1}^I + \mu_{H1}^I = \mu_{H0}^U + \mu_{H0}^I + \mu_{L0}^I$ . Combining this with (S.5) yet again, conclude that  $\mu_{L1}^U = \mu_{H0}^U \equiv \hat{\mu}^U$ . This, along with (S.6), implies  $(\mu_{H0}^I + \mu_{L0}^I) = (\mu_{L1}^I + \mu_{H1}^I)$ ; hence,  $\mu_{H0}^I = \mu_{L1}^I \equiv \hat{\mu}^I$ . Rewrite the inflow equals outflow conditions now as

$$\hat{\mu}^U(\eta + \lambda(\phi/2)) = \eta \frac{(1 - \phi - 2\hat{\mu}^U)}{2} \hat{\mu}^I, \quad (\eta + \lambda(\hat{\mu}^U + \hat{\mu}^I)) = \frac{\phi - 2\hat{\mu}^I}{2}(\eta + \lambda\hat{\mu}^U).$$

Solving these, I get

$$\hat{\mu}^U = \frac{1 - \phi}{4 + \frac{\lambda}{\eta}}, \quad \hat{\mu}^I = -\frac{\eta + \lambda, \hat{\mu}^U}{\lambda} + \sqrt{\left(\frac{\eta + \lambda\hat{\mu}^U}{\lambda}\right)^2 + \frac{\phi}{2} \frac{\eta + \lambda\hat{\mu}^U}{\lambda}}.$$

### S.3 Unflagged values

Again taking into account the guesses for  $\mathcal{I}$  and  $\beta$ , unflagged values are given by the system

$$\begin{aligned} rV_{H0}^I &= \eta(V_{H1}^I - V_{H0}^I) + \lambda\hat{\mu}^I \frac{V_{H1}^I - V_{H0}^I + V_{L0}^I - V_{L1}^I}{2} + \lambda\hat{\mu}^U \beta^I (V_{H1}^I - V_{H0}^I + V_{L0}^U - V_{L1}^U), \\ rV_{H1}^I &= \delta_H + \eta(V_{H0}^I - V_{H1}^I) + \lambda\hat{\mu}^U \beta^I (V_{H0}^I - V_{H1}^I + V_{H1}^U - V_{H0}^U), \\ rV_{L1}^I &= \delta_L + \eta(V_{L0}^I - V_{L1}^I) + \lambda\hat{\mu}^I \frac{V_{L0}^I - V_{L1}^I + V_{H1}^I - V_{H0}^I}{2} \\ &\quad + \lambda\hat{\mu}^U \beta^I (V_{L0}^I - V_{L1}^I + V_{H1}^U - V_{H0}^U), \\ rV_{L0}^I &= \eta(V_{L1}^I - V_{L0}^I) + \lambda\hat{\mu}^U \beta^I (V_{L1}^I - V_{L0}^I + V_{L0}^U - V_{L1}^U), \\ rV_{H0}^U &= \eta(V_{H1}^U - V_{H0}^U) + \lambda\hat{\mu}^I (1 - \beta^I) (V_{H1}^U - V_{H0}^U + V_{L0}^I - V_{L1}^I) \\ &\quad + \lambda\hat{\mu}^I (1 - \beta^I) (V_{H1}^U - V_{H0}^U + V_{H0}^I - V_{H1}^I), \\ rV_{H1}^U &= \delta_H + \eta(V_{H0}^U - V_{H1}^U), \\ rV_{L1}^U &= \delta_L + \eta(V_{L0}^U - V_{L1}^U) + \lambda\hat{\mu}^I (1 - \beta^I) (V_{L0}^U - V_{L1}^U + V_{H1}^I - V_{H0}^I) \\ &\quad + \lambda\hat{\mu}^I (1 - \beta^I) (V_{L0}^U - V_{L1}^U + V_{L1}^I - V_{L0}^I), \\ rV_{L0}^U &= \eta(V_{L1}^U - V_{L0}^U). \end{aligned}$$

In terms of the values of holding an asset, this system becomes

$$\begin{aligned} rS_H^I &= \delta_H - 2\eta S_H^I + \lambda\hat{\mu}^U \beta^I (S_H^U - S_H^I) + \lambda\hat{\mu}^I \frac{(S_H^I - S_H^U)}{2} + \lambda\hat{\mu}^U \beta^I (S_L^U - S_H^I), \\ rS_L^I &= \delta_L - 2\eta S_L^I + \lambda\hat{\mu}^U \beta^I (S_L^U - S_L^I) + \lambda\hat{\mu}^I \frac{(S_H^I - S_L^I)}{2} + \lambda\hat{\mu}^U \beta^I (S_H^U - S_L^I), \\ rS_H^U &= \delta_H - 2\eta S_H^U + \lambda\hat{\mu}^I (1 - \beta^I) (S_L^I - S_H^U) + \lambda \frac{\phi - 2\hat{\mu}^I}{2} (1 - \beta^I) (S_H^I - S_H^U), \end{aligned}$$

$$rS_L^U = \delta_L - 2\eta S_L^U + \lambda \hat{\mu}^I (1 - \beta^I) (S_H^I - S_L^U) + \lambda \frac{\phi - 2\hat{\mu}^I}{2} (1 - \beta^I) (S_L^I - S_L^U).$$

Add up the first two lines and the last two lines to get

$$\begin{aligned} (r + 2\eta)(S_H^I + S_L^I) &= \delta_H + \delta_L + 2\lambda \hat{\mu}^U \beta^I (S_H^U + S_L^U) - 2\lambda \hat{\mu}^U \beta^I (S_H^I + S_L^I), \\ (r + 2\eta)(S_H^U + S_L^U) &= \delta_H + \delta_L + \frac{\lambda\phi}{2} (1 - \beta^I) (S_H^I + S_L^I) - \frac{\lambda\phi}{2} (1 - \beta^I) (S_H^U + S_L^U). \end{aligned}$$

These imply  $(S_H^I + S_L^I) = (S_H^U + S_L^U) = \frac{\delta_H + \delta_L}{r + 2\eta}$ . Now from the original system, subtract the second equation from the first and the fourth from the third to find

$$\begin{aligned} (r + 2\eta)(S_H^I - S_L^I) &= \delta_H - \delta_L - \lambda(2\hat{\mu}^U \beta^I + \hat{\mu}^I)(S_H^I - S_L^I), \\ (r + 2\eta)(S_H^U - S_L^U) &= \delta_H - \delta_L - \frac{\lambda\phi}{2} (1 - \beta^I) (S_H^U - S_L^U) \\ &\quad + \left( \frac{\lambda\phi}{2} - \hat{\mu}^I \right) (1 - \beta^I) (S_H^I - S_L^I). \end{aligned}$$

Rearrange these to get the expressions

$$\begin{aligned} (S_H^I - S_L^I) &= \hat{\alpha}^I (\delta_H - \delta_L), \\ (S_H^U - S_L^U) &= \hat{\alpha}^U (\delta_H - \delta_L), \end{aligned} \tag{S.7}$$

where

$$\begin{aligned} \hat{\alpha}^I &= \frac{1}{2(r + 2\eta + \lambda(2\hat{\mu}^U \beta^I + \hat{\mu}^I))}, \\ \alpha^U = \hat{\alpha}^U &= \left[ \frac{r + 2\eta + \lambda \hat{\mu}^U + \frac{\lambda\phi}{4}}{r + 2\eta + \frac{\lambda\phi}{4}} \right] \hat{\alpha}^I, \end{aligned}$$

which finally implies

$$\begin{aligned} S_H^i &= \left[ \frac{1}{2(r + 2\eta)} \right] (\delta_H + \delta_L) + \frac{\hat{\alpha}^i}{2} (\delta_H - \delta_L), \\ S_L^i &= \left[ \frac{1}{2(r + 2\eta)} \right] (\delta_H + \delta_L) - \frac{\hat{\alpha}^i}{2} (\delta_H - \delta_L). \end{aligned}$$

#### S.4 Flagged values

I solve for  $D_{H1}^I$ ,  $D_{L1}^I$ , and  $D_{H1}^U$ . The system defining  $\{D_{va}^I\}$  is

$$\begin{aligned} rD_{H1}^I &= (\eta + \lambda \hat{\mu}^U)(D_{H0}^I - D_{H1}^I), \\ rD_{L1}^I &= (\eta + \lambda \hat{\mu}^U)(D_{L0}^I - D_{L1}^I) + \lambda \hat{\mu}^I \frac{S_H^I - S_L^I}{2}, \end{aligned}$$

$$\begin{aligned} rD_{L0}^I &= (\eta + \lambda\hat{\mu}^U)(D_{L1}^I - D_{L0}^I) - \lambda\hat{\mu}^U(\beta^I S_L^U + (1 - \beta^I)S_L^I), \\ rD_{H0}^I &= (\eta + \lambda\hat{\mu}^U)(D_{H1}^I - D_{H0}^I) + \lambda\hat{\mu}^I \frac{S_H^I - S_L^I}{2} - \lambda\hat{\mu}^U(\beta^I S_H^I + (1 - \beta^I)S_L^U). \end{aligned}$$

Combining the first equality with the fourth and the second equality with the third yields

$$\begin{aligned} r(D_{H1}^I - D_{H0}^I) &= -2(\eta + \lambda\hat{\mu}^U)(D_{H1}^I - D_{H0}^I) - \lambda\hat{\mu}^I \frac{S_H^I - S_L^I}{2} + \lambda\hat{\mu}^U(\beta^I S_H^I + (1 - \beta^I)S_L^U), \\ r(D_{L1}^I - D_{L0}^I) &= -2(\eta + \lambda\hat{\mu}^U)(D_{L1}^I - D_{L0}^I) + \lambda\hat{\mu}^I \frac{S_H^I - S_L^I}{2} + \lambda\hat{\mu}^U(\beta^I S_L^U + (1 - \beta^I)S_L^I). \end{aligned}$$

Solve to find

$$\begin{aligned} (D_{H1}^I - D_{H0}^I) &= -\frac{\lambda\hat{\mu}^I}{(r + 2\eta + 2\lambda\hat{\mu}^U)} \frac{S_H^I - S_L^I}{2} + \frac{\lambda\hat{\mu}^U}{(r + 2\eta + 2\lambda\hat{\mu}^U)}(\beta^I S_H^I + (1 - \beta^I)S_L^U), \\ (D_{L1}^I - D_{L0}^I) &= \frac{\lambda\hat{\mu}^I}{(r + 2\eta + 2\lambda\hat{\mu}^U)} \frac{S_H^I - S_L^I}{2} + \frac{\lambda\hat{\mu}^U}{(r + 2\eta + 2\lambda\hat{\mu}^U)}(\beta^I S_L^U + (1 - \beta^I)S_L^I). \end{aligned}$$

Plug this back into the original system to get

$$D_{H1}^I = \frac{\eta + \lambda\hat{\mu}^U}{r(r + 2\eta + 2\lambda\hat{\mu}^U)} \left[ \lambda\hat{\mu}^I \frac{S_H^I - S_L^I}{2} - \lambda\hat{\mu}^U(\beta^I S_H^I + (1 - \beta^I)S_L^U) \right], \quad (\text{S.8})$$

$$\begin{aligned} D_{L1}^I &= \frac{r + \eta + \lambda\hat{\mu}^U}{r(r + 2\eta + 2\lambda\hat{\mu}^U)} \lambda\hat{\mu}^I \frac{S_H^I - S_L^I}{2} \\ &\quad - \frac{\lambda\hat{\mu}^U(\eta + \lambda\hat{\mu}^U)}{r(r + 2\eta + 2\lambda\hat{\mu}^U)}(\beta^I S_L^U + (1 - \beta^I)S_L^I). \end{aligned} \quad (\text{S.9})$$

The system defining  $\{D_{\text{va}}^U\}$  is

$$\begin{aligned} rD_{H1}^U &= \eta(D_{H0}^U - D_{H1}^U), \\ rD_{H0}^U &= \eta(D_{H1}^U - D_{H0}^U) + \lambda\hat{\mu}^I(1 - \beta^I)(S_H^U - S_L^I) + \lambda(\phi/2 - \hat{\mu}^I)(1 - \beta^I)(S_H^U - S_H^I), \\ rD_{L1}^U &= \eta(D_{L0}^U - D_{L1}^U) + \lambda\hat{\mu}^I(1 - \beta^I)(S_H^U - S_L^I) + \lambda(\phi/2 - \hat{\mu}^I)(1 - \beta^I)(S_L^U - S_L^I), \\ rD_{L0}^U &= \eta(D_{L1}^U - D_{L0}^U). \end{aligned}$$

Subtract the second equality from the first to get

$$\begin{aligned} r(D_{H1}^U - D_{H0}^U) &= -2\eta(D_{H1}^U - D_{H0}^U) - \lambda\hat{\mu}^I(1 - \beta^I)(S_H^U - S_L^I) \\ &\quad - \lambda(\phi/2 - \hat{\mu}^I)(1 - \beta^I)(S_H^U - S_H^I) \\ \Rightarrow (D_{H1}^U - D_{H0}^U) &= -\frac{\lambda\hat{\mu}^I(1 - \beta^I)}{r + 2\eta}(S_H^U - S_L^I) - \frac{\lambda(\phi/2 - \hat{\mu}^I)(1 - \beta^I)}{r + 2\eta}(S_H^U - S_H^I). \end{aligned}$$

Finally plug back into the original system to get

$$D_{H1}^U = \frac{\eta\lambda\hat{\mu}^I(1-\beta^I)}{r(r+2\eta)}(S_H^U - S_L^I) + \frac{\eta\lambda(\phi/2 - \hat{\mu}^I)(1-\beta^I)}{r(r+2\eta)}(S_H^U - S_H^I). \quad (\text{S.10})$$

Co-editor Florian Scheuer handled this manuscript.

Manuscript received 11 December, 2019; final version accepted 8 March, 2021; available online 17 March, 2021.