

Baker, Julien S. et al.

## Article

# The preventive role of exercise on the physiological, psychological, and psychophysiological parameters of Coronavirus 2 (SARS-CoV-2): A mini review

Journal of Risk and Financial Management

## Provided in Cooperation with:

MDPI – Multidisciplinary Digital Publishing Institute, Basel

*Suggested Citation:* Baker, Julien S. et al. (2021) : The preventive role of exercise on the physiological, psychological, and psychophysiological parameters of Coronavirus 2 (SARS-CoV-2): A mini review, Journal of Risk and Financial Management, ISSN 1911-8074, MDPI, Basel, Vol. 14, Iss. 10, pp. 1-12, <https://doi.org/10.3390/jrfm14100476>

This Version is available at:

<https://hdl.handle.net/10419/258580>

### Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

### Terms of use:

*Documents in EconStor may be saved and copied for your personal and scholarly purposes.*

*You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.*

*If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.*



<https://creativecommons.org/licenses/by/4.0/>



Review

# The Preventive Role of Exercise on the Physiological, Psychological, and Psychophysiological Parameters of Coronavirus 2 (SARS-CoV-2): A Mini Review

Julien S. Baker <sup>1</sup>, Alistair Cole <sup>2</sup>, Dan Tao <sup>1</sup>, Feifei Li <sup>1</sup>, Wei Liang <sup>1</sup>, Jojo Jiao <sup>1</sup>, Yang Gao <sup>1</sup>  
and Rashmi Supriya <sup>1,\*</sup>

<sup>1</sup> Centre for Health and Exercise Science Research, Department of Sport, Physical Education and Health, Hong Kong Baptist University, Kowloon Tong 999077, Hong Kong; jsbaker@hkbu.edu.hk (J.S.B.); emma0514@hkbu.edu.hk (D.T.); lifeifei@hkbu.edu.hk (F.L.); wliang1020@hkbu.edu.hk (W.L.); jojojiao@hkbu.edu.hk (J.J.); gaoyang@hkbu.edu.hk (Y.G.)

<sup>2</sup> Department of Government and International Studies, Hong Kong Baptist University, Kowloon Tong 999077, Hong Kong; alistaircole@hkbu.edu.hk

\* Correspondence: rashmisupriya@hkbu.edu.hk

**Abstract:** The world has been severely challenged by the Coronavirus Disease (COVID-19) outbreak since the early 2020s. Worldwide, there have been more than 66 million cases of infection and over 3,880,450 deaths caused by this highly contagious disease. All sections of the population including those who are affected, those who are not affected and those who have recovered from this disease, are suffering physiologically, psychologically or psychophysiological. In this paper we briefly discuss the consequences of COVID-19 on physiological, psychological and psychophysiological vulnerability. We also attempt to provide evidence in support of exercise management as a prevention strategy for improving and minimizing the physiological, psychological and psychophysiological effects of COVID-19. Moderate exercise including walking, yoga and tai-chi to name but a few exercise regimes are critical in preventing COVID-19 and its complications. Governments, public health authorities and the general population should maintain physical activity during the COVID-19 pandemic to prevent additional physical and mental distress.

**Keywords:** COVID-19; coronavirus disease; physiological effects; psychological effects; psychophysiological effects; exercise; physical activity



**Citation:** Baker, Julien S., Alistair Cole, Dan Tao, Feifei Li, Wei Liang, Jojo Jiao, Yang Gao, and Rashmi Supriya. 2021. The Preventive Role of Exercise on the Physiological, Psychological, and Psychophysiological Parameters of Coronavirus 2 (SARS-CoV-2): A Mini Review. *Journal of Risk and Financial Management* 14: 476. <https://doi.org/10.3390/jrfm14100476>

Academic Editor: Thanasis Stengos

Received: 18 September 2021

Accepted: 5 October 2021

Published: 9 October 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.




**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

There has been a global pandemic of COVID-19 since 2019. The WHO reported 3,880,450 deaths associated with COVID-19 as of 23 June 2021 ([World Health Organization 2021](#)). COVID-19 and associated public health measures may negatively affect individuals with physical, behavioral health (mental illness, substance abuse), financial insecurity and socioeconomic vulnerability. Recent studies suggest that the measures taken to curb the spread of the COVID-19 outbreak have generated issues throughout the population ([Clemente-Suárez et al. 2020](#); [Adhikari et al. 2020](#)). It has been established and identified that the possible risk factors are related to physiological, psychological and psychophysiological vulnerability. Research suggests that impaired immunity may be the root cause of physiological, psychological and psychophysiological vulnerability. For instance, physiological complications due to the pandemic have been reported to be severe in immune-compromised people including the obese, overweight, elderly or subjects with metabolic syndrome or underlying pathologies. Similarly, anxiety and depression, the two major psychological complications reported during the COVID-19 pandemic are influenced by inflammatory pathways and neurotransmitters that are activated by an imbalance of the immune system. Furthermore, psychophysiological vulnerability during COVID-19 including a wide variety of health disorders, such as headaches, essential hypertension,

insomnia, asthma, gastrointestinal and dermatological conditions have been linked with modulation of cortisol levels and increases in pro-inflammatory cytokines. The purpose of this paper is to briefly discuss the consequences of COVID-19 on physiological, psychological and psychophysiological vulnerability; and to briefly outline the mechanisms of how exercise can help in modulating these factors. We have also attempted to provide evidence in support of exercise management as a prevention strategy for improving and minimizing the physiological, psychological and psychophysiological effects of COVID-19 (Figure 1).



**Figure 1.** Summary figure showing the preventive role of exercise on physiological, psychological and psychophysiological effects of COVID-19.

## 2. Search Strategy

This paper was prepared from the presentation on idea presented in the symposium “Transnational and Transdisciplinary Lessons of COVID-19 From the Perspective of Risk and Management”. A structured search strategy was conducted in PubMed and Google scholar to search for publications in English using the search term “COVID-19 OR SARS-CoV-2 OR CORONA” in combination with one of the following keywords: “physiological effects”, “psychological effects”, “psychophysiological effects”, “exercise”, “Yoga” and “tai chi”. We focused on clinical trials, meta-analyses and review articles. We did not include research related with athletes and psychosocial factors. The search was completed on 21 June 2021. When many similar articles were available, the most recent articles were used. Additional papers were identified from random searches and the reference lists of retrieved articles. No date restriction was put on the search so that a longitudinal ‘map’ of the subject area could be obtained

### 3. Physiological Effects of COVID-19 and Its Management by Exercise

**Physiological Effects of COVID-19:** The pandemic has caused physiological complications mostly in elderly populations including (age  $\geq 60$  years) the obese and overweight people with body mass indexes (BMI) over  $25 \text{ kg/m}^2$  or even higher, these individuals contribute to increased risk of infection scores for COVID-19. According to our previous review we have demonstrated that more than 50% of the population with obesity and/or overweight were admitted for critical care and had high mortality (Wang et al. 2020). The condition of obesity refers to excess body fat, low-grade chronic inflammation and impaired immunity that is associated with many debilitating and life-threatening diseases. The list includes respiratory dysfunction, cardiovascular disease, diabetes, some cancers, metabolic risk and co-morbidities, some of which have been linked to more severe COVID-19 infections (Stefan et al. 2020). Additionally, inactivity has now been identified as one of the biggest COVID-19 risk factors. Researchers studied 48,440 adults (a cross-section of the racially diverse Southern California population with an average age of 47 years) between 1 January and 21 October 2020 who were tested positive for COVID-19. The data showed that 6.4 percent of participants were consistently active while 14.4 percent were consistently inactive. Surprisingly, 79.2 percent of respondents were inconsistent exercisers. Study results found that a person who is consistently inactive has higher risk levels for COVID-19 complications. Their risk is comparable to that of people over the age of 60 and those who have undergone organ transplantation. Hospitalization rates for inactive people were twice as high as those for consistently active people. Additionally, these individuals have a 1.73 times greater chance of needing care in the intensive care unit (ICU) and a 2.49 times greater chance of dying from COVID-19 (Narici et al. 2021). Narrowing down to the molecular level, Angiotensin-converting enzyme 2 (ACE2) has been identified as a receptor for COVID-19 entry. ACE2 appears to be a receptor for COVID-19, and obesity may increase ACE2 expression in lung epithelial cells. Clearly, the more adipocytes present, the more ACE2 receptors there are to spread the virus. Additionally, COVID-19 exhibits hyperactive inflammatory responses. The immune system produces pro-inflammatory cytokines Interleukins (IL-6, IL-1 $\beta$ , IL-2, IL-15, IL-10, IL-13) and tumor necrosis factor-alpha (TNF-alpha) in response to an infection. When a virus attacks the immune system, it damages the parenchyma of the lungs and the bronchi and triggers respiratory discomfort resulting in adverse respiratory disorders. Further, it becomes serious enough to require medication (Jose and Manuel 2020; Stebbing et al. 2020; Gil-Etayo et al. 2021). Research also suggests that increased levels of IL-15 and a high T-helper 2 response are associated with a fatal outcome related to COVID-19 (Gil-Etayo et al. 2021). Research has established a significant reduction of %T helper 1 cells and %T helper 17 cells with higher activated %T helper 2 cells in COVID-19 patients compared with controls. Senescent T helper 2 cell percentage was observed to be an independent risk factor for death (Gil-Etayo et al. 2021).

**Exercise Benefits related to the Physiological Effects of COVID-19:** As per our previous review we established that there are multiple advantages of exercise in relation to human health (Wang et al. 2020). These include decreased adipose tissue, improved cardio-respiratory fitness (CRF) and enhancement of metabolic homeostasis and even suppression of inflammation. The most effective way to lose weight is to increase daily energy expenditure through exercise or physical activity. Exercise, in particular, accelerates the breakdown of glycogen in the muscles and the liver; this results in the breakdown of fat in adipose tissue and muscle; and facilitates the oxidation of fatty acids in the muscles (Poirier and Després 2001). A higher CRF corresponds to a lower accumulation of visceral adipose tissue (overweight and obesity) (Brock et al. 2011). In addition, exercise improves oxygen delivery and uptake in exercising muscles. In studies involving subjects with a low baseline CRF, exercise interventions using different modalities (aerobic and resistance) at moderate intensities significantly improved CRF in overweight, obese middle-aged women and men (Church et al. 2010). In contrast, during physical inactivity, metabolic homeostasis is disrupted, leading to insulin resistance, reduced lipid clearance post-prandially, muscle decline and an increase in visceral adiposity (Pedersen and Febbraio 2012). Moreover,

chronic systemic inflammation resulting from metabolic disruption has been linked to persistent physical inactivity (Nunn et al. 2010). Hence, increased exercise would help prevent metabolic derangements. It has been found that moderate-intensity exercise of at least 150 min per week can lower the prevalence of metabolic syndrome (Donnelly et al. 2009). Additionally, it has been observed in previous studies that inflammation and exercise have a negative relationship. Infection leads to the overproduction of pro-inflammatory cytokines such as interleukins (IL-6, IL-2, IL15 etc.) and TNF-alpha (Ostrowski et al. 1999). On the other hand, individuals who perform more frequent and intense physical activity have lower levels of inflammatory biomarkers (IL-6, IL-1 $\beta$ , and TNF-alpha). Through its unique cytokine and hormonal effects, exercise intensity may affect T helper 1/T helper 2 cell balance. Exercises such as walking, tai chi and restorative yoga stimulate the T helper 1 response (Shimizu et al. 2008; Nieman et al. 2005; Raghavendra et al. 2009; Esch et al. 2007), while more intense workouts and longer durations stimulate T helper 2. Cortisol is lowered by low-intensity exercise and relaxing activities, while it is raised by high-intensity exercise and long-duration workouts. There has been evidence that cortisol increases the development of T helper 2 cells and decreases that of T helper 1. Exercise may also push the body toward a more T helper 2 dominated state (as muscle-derived IL-6 behaves differently from TNF-alpha-associated IL-6) and associated IL-10. Muscle-derived IL-6 and associated IL-10 are anti-inflammatory (Narici et al. 2021) and enhances the body's immune response. However, low- to medium-intensity high-volume resistive exercise, that is easily implemented at home, will have positive effects, particularly if combined with a 15–25% reduction in daily energy intake. The combination of these regimens seems ideal for preserving neuromuscular, metabolic, and cardiovascular health (Narici et al. 2021). Furthermore, individuals have been advised to perform 150 min of moderate intensity exercise per week initially to obtain the necessary amount of exercise. This exercise intensity seems favorable in modifying the effects of COVID-19. A larger amount of exercise should be followed for significant weight loss and to prevent weight gain. Generally, 11,000 steps per day with 64–170 steps/minute and at least 10 min duration is sufficient for healthy adults (Wang et al. 2020).

#### 4. Psychological Effects of COVID-19 and Its Management by Exercise

**Psychological Effects of COVID-19:** A state of psychological distress is a state of emotional suffering caused by stressors and demands that people have difficulty in coping with. It is usually characterized by feelings of depression and anxiety (Arvidsdotter et al. 2016). Globally, adolescents of varying backgrounds experience higher levels of anxiety, depression, and stress due to the pandemic and have become frequent consumers of alcohol and recreational drugs (Jones et al. 2021). Based on US Census Bureau, around 56.2% of adults (age = 18–24 years), 48.9% of adults (age = 25–49 years), 39.1 % adults (age = 50–66 years) and 29.3% of adults (age = 65+ years) have been reported to have anxiety and/or depressive symptoms during the pandemic (Anon 2020). The most common psychological effects of COVID-19 on the general population are depression and anxiety. Among the general population, changing social dynamics, such as social distancing, restricted movement and lockdown, have led to severe behavioral and psychological changes. Due to the imposition of lockdown, people work from home, attend online classes and are restricted from visiting established social networks, such as religious meetings, which result in people suffering from stress problems and non-regulation of emotions, that help individuals develop resilience. In addition to social changes, people also experience severe mental and emotional changes due to their fear of getting the disease and uncertainty about their futures. This results from having been confined, abnormally reduced social/physical contact with others, and losing familiar habits are all directly related to feeling distressed, bored, socially isolated and frustrated (Reynolds et al. 2008; Hawryluck et al. 2004). Frustration and loneliness can be attributed to inhibitions from daily activities, interruption of social necessities, and not participating in social network activities (Jeong et al. 2016). The prevalence of loneliness may be associated with depression and suicidal behavior

(Serafini et al. 2020). An evaluation of depression and anxiety in Hong Kong during the COVID-19 pandemic revealed that 19% of the 500 respondents had depression and 14% had anxiety. Additionally, 25.4% reported that their mental health had deteriorated since the pandemic (Choi et al. 2020). Even people who are infected with the disease mainly suffer from anxiety and depression. A study conducted in Hubei on COVID-19 patients (n = 460) revealed that 304 people were diagnosed with somatization symptoms (66.09%), followed by depression (53.48%), anxiety (46.30%), insomnia (42.01%) and self-mutilating or suicidal thoughts (23.26%) (Wang et al. 2021). Existing studies have shown that those who have been exposed to the risk of infection may develop pervasive fears about their health, worries about infecting others, and fear of infecting family members. Cognitive, emotional, and behavioral disorders are reported to be associated with gender-based differences during the COVID-19 pandemic in many studies. Mostly women showed significant higher anxiety, depression, precautionary behavior and emotional responses and acute stress levels than men (Levkovich and Shinan-Altman 2021; García-Fernández et al. 2021; Hou et al. 2020). Focusing on the molecular level, research suggests that cortisol levels were significantly higher in patients with COVID-19 than in individuals who had not contracted the disease. They suggest that COVID-19 patients with highly elevated cortisol levels are more likely to quickly deteriorate in relation to health status. Patients with a cortisol baseline level of 744 or less survived on average for 36 days. Survival was only 15 days for patients with levels over 744 (Tan et al. 2020).

*Exercise Benefits on the Psychological Effects of COVID-19:* Numerous studies have shown the positive effects of exercise on anxiety and depression (Ströhle 2009; Stonerock et al. 2015; Stubbs et al. 2017), and exercise is widely accepted as a non-invasive means of treating or preventing those conditions (Rosa Rimes et al. 2015; Mikkelsen et al. 2017). Relaxation techniques and breathing exercises have been recommended as an intervention for COVID-19 patients to reduce acute anxiety, although more evidence is needed (Khawam et al. 2020). It was found that aerobic exercise was effective in treating anxiety and depression in a study of exercise treatment for clinical anxiety. Exercise programs with a high intensity gain more benefits than programs with a low intensity (Aylett et al. 2018). Some of the most helpful exercises in reducing anxiety include running, swimming, cycling, walking, dancing and even gardening (Hu et al. 2020). Exercise releases neurotransmitters in the body called endorphins, dopamine, and serotonin. They are natural chemicals produced in the brain that helps our body cope with stress. This boosts mood by releasing serotonin, improves appetite and sleep cycles (Aylett et al. 2018). The dysfunction of hypothalamic–pituitary–adrenal (HPA) axis, the increased secretion of corticotropin-releasing hormone (CRH), the impaired responsiveness to glucocorticoids, the increased size and activity of the pituitary were all observed in patients with depression. The ability of exercise on regulating HPA supported physical exercise as one of the methods to improve depressive symptoms (Stranahan et al. 2008). In contrast a study suggests that moderate-to-vigorous physical activities were negatively related to the symptoms of depression. The study proposed that increased light activity and reduced sedentary behavior might contribute to the decreased prevalence of depression (Kandola et al. 2020). At the molecular level, decreased brain-derived neurotrophic factor (BDNF) level is a vulnerability factor for anxiety (Janke et al. 2015). Numerous studies have found that physical exercise can increase the expression of BDNF in the dentate gyrus (area of the brain where all sensory parts come together to create unique representations and memories and is critical for memory and learning) (Adlard and Cotman 2004). Physical exercise was found to be able to restore BDNF to pre-stress levels, suggesting that exercise protects against stress-induced decreased levels of BDNF (Adlard and Cotman 2004). Exercise can also control inflammation, which is a possible mechanism to help reduce anxiety. As reported in a previous study, elevated levels of the pro-inflammatory cytokine C-reactive protein was associated with anxiety disorders (Vogelzangs et al. 2013). Intriguingly, exercise conferred its beneficial effect on anxiety by regulating inflammatory systems (Flynn et al. 2007). Yoga, an ancient eastern practice consisting of breath control, physical postures and meditation,

has demonstrated beneficial effects in patients with severe anxiety symptoms, although the effect was relatively mild (Saeed et al. 2019). A study examined the effect of Tai Chi, a traditional Chinese martial art, on anxiety in older adults and found that those receiving medical therapy could benefit from Tai Chi exercise, but those only receiving medical therapy in isolation, could not (Song et al. 2014). As far as mental health is concerned, aerobic exercise interventions have proven to successfully reduce depression in patients with low to moderate symptoms (Morres et al. 2019; Schuch et al. 2016). The alleviating effects of aerobic exercise interventions on self-reported depressive symptoms in clinical depression have been shown to be comparable to those of antidepressant treatments (Dinas et al. 2011; Blumenthal et al. 2007). Therefore, during the current pandemic, aerobic exercise can help to protect from an increase in depressive symptomatology in people with a diagnosis of clinical depression who already suffer from mild to moderate depressive symptoms. Exercise can also help in preventing depressive symptoms in people at risk for depression. Moreover, aerobic exercise interventions have been found to be effective in anxiety disorders (Generalized Anxiety Disorder, Panic Disorder, Obsessive-Compulsive Disorder, Social Phobia) (Aylett et al. 2018), and stress-related disorders such as post-traumatic stress disorder (Franklin and McCullough 2009), i.e., disorders that are expected to increase during the current pandemic. This psychological issue may not only occur in COVID survivors or health workers, but also the general population (Bohlken et al. 2020; Petzold et al. 2020). As far as clinical anxiety symptoms are concerned, aerobic exercise interventions do not achieve the same effects as psychopharmaceutical treatment in clinical anxiety patients (Carek et al. 2011), but it has proven itself as an adjunctive therapeutic treatment for anxiety disorders in several randomized controlled trial studies (Jayakody et al. 2014). In addition, as a means of primary prevention with respect to the current pandemic, already an acute bout of aerobic exercise of moderate intensity (>21 min) can successfully reduce anxiety (Rebar et al. 2015). Additionally, there is a mood-regulatory effect possible after just seven minutes of aerobic exercise (low to moderate intensity).

## 5. Psychophysiological Effects of COVID-19 and Its Management by Exercise

*Psychophysiological Effects of COVID-19:* Psychophysiology studies investigate the physiological effects of the psychological processes and behavior, as well as the impact of behavioral or psychological manipulations on the body. Psychophysiological measures include techniques designed to assess activity in a variety of bodily systems. As a result of psychological and psychosocial stressors/triggers, a person may experience depression, anxiety, panic attacks, somatic symptoms or posttraumatic stress disorder symptoms, delirium, psychosis or even suicidality. The term psychophysiology has been used to refer to a wide variety of disorders, such as headaches, essential hypertension, insomnia, asthma and gastrointestinal and dermatological conditions (Clemente-Suárez et al. 2020). Constrictions and restrictions mean a sudden end to citizens' normal lives. As a result, the individual begins to display symptoms of anxiety caused by social isolation, lack of mental hygiene habits and repeated exposure to negative news and information. Feelings of sadness, apathy, fear, uncertainty, frustration, irregular circadian cycles, insomnia, hyper vigilance and difficulty concentrating may accompany this illness. COVID symptomatology can trigger feelings of shame, social stigma and even guilt. The subject can mistake some symptoms, such as headaches, migraines or irritation in the throat, for COVID symptoms despite not being infected. The social isolation measures imposed by governments around the world led to a sharp decline in physical activity as recreation facilities, athletic centers, gyms, public parks, playgrounds and schools were forced to close. Furthermore, the gut microbiome may play a crucial role, since it is known that gut bacteria are responsible for mental disorders such as depression and anxiety (strongly present during the pandemic) (Anderson and Maes 2020; Keely et al. 2012). The gut microbiome may modulate the pathophysiology of COVID-19 by altering the immune system and triggering reactions (Anderson and Maes 2020). Additionally, many of the conditions that appear to increase the risk of COVID-19 fatalities are associated with altered gut permeability and dysbio-

sis, such as obesity, diabetes, cardiovascular disorders and lung and respiratory airway inflammation (Anderson and Maes 2020). Stress makes the population more vulnerable to viral infection by affecting the second brain, the gut microbiome (Narici et al. 2021). COVID-19 is predominantly a pulmonary condition, highlighting the importance of the gut, since previous studies have identified a gut-lung axis which suggests that respiratory infections are associated with changes in gut microbiota (Keely et al. 2012). In this way, microbial metabolites and certain endotoxins may have an impact on the lung through the bloodstream, and inflammation may have a reciprocal impact on the gut microbiota (Dumas et al. 2018). The increase in pro-inflammatory cytokines directly leads to a down regulation of hormones that maintain circadian rhythms (Markus et al. 2018). Narrowing to the molecular level there have been recent studies suggesting how psychological stress may also lead to higher levels of corticotropin-releasing hormone (CRH) in the brain, thereby contributing to cortisol production (Vanuytsel et al. 2014). CRH influences inflammatory cytokines and the production of TNF alpha, so its role in the emergence of COVID-19, and therefore its severity and fatalities, is likely to be linked to CRH. This raises the intriguing possibility that COVID-19 could also influence the gut microbiota.

**Exercise Benefits on the Psychophysiological Effects of COVID-19:** Exercise is not only likely to reduce stress, it also improves sleep as well, and it is a well-established fact that exercise has many benefits for overall health. General and mental health benefits of physical activity and regular aerobic exercise with respect to the functioning of the central nervous system, the peripheral autonomous nervous system, the immune system, mental health, stress and well-being have already been explored (Rosa Rimes et al. 2015). Regular aerobic exercise at a moderate level reduces respiratory tract infections because it stimulates the immune system (hormonal effects and cellular effects) (Krüger et al. 2016). Active lifestyles and regular aerobic exercise can save lives by preventing mental and physical health problems. Regarding the current SARS-CoV-2-Coronavirus pandemic, it is important to note that particularly aerobic exercise focused on endurance and cardiorespiratory fitness has positive effects on physical and mental health, particularly sports that involve large amounts of muscle. Besides stimulating the immune system (neuroimmunology and endocrine axis), it also improves cardiorespiratory functions, relieves anxiety and depression, and functions as a buffer against chronic stress by facilitating bodily recovery and reducing perceived stress symptoms. Mental and physical health are interdependent, as are physical activity's effects on both. The benefits of aerobic exercise have been observed in longitudinal and quasi-experimental studies on clinical patient groups (with neurological, somatic and mental disorders) and in healthy subjects without a history of chronic disease (Blair 1996; Franklin and McCullough 2009; Gerber and Pühse 2009; Haskell et al. 2007). During a pandemic lockdown, it is imperative for people with or without a history of lifestyle-related diseases or mental disorders to exercise regularly and stay physically active in order to avoid long-term harm to their physical and mental health. Fitness benefits are typically curvilinear and occur even with relatively low volumes of exercise. Furthermore, exercise positively impacts cognitive, daily, psychosocial and neuroendocrine functions. For instance, Tai-chi is beneficial for treating cognitive problems. Researchers have observed increased and reduced activity of the HPA axis in schizophrenia patients, though studies have shown both hyperfunction and hypofunction of the HPA axis which controls cortisol levels. They found that reducing salivary cortisol levels (a neuroendocrine indicator of stress and immunity, which can provide insights into the possible physiological mechanisms behind this effect) (Ho et al. 2014). Another study involved examining the physiological and psychological impacts of walking in urban parks throughout winter, spring, and early summer. A total of 12 middle-aged and older adults were involved in walking (11–15 min) in an urban park in Japan. To determine physiological responses, their heart rate and blood pressure were measured. The results show that walking in urban parks leads to physiological, psychological relaxation and varied landscape appreciation (Pratiwi et al. 2020). Exercise guidelines generally recommend that 150 min per week of moderate-intensity aerobic activity, such as brisk walking, and/or 75 min of vigorous-



intensity exercise, such as jogging, cycling, swimming, or aerobics. A combination of resistance training, strength-endurance training and yoga, targeting large muscle groups such as the legs or the entire body, is also highly recommended.

As stress is characterized by high levels of arousal and negative effects, mood regulation is a core function of stress recovery. Therefore, mood-regulating strategy involves redirecting cognitions and actions away from the stressor, the so-called diversion strategy. Music as a diversion strategy and results in psychophysiological recovery from stress. Music has been shown to lower anxiety and improve mood. There is some evidence that different music styles can impact psychophysiological recovery differently. Compared with listening to heavy metal, listening to classical music and music selected by the listener, helps in reducing stress. It decreases anxiety and increases relaxation. Listening to classical music after a stressful event decreased systolic blood pressure (Radstaak et al. 2014). Dance movement therapy (DMT) and physical exercise were examined in a study conducted for older adults with dementia. DMT participants showed significant reductions in depression, loneliness, and negative mood, as well as improving daily functioning and the diurnal cortisol slope. At 1-year follow-up, the effects on daily functioning and the cortisol slope remained unchanged. The study results suggest that DMT can be utilized as a multifaceted intervention to improve various aspects of functioning in older adults with declining cognitive abilities (Ho et al. 2018).

The use of artificial intelligence (AI) has increased in COVID-19 research, especially in Detection, classification, severity, and mortality risk assessment. Tracking coronavirus by using AI has already demonstrated its potential to stratify high-risk patients. This technique has also proven to be effective in predicting the infection rate in real time (Rogers 2021; Phillips 2021). Physical education (PE) can also be improved by incorporating AI. A practical understanding of education based on a technical understanding educational and research activities are taking place in all fields in order to consider new ideas pertinent to their respective fields. In case of physical education technology the new ideas are lacking in comparison to other academic fields (Lee and Lee 2021). Additionally, it is crucial to build an integrated research workforce and establish an effective research infrastructure. Similarly, in order to facilitate the comprehensive advancement in exercise, we must develop an environment for modern technology development in the field of physical education, and develop overall AI-related exercise applications.

## 6. Conclusions

Exercise has multiple benefits in combatting the vicious cycle of the physiological, psychological and psychophysiological impacts of COVID-19. Exercise could provide stimulation for the human constitution and could provide potential benefits on resisting the virus. These measures if implemented sooner may reduce underlying pathologies and contribute to the reduction of mortality associated with the COVID-19 pandemic. The health benefits of exercise, as well as its potential to save lives, should be promoted to future generations, governments, and policymakers. Additionally, future focus should be drawn on developing an environment for AI-related exercise applications. In addition, some important considerations should be considered. It is essential to realize the importance of mind relaxation techniques using music, dance and breathing exercises in conjunction with increases in physical activity. In addition, individuals should strictly follow social distancing and avoid mass gatherings in public areas while performing exercises during the COVID-19 pandemic.

**Author Contributions:** J.S.B. and R.S. wrote the manuscript; J.S.B. designed the study; A.C. and D.T. F.L., J.J., W.L. and Y.G. contributed to discussion and editing. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Acknowledgments:** We acknowledge all the presenters and participants at the conference “Translational and transdisciplinary lessons of COVID-19 from the perspective of risk and management”.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

- Adhikari, Sasmita Poudel, Sha Meng, Yu-Ju Wu, Yu-Ping Mao, Rui-Xue Ye, Qing-Zhi Wang, Chang Sun, Rozelle Scott, Hein Raat, Huan Zhou, and et al. 2020. Epidemiology, causes, clinical manifestation and diagnosis, prevention and control of coronavirus disease (COVID-19) during the early outbreak period: A scoping review. *Infectious Diseases of Poverty* 9: 29. [CrossRef]
- Adlard, P. A., and C. W. Cotman. 2004. Voluntary exercise protects against stress-induced decreases in brain-derived neurotrophic factor protein expression. *Neuroscience* 124: 985–92. [CrossRef]
- Anderson, George, and Michael Maes. 2020. Gut Dysbiosis Dysregulates Central and Systemic Homeostasis via Suboptimal Mitochondrial Function: Assessment, Treatment and Classification Implications. *Current Topics in Medicinal Chemistry* 20: 524–39. [CrossRef]
- Anon. 2020. US Census Bureau, Household Pulse Survey. Available online: <https://www.kff.org/coronavirus-covid-19/issue-brief/the-implications-of-covid-19-for-mental-health-and-substance-use/> (accessed on 3 September 2021).
- Arvidsdotter, Tina, Bertil Marklund, Sven Kylén, Charles Taft, and Inger Ekman. 2016. Understanding persons with psychological distress in primary health care. *Scandinavian Journal of Caring Sciences* 30: 687–94. [CrossRef]
- Aylett, Elizabeth, Nicola Small, and Peter Bower. 2018. Exercise in the treatment of clinical anxiety in general practice—a systematic review and meta-analysis. *BMC Health Services Research* 18: 559. [CrossRef] [PubMed]
- Blair, Steven N. 1996. Influences of Cardiorespiratory Fitness and Other Precursors on Cardiovascular Disease and All-Cause Mortality in Men and Women. *JAMA Journal of the American Medical Association* 276: 205. [CrossRef] [PubMed]
- Blumenthal, James A., Michael A. Babyak, P. Murali Doraiswamy, Lana Watkins, Benson M. Hoffman, Krista A. Barbour, Steve Herman, W. E. Craighead, A. L. Brosse, R. Waugh, and et al. 2007. Exercise and Pharmacotherapy in the Treatment of Major Depressive Disorder. *Psychosomatic Medicine* 69: 587–96. [CrossRef] [PubMed]
- Bohlken, Jens, Friederike Schömig, Matthias R. Lemke, Matthias Pumberger, and Steffi G. Riedel-Heller. 2020. COVID-19-Pandemie: Belastungen des medizinischen Personals. *Psychiatrische Praxis* 47: 190–97. [CrossRef]
- Brock, David W., Brian A. Irving, Barbara Gower, and Gary R. Hunter. 2011. Differences emerge in visceral adipose tissue accumulation after selection for innate cardiovascular fitness. *International Journal of Obesity* 35: 309–12. [CrossRef]
- Carek, Peter J., Sarah E. Laibstain, and Stephen M. Carek. 2011. Exercise for the Treatment of Depression and Anxiety. *The International Journal of Psychiatry in Medicine* 41: 15–28. [CrossRef]
- Choi, Edmond Pui Hang, Bryant Pui Hung Hui, and Eric Yuk Fai Wan. 2020. Depression and Anxiety in Hong Kong during COVID-19. *International Journal of Environmental Research and Public Health* 17: 3740. [CrossRef] [PubMed]
- Church, Timothy S., Conrad P. Earnest, Angela M. Thompson, Elisa Priest, Ruben Q. Rodarte, Travis Sanders, Robert Ross, and Steven N. Blair. 2010. Exercise without Weight Loss Does Not Reduce C-Reactive Protein. *Medicine and Science in Sports and Exercise* 42: 708–16. [CrossRef] [PubMed]
- Clemente-Suárez, Vicente Javier, Athanasios A. Dalamitros, Ana Isabel Beltran-Velasco, Juan Mielgo-Ayuso, and Jose Francisco Tornero-Aguilera. 2020. Social and Psychophysiological Consequences of the COVID-19 Pandemic: An Extensive Literature Review. *Frontiers in Psychology* 11: 3077. [CrossRef] [PubMed]
- Dinas, P. C., Y. Koutedakis, and A. D. Flouris. 2011. Effects of exercise and physical activity on depression. *Irish journal of Medical Science* 180: 319–25. [CrossRef] [PubMed]
- Donnelly, Joseph E., Steven N. Blair, John M. Jakicic, Melinda M. Manore, Janet W. Rankin, and Bryan K. Smith. 2009. Appropriate Physical Activity Intervention Strategies for Weight Loss and Prevention of Weight Regain for Adults. *Medicine and Science in Sports and Exercise* 41: 459–71. [CrossRef] [PubMed]
- Dumas, Alexia, Lucie Bernard, Yannick Poquet, Geanncarlo Lugo-Villarino, and Olivier Neyrolles. 2018. The role of the lung microbiota and the gut-lung axis in respiratory infectious diseases. *Cellular Microbiology* 20: e12966. [CrossRef] [PubMed]
- Esch, Tobias, Jorg Duckstein, Justus Welke, and Vittoria Braun. 2007. Mind/body techniques for physiological and psychological stress reduction: Stress management via Tai Chi training—A pilot study. *Medical Science Monitor* 13: CR488–CR497.
- Flynn, Michael G., Brian K. McFarlin, and Melissa M. Markofski. 2007. State of the Art Reviews: The Anti-Inflammatory Actions of Exercise Training. *American Journal of Lifestyle Medicine* 1: 220–35. [CrossRef]
- Franklin, Barry A., and Peter A. McCullough. 2009. Cardiorespiratory Fitness: An Independent and Additive Marker of Risk Stratification and Health Outcomes. *Mayo Clinic Proceedings* 84: 776–79. [CrossRef]
- García-Fernández, Lorena, Verónica Romero-Ferreiro, Sergio Padilla, Pedro David López-Roldán, María Monzó-García, and Roberto Rodríguez-Jimenez. 2021. Gender differences in emotional response to the COVID-19 outbreak in Spain. *Brain Behavior* 11: e01934. [CrossRef]
- Gerber, Markus, and Uwe Pühse. 2009. Review Article: Do exercise and fitness protect against stress-induced health complaints? A review of the literature. *Scandinavian Journal Public Health* 37: 801–19. [CrossRef]

- Gil-Etayo, Francisco Javier, Patricia Suárez-Fernández, Oscar Cabrera-Marante, Daniel Arroyo, Sara Garcinuño, Laura Naranjo, and Daniel E. Pleguezuelo. 2021. T-Helper Cell Subset Response Is a Determining Factor in COVID-19 Progression. *Frontiers in Cellular and Infection Microbiology* 11: 79. [\[CrossRef\]](#)
- Haskell, William L., I-Min Lee, Russell R. Pate, Kenneth E. Powell, Steven N. Blair, Barry A. Franklin, Caroline A. Macera, Gregory W. Heath, Paul D. Thompson, and Adrian Bauman. 2007. Physical Activity and Public Health. *Medicine and Science in Sports and Exercise* 39: 1423–34. [\[CrossRef\]](#) [\[PubMed\]](#)
- Hawryluck, Laura, Wayne L. Gold, Susan Robinson, Stephen Pogorski, Sandro Galea, and Rima Styra. 2004. SARS Control and Psychological Effects of Quarantine, Toronto, Canada. *Emerging Infectious Diseases* 10: 1206–12. [\[CrossRef\]](#)
- Ho, Rainbow Tin Hung, Adrian Ho Yin Wan, Friendly So Wah Au-Yeung, Phyllis Hau Yan Lo, Pantha Joey Chung Yue Siu, Cathy Pui Ki Wong, Winnie Yuen Han Ng, Irene Kit Man Cheung, Siu Man Ng, Cecilia Lai Wan Chan, and et al. 2014. The psychophysiological effects of Tai-chi and exercise in residential Schizophrenic patients: A 3-arm randomized controlled trial. *BMC Complementary and Alternative Medicine* 14: 364. [\[CrossRef\]](#) [\[PubMed\]](#)
- Ho, Rainbow Tin Hung, Ted C. T. Fong, Wai Chi Chan, Joseph S. K. Kwan, Patrick K. C. Chiu, Joshua C. Y. Yau, and Linda C. W. Lam. 2018. Psychophysiological Effects of Dance Movement Therapy and Physical Exercise on Older Adults With Mild Dementia: A Randomized Controlled Trial. *The Journals of Gerontology: Series B*. [\[CrossRef\]](#) [\[PubMed\]](#)
- Hou, Fengsu, Fengying Bi, Rong Jiao, Dan Luo, and Kangxing Song. 2020. Gender differences of depression and anxiety among social media users during the COVID-19 outbreak in China: a cross-sectional study. *BMC Public Health* 20: 1648. [\[CrossRef\]](#) [\[PubMed\]](#)
- Hu, Shaojuan, Lorelei Tucker, Chongyun Wu, and Luodan Yang. 2020. Beneficial Effects of Exercise on Depression and Anxiety During the COVID-19 Pandemic: A Narrative Review. *Frontiers in Psychiatry* 11: 1217. [\[CrossRef\]](#)
- Janke, Kellie L., Tara P. Cominski, Eldo V. Kuzhikandathil, Richard J. Servatius, and Kevin C. H. Pang. 2015. Investigating the Role of Hippocampal BDNF in Anxiety Vulnerability Using Classical Eyeblink Conditioning. *Frontiers in Psychiatry* 6. [\[CrossRef\]](#)
- Jayakody, Kaushadh, Shalmini Gunadasa, and Christian Hosker. 2014. Exercise for anxiety disorders: Systematic review. *British Journal of Sports Medicine* 48: 187–96. [\[CrossRef\]](#)
- Jeong, Hyunsuk, Hyeon Woo Yim, Yeong-Jun Song, Moran Ki, Jung-Ah Min, Juhee Cho, and Jeong-Ho Chae. 2016. Mental health status of people isolated due to Middle East Respiratory Syndrome. *Epidemiology and Health* 38: e2016048. [\[CrossRef\]](#)
- Jones, Elizabeth A. K., Amal K. Mitra, and Azad R. Bhuiyan. 2021. Impact of COVID-19 on Mental Health in Adolescents: A Systematic Review. *International Journal of Environmental Research and Public Health* 18: 2470. [\[CrossRef\]](#)
- Jose, Ricardo J., and Ari Manuel. 2020. COVID-19 cytokine storm: The interplay between inflammation and coagulation. *The Lancet Respiratory Medicine* 8: e46–e47. [\[CrossRef\]](#)
- Kandola, Aaron, Gemma Lewis, David P. J. Osborn, Brendon Stubbs, and Joseph F. Hayes. 2020. Depressive symptoms and objectively measured physical activity and sedentary behaviour throughout adolescence: A prospective cohort study. *The Lancet Psychiatry* 7: 262–71. [\[CrossRef\]](#)
- Keely, Simon, Nicholas J. Talley, and Philip M. Hansbro. 2012. Pulmonary-intestinal cross-talk in mucosal inflammatory disease. *Mucosal Immunology* 5: 7–18. [\[CrossRef\]](#) [\[PubMed\]](#)
- Khawam, Elias, Hassan Khouli, and Leo Pozuelo. 2020. Treating acute anxiety in patients with COVID-19. *Cleveland Clinic Journal of Medicine*. [\[CrossRef\]](#) [\[PubMed\]](#)
- Krüger, Karsten, Frank-Christoph Mooren, and Christian Pilat. 2016. The Immunomodulatory Effects of Physical Activity. *Current Pharmaceutical Design* 22: 3730–48. [\[CrossRef\]](#) [\[PubMed\]](#)
- Lee, Hyun Suk, and Junga Lee. 2021. Applying Artificial Intelligence in Physical Education and Future Perspectives. *Sustainability* 13: 351. [\[CrossRef\]](#)
- Levkovich, Inbar, and Shiri Shinan-Altman. 2021. The impact of gender on emotional reactions, perceived susceptibility and perceived knowledge about COVID-19 among the Israeli public. *International Health*. [\[CrossRef\]](#) [\[PubMed\]](#)
- Markus, Regina P., Pedro A. Fernandes, Gabriela S. Kinker, Sanseray da Silveira Cruz-Machado, and Marina Marçola. 2018. Immune-pineal axis—Acute inflammatory responses coordinate melatonin synthesis by pinealocytes and phagocytes. *British Journal of Pharmacology* 175: 3239–50. [\[CrossRef\]](#)
- Mikkelsen, K., L. Stojanovska, M. Polenakovic, M. Bosevski, and V. Apostolopoulos. 2017. Exercise and mental health. *Maturitas* 106: 48–56. [\[CrossRef\]](#) [\[PubMed\]](#)
- Morres, Ioannis D., Antonis Hatzigeorgiadis, Afroditi Stathi, Nikos Comoutos, Chantal Arpin-Cribbie, Charalampos Krommidas, and Yannis Theodorakis. 2019. Aerobic exercise for adult patients with major depressive disorder in mental health services: A systematic review and meta-analysis. *Depression Anxiety* 36: 39–53. [\[CrossRef\]](#) [\[PubMed\]](#)
- Narici, Marco, Giuseppe De Vito, Martino Franchi, Antonio Paoli, Tatiana Moro, Giuseppe Marcolin, Bruno Grassi, Lucrezia Zuccarelli, Biolo Gianni, Filippo Giorgio Di Girolamo, and et al. 2021. Impact of sedentarism due to the COVID-19 home confinement on neuromuscular, cardiovascular and metabolic health: Physiological and pathophysiological implications and recommendations for physical and nutritional countermeasures. *European Journal of Sport Science* 21: 614–35. [\[CrossRef\]](#)
- Nieman, David C., Dru A. Henson, Melanie D. Austin, and Victor A. Brown. 2005. Immune Response to a 30-Minute Walk. *Medicine and Science in Sports and Exercise* 37: 57–62. [\[CrossRef\]](#) [\[PubMed\]](#)
- Nunn, Alistair V., Geoffrey W. Guy, James S. Brodie, and Jimmy D. Bell. 2010. Inflammatory modulation of exercise salience: Using hormesis to return to a healthy lifestyle. *Nutrition & Metabolism* 7: 87. [\[CrossRef\]](#)

- Ostrowski, Kenneth, Thomas Rohde, Sven Asp, Peter Schjerling, and Bente Klarlund Pedersen. 1999. Pro- and anti-inflammatory cytokine balance in strenuous exercise in humans. *The Journal of physiology* 515: 287–91. [[CrossRef](#)] [[PubMed](#)]
- Pedersen, Bente K., and Mark A. Febbraio. 2012. Muscles, exercise and obesity: Skeletal muscle as a secretory organ. *Nature Reviews Endocrinology* 8: 457–65. [[CrossRef](#)] [[PubMed](#)]
- Petzold, Moritz Bruno, Jens Plag, and Andreas Ströhle. 2020. Umgang mit psychischer Belastung bei Gesundheitsfachkräften im Rahmen der COVID-19-Pandemie. *Nervenarzt* 91: 417–21. [[CrossRef](#)]
- Phillips, Ashley. 2021. Artificial Intelligence-enabled Healthcare Delivery and Digital Epidemiological Surveillance in the Remote Treatment of Patients during the COVID-19 Pandemic. *American Journal of Medical Research* 8: 30. [[CrossRef](#)]
- Poirier, Paul, and Jean-Pierre Després. 2001. Exercise In Weight Management of Obesity. *Cardiology Clinics* 19: 459–70. [[CrossRef](#)]
- Pratiwi, Prita Indah, Qiongying Xiang, and Katsunori Furuya. 2020. Physiological and Psychological Effects of Walking in Urban Parks and Its Imagery in Different Seasons in Middle-Aged and Older Adults: Evidence from Matsudo City, Japan. *Sustainability* 12: 4003. [[CrossRef](#)]
- Radstaak, Mirjam, Sabine A. E. Geurts, Jos F. Brosschot, and Michiel A. J. Kompier. 2014. Music and Psychophysiological Recovery from Stress. *Psychosomatic Medicine* 76: 529–37. [[CrossRef](#)] [[PubMed](#)]
- Raghavendra, Rao M., H. S. Vadiraja, Raghuram Nagarathna, H. R. Nagendra, M. Rekha, N. Vanitha, and K. S. Gopinath. 2009. Effects of a Yoga Program on Cortisol Rhythm and Mood States in Early Breast Cancer Patients Undergoing Adjuvant Radiotherapy: A Randomized Controlled Trial. *Integrative Cancer Therapies* 8: 37–46. [[CrossRef](#)] [[PubMed](#)]
- Rebar, Amanda L., Robert Stanton, David Geard, Camille Short, Mitch J. Duncan, and Corneel Vandelandotte. 2015. A meta-meta-analysis of the effect of physical activity on depression and anxiety in non-clinical adult populations. *Health Psychology Review* 9: 366–78. [[CrossRef](#)] [[PubMed](#)]
- Reynolds, Diane L., J. R. Garay, S. L. Deamond, Maura K. Moran, W. Gold, and R. Styra. 2008. Understanding, compliance and psychological impact of the SARS quarantine experience. *Epidemiology & Infection* 136: 997–1007. [[CrossRef](#)]
- Rogers, Rob. 2021. Internet of Things-based Smart Healthcare Systems, Wireless Connected Devices, and Body Sensor Networks in COVID-19 Remote Patient Monitoring. *American Journal of Medical Research* 8: 71. [[CrossRef](#)]
- Rosa Rimes, Ridson, Marcos de Souza Moura Antonio, Murilo Khede Lamego, Alberto Souza de Sa Filho, Joao Manochio, Flavia Paes, and Mauro Giovanni Carta. 2015. Effects of Exercise on Physical and Mental Health, and Cognitive and Brain Functions in Schizophrenia: Clinical and Experimental Evidence. *CNS Neurological Disorders-Drug Targets (Formerly Current Drug Targets-CNS & Neurological Disorders)* 14: 1244–54. [[CrossRef](#)]
- Saeed, Sy Ateaz, Karlene Cunningham, and Richard M. Bloch. 2019. Depression and Anxiety Disorders: Benefits of Exercise, Yoga, and Meditation. *American Family Physician* 99: 620–27.
- Schuch, Felipe B., Davy Vancampfort, Justin Richards, Simon Rosenbaum, Philip B. Ward, and Brendon Stubbs. 2016. Exercise as a treatment for depression: A meta-analysis adjusting for publication bias. *Journal of Psychiatric Research* 77: 42–51. [[CrossRef](#)]
- Serafini, Gianluca, Bianca Parmigiani, Andrea Amerio, Andrea Aguglia, Leo Sher, and Mario Amore. 2020. The psychological impact of COVID-19 on the mental health in the general population. *QJM An International Journal of Medicine* 113: 531–37. [[CrossRef](#)]
- Shimizu, Kazuhiro, Fuminori Kimura, Takayuki Akimoto, Takao Akama, Kai Tanabe, Takahiko Nishijima, Shinya Kuno, and Ichiro Kono. 2008. Effect of moderate exercise training on T-helper cell subpopulations in elderly people. *Exercise Immunology Review* 14: 24–37.
- Song, Qing-Hua, Guo-Qing Shen, Rong-Mei Xu, Quan-Hai Zhang, Ming Ma, Yan-Hua Guo, Xin-Ping Zhao, and Yu-Bing Han. 2014. Effect of Tai Chi exercise on the physical and mental health of the elder patients suffered from anxiety disorder. *International Journal of Physiology, Pathophysiology and Pharmacology* 6: 55–60. [[PubMed](#)]
- Stebbing, Justin, Anne Phelan, Ivan Griffin, Catherine Tucker, Olly Oechsle, Dan Smith, and Peter Richardson. 2020. COVID-19: Combining antiviral and anti-inflammatory treatments. *The Lancet Infectious Diseases* 20: 400–2. [[CrossRef](#)]
- Stefan, Norbert, Andreas L. Birkenfeld, Matthias B. Schulze, and David S. Ludwig. 2020. Obesity and impaired metabolic health in patients with COVID-19. *Nature Reviews Endocrinology* 16: 341–42. [[CrossRef](#)]
- Stonerock, Gregory L., Benson M. Hoffman, Patrick J. Smith, and James A. Blumenthal. 2015. Exercise as Treatment for Anxiety: Systematic Review and Analysis. *Annals of Behavioral Medicine* 49: 542–56. [[CrossRef](#)]
- Stranahan, Alexis M., Kim Lee, and Mark P. Mattson. 2008. Central Mechanisms of HPA Axis Regulation by Voluntary Exercise. *Neuromolecular Medicine* 10: 118–27. [[CrossRef](#)]
- Ströhle, Andreas. 2009. Physical activity, exercise, depression and anxiety disorders. *Journal of Neural Transmission* 116: 777–84. [[CrossRef](#)]
- Stubbs, Brendon, Davy Vancampfort, Simon Rosenbaum, Joseph Firth, Theodore Cosco, Nicola Veronese, Giovanni A. Salum, and Felipe B. Schuch. 2017. An examination of the anxiolytic effects of exercise for people with anxiety and stress-related disorders: A meta-analysis. *Psychiatry Research* 249: 102–8. [[CrossRef](#)]
- Tan, Tricia, Bernard Khoo, Edouard G. Mills, Maria Phylactou, Bijal Patel, Pei C. Eng, and Layla Thurston. 2020. Association between high serum total cortisol concentrations and mortality from COVID-19. *Lancet Diabetes Endocrinology* 8: 659–60. [[CrossRef](#)]
- Vanuytsel, Tim, Sander Van Wanrooy, Hanne Vanheel, Christophe Vanormelingen, Sofie Verschueren, Els Houben, Shadea Salim Rasool, J. Tóth, L. Holvoet, R. Farré, and et al. 2014. Psychological stress and corticotropin-releasing hormone increase intestinal permeability in humans by a mast cell-dependent mechanism. *Gut* 63: 1293–99. [[CrossRef](#)] [[PubMed](#)]

- Vogelzangs, N., A. T. F. Beekman, P. De Jonge, and B. W. J. H. Penninx. 2013. Anxiety disorders and inflammation in a large adult cohort. *Translational Psychiatry* 3: e249. [[CrossRef](#)] [[PubMed](#)]
- Wang, Meizi, Julien S. Baker, Wenjing Quan, Siqin Shen, Gusztáv Fekete, and Yaodong Gu. 2020. A Preventive Role of Exercise Across the Coronavirus 2 (SARS-CoV-2) Pandemic. *Frontiers in Physiology* 11: 572718. [[CrossRef](#)] [[PubMed](#)]
- Wang, Minghuan, Caihong Hu, Qian Zhao, Renjie Feng, Qing Wang, Hongbin Cai, and Zhenli Guo. 2021. Acute psychological impact on COVID-19 patients in Hubei: A multicenter observational study. *Translational Psychiatry* 11: 133. [[CrossRef](#)] [[PubMed](#)]
- World Health Organisation. 2021. Available online: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019> (accessed on 15 September 2021).