

Peña-Guerrero, Mayra Daniela; Umirbekov, Atabek; Tarasova, Larisa; Müller, Daniel

**Article — Published Version**

## Comparing the performance of high-resolution global precipitation products across topographic and climatic gradients of Central Asia

International Journal of Climatology

**Provided in Cooperation with:**

Leibniz Institute of Agricultural Development in Transition Economies (IAMO), Halle (Saale)

*Suggested Citation:* Peña-Guerrero, Mayra Daniela; Umirbekov, Atabek; Tarasova, Larisa; Müller, Daniel (2022) : Comparing the performance of high-resolution global precipitation products across topographic and climatic gradients of Central Asia, International Journal of Climatology, ISSN 1097-0088, Wiley, Hoboken, Vol. 42, Iss. 11, pp. 5554-5569, <https://doi.org/10.1002/joc.7548>, <https://rmets.onlinelibrary.wiley.com/doi/10.1002/joc.7548>

This Version is available at:

<https://hdl.handle.net/10419/264276>

**Standard-Nutzungsbedingungen:**

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

**Terms of use:**

*Documents in EconStor may be saved and copied for your personal and scholarly purposes.*

*You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.*

*If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.*



<http://creativecommons.org/licenses/by/4.0/>

## RESEARCH ARTICLE

# Comparing the performance of high-resolution global precipitation products across topographic and climatic gradients of Central Asia

Mayra Daniela Peña-Guerrero<sup>1,2,3,4</sup>  | Atabek Umirbekov<sup>1,2,3</sup>  |  
Larisa Tarasova<sup>4</sup>  | Daniel Müller<sup>1,2,5</sup> 

<sup>1</sup>Leibniz Institute of Agricultural Development in Transition Economies (IAMO), Halle (Saale), Germany

<sup>2</sup>Department of Geography, Humboldt-Universität zu Berlin, Berlin, Germany

<sup>3</sup>Tashkent Institute of Irrigation and Agricultural Mechanization Engineers, Tashkent, Uzbekistan

<sup>4</sup>Department of Catchment Hydrology, Helmholtz Centre for Environmental Research, Halle (Saale), Germany

<sup>5</sup>Integrative Research Institute on Transformations of Human-Environment System (IRI THESys), Humboldt-Universität zu Berlin, Berlin, Germany

## Correspondence

Mayra Daniela Peña-Guerrero, Leibniz Institute of Agricultural Development in Transition Economies (IAMO), Halle (Saale), Germany.

Email: [pena-guerrero@iamo.de](mailto:pena-guerrero@iamo.de)

## Funding information

Leibniz-Institut für Agrarentwicklung in Transformationsökonomien, Grant/Award Number: 687; Volkswagen Foundation, Grant/Award Number: 96264

## Abstract

Accurate and reliable precipitation data with high spatial and temporal resolution are essential in studying climate variability, water resources management, and hydrological forecasting. A range of global precipitation data are available to this end, but how well these capture actual precipitation remains unknown, particularly for mountain regions where ground stations are sparse. We examined the performance of three global high-resolution precipitation products for capturing precipitation over Central Asia, a hotspot of climate change, where reliable precipitation data are particularly scarce. Specifically, we evaluated MSWEP, CHIRPS, and GSMAP against independent gauging stations for the period 1985–2015. Our results show that MSWEP and CHIRPS outperformed GSMAP for wetter periods (i.e., winter and spring) and wetter locations (150–600 mm·year<sup>-1</sup>), lowlands, and mid-altitudes (0–3,000 m), and regions dominated by winter and spring precipitation. MSWEP performed best in representing temporal precipitation dynamics and CHIRPS excelled in capturing the volume and distribution of precipitation. All precipitation products poorly estimated precipitation at higher elevations (>3,000 m), in drier areas (<150 mm), and in regions characterized by summer precipitation. All products accurately detected dry spells, but their performance decreased for wet spells with increasing precipitation intensity. In sum, we find that CHIRPS and MSWEP provide the most reliable high-resolution precipitation estimates for Central Asia. However, the high spatial and temporal heterogeneity of the performance call for a careful selection of a suitable product for local applications considering the prevailing precipitation dynamics, climatic, and topographic conditions.

## KEYWORDS

climate change, green water, Kazakhstan, Kyrgyzstan, satellite imagery, Tajikistan, Turkmenistan, Uzbekistan, validation, water resources

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *International Journal of Climatology* published by John Wiley & Sons Ltd on behalf of Royal Meteorological Society.

## 1 | INTRODUCTION

The spatial and temporal variability of precipitation shape hydrological cycles (Michaelides *et al.*, 2009). Climate change alters these cycles through changes in precipitation frequency, intensity, and amount and by affecting evapotranspiration patterns (Trenberth, 2011; Tan *et al.*, 2020). These changes in turn impact freshwater availability for agriculture, hydropower, and socioeconomic development (Tan *et al.*, 2020). Most regions with rainfed or irrigated crops depend on precipitation totals during the peak months of plant growth to meet water demand (Funk *et al.*, 2015). Accurate and reliable precipitation records with high spatial and temporal resolutions are therefore essential to study climate variability, the management of water resources, and hydrological forecasting (Sun *et al.*, 2018).

Satellite precipitation sensors are currently the only instruments that can provide near real-time global coverage of precipitation with estimates from geosynchronous infrared sensors on geostationary satellites, which have a high sampling frequency, and polar-orbiting microwave sensors on low Earth-orbiting satellites with lower temporal resolution (Huffman *et al.*, 2007). Satellite-based estimates of precipitation are increasingly used to complement ground station observations, which are limited in areal coverage and density, particularly in inaccessible regions (e.g., mountainous areas), sparsely populated areas, and especially in developing countries (Zambrano-Bigiarini *et al.*, 2017; Rivera *et al.*, 2018; Sun *et al.*, 2018). The lack of station data in mountainous regions is worrisome because precipitation patterns in the mountains are crucial to assess changes in regional climate and in the cryosphere, which directly affect water availability in downstream regions (Unger-Shayesteh *et al.*, 2013; Immerzeel *et al.*, 2020; Viviroli *et al.*, 2020).

Merging satellite data with gauge measurements from ground stations and reanalysis estimations can improve the accuracy of global precipitation datasets (Tan *et al.*, 2020). Reanalysis-based estimates merge atmospheric measurements and climatic models encompassing physical and dynamical processes to produce consistent, accurate, and continuous meteorological data (Sun *et al.*, 2018; Tan *et al.*, 2020). However, varying availability of ground observations for calibrating the satellite algorithms and reanalysis estimates can compromise the quality of the merged global precipitation datasets (Hu *et al.*, 2018; Sun *et al.*, 2018; Zandler *et al.*, 2019). Therefore, quantitative validation of these global precipitation datasets against independent ground observations is critical to determine the accuracy and uncertainty of the global products at local and regional scales because misestimations can arise from sampling,

instrumental (e.g., sensor observations), and algorithmic errors (Nijssen, 2004; Ebert, 2007; Hu *et al.*, 2018). Validation of global precipitation data is hence a keystone for better understanding the impact of climate changes on regional hydrological cycles.

Station data for Central Asia are very scarce, especially in higher elevations and particularly since the dilapidation of much of the meteorological infrastructure following the collapse of the Soviet Union in 1991 and the independence of the Central Asian republics (Schiemann *et al.*, 2008; Unger-Shayesteh *et al.*, 2013). This is unfortunate because precipitation, glaciers, and snowmelt dominate the hydrological budget in the semiarid continental climate of Central Asia, where water fluxes in the mountainous areas play a crucial role in downstream hydrology and water availability (Schär *et al.*, 2004; Mannig *et al.*, 2013; Maussion *et al.*, 2013). The region's economy and ecology heavily rely on water from the two main endorheic rivers, the Amu Darya and Syr Darya, which originate in the headwater catchments of the Pamir and Tien Shan mountains, respectively (Schär *et al.*, 2004; Unger-Shayesteh *et al.*, 2013). In addition, the region is a hotspot of climate change with warming rates of up to 0.3°C per decade during the past half century (Teixeira *et al.*, 2013; Reyer *et al.*, 2017; Peng *et al.*, 2019).

Precipitation data sourced from global precipitation products are paramount for data-scarce or ungauged regions, such as Central Asia. Previous studies have evaluated precipitation products for the region with varied and sometimes contrasting results (Table S1, Supporting Information). Several studies have suggested that the gauge-based products from the Global Precipitation Climatology Centre (GPCC), with spatial resolutions ranging from 0.25° to 1°, were the most reliable precipitation data for the region but underestimated precipitation in the mountains (Malsy *et al.*, 2014; Hu *et al.*, 2018). In the Amu Darya basin, however, the gauge-based Climate Prediction Center (CPC) (0.5°) dataset performed best (Salehie *et al.*, 2021). In the Pamir mountains, the reanalysis product Modern-Era Retrospective analysis for Research Application (MERRA, 0.5°) stood out, although its performance deteriorated strongly for the period 1998–2012 due to the decline of station data availability (Zandler *et al.*, 2019).


The resolution of the abovementioned gauge-based and reanalysis products is not suitable for studies at regional and catchment scales in Central Asia because their spatial resolution is too coarse to capture precipitation gradients in the complex topography of the region (Hellwig *et al.*, 2018; Henn *et al.*, 2018). Here we evaluate global or near-global precipitation products with a spatial resolution higher than 12 × 12 km and for the period 1981–2015 for the Pamir and Tien Shan mountains and

adjacent lowlands of Central Asia. We only selected gauge-corrected products with proven reliability at local and regional scales with a sufficiently long time series available to support analysing climatic trends and variability (see Table S1 and references therein). Finally, we only consider products that are still operational. These criteria resulted in the selection of the Climate Hazard Group InfraRed Precipitation with Station Data (CHIRPS version 2, 0.05°) (Funk *et al.*, 2015), the Multi-Source Weighted-Ensemble Precipitation (MSWEP, 0.1°) (Beck *et al.*, 2017a), and the Global Satellite Mapping of Precipitation (GSMAP, 0.1°) (Ushio *et al.*, 2009).

The gauge-corrected versions of GSMAP and MSWEP products were previously considered the best-performing high-resolution precipitation data for the region (Guo *et al.*, 2015; Guo *et al.*, 2017; Lu *et al.*, 2021) and globally

(Beck *et al.*, 2017b). The gauge-calibrated CHIRPS product was ranked third in Central Asia when compared to six coarser precipitation products (Salehie *et al.*, 2021). However, the accuracy of MSWEP and CHIRPS has to date not been assessed for Central Asia with data from meteorological stations that were not used for gauge correction of the investigated products.

Here we aim to (a) identify the strengths and limitations of the three global precipitation products at daily, monthly, seasonal, and annual timescales, (b) determine the effect of topography and climate regimes on the performance of the precipitation products, (c) quantify the accuracy of the products for different precipitation intensities, and (d) based on these evaluations, propose which precipitation product is most appropriate for subsequent studies. Our analysis hence facilitates informed decisions



**FIGURE 1** Study area and location of the 30 precipitation stations used for validation. Bar plots represent the annual precipitation regimes (section 4.4) of the clusters shown in the map: cluster 1 (yellow) is characterized by winter and spring precipitation and long, dry summers and autumns; cluster 2 (blue) has winter and spring precipitation and a short, dry summer period; and summer precipitation dominates in cluster 3 (red) [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

for assessing climate variability, hydrological and agricultural studies, and water management in this heterogeneous and data-scarce region.

## 2 | STUDY AREA

The study area covers the Tien Shan and Pamir mountains, including the adjacent semi-arid lowlands of Kazakhstan, Kyrgyzstan, Tajikistan, Turkmenistan, and Uzbekistan (Figure 1). In the western and northern Tien Shan and Pamir mountains, most precipitation occurs during the winter and spring seasons (November–March) and falls primarily as snow (Barlow and Tippett, 2008; Sorg *et al.*, 2012). In contrast, parts of central and eastern Tien Shan and eastern Pamir receive most precipitation during summer months (Apel *et al.*, 2018). Precipitation amounts in the region range from 50 to 1,000 mm annually, primarily determined by orographic uplift and mid-latitude westerly cyclones (Mariotti, 2007; Barlow and Tippett, 2008).

Large-scale variation of extratropical westerlies, which transport moisture from the Atlantic Ocean, the Mediterranean and Caspian Sea, and the Persian Gulf, are the major moisture sources throughout the year in Central Asia; the central and eastern Tien Shan and southeastern Pamir are also affected by the Indian Monsoon in summer (Böhner, 2006; Meier *et al.*, 2013). Moisture fluxes from the Arabian Sea and tropical Africa during warm El Niño–Southern Oscillation events cause higher precipitation in autumn and spring in southwestern Central Asia (Mariotti, 2007).

## 3 | DATA

### 3.1 | Precipitation products

#### 3.1.1 | CHIRPS

CHIRPS provides daily blended gauge-satellite precipitation estimates covering most global land regions (50°N–50°S) with a resolution of 0.05° (about 5 km at the equator) from 1981 until present and with a low latency (updated roughly every 2 days, with a stable product released every 3 weeks) (Funk *et al.*, 2015). CHIRPS combines precipitation estimates based on observations of infrared cold cloud duration in which cold and bright clouds are related to convection and therefore rain (Sun *et al.*, 2018). CHIRPS incorporates station data from public data streams and private archives and uses reanalysis-based estimates of the Coupled Forecast System (CFS) to temporally disaggregate from 5-day to daily estimates

and when thermal infrared observations are missing (Shukla *et al.*, 2014; Funk *et al.*, 2015). Calibration of CHIRPS involves three main components: (a) the Climate Hazards group Precipitation climatology (CHPclim); (b) the satellite-only Climate Hazards group Infrared Precipitation (CHIRP); and (c) the station-blending procedure (Funk *et al.*, 2015). We downloaded the CHIRPS data from <https://data.chc.ucsb.edu/products/CHIRPS-2.0/>.

#### 3.1.2 | GSMAP

The GSMAP is a multisatellite algorithm developed by the Japan Science and Technology Agency (Okamoto *et al.*, 2005; Kubota *et al.*, 2007). The algorithm follows three main steps: (a) retrieval of precipitation rate from passive microwave data (precipitation-sized particles such as ice content are detected through clouds), provided by the CPC using a Kalman filter approach (Ushio *et al.*, 2009; Shige *et al.*, 2013; Yamamoto *et al.*, 2017); (b) propagation of the estimated precipitation rates using a backward- and forward-morphing technique (Joyce *et al.*, 2004); and (c) refinement of precipitation data based on the relationship between the infrared brightness temperature and surface precipitation rates. GSMAP has a spatial resolution of 0.1° (about 11 km at the equator) and near-global coverage (60°N–60°S). It provides hourly averaged rainfall (mm·hr<sup>-1</sup>). We used daily precipitation values (mm·day<sup>-1</sup>) of the GSMAP\_Gauge\_NRT (near real-time with gauge-calibration using the NOAA CPC Unified Gauge-Based Analysis of Global Precipitation dataset, 0.5°) that has the longest record, starting in 2000 up to present day. We downloaded the GSMAP data from <https://sharaku.eorc.jaxa.jp/GSMaP/>.

#### 3.1.3 | MSWEP

The MSWEP precipitation dataset provides 3-hourly and daily temporal resolution at 0.1°–0.25° spatial resolutions from 1979 to near present on a global scale (Beck *et al.*, 2017a; 2017b; Beck *et al.*, 2019). It merges gauge observations, satellite, and reanalysis estimates based on timescale and location (Beck *et al.*, 2019). The weight assigned to the gauge-based estimates is calculated from the gauge network density, and the weights assigned to the satellite- and reanalysis-based estimates are calculated from their comparative performance at surrounding gauges (Sun *et al.*, 2018). We used the latest version of MSWEP with a spatial resolution of 0.1° (about 11 km at the equator). This dataset relies on the reanalysis ERA5, the Multi-Satellite Retrievals from the Global (IMERG)

satellite constellation, and the Gridded Satellite (GridSat) thermal infrared imagery, with GridSat only used prior to 2000. Unlike previous versions of MSWEP, this version does not correct underestimation over mountainous and snow-dominated regions in order to match rain gauge observations as closely as possible (Beck *et al.*, 2021). We downloaded the MSWEP data from <http://www.gloh2o.org/>.

### 3.2 | Precipitation gauge data

We collected data for 30 stations within the five central Asian countries for the period 1981–2015 (Figure 1), which were not used to correct the global precipitation products. The selection of these stations was based on the list of station names and locations used for the CHIRPS product, the main public gauge sources of MSWEP (Funk *et al.*, 2015). We directly requested and collected the station data from the local research and governmental institutions for validation, and they are not available on open public archiving domains that are used for gauge correction of the global datasets. For some stations, the calibration of the global precipitation made only used a share of the available time series (see Table S2); in these cases, we used only the remaining data for the validation. For GSMAP, we used only the data from those 27 stations, which are available for the period April 2000 through December 2015.

## 4 | METHODS

### 4.1 | Evaluation of precipitation products at different timescales

We evaluated all products at daily, monthly, seasonal, and annual timescales to understand their value for applications that require precipitation data of various temporal resolutions (e.g., hydrological forecasting, water resource management, and agricultural drought monitoring) (Tobin and Bennett, 2014; Funk *et al.*, 2015). For the seasonal timescale, we used calendar seasons: December, January, February (DJF); March, April, May (MAM); June, July, August (JJA); and September, October, November (SON). We grouped stations by elevation bands, precipitation amount, and precipitation regime to evaluate the reliability of precipitation products in diverse environmental conditions and to determine the effect of topography and climate regimes on the performance of the products. Since different precipitation intensities challenge the accuracy of the precipitation estimates, we classified daily precipitation time series

into dry spells ( $<1 \text{ mm}\cdot\text{day}^{-1}$ ) and wet spells of various intensities ( $1\text{--}5$ ,  $5\text{--}20$ ,  $20\text{--}40$ , and  $>40 \text{ mm}\cdot\text{day}^{-1}$ ) (World Meteorological Organization, 2008; Zambrano-Bigiarini *et al.*, 2017).

### 4.2 | Evaluation of precipitation products at different spatial scales

We performed a point-to-pixel analysis to compare the time series of precipitation gauge data to the corresponding pixel of each product (Thiemig *et al.*, 2012; Zambrano-Bigiarini *et al.*, 2017; Baez-Villanueva *et al.*, 2018). To ensure a consistent comparison among the products, we upscaled CHIRPS to the coarser spatial resolution of MSWEP and GSMAP (i.e.,  $0.1^\circ$ ) using bilinear interpolation. To determine the effect of the upscaling, we performed the evaluation for both original and upscaled versions (hereafter termed CHIRPS upscaled).

### 4.3 | Evaluation metrics

We evaluated the performance of the products for continuous precipitation time series and for discrete precipitation events. For precipitation time series, we used the modified Kling–Gupta efficiency (KGE') (Gupta *et al.*, 2009; Kling *et al.*, 2012) (Equation S1), a dimensionless metric that measures the ability of the precipitation products to reproduce temporal dynamics (correlation coefficient  $r$ ) while preserving the volume (bias ratio  $\beta$ ) and the distribution of precipitation (variability ratio  $\gamma$ ). KGE',  $r$ ,  $\beta$ , and  $\gamma$  values of 1 indicate a perfect agreement between the precipitation estimates from the product and the ground observations. KGE' values range from  $-\infty$  to 1. To determine the product accuracy, we used the mean absolute error (MAE) (Equation S2), which measures the average magnitude of the difference between the estimated and observed values (Ebert, 2007).

We evaluated the ability of tested precipitation products measuring the correspondence between estimated and observed dry and wet spells of various intensity groups (section 4.1) using a standard contingency table (Ebert, 2007) that summarizes the frequency of correct and false predictions. We used three categorical measures—that is, the probability of detection (POD), the false alarm ratio (FAR), and frequency bias (fBias) (Equation S3)—that quantify various aspects of performance: POD measures the fraction of correctly identified observed events (“hit rate”), FAR gives the fraction of diagnosed events that were dry spells, and fBias

calculates the ratio of the estimated events to the observed precipitation (Ebert, 2007; Guo *et al.*, 2017; Baez-Villanueva *et al.*, 2018). Perfect values are fBias (no bias), POD (detection of all events) is 1, and FAR (no events are incorrectly identified) is 0.

#### 4.4 | Dominant precipitation regimes

To capture the heterogeneity of climatic conditions and precipitation seasonality, we determined the precipitation regimes of each gauge and its corresponding grid locations with a monthly sequence of the Pardé coefficients (Pardé, 1933) (Equation S4), which are dimensionless and can be used for interregional comparisons of precipitation regimes. We used the *k*-means clustering algorithm (Lloyd, 1982), which minimizes the sum of squares of distances between the gauging stations' values and the cluster with the nearest mean. In that way, we grouped the "shapes" of the seasonal precipitation regime according to membership in a cluster of precipitation with a similar shape (Weingartner *et al.*, 2013). We selected the optimal number of clusters (*k*) using the elbow method, a trade-off between the cluster sum of squared errors, and a

larger number of clusters (graphically) (Thorndike, 1953; Zhang *et al.*, 2016).

## 5 | RESULTS

### 5.1 | Performance at different timescales

At the seasonal scale, all products performed worst in summer (Figure 2). The overall performance of GSMAP was lower compared to the other products at all timescales, except summer, and especially in winter ( $KGE' < 0$ ). MSWEP, CHIRPS, and its upscaled version showed the best performance in winter. The second-best seasonal performance was spring for MSWEP and autumn for CHIRPS products. All products showed positive correlation coefficients (*r*) for all timescales (Figure S1). MSWEP best captures the temporal dynamics of precipitation in winter, followed by the two CHIRPS products. Moreover, MSWEP and both versions of CHIRPS performed similarly well in autumn and spring. In terms of bias values ( $\beta$ ), CHIRPS and CHIRPS upscaled showed the best performance at all timescales, except for the summer season, when it slightly




FIGURE 2 KGE' between the precipitation products and precipitation gauge data for six different timescales. The vertical blue line indicates the optimum value for KGE'. From left to right and up to bottom: monthly, winter (December, January, February), spring (March, April, May), summer (June, July, August), and autumn (September, October, November) [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

underestimated precipitation (Figure S2). GSMAP revealed higher overestimation in winter and underestimation in summer, whereas MSWEP overestimates precipitation in autumn and summer. Among all products, only GSMAP overestimated the variability ( $\gamma$ ) of the observed precipitation, especially in winter (Figure S3), whereas the other products underestimated it at all timescales but particularly during the summer season.

CHIRPS and its upscaled version performed best at monthly timescales and performed similar to MSWEP at the annual timescale. MSWEP showed the highest correlations but also the highest overestimation of precipitation at both scales. CHIRPS upscaled, followed by CHIRPS, performed best in terms of bias as well as in capturing the precipitation variability at monthly timescales, whereas MSWEP had a better performance in estimating the precipitation variability at the annual timescale.

Our results reveal distinct variations in MAE for different timescales (Figure 3). Regarding the lowest median MAE, both CHIRPS datasets showed it in autumn, GSMAP demonstrated it in summer, and MSWEP showed it in spring and winter. MSWEP in summer and GSMAP in spring, winter, and autumn exhibited the largest errors. CHIRPS products presented the lowest MAE at annual and monthly timescales.

## 5.2 | Spatial evaluation of the products' performance

The highest correlations for all products were found in the western part of the study area, where most of the precipitation occurs in winter and spring (Figure 4). GSMAP overestimated the variability of precipitation ( $\gamma$ ) in the southern Pamir and western Tien Shan, while the other products, especially MSWEP, underestimated the variability in this area. All of the datasets overestimated precipitation ( $\beta$ ) in the same region, with CHIRPS and its upscaled version performing slightly better. In the southeastern Pamir and western Tien Shan, the overall performance (measured with KGE') was poor for all products, but especially that of GSMAP. The precipitation products performed best in the stations located in the western Pamir and northern Tien Shan, with KGE' values of 0.92 for MSWEP, 0.87 for both CHIRPS products, and 0.83 for GSMAP. Overall, the MAE was lowest in the western region where MSWEP performed best, followed by GSMAP, CHIRPS upscaled, and CHIRPS. We found the highest MAE in the southwestern Tien Shan for all products, followed by the southern Pamir, especially for GSMAP and MSWEP.

To determine how topography and climate regime affected the products' performance, we grouped all






FIGURE 3 Mean absolute error (MAE) in mm of various global precipitation datasets and precipitation gauge data for six different temporal scales. The vertical blue line indicates the optimum value for MAE [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]



**FIGURE 4** KGE', its components ( $r$ ,  $\beta$ ,  $\gamma$ ), and MAE derived from a monthly timescale. The colours for KGE',  $r$ , and MAE range from light yellow (very poor performance) to dark red (best performance). For  $\beta$  and  $\gamma$ , white colours represent their best performance, while underestimation is depicted in dark purple and overestimation is depicted in dark green. Grey colour identifies stations not available for product evaluation (i.e., GSMAP, data before 2000). The black outlines correspond to the countries (see Figure 1 for a detailed map of the region) [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

stations by elevation bands, precipitation amount, and precipitation regime (Figure 5). All of the products, but especially GSMAP, had lower performance at high elevations (>3,000 m). MSWEP performed best in the lowlands (<1,000 m), while CHIRPS excelled at mid-altitudes (2,000–3,000 m). The MSWEP and CHIRPS products had similar performance between 1,000 m and 2,000 m elevation.

Considering annual precipitation, MSWEP performed best for wetter locations (>300 mm·year<sup>-1</sup>), while both CHIRPS performed best for moderately wet locations (150–300 mm·year<sup>-1</sup>) (Figure 5b). For drier locations (<150 mm·year<sup>-1</sup>), all of the products failed to capture the precipitation dynamics. With respect to precipitation regimes, all of the products performed best in cluster 1, where most of the precipitation falls



**FIGURE 5**  $KGE'$  at a monthly timescale between the precipitation products and precipitation gauge data for different (a) elevation bands: 0–1,000 m, 1,000–2,000 m, 2,000–3,000 m, and 3,000–4,000 m; (b) mean annual precipitation (2000–2015); (c) precipitation regimes with corresponding clusters (cluster 1 has winter–spring precipitation, long dry summers; cluster 2 has winter–spring precipitation, short dry summers, and cluster 3 is characterized by summer precipitation). The vertical blue line indicates the optimum value for  $KGE'$ .  $N$  indicates the number of stations in each group [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

in winter and spring, with long dry summers and autumns (Figures 1 and 5c). MSWEP showed the highest median values, followed by CHIRPS. In cluster 2, with precipitation in winter and spring but short, dry summers, CHIRPS products performed better than the other products did. However, in cluster 3 (summer precipitation), all of the products performed poorly, and only MSWEP had positive median KGE' values. CHIRPS and its upscaled version were unable to capture the summer precipitation regime of most of the stations (Figure 6e,f), and MSWEP performed better in representing the region's climatology but overestimated the precipitation amounts.

### 5.3 | Evaluation of dry and wet spells

All of the precipitation products were able to accurately detect dry spells with  $POD > 0.6$  (Figure 7). However, the ability to detect wet spells decreased proportionally with increasing precipitation intensity. MSWEP showed slightly better performance in terms of POD, except for the most intense precipitation class, for which CHIRPS upscaled performed better. The FAR values are consistent with the POD, and all of the products identified dry spells very well, with MSWEP having slightly better performance, which decreased with precipitation intensity, and GSMAP having better performance for moderate precipitation events. CHIRPS showed the closest agreement for all precipitation intensities in terms of fBias, with a slight overestimation (fBias  $> 1$ ) of light events and an underestimation (fBias  $< 1$ ) of heavy precipitation.


## 6 | DISCUSSION

We evaluated the performance of three precipitation products (CHIRPS, GSMAP, and MSWEP) with a spatial resolution higher than 12 km to capture local precipitation patterns over the heterogeneous topography and climate of Central Asia. To do so, we collected precipitation data from 30 independent gauging stations across the region. We accounted for elevation, precipitation regime, precipitation amount, intensity of wet spells, temporal dynamics, and different timescales. Overall, the products all performed best in (a) altitudes below 3,000 m; (b) regions dominated by winter and spring precipitation; and (c) wetter periods (i.e., winter and spring) and locations with between 150 and 600 mm of precipitation per year, and the products accurately detected dry spells. We found key differences between the products. MSWEP was best at capturing precipitation dynamics, CHIRPS was best at representing the volume and distribution of precipitation over


different timescales and locations, and GSMAP generally showed poorer performance. We also evaluated MSWEP v2.8 for the first time and found that, as compared to previous versions (the results of an earlier evaluation are presented in Figures S4 and S5), v2.8 improved the overall performance in the study region, especially for spring and winter, and did not overestimate precipitation as much.

MSWEP and CHIRPS also captured precipitation dynamics well for the Tibetan Plateau (MSWEP v2) (Liu *et al.*, 2019), Chile (MSWEP v1.1) (Zambrano-Bigiarini *et al.*, 2017), western Africa (MSWEPv2.2) (Satgé *et al.*, 2020), India (MSWEP v2.1) (Prakash, 2019), and the Bolivian Altiplano (MSWEP v2.1) (Satgé *et al.*, 2019). Because CHIRPS is intended to support agricultural drought monitoring, its best performance was expected at around the wettest months for each location (Funk *et al.*, 2015). This is supported by our results for Central Asia, where CHIRPS performed best for wetter periods and locations. Similar to our results, GSMAP also had the comparatively poor performance for the mountainous endorheic system of the Bolivian Altiplano (Satgé *et al.*, 2019) and for western Africa (Satgé *et al.*, 2020). The accuracy of GSMAP estimates may be affected by the lower number of stations in the source data CPC compared to MSWEP and CHIRPS (Satgé *et al.*, 2020).

Despite MSWEP and CHIRPS having the best overall performance, we found some limitations. Both products performed worst in summer (overestimation of precipitation), during the driest period in areas where winter and spring precipitation dominate (clusters 1 and 2; Figure 1), and for stations in areas with precipitation below  $150 \text{ mm} \cdot \text{year}^{-1}$ . Similar findings have been reported for CHIRPS in other drylands, such as northeast Brazil (Paredes-Trejo *et al.*, 2017), Sub-Saharan Africa (Harrison *et al.*, 2019), and Mainland China (Bai *et al.*, 2018), as well as for MSWEP (v2.1) in northeast India (Prakash, 2019). The low performance in these areas arguably is due to the very low precipitation, in that a single incorrectly identified rainfall event could lead to 100% overestimation or underestimation (Zambrano-Bigiarini *et al.*, 2017). Satellite-based precipitation estimates may be more suited to estimating convective tropical rainfall patterns than the isolated, highly localized, and short-lived convective rainfall typical in semi-arid to arid areas (Dinku *et al.*, 2010; Thiemi *et al.*, 2012; Beck *et al.*, 2017b). Our findings support this claim. In dry regions, detecting precipitation is difficult because space-born sensors (e.g., microwave and infrared sensors) can miss the subcloud evaporation of raindrops or rainfall suppression by desert aerosols (e.g., mineral dust) and be affected by the land's surface properties, such as a hot background (e.g., upwelling microwave radiation) (Dinku *et al.*, 2011; Beck *et al.*, 2017b).



**FIGURE 6** Interannual variation in precipitation and annual precipitation estimated by the precipitation products (dashed lines), as compared to the gauging stations (solid lines) for each cluster. Cluster 1 (winter/spring precipitation; long, dry summers); cluster 2 (winter/spring precipitation; short, dry summers), and cluster 3 (summer precipitation) [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]



**FIGURE 7** Median values of probability of detection (POD), false alarm ratio (FAR), and frequency of bias (fBias) performance for dry spells (<1 mm-day<sup>-1</sup>) and wet spells of different intensities (1–5, 5–20, 20–40, and >40 mm-day<sup>-1</sup>). The horizontal dashed line indicates the optimum POD, FAR, and fBIAS values [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

We found that the products overestimated precipitation at higher elevations (>3,000 m), possibly because the gauge network density in such areas is low (Harrison *et al.*, 2019). In complex mountainous terrains, precipitation can be falsely detected due to long-lasting orographic clouds or by the contrast between the temperature and the emissivity of rough land surfaces of water and snow-covered areas, which satellite sensors can misinterpret as precipitation (Gebregiorgis and Hossain, 2013; Guo *et al.*, 2015; Satgé *et al.*, 2019). In addition, in global evaluations, the reanalyses exhibited lower accuracy than the microwave- and infrared-based satellite datasets in the tropics did (Beck *et al.*, 2017b). In contrast, these products perform well in extratropical regions, probably linked to deficiencies in the subgrid convection parameterization schemes along with issues in the land surface parameterization (Beck *et al.*, 2017b). The coverage of the raw data sources, orographic correction, and interpolation techniques may compromise the accuracy of the precipitation products (Sun *et al.*, 2018). Considering the high dependency of the global precipitation products on local gauge calibration, more efforts are needed to increase the accessibility of local observations in order to improve the products' quality and reliability for hydrological, agricultural, and climate studies.

All of the products performed worse in southeastern Pamir and Tien Shan, where precipitation peaks in summer. Generally, such poor performance among all of the products for summer precipitation can be related to challenges in capturing the orographic uplift of warm clouds, false detection of very cold high clouds as precipitating by infrared products, and microwave products missing warm precipitation from shallow clouds (Gebregiorgis

and Hossain, 2013; Behrangi *et al.*, 2014; Satgé *et al.*, 2019). Moreover, for the reanalysis-based products (i.e., CHIRPS and MSWEP), poor summer performance might additionally have resulted from an unrealistic northward displacement of the monsoon cycle (Di Giuseppe *et al.*, 2013) and from the fact that atmospheric models in mid-latitudes can more reliably predict winter precipitation associated with synoptic systems such as fronts than it can summer precipitation, which is more often associated with convective systems such as thunderstorms (Haiden *et al.*, 2012; Zhu *et al.*, 2014).

Although the examined precipitation products were able to detect dry spells accurately, their performance decreased for wet spells as precipitation intensity increased. This lower performance for higher intensities can be associated with local storm events with a spatial extent smaller than the satellites' spatial resolution (Thiemig *et al.*, 2012). The weaker detection ability of CHIRPS might be related to its fixed threshold for detecting precipitation from cloud temperatures that might not be appropriate for this region as well as its dependency on the 0.25° TRMM training data, which contributes to the false detection of rainfall events when averaged over larger areas (Dinku *et al.*, 2010; Toté *et al.*, 2015; Paredes-Trejo *et al.*, 2017; Dinku *et al.*, 2018). The reported duplication and inconsistency in the gauge sources used to calibrate CHIRPS could be additional sources of uncertainty (Rivera *et al.*, 2018).

Finally, our results suggest that the precipitation product's selection depends on the specific user needs or application and the regional characteristics. For example, CHIRPS and MSWEP perform best during winter and spring, which makes them suitable to assess terrestrial

water storage prior to the irrigation season. While CHIRPS provides daily precipitation amounts, MSWEP has a higher temporal resolution (3-hr resolution), making it more appropriate for subseasonal hydrological monitoring and forecasting (Beck *et al.*, 2017b). Both products have had a long temporal record, from 1979 (MSWEP) and 1981 (CHIRPS) to near the present, with a delay of several days (CHIRPS) or several hours (MSWEP). CHIRPS has a higher spatial resolution ( $0.05^\circ$ ) as compared to MSWEP ( $0.1^\circ$ ) and more suitable for smaller catchments in elevations below 3,000 m. The products' best performance was achieved in the western, central, and northern Tien Shan and Pamir mountains and in adjacent regions. Although we did not find considerable differences between the original and upscaled version of CHIRPS, a lower performance of the upscaled version can arise from the resampling method used for upscaling; hence, we advise the use of CHIRPS in its native resolution.

## 7 | CONCLUSION

We presented the first evaluation of three global high-resolution precipitation products over the heterogeneous topography and climate of Central Asia using independent station data. We quantified the products' ability to reproduce temporal dynamics while preserving the volume and distribution of precipitation, evaluated the products' accuracy, and assessed the products' ability to detect dry and wet spells of different intensities accurately.

We found that CHIRPS and MSWEP were the most reliable global products for obtaining high-resolution precipitation estimates in Central Asia, especially for wet seasons. Nevertheless, our results highlight high spatial and temporal heterogeneity of the performance, which indicates that the final product for a local application must be selected with care, based on the guidelines provided above. This is particularly relevant for regions with low precipitation levels and in complex terrain where ground station data are sparse.

## ACKNOWLEDGEMENTS

This research is supported by the Volkswagen Foundation within the Sustainable Agricultural Development in Central Asia (SUSADICA) project (Grant No. 96264). We would like to thank the Hydrometeorological Service of Tajikistan, Dr. Stephan Weise, Dr. Heiko Apel, Dr. Abror Gafurov, and Dr. Zafar Gafurov for their help in acquiring the station data. We also would like to thank the two anonymous reviewers and the Editor Dr. Jose Marengo for their valuable comments.

## AUTHOR CONTRIBUTIONS

**Mayra Daniela Peña-Guerrero:** Conceptualization; data curation; formal analysis; investigation; methodology; software; visualization; writing – original draft; writing – review and editing. **Atabek Umirbekov:** Conceptualization; writing – review and editing. **Larisa Tarasova:** Conceptualization; methodology; visualization; writing – review and editing. **Daniel Müller:** Conceptualization; supervision; visualization; writing – review and editing.

## ORCID

Mayra Daniela Peña-Guerrero  <https://orcid.org/0000-0001-8974-2956>

Atabek Umirbekov  <https://orcid.org/0000-0001-5567-2574>

Larisa Tarasova  <https://orcid.org/0000-0002-7675-0751>

Daniel Müller  <https://orcid.org/0000-0001-8988-0718>

## REFERENCES

- Apel, H., Abdykerimova, Z., Agalhanova, M., Baimaganbetov, A., Gavrilenko, N., Gerlitz, L., Kalashnikova, O., Unger-Shayesteh, K., Vorogushyn, S. and Gafurov, A. (2018) Statistical forecast of seasonal discharge in central Asia using observational records: development of a generic linear modelling tool for operational water resource management. *Hydrology and Earth System Sciences*, 22, 2225–2254.
- Baez-Villanueva, O.M., Zambrano-Bigiarini, M., Ribbe, L., Nauditt, A., Giraldo-Osorio, J.D. and Thinh, N.X. (2018) Temporal and spatial evaluation of satellite rainfall estimates over different regions in Latin-America. *Atmospheric Research*, 213, 34–50.
- Bai, L., Shi, C., Li, L., Yang, Y. and Wu, J. (2018) Accuracy of CHIRPS satellite-rainfall products over mainland China. *Remote Sensing*, 10, 362.
- Barlow, M.A. and Tippett, M.K. (2008) Variability and predictability of central Asia river flows: antecedent winter precipitation and large-scale teleconnections. *Journal of Hydrometeorology*, 9, 1334–1349.
- Beck, H.E., Van Dijk, A.I.J.M., Levizzani, V., Schellekens, J., Miralles, D.G., Martens, B. and De Roo, A. (2017a) MSWEP: 3-hourly  $0.25^\circ$  global gridded precipitation (1979–2015) by merging gauge, satellite, and reanalysis data. *Hydrology and Earth System Sciences*, 21, 589–615.
- Beck, H.E., Vergopolan, N., Pan, M., Levizzani, V., Van Dijk, A.I.J.M., Weedon, G.P., Brocca, L., Pappenberger, F., Huffman, G.J. and Wood, E.F. (2017b) Global-scale evaluation of 22 precipitation datasets using gauge observations and hydrological modeling. *Hydrology and Earth System Sciences*, 21, 6201–6217.
- Beck, H.E., Wood, E.F., Pan, M., Fisher, C.K., Miralles, D.G., Van Dijk, A.I.J.M., Mcvicar, T.R. and Adler, R.F. (2019) MSWEP V2 global 3-hourly  $0.1^\circ$  precipitation: methodology and quantitative assessment. *Bulletin of the American Meteorological Society*, 100, 473–500.
- Beck, H. E., Wood, E. F., Pan, M., Fisher, C. K., Miralles, D. G., van Dijk, A. I. J. M., Mcvicar, T. R., and Adler, R. F. (2021) MSWEP

- V2.8 technical documentation. Available at: <http://www.gloh2o.org/mswep/> [Accessed 7th April, 2021].
- Behrangi, A., Tian, Y., Lambrigtsen, B.H. and Stephens, G.L. (2014) What does CloudSat reveal about global land precipitation detection by other spaceborne sensors? *Water Resources Research*, 50, 4893–4905.
- Böhner, J. (2006) General climatic controls and topoclimatic variations in central and high Asia. *Boreas*, 35, 279–295.
- Di Giuseppe, F., Molteni, F. and Dutra, E. (2013) Real-time correction of ERA-Interim monthly rainfall. *Geophysical Research Letters*, 40, 3750–3755.
- Dinku, T., Ceccato, P. and Connor, S.J. (2011) Challenges of satellite rainfall estimation over mountainous and arid parts of East Africa. *International Journal of Remote Sensing*, 32, 5965–5979.
- Dinku, T., Ceccato, P., Cressman, K. and Connor, S.J. (2010) Evaluating detection skills of satellite rainfall estimates over desert locust recession regions. *Journal of Applied Meteorology and Climatology*, 49, 1322–1332.
- Dinku, T., Funk, C., Peterson, P., Maidment, R., Tadesse, T., Gadain, H. and Ceccato, P. (2018) Validation of the CHIRPS satellite rainfall estimates over eastern Africa. *Quarterly Journal of the Royal Meteorological Society*, 144, 292–312.
- Ebert, E.E. (2007) Methods for verifying satellite precipitation estimates. In: Levizzani, V. and Turk, F.J. (Eds.) *Measuring Precipitation From Space*. Dordrecht: Springer.
- Funk, C., Peterson, P., Landsfeld, M., Pedreros, D., Verdin, J., Shukla, S., Husak, G., Rowland, J., Harrison, L., Hoell, A. and Michaelsen, J. (2015) The climate hazards infrared precipitation with stations—a new environmental record for monitoring extremes. *Scientific Data*, 2, 150066.
- Gebregiorgis, A.S. and Hossain, F. (2013) Understanding the dependence of satellite rainfall uncertainty on topography and climate for hydrologic model simulation. *IEEE Transactions on Geoscience and Remote Sensing*, 51, 704–718.
- Guo, H., Bao, A., Liu, T., Kurban, A. and De Maeyer, P. (2017) Systematic evaluation of satellite precipitation estimates over central Asia using an improved error-component procedure. *Journal of Geophysical Research: Atmospheres*, 122, 10906–11927.
- Guo, H., Chen, S., Bao, A., Hu, J., Gebregiorgis, A., Xue, X. and Zhang, X. (2015) Inter-comparison of high-resolution satellite precipitation products over central Asia. *Remote Sensing*, 7, 7181–7211.
- Gupta, H.V., Kling, H., Yilmaz, K.K. and Martinez, G.F. (2009) Decomposition of the mean squared error and NSE performance criteria: implications for improving hydrological modeling. *Journal of Hydrology*, 377, 80–91.
- Haiden, T., Richardson, D.S., Rodwell, M.J., Okagaki, A., Robinson, T. and Hewson, T. (2012) Intercomparison of global model precipitation forecast skill in 2010/11 using the SEEPS score. *Monthly Weather Review*, 140, 2720–2733.
- Harris, I., Jones, P.D., Osborn, T.J. and Lister, D.H. (2014) Updated high-resolution grids of monthly climatic observations—the CRU TS3.10 dataset. *International Journal of Climatology*, 34, 623–642.
- Harrison, L., Funk, C. and Peterson, P. (2019) Identifying changing precipitation extremes in sub-Saharan Africa with gauge and satellite products. *Environmental Research Letters*, 14, 085007.
- Hellwig, J., Stahl, K., Ziese, M. and Becker, A. (2018) The impact of the resolution of meteorological data sets on catchment-scale precipitation and drought studies. *International Journal of Climatology*, 38, 3069–3081.
- Henn, B., Newman, A.J., Livneh, B., Daly, C. and Lundquist, J.D. (2018) An assessment of differences in gridded precipitation datasets in complex terrain. *Journal of Hydrology*, 556, 1205–1219.
- Hu, Z., Zhou, Q., Chen, X., Li, J., Li, Q., Chen, D., Liu, W. and Yin, G. (2018) Evaluation of three global gridded precipitation data sets in central Asia based on rain gauge observations. *International Journal of Climatology*, 38, 3475–3493.
- Huffman, G.J., Adler, R.F., Gu, G., Nelkin, E.J., Bolvin, D.T., Gu, G., Hong, Y., Bowman, K.P., Stocker, E.F. and Wolff, D.B. (2007) The TRMM multisatellite precipitation analysis (TMPA): quasi-global, multiyear, combined-sensor precipitation estimates at fine scales. *Journal of Hydrometeorology*, 8, 38–55.
- Immerzeel, W.W., Lutz, A.F., Andrade, M., Bahl, A., Biemans, H., Bolch, T., Hyde, S., Brumby, S., Davies, B.J., Elmore, A.C., Emmer, A., Feng, M., Fernandez, A., Haritashya, U., Kargel, J. S., Koppes, M., Kraaijenbrink, P.D.A., Kulkarni, A.V., Mayewski, P.A., Nepal, S., Pacheco, P., Painter, T.H., Pellicciotti, F., Rajaram, H., Rupper, S., Sinisalo, A., Shrestha, A.B., Viviroli, D., Wada, Y., Xiao, C., Yao, T. and Baillie, J.E.M. (2020) Importance and vulnerability of the world's water towers. *Nature*, 577, 364–369.
- Joyce, R.J., Janowiak, J.E., Arkin, P.A. and Xie, P. (2004) CMORPH: a method that produces global precipitation estimates from passive microwave and infrared data at high spatial and temporal resolution. *Journal of Hydrometeorology*, 5, 487–503.
- Kling, H., Fuchs, M. and Paulin, M. (2012) Runoff conditions in the upper Danube basin under an ensemble of climate change scenarios. *Journal of Hydrology*, 424–425, 264–277.
- Kubota, T., Shige, S., Hashizume, H., Aonashi, K., Takahashi, N., Seto, S., Hirose, M., Takayabu, Y.N., Ushio, T., Nakagawa, K., Iwanami, K., Kachi, M. and Okamoto, K.I. (2007) Global precipitation map using satellite-borne microwave radiometers by the GSMaP project: production and validation. *IEEE Transactions on Geoscience and Remote Sensing*, 45, 2259–2275.
- Liu, J., Shangguan, D., Liu, S., Ding, Y., Wang, S. and Wang, X. (2019) Evaluation and comparison of CHIRPS and MSWEP daily-precipitation products in the Qinghai–Tibet Plateau during the period of 1981–2015. *Atmospheric Research*, 230, 104634.
- Lloyd, S.P. (1982) Least squares quantization in PCM. *IEEE Transactions on Information Theory*, 28, 129–137.
- Lu, X., Tang, G., Liu, X., Wang, X., Liu, Y. and Wei, M. (2021) The potential and uncertainty of triple collocation in assessing satellite precipitation products in central Asia. *Atmospheric Research*, 252, 105452.
- Malsy, M., Aus Der Beek, T. and Flörke, M. (2014) Evaluation of large-scale precipitation data sets for water resources modelling in central Asia. *Environmental Earth Sciences*, 73, 787–799.
- Mannig, B., Müller, M., Starke, E., Merckenschlager, C., Mao, W., Zhi, X., Podzun, R., Jacob, D. and Paeth, H. (2013) Dynamical downscaling of climate change in central Asia. *Global and Planetary Change*, 110, 26–39.
- Mariotti, A. (2007) How ENSO impacts precipitation in southwest central Asia. *Geophysical Research Letters*, 34(16), 1–5.
- Maussion, F., Scherer, D., Mölg, T., Collier, E., Curio, J. and Finkelnburg, R. (2013) Precipitation seasonality and variability

- over the Tibetan Plateau as resolved by the High Asia reanalysis. *Journal of Climate*, 27, 1910–1927.
- Meier, C., Knoche, M., Merz, R. and Weise, S.M. (2013) Stable isotopes in river waters in the Tajik Pamirs: regional and temporal characteristics. *Isotopes in Environmental and Health Studies*, 49, 542–554.
- Michaelides, S., Levizzani, V., Anagnostou, E., Bauer, P., Kasparis, T. and Lane, J.E. (2009) Precipitation: measurement, remote sensing, climatology and modeling. *Atmospheric Research*, 94, 512–533.
- Nijssen, B. (2004) Effect of precipitation sampling error on simulated hydrological fluxes and states: anticipating the global precipitation measurement satellites. *Journal of Geophysical Research*, 109, D02103.
- Okamoto, K.I., Ushio, T., Iguchi, T., Takahashi, N. and Iwanami, K. (2005) The global satellite mapping of precipitation (GSMaP) project. In: *IEEE International Geoscience and Remote Sensing Symposium*. Seoul: IEEE, pp. 3414–3416.
- Parajka, J., Merz, R., Szolgay, J., Blöschl, G., Kohnova, S. and Hlavčová, K. (2008) A comparison of precipitation and runoff seasonality in Slovakia and Austria. *Meteorologický časopis*, 11, 9–14.
- Pardé, M. (1933) *Fleuves et Rivières*. Paris: Armand Colin.
- Paredes-Trejo, F.J., Barbosa, H.A. and Lakshmi Kumar, T.V. (2017) Validating CHIRPS-based satellite precipitation estimates in Northeast Brazil. *Journal of Arid Environments*, 139, 26–40.
- Peng, D., Zhou, T., Zhang, L. and Zou, L. (2019) Detecting human influence on the temperature changes in central Asia. *Climate Dynamics*, 53, 4553–4568.
- Prakash, S. (2019) Performance assessment of CHIRPS, MSWEP, SM2RAIN-CCI, and TMPA precipitation products across India. *Journal of Hydrology*, 571, 50–59.
- Reyer, C.P.O., Otto, I.M., Adams, S., Albrecht, T., Baarsch, F., Carlsburg, M., Coumou, D., Eden, A., Ludi, E., Marcus, R., Mengel, M., Mosello, B., Robinson, A., Schleussner, C.-F., Serdeczny, O. and Stagl, J. (2017) Climate change impacts in central Asia and their implications for development. *Regional Environmental Change*, 17, 1639–1650.
- Rivera, J.A., Marianetti, G. and Hinrichs, S. (2018) Validation of CHIRPS precipitation dataset along the Central Andes of Argentina. *Atmospheric Research*, 213, 437–449.
- Salehie, O., Ismail, T., Shahid, S., Ahmed, K., Adarsh, S., Asaduzzaman, M. and Dewan, A. (2021) Ranking of gridded precipitation datasets by merging compromise programming and global performance index: a case study of the Amu Darya basin. *Theoretical and Applied Climatology*, 144, 985–999.
- Satgé, F., DeFrance, D., Sultan, B., Bonnet, M.-P., Seyler, F., Rouché, N., Pierron, F. and Paturel, J.-E. (2020) Evaluation of 23 gridded precipitation datasets across West Africa. *Journal of Hydrology*, 581, 124412.
- Satgé, F., Ruelland, D., Bonnet, M.-P., Molina, J. and Pillco, R. (2019) Consistency of satellite-based precipitation products in space and over time compared with gauge observations and snow-hydrological modelling in the Lake Titicaca region. *Hydrology and Earth System Sciences*, 23, 595–619.
- Schär, C., Vasilina, L., Pertziger, F. and Dirren, S. (2004) Seasonal runoff forecasting using precipitation from meteorological data assimilation systems. *Journal of Hydrometeorology*, 5, 959–973.
- Schiemann, R., Lüthi, D., Vidale, P.L. and Schär, C. (2008) The precipitation climate of central Asia—intercomparison of observational and numerical data sources in a remote semi-arid region. *International Journal of Climatology*, 28, 295–314.
- Shige, S., Kida, S., Ashiwake, H., Kubota, T. and Aonashi, K. (2013) Improvement of TMI rain retrievals in mountainous areas. *Journal of Applied Meteorology and Climatology*, 52, 242–254.
- Shukla, S., McNally, A., Husak, G. and Funk, C. (2014) A seasonal agricultural drought forecast system for food-insecure regions of East Africa. *Hydrology and Earth System Sciences*, 18, 3907–3921.
- Sorg, A., Bolch, T., Stoffel, M., Solomina, O. and Beniston, M. (2012) Climate change impacts on glaciers and runoff in Tien Shan (central Asia). *Nature Climate Change*, 2, 725–731.
- Sun, Q., Miao, C., Duan, Q., Ashouri, H., Sorooshian, S. and Hsu, K.L. (2018) A review of global precipitation data sets: data sources, estimation, and intercomparisons. *Reviews of Geophysics*, 56, 79–107.
- Tan, X., Wu, Y., Liu, B. and Chen, S. (2020) Inconsistent changes in global precipitation seasonality in seven precipitation datasets. *Climate Dynamics*, 54, 3091–3108.
- Teixeira, E.I., Fischer, G., Van Velthuisen, H., Walter, C. and Ewert, F. (2013) Global hot-spots of heat stress on agricultural crops due to climate change. *Agricultural and Forest Meteorology*, 170, 206–215.
- Thiemig, V., Rojas, R., Zambrano-Bigiarini, M., Levizzani, V. and De Roo, A. (2012) Validation of satellite-based precipitation products over sparsely gauged African River basins. *Journal of Hydrometeorology*, 13, 1760–1783.
- Thorndike, R.L. (1953) Who belongs in the family? *Psychometrika*, 18, 267–276.
- Tobin, K.J. and Bennett, M.E. (2014) Satellite precipitation products and hydrologic applications. *Water International*, 39, 360–380.
- Toté, C., Patricio, D., Boogaard, H., Van Der Wijngaart, R., Tarnavsky, E. and Funk, C. (2015) Evaluation of satellite rainfall estimates for drought and flood monitoring in Mozambique. *Remote Sensing*, 7, 1758–1776.
- Trenberth, K.E. (2011) Changes in precipitation with climate change. *Climate Research*, 47, 123–138.
- Unger-Shayesteh, K., Vorogushyn, S., Farinotti, D., Gafurov, A., Duethmann, D., Mandychev, A. and Merz, B. (2013) What do we know about past changes in the water cycle of central Asian headwaters? A review. *Global and Planetary Change*, 110, 4–25.
- Uppala, S.M., Kållberg, P.W., Simmons, A.J., Andrae, U., Bechtold, V.D.C., Fiorino, M., Gibson, J.K., Haseler, J., Hernandez, A., Kelly, G.A., Li, X., Onogi, K., Saarinen, S., Sokka, N., Allan, R.P., Andersson, E., Arpe, K., Balmaseda, M. A., Beljaars, A.C.M., Berg, L.V.D., Bidlot, J., Bormann, N., Caires, S., Chevallier, F., Dethof, A., Dragosavac, M., Fisher, M., Fuentes, M., Hagemann, S., Hólm, E., Hoskins, B.J., Isaksen, I., Janssen, P.A.E.M., Jenne, R., McNally, A.P., Mahfouf, J.F., Morcrette, J.J., Rayner, N.A., Saunders, R.W., Simon, P., Sterl, A., Trenberth, K.E., Untch, A., Vasiljevic, D., Viterbo, P. and Woollen, J. (2005) The ERA-40 re-analysis. *Quarterly Journal of the Royal Meteorological Society*, 131, 2961–3012.
- Ushio, T., Sasashige, K., Kubota, T., Shige, S., Okamoto, K.I., Aonashi, K., Inoue, T., Takahashi, N., Iguchi, T., Kachi, M., Oki, R., Morimoto, T. and Kawasaki, Z.-I. (2009) A Kalman

- filter approach to the global satellite mapping of precipitation (GSMaP) from combined passive microwave and infrared radiometric data. *Journal of the Meteorological Society of Japan*, 87A, 137–151.
- Viviroli, D., Kummu, M., Meybeck, M., Kallio, M. and Wada, Y. (2020) Increasing dependence of lowland populations on mountain water resources. *Nature Sustainability*, 3, 917–928.
- Weingartner, R., Blöschl, G., Hannah, D.M., Marks, D.G., Parajka, J., Pearson, C.S., Rogger, M., Salinas, J.L., Sauquet, E., Srikanthan, R., Thompson, S.E., Viglione, A., Blöschl, G., Sivapalan, M., Wagener, T., Viglione, A. and Savenije, H. (2013) Prediction of seasonal runoff in ungauged basins. In: *Runoff Prediction in Ungauged Basins*. pp. 132–134. Cambridge: Cambridge University Press.
- World Meteorological Organization. (2008) Present and past weather, state of the ground. In: *Guide to Meteorological Instruments and Methods of Observation*, 7th edition. Geneva: WMO.
- Yamamoto, M.K., Shige, S., Yu, C.-K. and Cheng, L.-W. (2017) Further improvement of the heavy orographic rainfall retrievals in the GSMaP algorithm for microwave radiometers. *Journal of Applied Meteorology and Climatology*, 56, 2607–2619.
- Zambrano-Bigiarini, M., Nauditt, A., Birkel, C., Verbist, K. and Ribbe, L. (2017) Temporal and spatial evaluation of satellite-based rainfall estimates across the complex topographical and climatic gradients of Chile. *Hydrology and Earth System Sciences*, 21, 1295–1320.
- Zandler, H., Haag, I. and Samimi, C. (2019) Evaluation needs and temporal performance differences of gridded precipitation products in peripheral mountain regions. *Scientific Reports*, 9, 15118.
- Zhang, Y., Moges, S. and Block, P. (2016) Optimal cluster analysis for objective regionalization of seasonal precipitation in regions of high spatial–temporal variability: application to western Ethiopia. *Journal of Climate*, 29, 3697–3717.
- Zhu, H., Wheeler, M.C., Sobel, A.H. and Hudson, D. (2014) Seamless precipitation prediction skill in the Tropics and extratropics from a global model. *Monthly Weather Review*, 142, 1556–1569.

## SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

**How to cite this article:** Peña-Guerrero, M. D., Umirbekov, A., Tarasova, L., & Müller, D. (2022). Comparing the performance of high-resolution global precipitation products across topographic and climatic gradients of Central Asia. *International Journal of Climatology*, 42(11), 5554–5569. <https://doi.org/10.1002/joc.7548>