

Baumann, Matthias et al.

Article — Published Version

Frontier metrics for a process-based understanding of deforestation dynamics

Environmental Research Letters

Provided in Cooperation with:

Leibniz Institute of Agricultural Development in Transition Economies (IAMO), Halle (Saale)

Suggested Citation: Baumann, Matthias et al. (2022) : Frontier metrics for a process-based understanding of deforestation dynamics, Environmental Research Letters, ISSN 1748-9326, IOP Publishing, Bristol, Vol. 17, Iss. 9, <https://doi.org/10.1088/1748-9326/ac8b9a> , <https://iopscience.iop.org/article/10.1088/1748-9326/ac8b9a>

This Version is available at:

<https://hdl.handle.net/10419/264381>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.



<http://creativecommons.org/licenses/by/4.0>

ENVIRONMENTAL RESEARCH
LETTERS

LETTER

Frontier metrics for a process-based understanding
of deforestation dynamics











OPEN ACCESS

RECEIVED
21 January 2022REVISED
17 August 2022ACCEPTED FOR PUBLICATION
22 August 2022PUBLISHED
7 September 2022

Original content from
this work may be used
under the terms of the
[Creative Commons
Attribution 4.0 licence](#).

Any further distribution
of this work must
maintain attribution to
the author(s) and the title
of the work, journal
citation and DOI.



Matthias Baumann^{1,*} , Ignacio Gasparri² , Ana Buchadas^{1,8} , Julian Oeser¹ , Patrick Meyfroidt^{3,4} ,
Christian Levers⁵ , Alfredo Romero-Muñoz¹ , Yann le Polain de Waroux⁶ , Daniel Müller^{1,7,8} 
and Tobias Kuemmerle^{1,8} 

¹ Geography Department, Humboldt-Universität zu Berlin, Unter den Linden 6, 10099 Berlin, Germany

² Instituto de Ecología Regional (IER), Universidad Nacional de Tucumán (UNT)—Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET). CC:34, CP 4107 Yerba Buena, Tucumán, Argentina & Facultad de Ciencias Naturales e Instituto Miguel Lillo, Universidad Nacional de Tucumán (UNT), Tucumán, Argentina

³ Earth and Life Institute, UCLouvain, Louvain-la-Neuve, Belgium

⁴ XX. Fonds de la Recherche Scientifique F.R.S.-FNRS, 1000 Brussels, Belgium

⁵ Department of Environmental Geography, Institute for Environmental Studies (IVM), Vrije Universiteit Amsterdam, De Boelelaan 1087, 1081 HV Amsterdam, The Netherlands

⁶ Institute for the Study of International Development and Department of Geography, McGill University, Montreal, QC, Canada

⁷ Leibniz Institute of Agricultural Development in Transition Economies (IAMO), Theodor-Lieser-Str. 2, 06120 Halle (Saale), Germany

⁸ Integrative Research Institute on Transformations of Human-Environment Systems (IRI THESys), Humboldt-Universität zu Berlin, Unter den Linden 6, Berlin 10099, Germany

* Author to whom any correspondence should be addressed.

E-mail: matthias.baumann@hu-berlin.de

Keywords: commodity frontiers, deforestation, tropical dry forests and savannahs, agricultural expansion, social-ecological archetypes, Landsat time series

Supplementary material for this article is available [online](#)

Abstract

Agricultural expansion into tropical and subtropical forests often leads to major social-ecological trade-offs. Yet, despite ever-more detailed information on where deforestation occurs, how agriculture expands into forests remains unclear, which is hampered by a lack of spatially and temporally detailed reconstruction of agricultural expansion. Here, we developed and mapped a novel set of metrics that quantify agricultural frontier processes at unprecedented spatial and temporal detail. Specifically, we first derived consistent annual time series of land-use/cover to, second, describe archetypical patterns of frontier expansion, pertaining to the speed, the diffusion and activity of deforestation, as well as post-deforestation land use. We exemplify this approach for understanding agricultural frontier expansion across the entire South American Chaco (1.1 million km²), a global deforestation hotspot. Our study provides three major insights. First, agricultural expansion has been rampant in the Chaco, with more than 19.3 million ha of woodlands converted between 1985 and 2020, including a surge in deforestation after 2019. Second, land-use trajectories connected to frontier processes have changed in major ways over the 35 year study period we studied, including substantial regional variations. For instance, while ranching expansion drove most of the deforestation in the 1980s and 1990s, cropland expansion dominated during the mid-2000s in Argentina, but not in Paraguay. Similarly, 40% of all areas deforested were initially used for ranching, but later on converted to cropping. Accounting for post-deforestation land-use change is thus needed to properly attribute deforestation and associated environmental impacts, such as carbon emissions or biodiversity loss, to commodities. Finally, we identified major, recurrent frontier types that may be a useful spatial template for land governance to match policies to specific frontier situations. Collectively, our study reveals the diversity of frontier processes and how frontier metrics can capture and structure this diversity to uncover major patterns of human–nature interactions, which can be used to guide spatially-targeted policies.

1. Introduction

Agricultural expansion into natural areas has helped to meet the growing global demand for food, feed, and fiber (Godfray *et al* 2010), but has also produced unsustainable land-use outcomes. Agricultural expansion causes widespread deforestation in tropical and subtropical forests, triggering globally-relevant greenhouse gas emissions (Carlson *et al* 2017), biodiversity losses (Chaplin-Kramer *et al* 2015), and major livelihood impacts on forest-dependent people (Andersson and Agrawal 2011, Oldekop *et al* 2020). Much of the agricultural expansion during the past decades has taken place in the tropics (Gibbs *et al* 2010), where most of the last uncultivated, productive lands are found (Ramankutty *et al* 2002, Lambin *et al* 2013). Sustainability planning to prevent or minimize undesirable social-ecological outcomes in regions where agriculture expands is thus needed.

This, first and foremost, requires a robust understanding of where and how frontiers expand. Considerable progress has been made on the prior, that is mapping where deforestation takes place (Hansen *et al* 2013, Turubanova *et al* 2018, Vancutsem *et al* 2021, Zalles *et al* 2021). Yet, mapping *how* agricultural frontiers progress is more complex, and requires a thorough understanding of the underlying processes inherent in frontier dynamics. For example, some frontiers advance slowly while others erupt rapidly (Kröger and Nygren 2020), some frontiers grow outward while others leap-frog to remote places (Bowman *et al* 2012), and some frontiers accelerate while others consolidate and slow down (Bonilla-Moheno and Aide 2020). Likewise, a wide range of land-use-actors drive frontier expansion, such as swidden cultivators (Vieilledent *et al* 2018), forest smallholders (Tyukavina *et al* 2018, Phiri *et al* 2019), or agribusinesses (Klink and Machado 2005). Further, in some regions, frontiers may be considered old or suspended, whereas in other regions new frontiers emerge. Lastly, land-use trajectories after initial deforestation are diverse (Hosonuma *et al* 2012, De Sy *et al* 2019, Souza *et al* 2020, Song *et al* 2021). Although methods to quantitatively characterize such complexity across large agricultural frontiers are still underdeveloped, there is a need to capture and describe complex frontier dynamics to support context-specific land governance and address sustainability challenges in frontier regions (Pacheco *et al* 2021). Archetype analyses aimed at identifying major patterns of human–environment interactions (Lambin *et al* 2003, Geist and Lambin 2004, Eisenack *et al* 2006, 2019, Oberlack *et al* 2019, Sietz *et al* 2019, Rocha *et al* 2020), such as typical land systems (Vaclavik *et al* 2013, Levers *et al* 2018), land-use change trajectories (Levers *et al* 2018, Meyfroidt *et al* 2018), or land-use outcomes (Cumming *et al* 2014, Pacheco-Romero *et al* 2021), are a potentially

powerful way to structure diversity and complexity of land-use dynamics for that purpose.

Mapping archetypical patterns of frontier processes and better identifying what drives them could enable more nuanced land governance. For example, identifying emerging frontiers would allow for proactive land-use and conservation planning (e.g. zoning), whereas reactive interventions (e.g. forest protection) would be needed where frontiers are particularly active (Hansen *et al* 2020). Likewise, where frontiers consolidate, restoration opportunities might unfold, as land-use actors are more interested in long-term sustainability (Latawiec *et al* 2015, Lerner *et al* 2015, Strassburg *et al* 2017). Disentangling frontier dynamics can furthermore help to identify actor-specific governance interventions. For example, historically, frontiers have mainly been driven by smallholders (Barbier 2012, Pacheco 2012, Godar *et al* 2014), but since the late 1990s, capital-intensive, influential actors have been driving frontiers to produce commodities for global markets (Rudel 2007, Kröger and Nygren 2020). Such commodity frontiers are typically characterized by agglomeration effects (Garrett *et al* 2013, Austin *et al* 2017, Richards 2018) and are sensitive to macroeconomic and trade signals, which can produce abrupt accelerations of frontier dynamics. In addition, land-use actors in commodity frontiers are potentially responsive to market-based interventions (Zu Ermgassen *et al* 2020), for example through supply-chain governance interventions or certification systems (Baynes *et al* 2015). Finally, identifying recurring patterns of frontier will allow for assessments of what explains these key dynamics, which directly contributes to building theories in land system science (Meyfroidt *et al* 2018, Turner *et al* 2020). Yet, we lack a robust understanding and a set of quantitative indicators that capture how frontiers unfold.

Increasing access to satellite images along with new processing capabilities offer new opportunities for understanding frontier dynamics at unprecedented temporal and spatial resolution (Gorelick *et al* 2017, Wulder *et al* 2019, Woodcock *et al* 2020), yet these opportunities have so far not been explored. Prior work on assessing frontiers has mostly focused on mapping deforestation (Hansen *et al* 2013, Müller *et al* 2016, Griffiths *et al* 2018, Vancutsem *et al* 2021), what follows deforestation (Zalles *et al* 2019, 2021, Souza *et al* 2020, Song *et al* 2021) or, most recently, who drives deforestation frontiers (Curtis *et al* 2018, Pacheco *et al* 2021). The question of how frontier dynamics unfold, beyond identifying hotspots of deforestation (Harris *et al* 2017, Tyukavina *et al* 2018, Potapov *et al* 2019), remains largely unexplored. Specifically, remote-sensing time series should allow to describe speed at which frontiers expand (e.g. slow vs. fast progressing), frontier stage (e.g. emerging, active, consolidated) or the frontier diffusion process

(e.g. gradually progressing vs. leap-frogging frontiers). Translating land-cover time series into such process-based system metrics requires overcoming two key challenges. First, it requires deriving consistent time series that do not solely represent collections of individual maps, as error propagation makes analyzing changes between these maps and the derivation of process-based metrics difficult to interpret (Friedl *et al* 2010, Sulla-Menashe *et al* 2019, Liu *et al* 2020). Second, it requires separating human land-use change or management changes (e.g. fallow periods, logging) from other land-cover change such as natural disturbances (e.g. fire) (Gómez *et al* 2016). Overcoming these challenges and establishing consistent time series that represent land-use change would mean a step-change towards translating process-based metrics for better understanding deforestation frontier dynamics.

A better understanding of frontier dynamics is particularly urgent for the world's subtropical tropical dry forests and savannas (hereafter: dry forests). Frontiers have expanded particularly rapidly in these forests over the last decades, but dry forests have received much less attention than rainforests (Miles *et al* 2006, Pennington *et al* 2018). This is surprising, given that dry forests account for nearly 40% of all tropical forests (Murphy and Lugo 1986), harbor high biodiversity (Mayle *et al* 2007), and account for about 30% of the terrestrial primary productivity (Grace *et al* 2006). Dry forest loss has been particularly widespread in South America where agricultural expansion since the early 2000s has turned several dry forest regions into a global deforestation hotspots (Hansen *et al* 2013, Pacheco *et al* 2021, Buchadas *et al* 2022). One of these hotspots is the Gran Chaco in South America shared by Argentina, Bolivia, and Paraguay, where agricultural expansion has been rampant (Hansen *et al* 2013) mostly for beef and cash crop production (Gasparri and Baldi 2013, Fehlenberg *et al* 2017). Where deforestation has occurred in the Chaco is relatively well-understood (Killeen *et al* 2007, Gasparri and Grau 2009, Vallejos *et al* 2015), including post-deforestation land-uses (Boletta *et al* 2006, Campos-Krauer and Wisely 2011, Volante *et al* 2012, Caldas *et al* 2015, Baumann *et al* 2017), and the importance of actors in shaping these pattern (le Polain de Waroux *et al* 2018, Levers *et al* 2021). Yet, how the diversity of actors and social-ecological conditions has produced different types of frontier patterns remains unclear.

Our overarching goal was to develop and test a novel set of frontier metrics that quantitatively describe frontier processes across space and over time. We demonstrate the value of these metrics by deriving archetypical pattern of frontier dynamics driven by agricultural expansion for the Chaco, across the entire history of modern agricultural expansion (1985–2020). Doing so required us to develop the first consistent, spatio-temporally detailed land-cover

reconstruction for this global deforestation hotspot. Specifically, we asked the following questions:

- (a) How can frontier processes and dynamics be described using satellite-based time-series of land cover?
- (b) Where and how have agricultural frontiers expanded into the Chaco's forests since 1985?
- (c) What are archetypical frontier dynamics, including post-deforestation land use change?

2. Methods

2.1. Study area

The Chaco is a 1.1 million km² ecoregion in South America, extending into Argentina, Bolivia, and Paraguay. Mean annual temperature in the Chaco is 22 °C, and annual precipitation shows a pronounced east-west-gradient from 1200 mm in the humid Chaco to 400 mm in the driest regions in the southwest (Bucher 1982). The Chaco is subdivided into the dry Chaco in the west, where xerophyllous forests are dominant and cropping and intensive ranching are dominant, and the wet Chaco in the east where widespread wetlands together with palm savannas and intermixed grasslands for a diverse mosaic of vegetation (Bucher 1982). Historically, land use in the Chaco was dominated by small-scale producers, such as the Eastern European colonies in the Chaco province, or forest smallholders who used a few hectares of land for subsistence cropping to sell on local markets, and the surrounding woodlands to gather firewood and material for rural construction, as well as forest grazing of roaming livestock (Fatecha 1989, Bucher and Huszar 1999). While smallholders continue to be important in parts of the Chaco (Levers *et al* 2021), the emergence and rapid expansion of large-scale agribusinesses has happened over wide areas since the 1990s. These actors have substantial capital and knowledge on the existence and the amount of expected land rent, allowing them to quickly and efficiently capitalize on opportunities that frontier situations entail (le Polain de Waroux 2019). Together with the liberalization of genetically modified soybean variants in the Chaco during the 1990s (Reenberg and Fenger 2011), the introduction of highly productive pasture grasses (e.g. Gatton panic (*Panicum maximum*)) (Vazquez 2013), and the changing export policies of Argentina in reaction to the peso devaluation in 2001 (Leguizamón 2014), this has converted the Chaco into a global deforestation hotspot in the 2000s and 2010s (Hansen *et al* 2013, Baumann *et al* 2017).

2.2. Overview of methodology

Our analytical framework contains three main steps (figure 1). We provide a summary of our




Figure 1. Framework for identifying and characterizing deforestation frontiers. In step 1, we used the Landsat archive to derive consistent land-cover/use time series. Step 2 then derived and mapped six frontier metrics, and sub-categorized each metric (e.g. the metric ‘Frontier speed’ into ‘Slow’, ‘Medium’ and ‘Fast’). Finally, step 3 uses the frontier metrics to identify archetypal frontier dynamics and analyzes them.

methodology here, and a detailed, step-by-step description in the supporting information (text S1–S3). In step 1, we re-constructed land cover across the entire Chaco, annually and consistently for the period 1985–2020. To do so, we made full use of the Landsat satellite archive (>80 000 images) and derived an annual time series of spectral-temporal

metrics (STM) (Oeser *et al* 2020), which we combined with a comprehensive set of training data in a random forest regression framework to derive annual classification probabilities for the classes: (a) woodlands, (b) other vegetation (i.e. natural grasslands, palm savannas), (c) croplands, (d) pastures, and (e) other land covers (i.e. water, bare land, urban

areas). Using these probabilities, we then mapped six land-cover transitions (table S1). We validated each individual map (i.e. 1985–2020) using an independent set of random points, which we evaluated through on-screen interpretation of spectral profiles (e.g. pastures exhibit a different phenological profile throughout the year compared to croplands), mean STM composites of the dry and wet season, and, where available, high-resolution imagery in Google Earth and Bing maps as well as Planet Labs. Based on these points we generated error matrices, calculated overall accuracies as well as class-wise user's and producer's accuracies, and calculated area estimates including confidence intervals of each land-cover class (Olofsson *et al* 2014). Lastly, we aggregated the $30 \times 30 \text{ m}^2$ land-cover maps into two datasets at $1.5 \times 1.5 \text{ km}^2$ resolution: a time series of fractional woodland cover 1985–2020, and a time series of dominant agricultural land cover (i.e. pasture or cropland).

In step 2, we identified frontier areas (i.e. areas with at least 0.5% woodland loss (Rodrigues *et al* 2009, Buchadas *et al* 2022) during three consecutive years and where the final land cover was either cropland or pasture; more information in text S1) and derived six frontier metrics for these areas, which reflect key aspects of frontier processes (table 1): (a) we defined *frontier timing*, describing woodland change 2016–2020 (i.e. the last five years of the time series) relative to 1985–2015. The rationale for this metric was that the temporal course of the frontier determines the type of intervention. (b) We assessed *frontier speed*, representing the strongest annual woodland loss, which determines the focus of regional or national policies aiming at conserving remaining woodlands. (c) *Frontier naturalness* refers to woodlands left relative to the baseline woodlands, which influences the balances of priorities between conservation and restoration. (d) *Frontier diffusion* distinguished between gradual and leap-frogging frontiers, serving as an indicator for the type of actor being dominant in these areas. (e) *Frontier onset*, described the starting year of frontier development and allows for a temporal evaluation of the emergence of different frontier types during our study period. (f) *Frontier land use*, describing land use after woodland loss.

In step 3, we reconstructed how frontiers have unfolded across the region by characterizing the spatio-temporal pattern of our frontier metrics for the time period 1985–2020. First, we assessed frontier dynamics by relating our metric frontier onset (i.e. the year of emergence of a frontier pixel) to the other five frontier metrics in order to find temporal patterns of frontier emergence, and summarized each frontier type for each year of our analysis 1985–2020. We did this for the whole Chaco, the Chaco sections in the three countries, as well as the dry and wet Chaco separately. Second, we identified archetypical

frontier dynamics, by (a) identifying typical combinations of frontier metrics across the entire Chaco, and (b) by quantitatively evaluating our metrics in frontier regions identified in previous research (le Polain de Waroux *et al* 2018). To do so, we used the three most common metric combinations per region and assigned the majority of a category.

3. Results

Forest loss has been rampant in the Chaco with a total of $193\,321 \text{ km}^2$ (28%) of woodlands lost since 1985. Woodland loss increased steadily until 2009/10, when we found highest annual loss rates (1.7%, equaling $10\,167 \text{ km}^2$ in 2009 and 9507 km^2 in 2010), with loss rates declining thereafter (1.1% on average 2011–2019). Most of the woodland loss in 1985–2020 occurred in Argentina ($103\,480 \text{ km}^2$; average annual loss rate of 0.9%), followed by Paraguay ($77\,850 \text{ km}^2$, 1.3%), and Bolivia ($11\,989 \text{ km}^2$, 0.35%). Alarmingly, our analyses revealed a recent surge in woodland loss, in 2019/20, with the highest woodland loss rate registered since the previous high in 2009/2010 (1.7%). The highest rates of woodland loss occurred in the wet Chaco (3.3%, figures 2 and 3).

Our classifications had high overall accuracies, on average 86.1% (max: 93.9%, min: 77.1%). Average user's and producer's accuracy of the woodland class were also high and ranged between 90.6% and 96.9%, whereas accuracy for the cropland class (73.6%–61.5%) and pasture class (74.1%–81.5%) were somewhat lower (see supplementary material for more detailed information on class-wise accuracies).

Of the total woodlands loss we identified, the dominant initial proximate cause was pasture expansion (47%) followed by cropland expansion (2.5%), while 50.5% were disturbed but did convert to cropland or pasture right after woodland loss had occurred. These patterns varied slightly across countries, as well as for the dry and wet Chaco. In Argentina, pasture expansion was the dominant proximate cause of deforestation (34.4%), whereas only 3.6% were deforested for being immediately used as cropland. An additional $64\,000 \text{ km}^2$ of woodlands were disturbed (62%). In Bolivia and Paraguay, pasture expansion was the dominant proximate cause of deforestation (57.1% and 61.4% of all woodland loss, respectively), whereas cropland expansion (2.0% and 0.7%, respectively) only had a minor importance as a proximate cause. An additional 4908 km^2 (40.9%) and $29\,570 \text{ km}^2$ (37.9%) of woodlands were disturbed in Bolivia and Paraguay, respectively. In the dry Chaco, pasture expansion was the most dominant proximate cause of deforestation (51.8%), followed by cropland expansion (2.9%). Contrary, only 28.0% and 0.4% of woodland loss in the wet Chaco was due to pasture or cropland expansion, respectively (figure 3(B)).

Table 1. Rationale and relevance of the six metrics describing processes in agricultural frontiers.

Metric	Variable	Types and explanation	Rationale and relevance
Frontier timing	Temporal course of the frontier	<ol style="list-style-type: none"> Active (frontiers that are active fronts) Suspended (frontiers that were active but then appear inactive) Emerging (frontiers that are newly appearing) 	<p>Different types of activity require different interventions:</p> <ul style="list-style-type: none"> Active frontiers require urgent stop-gap measures (e.g. strengthening law enforcement, moratoria etc). Suspended frontiers require monitoring and measures of land consolidation and intensification. Emerging frontiers might be targets for various long-term interventions including the development of sustainable production (e.g. certification systems) or community-based natural resource management.
Frontier land use	Post-deforestation land-use trajectories	<ol style="list-style-type: none"> Pasture Cropland Transition (frontier that was dominated by pasture first, but shifted to croplands) 	<p>Different land uses are operated by differently to incentives and interventions. Supply chain interventions need to target the key commodities in a frontier.</p> <ul style="list-style-type: none"> Pasture frontiers may be target for implementing more sustainable production systems (e.g. silvopastures). Transition frontiers may represent focus regions for policies that focus on limiting the further expansion of intensive cropping systems.
Frontier speed	Rate of fastest woodland loss	<ol style="list-style-type: none"> Slow Medium Fast 	<p>The speed with which frontiers progress determines the focus of regional/national policies aiming at conserving remaining woodlands.</p> <ul style="list-style-type: none"> Fast frontiers can be hotspots of policy focus. Slow frontiers might be places to develop longer-term interventions.
Frontier diffusion	Spatial distance to other frontiers	<ol style="list-style-type: none"> Progressing (frontiers, that diffuse through spatial contagion) Leapfrogging (new nexus of frontiers that can then diffuse by contagion) 	<p>How frontiers diffuse represent the group of actors in these areas and require different types of interventions.</p> <ul style="list-style-type: none"> Progressing frontiers might be contained by networks of protected areas and land-use zoning (as in the Brazilian Amazon). Leapfrogging frontiers require an understanding of the mechanisms through which these frontiers diffuse to be governed efficiently (social networks, etc).
Frontier naturalness	Remaining woodland	<ol style="list-style-type: none"> High Medium Low 	<p>The level of remaining woodland cover can influence the balance of priorities between conservation and restoration.</p> <ul style="list-style-type: none"> In high woodland frontiers conservation interventions may be more suitable to avoid tipping points in woody cover below which biodiversity may be lost rapidly. In low woodland frontiers, restoration efforts in degraded areas may be more suitable.
Frontier onset	Year of start of woodland loss	Year	The year of onset represents the timing of frontier dynamics; normally precedes maximum woodland loss.





Figure 2. Agricultural expansion into Chaco woodlands. The map shows the extent of natural vegetation and agriculture in 2020; the two times series (Landsat images and classification) show frontier evolution in the Argentinian Chaco (pink marker, left) and the Paraguayan Chaco (blue marker, right). The white line marks the border between the dry (west) and wet (east) Chaco.

Land cover in 2020 often differed compared to the initial post-deforestation land cover. Across the Chaco, nearly 37% of all woodlands that were not converted into agriculture immediately, (i.e. were classified as disturbed forest) were later converted to pastures (29 635 km²) or cropland (6707 km²), and 17% of all areas initially converted into pastures became cropland later on (15 279 km²). This trend was strongest in Paraguay, where 43.2% of all deforested areas became agriculture by 2020, from which 98.2% became pasture (42.5%), and 1.8% cropland, followed by Bolivia (35.35% of all deforestation, 94.8% of these became pasture and 5.2% cropland) and Argentina (34.1% of all deforestation, of which 70.1% for pasture and 29.9% for cropland). In Argentina, 40.1% of all areas where post-deforestation land use was pastures later became cropland (14 244 km²), whereas in Paraguay (1.3%) and Bolivia (6.0%) this trend was weaker (figure 3).

Our six frontier metrics provided further insight into the dynamics of agricultural expansion in the Chaco, revealing typical frontiers patterns (figures 4 and 5). Most frontier areas were identified as old frontiers, classified as either suspended (48.0%) or active (51.2%), whereas we classified only a minor proportion of the Chaco as emerging frontiers (0.7%, primarily in Paraguay). As highlighted above, only a minor proportion of the frontier areas were classified as cropland frontiers (2.2%, direct conversion

from woodlands to croplands), whereas most frontiers were due to pasture expansion, either directly (80.3%) or with a time lag (e.g. 17.5%, with a time lag of >3 years; figure 4). Most frontiers in the Chaco were characterized as slow (63.2%), with fast (28.2%) and medium frontiers (8.6%) less common. As can be expected, progressing frontiers formed the overwhelming type of frontier expansion (98.8%) compared to leapfrogging frontiers (1.2%; primarily in Argentina and Paraguay). Lastly, remaining woodlands in frontiers were either low (45.6%) or medium (32.8%), whereas in only 21.6% woodlands were high.


The temporal evolution of our frontier metrics varied substantially between the three countries. For example, while Argentina experienced its peak of new frontier areas in 2002, in Paraguay this peak occurred much later (2008) and earlier in Bolivia (1995). In all three countries the peak of new frontier areas was also associated with the largest amount of fast progressing frontiers. Interestingly, with regards to frontier land use Argentina features distinct periods of frontier emergence that is not visible in Bolivia or Paraguay: while pasture was the dominant frontier land use around 2002 (i.e. during the peak of new frontier emergence), we found that transition frontiers occurred earlier (i.e. ~1995–1998) whereas cropland frontiers emerged only during a short period of time (~2005–2007, figure 4).



Integrating our frontier metrics across the Chaco showed that the Chaco is dominated by a set of archetypal frontier types. Out of 162 possible combinations of metric levels, 59.8% all frontier areas fell into ten distinct combinations (hereafter: frontier types, figure 5). Across the entire Chaco, these ten most prevalent frontier types were all characterized as *progressing* frontiers, though with differences regarding frontier timing (39.8% are *active* vs. 19.9% *suspended*) and frontier naturalness (13.5% *high*, 17.8% *medium*, 27.9% *low*). The most common frontier type comprised 17.8% of the study area and was an active pasture frontier that is slowly progressing with medium naturalness. At the country-level we also found distinct differences. For example, while in Bolivia and Paraguay the ten most common frontier types were all *pasture* frontiers, this was

not the case in Argentina, where four out of the ten most common frontier types were *transition* frontiers. Likewise, naturalness in Argentina was generally medium, whereas it was generally high and low in Bolivia and Paraguay, respectively. Contrary, all three countries had a large proportion of active frontiers (Argentina: 55.6% of the total area, Bolivia: 71.8%, Paraguay: 48.9%) that were generally progressing slowly (Argentina 76.3%, Bolivia 82.4%, Paraguay 60.4%; figure 6). With regards to the dry and wet Chaco, we found that while the wet Chaco exclusively was dominated by active pasture frontiers, we found in the dry Chaco a mix of pasture and transition frontiers.


Associating our metrics with qualitatively outlined frontier regions (le Polain de Waroux *et al* 2018) suggested four clear groups of frontier types. Group



I (blue color, figure 6) was characterized as *suspended* frontiers, with *low* naturalness and where frontier land use was either *transition* or *cropland* (i.e. Anta I, Córdoba, San Luís, Bandera, and Chaco-Santiago I). Group II (yellow) was similar to group I, except that the frontier land use was *pasture* (i.e. Tartagal, Chaco-Pantanal, and Central Chaco). Group III (green) were *active* frontiers with either *high* or *medium* frontier naturalness (i.e. Andean Foothills, Anta II, Chaco-Santiago I, Corrientes, Formosa). Lastly, group IV (red) encompassed all frontier regions, where naturalness was already *low*, but which were identified as *active*, independently from the frontier land use (i.e. Santa Cruz, Tucumán, Semi-arid Chaco, figure 6).

4. Discussion

Better understanding how agriculture expands into tropical and subtropical forests is important for addressing the major sustainability challenges associated with frontier expansion. This is particularly urgent for the world’s tropical dry forests, many of which are hotspots of deforestation, carbon emissions, and biodiversity loss. Here, we developed a novel set of frontier metrics, and demonstrated how these can be used to identify typical frontier dynamics. We demonstrate this approach for the entire South American Chaco, highlighting three key insights. First, reconstructing frontier dynamics since



1985 revealed rampant agricultural expansion, with 193 321 km² of Chaco woodlands being converted. Importantly, for the first time we document a recent surge in woodland loss (after 2019). Second, translating our land-cover time series into frontier metrics uncovered distinct frontier processes. For example, whereas ranching expansion drove woodland loss in Paraguay and Bolivia, cropland expansion remained the primary driver of woodland loss since the mid-2000s in Argentina. Similarly, we uncover typical land-cover trajectories following woodland loss, such as initial conversion for pasture and a later shift to cropping, or a considerable fallow period before agriculture is established. Fourth, the multidimensionality of our metrics allowed us to identify groups of frontiers with similar characteristics and development stages that are likely characterized by similar underlying processes and sustainability outcomes. Our metrics hence provide a deeper understanding of frontier processes while allowing to better target land governance policies to sustainably manage frontier regions.

Land-cover change in the Chaco had previously been mapped (Hansen *et al* 2013, Vallejos *et al* 2015, Guyra 2018, Song *et al* 2021, Zalles *et al* 2021), but never with the spatial, temporal and thematic detail that we provide here. Specifically, our mapping goes beyond prior efforts in at least four ways. First, our analysis reconstructs land-cover change back to 1985 at annual resolution, covering the entire history of modern agricultural expansion in this deforestation hotspot. Importantly, we developed an approach that ensures consistent, logical trajectories, avoiding pseudo-change. Second, our analysis, for the first time, separates agricultural expansion from forest disturbances, which constituted a substantial share of the woodland loss in the Chaco (34%, figure 3). Third, because our assessment was validated, we were able to derive the first robust area estimates of frontier dynamics in the Chaco. Fourth, our approach is novel in disentangling post-deforestation land-use transitions, including multiple, subsequent land-cover transitions. This revealed, for example, that deforested areas in Argentina are often eventually used




Figure 6. The ten most common combinations of the frontier metric levels (i.e. frontier types), and their relative share on all frontier areas. Results are presented for the entire Chaco and separated by the three countries as well as for the dry and wet Chaco. For a complete list of frontier types and their area share please refer to table S1 in the supplementary material.

for cropping, although initial deforestation occurs for ranching. It is important to highlight that our land-cover reconstruction is solely based on satellite imagery, allowing for subsequent analyses (e.g. statistical analyzing of drivers of change). Likewise, our approach is easily transferable, can be scaled up to even larger regions, and can be updated as satellite image archives grow. This, we humbly suggest, constitutes a step-change in our ability to monitor land-cover change and forms the basis for a deeper process-understanding of frontiers.

The patterns and trends of agricultural expansion we derived here are highly plausible. For example, our results suggest that frontiers expanded particularly rapidly in the 2000s in Argentina but slowed down after 2010. The agricultural expansion boom in the 2000s was the result of several factors, most importantly the currency devaluation in 2001, which strongly increased profits from soy exports (Gasparri

and Baldi 2013) and the introduction of genetically modified soybean in the Chaco (Reenberg and Fenger 2011, le Polain de Waroux 2019). Indeed, most of the cropland frontiers emerged during that time (figure 4). Later, increasing taxation, economic instability, an outflow of capital (le Polain de Waroux *et al* 2019), increasing land-use restrictions through Argentina's zoning law (Marinaro *et al* 2020), and the increasingly more marginal conditions for sites on which remaining forests are found (Houspanossian *et al* 2016) lowered cropland expansion rates after 2010. In contrast, capital that accumulated in the soybean boom (in the Chaco or elsewhere, such as Brazil), combined with evolving know-how and infrastructure to optimize cattle ranching in the Chaco (le Polain de Waroux 2019) explains surging woodland conversion we found in the Paraguayan Chaco after 2010. As a final example, the recent, more than two-fold surge in deforestation after 2019

(figure 3(A)) that we here document for the first time converges well with reports of increasing forest conversion, during the lockdown situation—in the Chaco and other deforestation frontiers globally (Fair 2020, Price 2020).


A major surprise in our findings was that most converted woodlands did not transition to agriculture right away, and many never. Four complementary explanations for this finding are plausible. First, natural disturbances, such as from fires or river-bed migrations are common in the Chaco (Adamoli *et al* 1990, Bravo *et al* 2001, de Marzo *et al* 2021). However, disturbance attribution is not always straightforward. For example, fires occur naturally, are used as a management tool to control woody encroachment, or are associated with the deforestation process (Boletta *et al* 2006). Second, woodland conversion may not be driven by the goal to immediately produce agricultural commodities, but might happen to secure land, to prepare land for resale, or simply in fear of tightening regulations (Seghezze *et al* 2011). Third, given that removing woodland and preparing land for agriculture requires capital (e.g. sowing with productive pasture grasses), there may be a time lag between deforestation and agricultural use, which we found for 34% of all woodlands converted to agriculture (figure 3(B)). Finally, silvopastoral systems, where parts of the tree canopy remain, are becoming more common in Argentina (Baldassini *et al* 2018, Fernández *et al* 2020), and these areas would fall outside of our pasture class. All of these factors point towards the importance of adopting approaches that deliver information at high spatial, temporal, and, importantly, thematic resolutions to quantify agricultural expansion in the tropics and to understand the causes and mechanisms of deforestation. This, in turn, is critical for properly attributing environmental trade-offs properly to commodities, which is a key research frontier for achieving supply chain sustainability (Gardner *et al* 2019, Pendrill *et al* 2019, Zu Ermgassen *et al* 2020).

Translating our land-cover time series into a consistent set of frontier metrics, allowed us to move beyond land cover to characterizing land-use change processes. In our case, this enabled us to identify distinct frontier types, characterized by similar land-use and woodland loss dynamics in space and time. Such archetypical, high-level patterns and outcomes of human–environment interactions can help to structure complexity in land-use change (Vaclavik *et al* 2013, Levers *et al* 2018, Pacheco-Romero *et al* 2021), foster a more mechanistic understanding of land-use change (Magliocca *et al* 2018), and contribute to developing theories of the middle range (Meyfroidt *et al* 2018). Importantly though, identifying archetypes, such as recurring frontier types, allows for the more context-specific, regionally-targeted land governance increasingly asked for (Kuemmerle *et al* 2016, Thomson *et al* 2019, Christie *et al* 2020, Pacheco

et al 2021). For example, *cropland* or *transition* frontiers with *low remaining* naturalness (i.e. group I (blue), figure 7) are regions where restoration efforts in degraded lands are most suitable. Likewise, *pasture* frontiers that are *suspended* (i.e. group II (yellow)) may increasingly experience pasture to cropland conversions in the future, and hence actor-focused interventions, such as building incentives towards more sustainable production systems (e.g. silvopastures), may be most effective. Contrary, *active* and *fast* frontiers with *high* or *medium* naturalness (e.g. group III (green)) should become hotspots of policies with the goal to avoid tipping points in woody cover, for example through the identification of biodiversity-rich areas and their subsequent protection through new protected areas or zonation (e.g. Ley de Protección Ambiental de Bosques 2007) to restrict further agricultural expansion.

Our analyses provide the most detailed reconstruction of woodland and agricultural dynamics for the Chaco, including novel insights into how agricultural frontiers have expanded. A few limitations still need to be mentioned. First, we only mapped agricultural expansion and intensification, but not agricultural abandonment. Abandonment is not (yet) a widespread process in the Chaco and vegetation recovery on abandoned fields takes time (Basualdo *et al* 2019). Still, adding de-intensification and abandonment processes would be a useful expansion of our approach in future work. Second, we describe frontier expansion related to intensified, large-scale agriculture but did not explicitly address forest smallholders practicing subsistence agriculture inside forests. While these actors are important in the Chaco, dynamics in forest smallholders mainly are due to agribusiness expansion (Levers *et al* 2021), and so are indirectly captured here. Third, our classification contains remaining uncertainty. For example, some class confusions occur between natural vegetation after woodland loss and pastures, which might represent silvopastures. Likewise, some accuracies for the cropland and pasture classes were relatively low (~70%, figure SI-3) for some years, likely due to missing observations. Yet, changes not captured in that year (e.g. cropland expansion) were likely captured in subsequent years, so that the overall trajectory of land-cover changes should remain unaffected.

Agricultural expansion into tropical and subtropical forests contributes heavily to many global sustainability challenges. Steering these frontiers towards more sustainable outcomes requires a better understanding of the dynamics of frontier processes. Here, we developed and demonstrated a novel approach to generate such understanding on the basis of frontier metrics derived from freely available, high-resolution satellite imagery. For the Chaco, our frontier metrics characterize and structure the complexity of frontier dynamics, for example revealing slow vs. rampant frontiers, where frontiers are emerging, or



when frontiers were particularly active. This allows for exploring the underlying drivers of these frontier processes, including testing hypotheses about causal mechanisms. For example, the emergence of cropland frontiers could be associated with the introduction of genetically modified soybeans, and through our metrics one could explore the causal links between the two. Further, our study revealed that about 34% of the deforestation in the Chaco that might be wrongfully attributed to commodity agriculture, and another 17% that might be attributed to the wrong commodity depending on which baseline is chosen, providing important insights for attributing environmental trade-offs properly to commodities. Our transferable, repeatable, scalable, and extendable approach allows for comparative research across regions to find rules

governing frontiers in many situations, as well as to identify generalizable patterns and processes that shape frontiers in different regions. In the Chaco and elsewhere this can enable cross-regional learning and the more regionally targeted, context-specific policy-interventions that are often asked for. More broadly, our study highlights the opportunities of the big data era of remote sensing for creating a step change in our understanding of land-use change and for uncovering patterns of human–environment interactions at the ecoregional and national scale.

Data availability statement

The data that support the findings of this study are available upon reasonable request from the authors.

Acknowledgments

We gratefully acknowledge support by the German Research Foundation (DFG, Project KUIJ 2458/5-1), the German Federal Ministry of Education and Science (BMBF, Project PASANOVA, 031B0034A), and the Belgian Science Policy Office Research Program for Earth Observation (belspo-STEREO-III, Project REFORCHA, SR/00/338). This work was supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (Grant Agreement No. 101001239 SYSTEMSHIFT). This research contributes to the Global Land Program (glp.earth). We are grateful for the very helpful, constructive and encouraging comments made by two reviewers and the handling editor.

ORCID iDs

Matthias Baumann  <https://orcid.org/0000-0003-2375-3622>
 Ignacio Gasparri  <https://orcid.org/0000-0001-8389-1379>
 Ana Buchadas  <https://orcid.org/0000-0003-4219-108X>
 Julian Oeser  <https://orcid.org/0000-0003-3216-6817>
 Patrick Meyfroidt  <https://orcid.org/0000-0002-1047-9794>
 Christian Levers  <https://orcid.org/0000-0003-4810-9024>
 Alfredo Romero-Muñoz  <https://orcid.org/0000-0002-7905-6087>
 Yann le Polain de Waroux  <https://orcid.org/0000-0001-9887-7270>
 Daniel Müller  <https://orcid.org/0000-0001-8988-0718>
 Tobias Kuemmerle  <https://orcid.org/0000-0002-9775-142X>

References

- Adamoli J, Sennhauser E, Acero J M and Rescia A 1990 Stress and disturbance—vegetation dynamics in the dry Chaco region of Argentina *J. Biogeogr.* **17** 491–500
- Andersson K and Agrawal A 2011 Inequalities, institutions, and forest commons *Glob. Environ. Change* **21** 866–75
- Austin K G, González-Roglich M, Schaffer-Smith D, Schwantes A M and Swenson J J 2017 Trends in size of tropical deforestation events signal increasing dominance of industrial-scale drivers *Environ. Res. Lett.* **12** 054009
- Baldassini P, Despósito C, Piñeiro G and Paruelo J M 2018 Silvopastoral systems of the Chaco forests: effects of trees on grass growth *J. Arid Environ.* **156** 87–95
- Barbier E B 2012 Scarcity, frontiers and development *Geogr. J.* **178** 110–22
- Basualdo M, Huykman N, Volante J N, Paruelo J M and Piñeiro G 2019 Lost forever? Ecosystem functional changes occurring after agricultural abandonment and forest recovery in the semiarid Chaco forests *Sci. Total Environ.* **650** 1537–46
- Baumann M, Gasparri I, Piquer-Rodríguez M, Gavier Pizarro G, Griffiths P, Hostert P and Kuemmerle T 2017 Carbon emissions from agricultural expansion and intensification in the Chaco *Glob. Change Biol.* **23** 1902–16
- Baynes J, Herbohn J, Smith C, Fisher R and Bray D 2015 Key factors which influence the success of community forestry in developing countries *Glob. Environ. Change* **35** 226–38
- Boletta P E, Ravelo A C, Planchuelo A M and Grilli M 2006 Assessing deforestation in the Argentine Chaco *For. Ecol. Manage.* **228** 108–14
- Bonilla-Moheno M and Aide T M 2020 Beyond deforestation: land cover transitions in Mexico *Agric. Syst.* **178** 102734
- Bowman M S, Soares-Filho B S, Merry F D, Nepstad D C, Rodrigues H and Almeida O T 2012 Persistence of cattle ranching in the Brazilian Amazon: a spatial analysis of the rationale for beef production *Land Use Policy* **29** 558–68
- Bravo S, Kunst C, Gimenez A and Moglia G 2001 Fire regime of a *Elionorus muticus* Spreng. savanna, western Chaco region, Argentina *Int. J. Wildland Fire* **10** 65
- Buchadas A, Baumann M, Meyfroidt P and Kuemmerle T 2022 Uncovering major types of deforestation frontiers across the world's tropical dry woodlands *Nat. Sustain.* **5** 619–27
- Bucher E H 1982 Chaco and Caatinga—South American arid savannas, woodlands and thickets *Ecology of Tropical Savannas* ed B J Huntley and B H Walker (Berlin: Springer) pp 48–79
- Bucher E H and Huszar P C 1999 Sustainable management of the Gran Chaco of South America *J. Environ. Manage.* **57** 99–108
- Caldas M M, Goodin D, Sherwood S, Campos Krauer J M and Wisely S M 2015 Land-cover change in the Paraguayan Chaco *J. Land Use Sci.* **10** 1–18
- Campos-Krauer J M and Wisely S M 2011 Deforestation and cattle ranching drive rapid range expansion of capybara in the Gran Chaco ecosystem *Glob. Change Biol.* **17** 206–18
- Carlson K M et al 2017 Greenhouse gas emissions intensity of global croplands *Nat. Clim. Change* **7** 63–68
- Chaplin-Kramer R et al 2015 Spatial patterns of agricultural expansion determine impacts on biodiversity and carbon storage *Proc. Natl Acad. Sci.* **112** 7402–7
- Christie A P, Amano T, Martin P A, Petrovan S O, Shackelford G E, Simmons B I, Smith R K, Williams D R, Wordley C F R and Sutherland W J 2020 Poor availability of context-specific evidence hampers decision-making in conservation *Biol. Conserv.* **248** 108666
- Cumming G S, Buerkert A, Hoffmann E M, Schlecht E, von Cramon-taubadel S and Tschardt T 2014 Implications of agricultural transitions and urbanization for ecosystem services *Nature* **515** 50–57
- Curtis P G, Slay C M, Harris N L, Tyukavina A and Hansen M C 2018 Classifying drivers of global forest loss *Science* **361** 1108–11
- de Marzo T, Pflugmacher D, Baumann M, Lambin E F, Gasparri I and Kuemmerle T 2021 Characterizing forest disturbances across the Argentine Dry Chaco based on Landsat time series *Int. J. Appl. Earth Obs. Geoinf.* **98** 102310
- De Sy V, Herold M, Achard F, Avitabile V, Baccini A, Carter S, Clevers J G P W, Lindquist E, Pereira M and Verchot L 2019 Tropical deforestation drivers and associated carbon emission factors derived from remote sensing data *Environ. Res. Lett.* **14** 094022
- Eisenack K, Lüdeke M and Kropp J 2006 Construction of archetypes as a formal method to analyze social-ecological systems, *IDGEC Synthesis Conf. Institutional Dimensions of Global Environmental Change (Bali, Indonesia, 6–9 December 2006)* (International Human Dimensions Programme of Global Environmental Change)
- Eisenack K, Villamayor-Tomas S, Epstein G, Kimmich C, Magliocca N, Manuel-Navarrete D, Oberlack C, Roggero M and Sietz D 2019 Design and quality criteria for archetype analysis *Ecol. Soc.* **24** 6
- Fair J 2020 COVID-19 lockdown precipitates deforestation across Asia and South America (available at: <https://news.mongabay.com/2020/07/covid-19-lockdown-precipitates-deforestation-across-asia-and-south-america/>)

- Fatecha A 1989 Present and potential area for agricultural use in the arid Chaco of Paraguay *Agricultural Production under Semi-arid Conditions with Special Reference to the Paraguayan Chaco: Strategies and Appropriate Technologies* ed M Hamp and M A Tiefert (Feldafing: Deutsche Stiftung für Internationale Entwicklung (DSE)) pp 26–49
- Fehlenberg V, Baumann M, Gasparri N I, Piquer-Rodríguez M, Gavier-Pizarro G and Kuemmerle T 2017 The role of soybean production as an underlying driver of deforestation in the South American Chaco *Glob. Environ. Change* **45** 24–34
- Fernández P D, Kuemmerle T, Baumann M, Grau H R, Nasca J A, Radrizzani A and Gasparri N I 2020 Understanding the distribution of cattle production systems in the South American Chaco *J. Land Use Sci.* **17** 1–17
- Friedl M A, Sulla-Menashe D, Tan B, Schneider A, Ramankutty N, Sibley A and Huang X 2010 MODIS Collection 5 global land cover: algorithm refinements and characterization of new datasets *Remote Sens. Environ.* **114** 168–82
- Gardner T A et al 2019 Transparency and sustainability in global commodity supply chains *World Dev.* **121** 163–77
- Garrett R D, Lambin E F and Naylor R L 2013 The new economic geography of land use change: supply chain configurations and land use in the Brazilian Amazon *Land Use Policy* **34** 265–75
- Gasparri N I and Baldi G 2013 Regional patterns and controls of biomass in semiarid woodlands *Reg. Environ. Change* **13** 1131–44
- Gasparri N I and Grau H R 2009 Deforestation and fragmentation of Chaco dry forest in NW Argentina (1972–2007) *For. Ecol. Manage.* **258** 913–21
- Geist H J and Lambin E F 2004 Dynamic causal patterns of desertification *BioScience* **54** 817–29
- Gibbs H K, Ruesch A S, Achard F, Clayton M K, Holmgren P, Ramankutty N and Foley J A 2010 Tropical forests were the primary sources of new agricultural land in the 1980s and 1990s *Proc. Natl Acad. Sci.* **107** 16732–7
- Godar J, Gardner T A, Tizado E J and Pacheco P 2014 Actor-specific contributions to the deforestation slowdown in the Brazilian Amazon *Proc. Natl Acad. Sci.* **111** 15591–6
- Godfray H C J, Beddington J R, Crute I R, Haddad L, Lawrence D, Muir J F, Pretty J, Robinson S, Thomas S M and Toulmin C 2010 Food security *Science* **327** 812–8
- Gómez C, White J C and Wulder M A 2016 Optical remotely sensed time series data for land cover classification *ISPRS J. Photogramm. Remote Sens.* **116** 55–72
- Gorelick N, Hancher M, Dixon M, Ilyushchenko S, Thau D and Moore R 2017 Google earth engine *Remote Sens. Environ.* **202** 18–27
- Grace J, San José J, Meir P, Miranda H S and Montes R A 2006 Productivity and carbon fluxes of tropical savannas *J. Biogeogr.* **33** 387–400
- Griffiths P, Jakimow B and Hostert P 2018 Reconstructing long term annual deforestation dynamics in Pará and Mato Grosso using the Landsat archive *Remote Sens. Environ.* **216** 497–513
- Guyra P 2018 Informe deforestación (available at: <http://guyra.org.py/informe-deforestacion/M4-Citavi>)
- Hansen A J et al 2020 A policy-driven framework for conserving the best of Earth's remaining moist tropical forests *Nat. Ecol. Evol.* **4** 1377–84
- Hansen M C et al 2013 High-resolution global maps of 21st-century forest cover change *Science* **342** 850–3
- Harris N L et al 2017 Using spatial statistics to identify emerging hot spots of forest loss *Environ. Res. Lett.* **12** 024012
- Hosonuma N, Herold M, Sy V, Fries R S, Brockhaus M, Verchot L, Angelsen A and Romijn E 2012 An assessment of deforestation and forest degradation drivers in developing countries *Environ. Res. Lett.* **7** 044009
- Houspanossian J, Giménez R, Baldi G and Noretto M 2016 Is aridity restricting deforestation and land uses in the South American Dry Chaco? *J. Land Use Sci.* **11** 369–83
- Killeen T J, Calderon V, Soria L, Quezada B, Steininger M K, Harper G, Solórzano L A and Tucker C J 2007 Thirty years of land-cover change in Bolivia *Ambio* **36** 600–6
- Klink C A and Machado R B 2005 Conservation of the Brazilian Cerrado *Conserv. Biol.* **19** 707–13
- Kröger M and Nygren A 2020 Shifting frontier dynamics in Latin America *J. Agrar. Change* **424** 3
- Kuemmerle T et al 2016 Hotspots of land use change in Europe *Environ. Res. Lett.* **11** 064020
- Lambin E F, Geist H J and Lepers E 2003 Dynamics of land-use and land-cover change in tropical regions *Annu. Rev. Environ. Resour.* **28** 205–41
- Lambin E F, Gibbs H K, Ferreira L, Grau R, Mayaux P, Meyfroidt P, Morton D C, Rudel T K, Gasparri I and Munger J 2013 Estimating the world's potentially available cropland using a bottom-up approach *Glob. Environ. Change* **23** 892–901
- Latawiec A E, Strassburg B B N, Brancalion P H S, Rodrigues R R and Gardner T 2015 Creating space for large-scale restoration in tropical agricultural landscapes *Front. Ecol. Environ.* **13** 211–8
- le Polain de Waroux Y 2019 Capital has no homeland *Geoforum* **105** 131–44
- le Polain de Waroux Y, Baumann M, Gasparri N I, Gavier-Pizarro G, Godar J, Kuemmerle T, Müller R, Vázquez F, Volante J N and Meyfroidt P 2018 Rents, actors, and the expansion of commodity frontiers in the Gran Chaco *Ann. Am. Assoc. Geogr.* **108** 204–25
- le Polain de Waroux Y, Garrett R D, Graesser J, Nolte C, White C and Lambin E F 2019 The restructuring of South American soy and beef production and trade under changing environmental regulations *World Dev.* **121** 188–202
- Leguizamón A 2014 Modifying Argentina *Geoforum* **53** 149–60
- Lerner A M, Rudel T K, Schneider L C, McGroddy M, Burbano D V and Mena C F 2015 The spontaneous emergence of silvo-pastoral landscapes in the Ecuadorian Amazon: patterns and processes *Reg. Environ. Change* **15** 1421–31
- Levers C et al 2018 Archetypical patterns and trajectories of land systems in Europe *Reg. Environ. Change* **18** 715–32
- Levers C et al 2021 Agricultural expansion and the ecological marginalization of forest-dependent people *Proc. Natl Acad. Sci. USA* **118** e2100436118
- Ley de Protección Ambiental de Bosques N 2007 *Ley No 26331, primera sección* (Argentina) p 3 (available at: www.infoleg.gov.ar/infolegInternet/anexos/135000-139999/136125/norma.htm)
- Liu H, Gong P, Wang J, Clinton N, Bai Y and Liang S 2020 Annual dynamics of global land cover and its long-term changes from 1982 to 2015 *Earth Syst. Sci. Data* **12** 1217–43
- Magliocca N R, Ellis E C, Allington G R H, de Bremond A, Dell'Angelo J, Mertz O, Messerli P, Meyfroidt P, Seppelt R and Verburg P H 2018 Closing global knowledge gaps: producing generalized knowledge from case studies of social-ecological systems *Glob. Environ. Change* **50** 1–14
- Marinaro S, Gasparri N I and Piriz-Carrillo V 2020 Private-land control and deforestation dynamics in the context of implementing the Native Forest Law in the Northern Argentinian Dry Chaco *Environ. Conserv.* **47** 1–7
- Mayle F E, Langstroth R P, Fisher R A and Meir P 2007 Long-term forest-savannah dynamics in the Bolivian Amazon *Phil. Trans. R. Soc. B* **362** 291
- Meyfroidt P et al 2018 Middle-range theories of land system change *Glob. Environ. Change* **53** 52–67
- Miles L, Newton A C, DeFries R S, Ravilious C, May I, Blyth S, Kapos V and Gordon J E 2006 A global overview of the conservation status of tropical dry forests *J. Biogeogr.* **33** 491–505
- Müller H, Griffiths P and Hostert P 2016 Long-term deforestation dynamics in the Brazilian Amazon—uncovering historic frontier development along the Cuiabá-Santarém highway *Int. J. Appl. Earth Obs. Geoinf.* **44** 61–69
- Murphy P G and Lugo A E 1986 Ecology of tropical dry forest *Annu. Rev. Ecol. Syst.* **17** 67–88

- Oberlack C et al 2019 Archetype analysis in sustainability research: meanings, motivations, and evidence-based policy making *Ecol. Soc.* **24** 26
- Oeser J, Heurich M, Senf C, Pflugmacher D, Belotti E, Kuemmerle T, Pettorelli N and Armenteras D 2020 Habitat metrics based on multi-temporal Landsat imagery for mapping large mammal habitat *Remote Sens. Ecol. Conserv.* **6** 52–69
- Oldekop J A et al 2020 Forest-linked livelihoods in a globalized world *Nat. Plants* **6** 1400–7
- Olofsson P, Foody G M, Herold M, Stehman S V, Woodcock C E and Wulder M A 2014 Good practices for estimating area and assessing accuracy of land change *Remote Sens. Environ.* **148** 42–57
- Pacheco P 2012 Actor and frontier types in the Brazilian Amazon *Geoforum* **43** 864–74
- Pacheco P, Mo K, Dudley N, Shapiro A, Aguilar-Amuchastegui N, Ling P Y, Anderson C and Marx A 2021 *Deforestation Frontiers: Drivers and Responses in a Changing World* (Gland: WWF) p 125
- Pacheco-Romero M, Kuemmerle T, Levers C, Alcaraz-Segura D and Cabello J 2021 Integrating inductive and deductive analysis to identify and characterize archetypical social-ecological systems and their changes *Landsc. Urban Plan.* **215** 104199
- Pendrill F, Persson U M, Godar J, Kastner T, Moran D, Schmidt S and Wood R 2019 Agricultural and forestry trade drives large share of tropical deforestation emissions *Glob. Environ. Change* **56** 1–10
- Pennington R T, Lehmann C E R and Rowland L M 2018 Tropical savannas and dry forests *Curr. Biol.* **28** R541–45
- Phiri D, Morgenroth J and Xu C 2019 Long-term land cover change in Zambia: an assessment of driving factors *Sci. Total Environ.* **697** 134206
- Potapov P et al 2019 Annual continuous fields of woody vegetation structure in the Lower Mekong region from 2000–2017 Landsat time-series *Remote Sens. Environ.* **232** 111278
- Price K 2020 Poaching, deforestation reportedly on the rise since COVID-19 lockdowns (available at: www.conservation.org/blog/poaching-deforestation-reportedly-on-the-rise-since-covid-19-lockdowns)
- Ramankutty N, Foley J A and Olejniczak N J 2002 People on the land *Ambio* **31** 251–7
- Reenberg A and Fenger N A 2011 Globalizing land use transitions *Geogr. Tidsskr.* **111** 85–92
- Richards P 2018 It's not just where you farm; it's whether your neighbor does too. How agglomeration economies are shaping new agricultural landscapes *J. Econ. Geogr.* **18** 87–110
- Rocha J, Malmborg K, Gordon L, Brauman K and DeClerck F 2020 Mapping social-ecological systems archetypes *Environ. Res. Lett.* **15** 034017
- Rodrigues A S L, Ewers R M, Parry L, Souza C, Verissimo A and Balmford A 2009 Boom-and-bust development patterns across the Amazon deforestation *Frontier* **324** 1435–7
- Rudel T K 2007 Changing agents of deforestation *Land Use Policy* **24** 35–41
- Seghezzo L, Volante J N, Paruelo J M, Somma D J, Buliubasich E C, Rodríguez H E, Gagnon S and Hufty M 2011 Native forests and agriculture in Salta (Argentina) *J. Environ. Dev.* **20** 251–77
- Sietz D, Frey U, Roggero M, Gong Y, Magliocca N, Tan R, Janssen P and Václavík T 2019 Archetype analysis in sustainability research: methodological portfolio and analytical frontiers *Ecol. Soc.* **24** 34
- Song X-P et al 2021 Massive soybean expansion in South America since 2000 and implications for conservation *Nat. Sustain.* **4** 784–92
- Souza C M et al 2020 Reconstructing three decades of land use and land cover changes in Brazilian biomes with Landsat archive and earth engine *Remote Sens.* **12** 2735
- Strassburg B B N et al 2017 Moment of truth for the Cerrado hotspot *Nat. Ecol. Evol.* **1** 0099
- Sulla-Menashe D, Gray J M, Abercrombie S P and Friedl M A 2019 Hierarchical mapping of annual global land cover 2001 to present: the MODIS Collection 6 Land Cover product *Remote Sens. Environ.* **222** 183–94
- Thomson A M, Ellis E C, Grau H R, Kuemmerle T, Meyfroidt P, Ramankutty N and Zeleke G 2019 Sustainable intensification in land systems *Curr. Opin. Environ. Sustain.* **38** 37–43
- Turner B L, Meyfroidt P, Kuemmerle T, Müller D and Roy Chowdhury R 2020 Framing the search for a theory of land use *J. Land Use Sci.* **15** 489–508
- Turubanova S, Potapov P V, Tyukavina A and Hansen M C 2018 Ongoing primary forest loss in Brazil, Democratic Republic of the Congo, and Indonesia *Environ. Res. Lett.* **13** 074028
- Tyukavina A, Hansen M C, Potapov P, Parker D, Okpa C, Stehman S V, Kommareddy I and Turubanova S 2018 Congo Basin forest loss dominated by increasing smallholder clearing *Sci. Adv.* **4** eaat2993
- Vaclavik T, Lautenbach S, Kuemmerle T and Seppelt R 2013 Mapping global land system archetypes *Glob. Environ. Change* **23** 1637–47
- Vallejos M, Volante J N, Mosciaro M J, Vale L M, Bustamante M L and Paruelo J M 2015 Transformation dynamics of the natural cover in the Dry Chaco ecoregion *J. Arid Environ.* **123** 3–11
- Vancutsem C, Achard F, Pekel J-F, Vieilledent G, Carboni S, Simonetti D, Gallego J, Aragão L E O C and Nasi R 2021 Long-term (1990–2019) monitoring of forest cover changes in the humid tropics *Sci. Adv.* **7** eab1603
- Vazquez F 2013 Geografía humana del Chaco paraguayo—transformaciones territoriales y desarrollo regional (Asunción: Ediciones ADEPO)
- Vieilledent G, Grinand C, Rakotomalala F A, Ranaiivosoa R, Rakotoarijaona J-R, Allnut T F and Achard F 2018 Combining global tree cover loss data with historical national forest cover maps to look at six decades of deforestation and forest fragmentation in Madagascar *Biol. Conserv.* **222** 189–97
- Volante J N, Alcaraz-Segura D, Mosciaro M J, Viglizzo E F and Paruelo J M 2012 Ecosystem functional changes associated with land clearing in NW Argentina *Agric. Ecosyst. Environ.* **154** 12–22
- Woodcock C E, Loveland T R, Herold M and Bauer M E 2020 Transitioning from change detection to monitoring with remote sensing: a paradigm shift *Remote Sens. Environ.* **238** 111558
- Wulder M A et al 2019 Current status of Landsat program, science, and applications *Remote Sens. Environ.* **225** 127–47
- Zalles V et al 2019 Near doubling of Brazil's intensive row crop area since 2000 *Proc. Natl Acad. Sci. USA* **116** 428–35
- Zalles V et al 2021 Rapid expansion of human impact on natural land in South America since 1985 *Sci. Adv.* **7** eabg1620
- Zu Ermgassen E K H J, Godar J, Lathuilière M J, Löfgren P, Gardner T, Vasconcelos A and Meyfroidt P 2020 The origin, supply chain, and deforestation risk of Brazil's beef exports *Proc. Natl Acad. Sci.* **117** 31770–9