

Maki, Daiki; Ota, Yasushi

Article

Robust tests for ARCH in the presence of a misspecified conditional mean: A comparison of nonparametric approaches

Cogent Economics & Finance

Provided in Cooperation with:

Taylor & Francis Group

Suggested Citation: Maki, Daiki; Ota, Yasushi (2021) : Robust tests for ARCH in the presence of a misspecified conditional mean: A comparison of nonparametric approaches, Cogent Economics & Finance, ISSN 2332-2039, Taylor & Francis, Abingdon, Vol. 9, Iss. 1, pp. 1-18, <https://doi.org/10.1080/23322039.2020.1862445>

This Version is available at:

<https://hdl.handle.net/10419/270029>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.



<https://creativecommons.org/licenses/by/4.0/>

Robust tests for ARCH in the presence of a misspecified conditional mean: A comparison of nonparametric approaches

Daiki Maki & Yasushi Ota |

To cite this article: Daiki Maki & Yasushi Ota | (2021) Robust tests for ARCH in the presence of a misspecified conditional mean: A comparison of nonparametric approaches, Cogent Economics & Finance, 9:1, 1862445, DOI: [10.1080/23322039.2020.1862445](https://doi.org/10.1080/23322039.2020.1862445)

To link to this article: <https://doi.org/10.1080/23322039.2020.1862445>



© 2021 The Author(s). This open access article is distributed under a Creative Commons Attribution (CC-BY) 4.0 license.



Published online: 06 Jan 2021.



Submit your article to this journal [↗](#)



Article views: 1455



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 1 View citing articles [↗](#)



Received: 17 June 2020
Accepted: 04 December 2020

*Corresponding author: Daiki Maki,
Doshisha University, Kyoto 602-8580,
Japan
E-mail: dmaki@mail.doshisha.ac.jp

Reviewing editor:
Xibin Zhang, Econometrics and
Business Statistics, Monash
University, Caulfield East, Australia

Additional information is available at
the end of the article

GENERAL & APPLIED ECONOMICS | RESEARCH ARTICLE

Robust tests for ARCH in the presence of a misspecified conditional mean: A comparison of nonparametric approaches

Daiki Maki^{1*} and Yasushi Ota²

Abstract: This study compares the size and power of autoregressive conditional heteroskedasticity (ARCH) tests that are robust to the presence of a misspecified conditional mean. The approaches employed are based on two nonparametric regressions for the conditional mean: an ARCH test with a Nadaraya-Watson kernel regression and an ARCH test using a polynomial approximation regression. The two approaches do not require the specification of a conditional mean and can adapt to various nonlinear models, which are unknown a priori. The results reveal that the ARCH tests are robust to the misspecified conditional mean models. The simulation results show that the ARCH tests based on the polynomial approximation regression approach have better properties of the size and power than those using the Nadaraya-Watson kernel regression approach for various nonlinear models.

Subjects: Econometrics; Financial Econometrics

Keywords: ARCH test; nonparametric regression; misspecified models; size and power
Jel classification: C12; C14; C58

ABOUT THE AUTHORS

Daiki Maki is a professor of Econometrics at the Faculty of Commerce, Doshisha University. He is an expert at time series analysis. His research interests cover nonlinear time series analysis, nonstationary time series analysis, and financial econometrics.

Yasushi Ota is an associate professor of Finance at the Faculty of Management, Okayama University of Science. His field of expertise is financial theory. His research interests cover option pricing and mathematical finance.

PUBLIC INTEREST STATEMENT

The presence of heteroskedasticity significantly impacts estimations and inferences in a time series analysis. Autoregressive conditional heteroskedasticity (ARCH) model is a simple and popular volatility model and continues to be widely used in the literature. Many studies evidence the overrejection of no ARCH effects when the nonlinearity of the conditional mean regression model is neglected. To appropriately test for ARCH, it is necessary to avoid the misspecified model of the conditional mean. This study compares the size and power of ARCH tests that do not depend on the conditional mean model. The tests are applicable to various nonlinear conditional mean models and are robust to the misspecified conditional mean model. The Monte Carlo simulation results evidence that ARCH tests that are based on the polynomial approximation regression approach have better size and power properties than those using the Nadaraya-Watson kernel regression approach when DGPs are various nonlinear models.

1. Introduction

The presence of heteroskedasticity significantly impacts estimations and inferences in a time series analysis. Becker and Hurn (2009) and Pavlidis et al. (2010), for example, demonstrate that the presence of heteroskedasticity frequently leads to over-rejections of the null hypothesis when testing the null for the linearity of a conditional mean model against the alternative hypothesis of nonlinear time series models. Pavlidis et al. (2013) show that causality tests on the conditional mean demonstrate spurious causality relationships in the presence of multivariate heteroskedasticity. These facts indicate that tests for heteroskedasticity in data-generating processes (DGP) play an important role in time-series analyses.

The most representative model for heteroskedasticity is Engle's (1982) autoregressive conditional heteroskedasticity (ARCH) model. ARCH is a simple and popular volatility model and continues to be widely used in the literature. When testing for heteroskedasticity, a regression model for the assumed conditional mean is first estimated. Next, ARCH is examined to use statistics such as the Lagrange multiplier (LM). If the conditional mean regression model is correctly specified, the ARCH test performs well. However, a misspecified conditional mean severely impedes the ARCH tests. Lumsdaine and Ng (1999) examine the properties of ARCH tests under a misspecified conditional mean. They show that the misspecification of the conditional mean over-rejects the null hypothesis for homoskedasticity. Similarly, Balke and Kapetanios (2007) clarify the influence of the neglected nonlinearity of the conditional mean on ARCH tests. Their analysis evidences the over-rejection of no ARCH effects when the nonlinearity of the conditional mean regression model is neglected. Erroneous ARCH tests based on misspecified conditional mean models and the failure to obtain sufficient reliability for the derived results increasingly impede model constructions and statistical evaluation. To appropriately test for ARCH, it is necessary to avoid the misspecified model of the conditional mean. Thus, it is important to investigate which ARCH tests are robust to the misspecification of the conditional mean.

This study clarifies which ARCH tests that do not depend on the conditional mean model have better size and power. To this end, we first introduce ARCH tests using nonparametric regression approaches to avoid the misspecification of the conditional mean and next investigate the size and power of the introduced tests in various linear and nonlinear models with/without ARCH effects. The tests are applicable to various nonlinear conditional mean models and are robust to the misspecified conditional mean model. We employ two nonparametric approaches to avoid the misspecification of the conditional mean model. First is a regression using the Nadaraya-Watson kernel estimator, which is a representative nonparametric method. Nadaraya (1964) and Watson (1964) propose the method using a kernel density function in a regression analysis that does not depend on the model. McMillan (2001) and Exterkate et al. (2016) show that the Nadaraya-Watson estimator is useful under various nonlinear models. Second is the regression analysis using a polynomial approximation of a general unknown nonlinear model. Stone (1977) and Katkovnik (1979) propose the local polynomial estimator on the basis of a polynomial approximation. Balke and Kapetanios (2007) develop a method to approximate unknown models using a neural network. Péguin-Feissolle et al. (2013) introduce a causality test that is based on a Taylor approximation of a general nonlinear model and is applicable to various nonlinear models. These approaches are relevant from the viewpoint of a polynomial approximation.

In this study, we examine rejection frequencies under the null and alternative hypotheses for the introduced ARCH tests using Monte Carlo simulations. The simulation analyzes the influence of the lag length, the bandwidth selection for the Nadaraya-Watson estimator, and the approximation order for the polynomial approximation method on the results. The conditional mean models investigated in this study are linear autoregressive, threshold autoregressive, smooth transition autoregressive, Markov switching, and bi-linear models. These are popular nonlinear models used for empirical analysis and tend to cause spurious ARCH effects because as shown by Lumsdaine and Ng (1999), Van Dijk et al. (1999), and Balke and Kapetanios (2007), it is difficult to distinguish between nonlinear models with homoskedastic variance and linear models with an ARCH effect. The Monte Carlo simulation results evidence that ARCH tests that are based on the polynomial approximation regression approach have better size and power properties than those using the

Nadaraya-Watson kernel regression approach when DGPs are various nonlinear models. Empirical applications to economic variables support the simulation results.

The remainder of this paper is organized as follows: Section 2 presents the influence of a misspecified conditional mean on the ARCH tests and proposes ARCH tests using nonparametric regression approaches for the conditional mean. Section 3 presents the size and power of the tests under nonlinear models. Section 4 illustrates empirical applications. Section 5 concludes the paper.

2. ARCH tests using nonparametric regression approaches for conditional mean

We consider the following DGP with lag order m :

$$y_t = f(y_{t-1}, \dots, y_{t-m}; \beta) + u_t, \quad t = 1, \dots, T \quad (1)$$

where $f(\cdot; \cdot)$ is an unknown function and β is a parameter vector. u_t is a disturbance term with mean zero and variance denoted by

$$u_t = \sigma_t \varepsilon_t; \quad \sigma_t^2 = \gamma_0 + \sum_{i=1}^p \gamma_i u_{t-i}^2, \quad (2)$$

where ε_t are independently and identically distributed (iid) random variables with mean zero and variance equal to one. Although the conditional variance could have model misspecification similar to the conditional mean, standard heteroskedastic tests have the ability to find linear ARCH effects even if the true conditional variance is generalized ARCH (GARCH) with or without nonlinear parts.¹ On the other hand, spurious ARCH effects tend to be observed when the conditional mean has model misspecifications. The misspecification of the conditional mean has clear impacts on the inference of variance, as shown by Lumsdaine and Ng (1999) and Balke and Kapetanios (2007). Thus, we focus on investigating the influence that the model misspecification of the conditional mean has on ARCH effects.

The null hypothesis of homoskedasticity to test for the ARCH effect is denoted by

$$H_0 : \gamma_1 = \dots = \gamma_p = 0, \quad (3)$$

and the alternative hypothesis is

$$H_1 : \text{at least one } \gamma_i \neq 0, \quad i = 1, \dots, p. \quad (4)$$

Even if we assume a GARCH model to be heteroskedastic, the testing procedure is the same as that in by Lee (1991) and Gel and Chen (2012). Therefore, we focus only on the ARCH test. Engle's (1982) standard ARCH test uses the auxiliary regression of squared residuals:

$$\hat{u}_t^2 = \gamma_0 + \gamma_1 \hat{u}_{t-1}^2 + \dots + \gamma_p \hat{u}_{t-p}^2 + \eta_t, \quad (5)$$

where η_t is an error term. The LM test statistics is given by

$$LM = \frac{T \hat{d}' \hat{W} (\hat{W}' \hat{W})^{-1} \hat{W} \hat{d}}{\hat{d}' \hat{d}}, \quad (6)$$

where $\hat{d} = (\hat{d}_1, \dots, \hat{d}_T)$, $\hat{d}_t = (\hat{u}_t^2 / \hat{\sigma}_u^2 - 1)$, $\hat{\sigma}_u^2 = (1/T) \sum_{t=1}^T \hat{u}_t^2$, $\hat{W} = (\hat{w}_1, \dots, \hat{w}_T)$, and $\hat{w}_t = (1, \hat{u}_{t-1}^2, \dots, \hat{u}_{t-p}^2)$. The LM test statistic (6) is equivalent to TR^2 , where R^2 is the coefficient for the determination of (5)². Under the null hypothesis of no ARCH effects, the asymptotic distribution of (6) is $\chi^2(p)$.

When true DGP are denoted by (1), suppose that we estimate the following misspecified model:

$$y_t = g(y_{t-1}, \dots, y_{t-\tilde{m}}; \alpha) + u_t, \quad (7)$$

where $g(\cdot; \cdot)$ is a misspecified function, \tilde{m} is the lag length, and α is a parameter vector for the misspecified model. Accordingly, the residual is denoted by

$$\hat{u}_t = u_t + f(y_{t-1}, \dots, y_{t-m}; \beta) - \hat{g}(y_{t-1}, \dots, y_{t-\tilde{m}}; \alpha) = u_t + e_t, \quad (8)$$

where $e_t = f(y_{t-1}, \dots, y_{t-m}; \beta) - \hat{g}(y_{t-1}, \dots, y_{t-\tilde{m}}; \alpha)$. The squared residual for \hat{u}_t is

$$\hat{u}_t^2 = u_t^2 + 2u_te_t + e_t^2. \quad (9)$$

Equation (9) means that the ARCH test correctly performs when $e_t \xrightarrow{P} 0$, whereas the ARCH test is subject to a model misspecification and leads to unreliable results when $e_t \xrightarrow{P} 0$ does not hold. For example, when true DGP (1) are a threshold autoregressive (TAR) model and misspecified estimation model (7) is a linear AR model, e_t includes nonlinearity. As highlighted by Lumsdaine and Ng (1999) and Balke and Kapetanios (2007), such a misspecification results in a spurious ARCH effect. Therefore, a regression approach that does not depend on a specific model is necessary to avoid model misspecification and spurious ARCH effects.

The first approach that is robust to model misspecification is a nonparametric regression that is based on the Nadaraya-Watson kernel estimator. We consider the following conditional mean regression model:

$$y_t = m(y_{t-1}, \dots, y_{t-s}) + u_t, \quad t = 1, \dots, T, \quad (10)$$

where $m(\cdot)$ is the unknown regression function without any parametric form. The regression function for y_t on $Y_t = (y_{t-1}, \dots, y_{t-s})$ is

$$z(y_{t-1}, \dots, y_{t-s}) = E(y_t | Y_t = y). \quad (11)$$

The most representative method to estimate the function is the Nadaraya-Watson estimator. The estimator is denoted by

$$\hat{z}(y_{t-1}, \dots, y_{t-s}) = \frac{\sum_{t=1}^T K\left(\frac{y_{t-1}-y_1}{h_1}, \dots, \frac{y_{t-s}-y_s}{h_s}\right) y_t}{\sum_{t=1}^T K\left(\frac{y_{t-1}-y_1}{h_1}, \dots, \frac{y_{t-s}-y_s}{h_s}\right)} \quad (12)$$

where h_s denotes the bandwidth to determine the smoothness of the kernel function. The kernel function $K(y)$ satisfies the following:

$$\int K(y) dy = 1, \quad \int yK(y) dy = 0, \quad \int y^2 K(y) dy > 0. \quad (13)$$

This study uses the Gaussian kernel denoted by³:

$$K(\cdot) = \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{y^2}{2}\right). \quad (14)$$

We use two bandwidth selections for h_s that are derived by minimizing the integrated mean squared error (IMSE). First is Silverman's (1986) plug-in method. The bandwidth obtained using the plug-in method is based on the following equation:

$$h_s = c_0 T^{-1/(1+s)}, \quad (15)$$

where c_0 is a constant that depends on the kernel function. The optimal bandwidth selection for the Gaussian kernel is denoted by

$$h_s^{opt} \approx 1.06 \sigma_s T^{-1/(s+4)}, \quad (16)$$

where σ_s is the standard deviation for y_{t-s} . The modified h_s^{opt} that is robust to outliers is written as

$$\tilde{h}_s^{opt} = 1.06 \min(\hat{\sigma}_s, \hat{Q}_s/1.34) T^{-1/(s+4)}, \quad (17)$$

where \hat{Q}_s is the estimate for the interquartile range of y_{t-s} .⁴

Second is the cross-validation procedure developed by Rudemo (1982). When using the Gaussian kernel, we consider the following mean squared error called the cross-validation criterion:

$$CV(\mathbf{h}) = \frac{1}{T} \sum_{i=1}^T (y_i - \hat{z}(Y_{-i}))^2, \quad (18)$$

where $\mathbf{h} = (h_1, \dots, h_s)$ and $\hat{z}(Y_{-i})$ is a leave-one-out estimator that excludes i th observation. The optimal bandwidth \mathbf{h} for the cross-validation procedure is determined by minimizing $CV(\mathbf{h})$. Stone (1984) shows that bandwidth \mathbf{h} for the cross-validation can asymptotically select the optimal bandwidth from an IMSE viewpoint and has probability convergence to the bandwidth for the plug-in method. While bandwidth h_s for the plug-in method depends on the assumed kernel density function, the cross-validation is not required to assume the kernel density function and can obtain a consistent estimator for the bandwidth that minimizes IMSE. It is possible that the residuals obtained using Nadaraya-Watson estimator (12) with bandwidth selection (17) or (18) have similar properties. Accordingly, the above-mentioned nonparametric regression approach is robust to the model misspecification of the conditional mean and thus, the ARCH test is correctly performed.⁵

The next approach adopted to avoid misspecification is a polynomial approximation of a general unknown nonlinear model. When we apply a k th-order Taylor approximation to true model (1), the regression model is denoted by

$$y_t = \beta_0 + \sum_{j=1}^l \beta_j y_{t-j} + \sum_{j_1=1}^l \sum_{j_2=1}^l \beta_{j_1 j_2} y_{t-j_1} y_{t-j_2} + \dots + \sum_{j_1=1}^l \sum_{j_2=1}^l \dots \sum_{j_k=1}^l \beta_{j_1 \dots j_k} y_{t-j_1} \dots y_{t-j_k} + \epsilon_t, \quad (19)$$

where l is the lag length and ϵ_t is an error term that includes the remainder term of the Taylor series approximation. We assume $l \leq k$ as a simple notation. If the true model is a linear AR model, all $\beta_{j_1 j_2}$ and $\beta_{j_1 \dots j_k}$ are zero. In contrast, if the true model is nonlinear, one $\beta_{j_1 j_2}$ or $\beta_{j_1 \dots j_k}$ is not zero at least. We investigate this using a standard Wald test. For example, (19) with $l = 2$ and $k = 2$ can be written as

$$y_t = \beta_0 + \sum_{j=1}^2 \beta_j y_{t-j} + \sum_{j_1=1}^2 \sum_{j_2=1}^2 \beta_{j_1 j_2} y_{t-j_1} y_{t-j_2} + \epsilon_t. \quad (20)$$

The difference between the true model and the polynomial approximation regression model reduces because the polynomial regression can approximate various nonlinear models including

the TAR and Markov switching models. When testing for ARCH effects under an unknown (true) model, using residuals obtained from polynomial approximation regression (19) can be advantageous since it is possible that they show size and power properties similar to those of the true model. Therefore, the ARCH test using the residuals from the polynomial approximation regression does not appear to be influenced by model misspecification.

3. Size and power properties of ARCH tests using nonparametric regression models

This section examines the size and power of the ARCH tests using nonparametric regression models for the conditional mean presented in Section 2. We conduct Monte Carlo simulations to compare the rejection frequencies of the test statistics under various conditional mean models with and without ARCH effects. The simulations are based on 10,000 replications; a significance level of 5%; and sample sizes with $T = 100, 250$, and 500 .⁶ To avoid the effect of initial conditions, data with $T + 100$ are generated. We discard the initial 100 samples and use the data with sample size T . We compare ARCH tests (6) using the following regression models for the conditional mean: the AR model denoted $AR(q)$, polynomial approximation model (19) with second- and third-order Taylor approximation denoted as $T2(q)$ and $T3(q)$, and nonparametric regression model (12) with plug-in method (17) and cross-validation method (18) denoted as $NP_{pl}(q)$ and $NP_{cv}(q)$. We set lag length q to $q = 1$ or $q = 2$.⁷ The AR model is used as a benchmark for comparison.

First, we consider the following AR processes to examine the influence of lag length on the tests' performance.

$$y_t = \beta_0 + \beta_1 y_{t-1} + \beta_2 y_{t-2} + u_t, \quad (21)$$

$$u_t = \sigma_t \epsilon_t, \quad (22)$$

$$\sigma_t^2 = \gamma_0 + \gamma_1 u_{t-1}^2, \quad (23)$$

where $u_t \sim \text{i.i.d.} N(0, 1)$. β_0 is set to $\beta_0 = 0$. Table 1 presents the rejection frequencies for the ARCH tests obtained from each regression model for the conditional mean. We use the following DGP:

$$\text{DGP1-1: } y_t = 0.2y_{t-1} + u_t,$$

$$\text{DGP1-2: } y_t = 0.7y_{t-1} + u_t,$$

$$\text{DGP1-3: } y_t = 0.7y_{t-1} - 0.2y_{t-2} + u_t,$$

$$\text{DGP1-4: } y_t = 0.7y_{t-1} - 0.5y_{t-2} + u_t.$$

These DGP have homoskedastic errors with $\gamma_0 = 1$ and $\gamma_1 = 0$ for (23). The rejection frequencies presented in Table 1 indicate the empirical size of the ARCH tests on the basis of each regression model.

For DGP1-1 and DGP1-2, which have lag order one, most of the tests have a small under-rejection but reasonable size performance, except for $NP_{pl}(2)$ and $NP_{cv}(2)$. $NP_{pl}(2)$ and $NP_{cv}(2)$ report over-rejections for DGP1-1 and DGP1-2. The rejection frequencies of $NP_{pl}(2)$ for DGP1-1 with $T = 500$ and of $NP_{cv}(2)$ for DGP1-2 with $T = 500$ are 0.143 and 0.101. An additional lag for the nonparametric regression of the conditional mean using the Nadaraya-Watson estimator leads to size distortions in the ARCH tests. In contrast, $AR(2)$, $T2(2)$, and $T3(2)$ do not report overrejections for DGP1-1 and DGP1-2. The results show that the additional lag for AR and polynomial approximation regression does not impact the size of the ARCH tests. However, a lower lag length clearly influences the empirical size of all the tests. We see that the ARCH tests based on $AR(1)$, $T2(1)$, $T3(1)$, $NP_{pl}(1)$,

Table 1. Rejection frequencies under AR models

	AR(1)	AR(2)	T2(1)	T2(2)	T3(1)	T3(2)	$NP_{pl}(1)$	$NP_{pl}(2)$	$NP_{cv}(1)$	$NP_{cv}(2)$
DGP1-1										
$T = 100$	0.035	0.035	0.035	0.025	0.032	0.028	0.031	0.041	0.040	0.034
$T = 250$	0.044	0.040	0.038	0.040	0.043	0.037	0.046	0.093	0.051	0.042
$T = 500$	0.041	0.043	0.046	0.041	0.046	0.046	0.051	0.143	0.052	0.047
DGP1-2										
$T = 100$	0.036	0.032	0.035	0.029	0.029	0.034	0.034	0.026	0.030	0.031
$T = 250$	0.039	0.040	0.038	0.038	0.041	0.043	0.037	0.025	0.039	0.066
$T = 500$	0.042	0.045	0.041	0.039	0.042	0.044	0.041	0.029	0.044	0.101
DGP1-3										
$T = 100$	0.042	0.031	0.032	0.027	0.032	0.031	0.035	0.018	0.042	0.030
$T = 250$	0.052	0.042	0.046	0.035	0.047	0.040	0.049	0.026	0.046	0.049
$T = 500$	0.063	0.044	0.061	0.040	0.054	0.045	0.064	0.037	0.052	0.075
DGP1-4										
$T = 100$	0.063	0.034	0.060	0.026	0.048	0.028	0.051	0.019	0.072	0.029
$T = 250$	0.127	0.043	0.116	0.037	0.097	0.038	0.115	0.027	0.113	0.048
$T = 500$	0.213	0.041	0.201	0.042	0.189	0.040	0.208	0.032	0.181	0.069

and $NP_{cv}(1)$ over-reject the null hypothesis of homoskedastic variance under DGP1-3 or DGP1-4, which have a lag order of two. For example, the rejection frequencies of $AR(1)$, $T2(1)$, $T3(1)$, $NP_{pl}(1)$, and $NP_{cv}(1)$ for DGP1-4 with $T = 250$ are 0.127, 0.116, 0.097, 0.115, and 0.113, respectively. The size distortions in DGP1-4 are greater than those in DGP1-3. The influence of the lower lag length on the empirical size depends on the persistence parameter of DGP. Compared with the size distortions for the model with a lower lag length, those for the model with an additional lag length are smaller. The lower lag length tends to cause the size distortions because the conditional variance includes the influence of extra lags and leads to spurious ARCH effects. In contrast, when the conditional mean regression includes additional lags, they are not significant and the conditional variance does not include the influence of additional lag length. Accordingly, we present size and power properties for the models with two lags below.

We examine the empirical size of the ARCH tests under the following conditional mean generated by the TAR models.

$$\text{DGP2-1: } y_t = (0.7y_{t-1} - 0.2y_{t-2})I(y_{t-1} \geq 0) + (0.1y_{t-1} - 0.2y_{t-2})I(y_{t-1} < 0) + u_t,$$

$$\text{DGP2-2: } y_t = (0.7y_{t-1} - 0.2y_{t-2})I(y_{t-1} \geq 0) + (-0.5y_{t-1} - 0.2y_{t-2})I(y_{t-1} < 0) + u_t,$$

$$\text{DGP2-3: } y_t = (0.7y_{t-1} + 0.2y_{t-2})I(y_{t-1} \geq 0) + (0.7y_{t-1} - 0.7y_{t-2})I(y_{t-1} < 0) + u_t,$$

$$\text{DGP2-4: } y_t = (0.7y_{t-1} - 0.2y_{t-2})I(\Delta y_{t-1} \geq 0) + (0.1y_{t-1} - 0.2y_{t-2})I(\Delta y_{t-1} < 0) + u_t,$$

$$\text{DGP2-5: } y_t = (0.7y_{t-1} - 0.2y_{t-2})I(\Delta y_{t-1} \geq 0) + (-0.5y_{t-1} - 0.2y_{t-2})I(\Delta y_{t-1} < 0) + u_t,$$

$$\text{DGP2-6: } y_t = (0.7y_{t-1} + 0.2y_{t-2})I(\Delta y_{t-1} \geq 0) + (0.7y_{t-1} - 0.7y_{t-2})I(\Delta y_{t-1} < 0) + u_t,$$

where $I(\cdot)$ is an indicator function that takes the value of 1 if $I(\cdot)$ is true and 0 if $I(\cdot)$ is not true. u_t denotes a homoskedastic error similar to that from DGP1-1 to 1-4. While DGP2-1, 2-2, and 2-3 are standard TAR models whose indicator functions depend on y_{t-1} , DGP2-4, 2-5, and 2-6 are

momentum threshold autoregressive (MTAR) models wherein the threshold is the difference Δy_{t-1} . These TAR models allow for asymmetric adjustments. In addition, MTAR can capture the spiky properties of the process.

Figures 1 and 2 illustrate the sample path for DGP2-1 with homoskedastic errors and the ARCH effect $\gamma_0 = 0.3$ for (23). Figure 2 clearly shows the volatile behavior generated by the ARCH effect. However, Figure 3 illustrates that the sample path for DGP2-3 demonstrates a similar volatile movement even if the error is homoskedastic. As shown in Figures 2 and 3, it is generally difficult to distinguish between the nonlinear conditional mean model with the homoskedastic error and the linear AR model with ARCH effect. Such a similarity between the TAR model with homoskedastic errors and the linear AR model with ARCH effects may produce spurious size and power properties.

Table 2 tabulates the simulation results. $AR(2)$ reports over-rejections for the null hypothesis of no ARCH effects. For DGP2-2 and DGP2-5, which have strong asymmetry, the size distortions of $AR(2)$ are significantly large. These results indicate that the use of the AR model for the conditional mean leads to spurious ARCH effects when the true DGP are based on the TAR or MTAR model. In addition, the over-rejections increase with large sample size. Unlike the performance of $AR(2)$, the polynomial approximation regression models $T2(2)$ and $T3(2)$ and nonparametric regression models $NP_{pl}(2)$ and $NP_{cv}(2)$ perform better. For example, the rejection frequencies of $AR(2)$, $T2(2)$, $T3(2)$, $NP_{pl}(2)$, and $NP_{cv}(2)$ for DGP2-2 with $T = 250$ are 0.373, 0.040, 0.033, 0.042, and 0.051, respectively. $T3(2)$ has a more reasonable size compared with those for $T2(2)$, $NP_{pl}(2)$, and $NP_{cv}(2)$. $T2(2)$, $NP_{pl}(2)$, and $NP_{cv}(2)$ report size distortions in certain cases. The rejection frequencies of $T2(2)$, $NP_{pl}(2)$, and $NP_{cv}(2)$ for DGP2-3 with $T = 500$ are 0.096, 0.139, and 0.104, respectively. Thus, the polynomial approximation regression model $T3(2)$ is a more appropriate approach to test for ARCH than other approaches under the TAR or MTAR model.

Table 3 presents the rejection frequencies for each test under smooth transition autoregressive (STAR) models generated by the followings:

$$\text{DGP3-1: } y_t = 0.7y_{t-1} - 0.2y_{t-2} + (-0.5y_{t-1} - 0.2y_{t-2})(1 - \exp(-0.1y_{t-1}^2)) + u_t,$$

Figure 1. Sample path for DGP1-1.




Figure 2. Sample path for DGP1-1 with $\gamma_1 = 0.3$.





Figure 3. Sample path for DGP2-3.



$$\text{DGP3-2: } y_t = 0.7y_{t-1} - 0.2y_{t-2} + (-y_{t-1} - 0.2y_{t-2})(1 - \exp(-0.1y_{t-1}^2)) + u_t,$$

$$\text{DGP3-3: } y_t = 0.7y_{t-1} - 0.2y_{t-2} + (-y_{t-1} - 0.2y_{t-2})(1 - \exp(-y_{t-1}^2)) + u_t,$$

$$\text{DGP3-4: } y_t = 0.7y_{t-1} - 0.2y_{t-2} + (-0.5y_{t-1} - 0.2y_{t-2})(1 + \exp(-0.1y_{t-1}))^{-1} + u_t,$$

$$\text{DGP3-5: } y_t = 0.7y_{t-1} - 0.2y_{t-2} + (-y_{t-1} - 0.2y_{t-2})(1 + \exp(-0.1y_{t-1}))^{-1} + u_t,$$

$$\text{DGP3-6: } y_t = 0.7y_{t-1} - 0.2y_{t-2} + (-y_{t-1} - 0.2y_{t-2})(1 + \exp(-y_{t-1}))^{-1} + u_t,$$

Table 2. Rejection frequencies under TAR models

	$AR(2)$	$T2(2)$	$T3(2)$	$NP_{pl}(2)$	$NP_{cv}(2)$
DGP2-1					
$T = 100$	0.049	0.029	0.029	0.028	0.030
$T = 250$	0.080	0.036	0.040	0.051	0.041
$T = 500$	0.116	0.046	0.041	0.081	0.071
DGP2-2					
$T = 100$	0.150	0.029	0.025	0.021	0.033
$T = 250$	0.373	0.040	0.033	0.042	0.051
$T = 500$	0.658	0.055	0.040	0.069	0.073
DGP2-3					
$T = 100$	0.083	0.038	0.026	0.104	0.026
$T = 250$	0.144	0.064	0.040	0.132	0.065
$T = 500$	0.224	0.096	0.058	0.139	0.104
DGP2-4					
$T = 100$	0.063	0.027	0.027	0.034	0.030
$T = 250$	0.117	0.040	0.038	0.086	0.046
$T = 500$	0.188	0.045	0.041	0.135	0.073
DGP2-5					
$T = 100$	0.359	0.039	0.026	0.027	0.053
$T = 250$	0.748	0.083	0.051	0.059	0.076
$T = 500$	0.962	0.150	0.090	0.094	0.144
DGP2-6					
$T = 100$	0.081	0.036	0.028	0.030	0.040
$T = 250$	0.176	0.050	0.044	0.037	0.075
$T = 500$	0.277	0.069	0.052	0.044	0.142

where u_t denotes homoskedastic errors similar to those in Tables 1 and 2. STAR models have the time-varying properties of the conditional mean. DGP3-1, 3-2, and 3-3 impose symmetry constraints on the time-varying properties, whereas DGP3-4, 3-5, and 3-6, which are logistic STAR models, allow asymmetry. DGP3-2 and 3-5 produce a smoother and more marginal change than DGP3-3 and 3-6. We observe that $AR(2)$, $T2(2)$, and $NP_{pl}(2)$ partially reject the null hypothesis of no ARCH effects. The rejection frequencies of $AR(2)$ is higher than those of the other regression models for DGP3-2 and 3-6. $T2(2)$ shows size distortions for DGP3-2. $NP_{pl}(2)$ reports a slight over-rejection with $T = 500$. In contrast, the shape of the transition function does not have a clear impact on the empirical size of $T3(2)$ and $NP_{cv}(2)$. $T3(2)$ and $NP_{cv}(2)$ can capture the properties of STAR models and allows the ARCH test to perform well.

In addition, we present the results of each test for the other nonlinear processes:

$$\text{DGP4-1: } y_t = (0.7y_{t-1} - 0.2y_{t-2})s_t + (0.3y_{t-1} - 0.2y_{t-2})(1 - s_t) + u_t, p_{00} = p_{11} = 0.7,$$

$$\text{DGP4-2: } y_t = (0.7y_{t-1} - 0.2y_{t-2})s_t + (0.3y_{t-1} - 0.2y_{t-2})(1 - s_t) + u_t, p_{00} = p_{11} = 0.98,$$

$$\text{DGP4-3: } y_t = (0.7y_{t-1} + 0.2y_{t-2})s_t + (0.3y_{t-1} - 0.2y_{t-2})(1 - s_t) + u_t, p_{00} = p_{11} = 0.98,$$

$$\text{DGP4-4: } y_t = 0.1y_{t-1}u_{t-1} + 0.1y_{t-2}u_{t-2} + u_t,$$

$$\text{DGP4-5: } y_t = 0.3y_{t-1}u_{t-1} + 0.1y_{t-2}u_{t-2} + u_t,$$

Table 3. Rejection frequencies under STAR models

	AR(2)	T2(2)	T3(2)	$NP_{pl}(2)$	$NP_{cv}(2)$
DGP3-1					
$T = 100$	0.053	0.032	0.033	0.023	0.027
$T = 250$	0.068	0.045	0.038	0.053	0.043
$T = 500$	0.094	0.068	0.043	0.088	0.063
DGP3-2					
$T = 100$	0.088	0.043	0.029	0.028	0.028
$T = 250$	0.184	0.107	0.038	0.061	0.042
$T = 500$	0.303	0.238	0.044	0.088	0.054
DGP3-3					
$T = 100$	0.042	0.024	0.026	0.027	0.031
$T = 250$	0.048	0.038	0.030	0.054	0.047
$T = 500$	0.057	0.046	0.042	0.085	0.069
DGP3-4					
$T = 100$	0.035	0.030	0.031	0.027	0.031
$T = 250$	0.044	0.034	0.040	0.055	0.043
$T = 500$	0.045	0.040	0.043	0.082	0.049
DGP3-5					
$T = 100$	0.032	0.029	0.025	0.034	0.028
$T = 250$	0.042	0.039	0.040	0.067	0.043
$T = 500$	0.053	0.041	0.042	0.111	0.047
DGP3-6					
$T = 100$	0.109	0.029	0.029	0.028	0.032
$T = 250$	0.273	0.042	0.037	0.049	0.042
$T = 500$	0.483	0.048	0.042	0.073	0.055

$$\text{DGP4-6: } y_t = 0.1y_{t-1}u_{t-1} - 0.1y_{t-2}u_{t-2} + u_t,$$

where $u_t \sim \text{i.i.d.} N(0, 1)$ and s_t is a random variable that takes the value of 0 or 1. DGP4-1, 4-2, and 4-3 are Markov switching processes and s_t determines the behavior. Whether s_t takes the value of 0 or 1 depends on the transition probabilities p_{11} and p_{00} . $p_{11} = P(s_{t+1} = 1 | s_t = 1)$ denotes the change probability from state $s_t = 1$ to state $s_{t+1} = 1$. Similarly, the transition probabilities are denoted by $p_{00} = P(s_{t+1} = 0 | s_t = 0)$, $p_{10} = 1 - p_{00} = P(s_{t+1} = 1 | s_t = 0)$, and $p_{01} = 1 - p_{11} = P(s_{t+1} = 0 | s_t = 1)$, respectively. They are set to $p_{11} = p_{00} = 0.7$ for DGP4-1 and $p_{11} = p_{00} = 0.98$ for DGP4-2 and 4-3. While DGP4-1 has frequent switches in the AR parameters, DGP4-2 and 4-3 show persistent switches. DGP4-4, 4-5, and 4-6 are bilinear models that are used to model rare, volatile, or outburst processes.

AR(2) that neglects nonlinearity causes the spurious ARCH effect, which is similar to the results in Tables 2 and 3. The results for the nonparametric regression models using the Nadaraya-Watson estimator depend on the bandwidth selection. $NP_{pl}(2)$ under-rejects the null hypothesis for DGP4-1, DGP4-2, and DGP4-5 and over-rejects that for DGP4-3, 4-4, and 4-6. $NP_{cv}(2)$ performs well for DGP4-2, DGP4-4, and DGP4-6 and over-rejects the null hypothesis for DGP4-1, 4-3, and 4-5. T2(2) has relatively reasonable empirical sizes for $T = 100$ and 200, but reports size distortions for DGP4-1, 4-2, 4-3, 4-5, and 4-6 with $T = 500$. Here as well, we find that T3(2) generally performs better.

The simulation results from Tables 1–4 evidence that the model misspecification of the conditional mean causes size distortions for the null hypothesis of no ARCH effects. The ARCH tests using

Table 4. Rejection frequencies under MS and bilinear models

	AR(2)	T2(2)	T3(2)	NP _{pl} (2)	NP _{cv} (2)
DGP4-1					
T = 100	0.051	0.034	0.029	0.021	0.049
T = 250	0.092	0.064	0.040	0.025	0.065
T = 500	0.137	0.101	0.068	0.029	0.077
DGP4-2					
T = 100	0.047	0.029	0.026	0.023	0.040
T = 250	0.076	0.053	0.037	0.026	0.051
T = 500	0.108	0.084	0.055	0.034	0.057
DGP4-3					
T = 100	0.048	0.035	0.031	0.112	0.051
T = 250	0.084	0.054	0.044	0.156	0.073
T = 500	0.116	0.087	0.057	0.141	0.100
DGP4-4					
T = 100	0.066	0.026	0.028	0.034	0.049
T = 250	0.112	0.036	0.038	0.076	0.051
T = 500	0.170	0.043	0.044	0.129	0.052
DGP4-5					
T = 100	0.456	0.035	0.032	0.027	0.107
T = 250	0.851	0.071	0.040	0.033	0.110
T = 500	0.987	0.100	0.046	0.036	0.133
DGP4-6					
T = 100	0.053	0.029	0.030	0.037	0.040
T = 250	0.090	0.040	0.039	0.078	0.051
T = 500	0.134	0.040	0.046	0.133	0.056

the AR regression model are sensitive to the presence of the nonlinear conditional mean and show high over-rejections. This can be attributed by neglected nonlinearity and difficulties in distinguishing between the nonlinearity of the conditional mean and the ARCH effects. While the nonparametric regression models using the Nadaraya-Watson estimator partially perform well, the rejection frequencies strongly depend on DGP and the bandwidth selection. By contrast, the size properties of T3(2) outperform those of other models and are close to the nominal size at 5%. Therefore, T3(2) can approximate the (unknown) linear and nonlinear conditional mean models well and produce reliable ARCH tests.

Tables 5 and 6 report the nominal power and size-corrected power properties for the ARCH tests. We use DGP1-3, DGP2-1, 2-4, 3-1, 3-4, 4-1, and 4-4 for power comparison. Each DGP has an ARCH effect denoted by

$$u_t = \sigma_t \epsilon_t, \quad (24)$$

$$\sigma_t^2 = \gamma_0 + \gamma_1 u_{t-1}^2, \quad (25)$$

where γ_0 and γ_1 are set to $\gamma_0 = 1$ and $\gamma_1 = (0.1, 0.3)$. The powers of AR(2) are clearly higher than those of other models in Table 5. We have a relatively reasonable evaluation of the power for DGP1-3 because the

Table 5. Nominal power properties for ARCH tests

	$\gamma_1 = 0.1$					$\gamma_1 = 0.3$				
	AR(2)	T2(2)	T3(2)	NP _{pl} (2)	NP _{cv} (2)	AR(2)	T2(2)	T3(2)	NP _{pl} (2)	NP _{cv} (2)
DGP1-3										
$T = 100$	0.119	0.070	0.032	0.043	0.055	0.440	0.294	0.138	0.218	0.205
$T = 250$	0.267	0.197	0.122	0.082	0.089	0.848	0.777	0.646	0.589	0.493
$T = 500$	0.486	0.413	0.309	0.182	0.164	0.990	0.982	0.959	0.913	0.845
DGP2-1										
$T = 100$	0.146	0.065	0.032	0.027	0.065	0.444	0.290	0.143	0.159	0.243
$T = 250$	0.364	0.213	0.119	0.046	0.102	0.868	0.787	0.673	0.484	0.552
$T = 500$	0.624	0.429	0.325	0.108	0.192	0.991	0.980	0.963	0.877	0.879
DGP2-4										
$T = 100$	0.164	0.068	0.038	0.018	0.072	0.446	0.284	0.147	0.122	0.239
$T = 250$	0.400	0.218	0.127	0.029	0.099	0.853	0.774	0.642	0.408	0.516
$T = 500$	0.677	0.446	0.337	0.065	0.158	0.989	0.980	0.958	0.792	0.844
DGP3-1										
$T = 100$	0.164	0.081	0.029	0.027	0.057	0.474	0.315	0.150	0.142	0.245
$T = 250$	0.363	0.275	0.119	0.047	0.111	0.879	0.817	0.647	0.477	0.609
$T = 500$	0.617	0.540	0.320	0.105	0.224	0.992	0.986	0.964	0.858	0.917
DGP3-4										
$T = 100$	0.115	0.062	0.033	0.024	0.072	0.440	0.287	0.143	0.159	0.282
$T = 250$	0.275	0.203	0.116	0.051	0.141	0.850	0.777	0.655	0.486	0.647
$T = 500$	0.490	0.418	0.325	0.109	0.258	0.988	0.982	0.964	0.859	0.933
DGP4-1										
$T = 100$	0.154	0.083	0.038	0.034	0.113	0.456	0.297	0.143	0.178	0.323
$T = 250$	0.379	0.294	0.180	0.086	0.235	0.863	0.797	0.662	0.532	0.696
$T = 500$	0.654	0.586	0.466	0.213	0.431	0.990	0.985	0.973	0.891	0.948
DGP4-4										
$T = 100$	0.189	0.067	0.030	0.023	0.117	0.481	0.283	0.135	0.140	0.369
$T = 250$	0.425	0.201	0.111	0.036	0.215	0.884	0.781	0.639	0.430	0.715
$T = 500$	0.693	0.419	0.310	0.067	0.354	0.991	0.983	0.964	0.828	0.950

size properties of AR(2) and other tests are close to the nominal level 0.05 (Table 1). However, we cannot correctly evaluate the high nominal powers of AR(2) for other DGP. The higher powers of AR(2) are influenced by size distortions presented in tables from 2 to 4. The power properties of the nonparametric models are more appropriately interpreted because T2(2) and T3(3) do not overreject the null hypothesis for DGP in Table 5 and the size distortions of NP_{pl}(2) and NP_{cv}(2) are smaller than those of AR(2). In comparison, we observe that the polynomial approximation models T2(2) and T3(2) perform better than NP_{pl}(2) and NP_{cv}(2). Note that the powers of NP_{pl}(2) are quite small when the ARCH effect is $\gamma_1 = 0.1$. For $\gamma_1 = 0.3$, the nonparametric regression models report sufficient power to identify the ARCH effects.

We compare the power properties among the models without the influences of size distortions. Table 6 demonstrates the size-corrected power. The powers of AR(2) in Table 6 are lower than those in Table 5 because the size distortions are corrected. AR(2) still performs well even if the size is corrected. The ability to detect ARCH effects in the nonlinear models for T2(2) is high, similar to that of AR(2). While the powers of T3(2) is slightly smaller than those of T2(2) because T3(2) has additional regression parameters for the conditional mean, it has sufficient power to find the ARCH

Table 6. Size-corrected power properties for ARCH tests

	$\gamma_1 = 0.1$					$\gamma_1 = 0.3$				
	AR(2)	T2(2)	T3(2)	NP _{pl} (2)	NP _{cv} (2)	AR(2)	T2(2)	T3(2)	NP _{pl} (2)	NP _{cv} (2)
DGP1-3										
$T = 100$	0.145	0.088	0.047	0.075	0.081	0.479	0.325	0.165	0.280	0.235
$T = 250$	0.285	0.235	0.127	0.114	0.092	0.862	0.808	0.665	0.644	0.494
$T = 500$	0.501	0.429	0.321	0.191	0.127	0.990	0.982	0.967	0.919	0.807
DGP2-1										
$T = 100$	0.162	0.094	0.048	0.041	0.080	0.460	0.343	0.177	0.203	0.261
$T = 250$	0.296	0.235	0.138	0.044	0.107	0.820	0.811	0.678	0.472	0.571
$T = 500$	0.476	0.453	0.342	0.078	0.168	0.980	0.984	0.968	0.819	0.864
DGP2-4										
$T = 100$	0.148	0.089	0.058	0.028	0.086	0.417	0.331	0.185	0.133	0.271
$T = 250$	0.279	0.240	0.149	0.020	0.088	0.763	0.786	0.669	0.348	0.500
$T = 500$	0.412	0.457	0.352	0.032	0.120	0.951	0.979	0.966	0.719	0.803
DGP3-1										
$T = 100$	0.165	0.107	0.046	0.041	0.098	0.470	0.366	0.179	0.188	0.293
$T = 250$	0.315	0.262	0.129	0.041	0.156	0.849	0.815	0.668	0.472	0.620
$T = 500$	0.502	0.483	0.342	0.075	0.246	0.984	0.982	0.971	0.820	0.908
DGP3-4										
$T = 100$	0.193	0.106	0.044	0.043	0.098	0.477	0.346	0.176	0.192	0.315
$T = 250$	0.387	0.295	0.132	0.046	0.150	0.865	0.808	0.674	0.475	0.674
$T = 500$	0.625	0.545	0.327	0.080	0.259	0.988	0.984	0.966	0.819	0.936
DGP4-1										
$T = 100$	0.153	0.115	0.059	0.063	0.117	0.448	0.345	0.182	0.236	0.326
$T = 250$	0.287	0.263	0.207	0.119	0.219	0.803	0.773	0.690	0.599	0.671
$T = 500$	0.451	0.452	0.415	0.267	0.391	0.972	0.967	0.960	0.919	0.937
DGP4-4										
$T = 100$	0.151	0.086	0.043	0.028	0.124	0.440	0.332	0.166	0.152	0.371
$T = 250$	0.287	0.227	0.133	0.025	0.211	0.793	0.804	0.673	0.383	0.724
$T = 500$	0.457	0.431	0.327	0.032	0.341	0.961	0.983	0.967	0.733	0.947

effect. The rejection frequencies of $NP_{pl}(2)$ and $NP_{cv}(2)$ for $\gamma_1 = 0.1$ are inferior to those of other models in Table 6. While they relatively perform well for $\gamma_1 = 0.3$ with $T = 100$, other models have better power properties, particularly for $T = 250$ and 500.

The comparison of the ARCH tests using each regression model for the conditional mean indicates that the presence of the nonlinear conditional mean has influences of size and power properties on the ARCH tests. The AR regression models have higher over-rejection of the null hypothesis of no ARCH effects for the nonlinear conditional mean models. The ARCH tests based on AR models for the nonlinear conditional mean are not effective from the viewpoints of size and power. This is because size-corrected tests are needed and the true model is generally unknown a priori. The nonparametric regression models using the Nadaraya-Watson estimator tend to have slight size distortions and low power. The polynomial approximation model $T2(2)$ shows slight over-rejection depending on the nonlinear conditional mean and sample size, although it has better power properties for the ARCH effect with the nonlinear conditional mean. $T3(2)$ has a reasonable size and power properties and yields reliable results for the ARCH tests irrespective of the conditional mean models.

4. Empirical applications

In this section, we apply the tests to economic variables. We employ six economic data including those of Real Gross Domestic Product (GDP) for U.S., M2 for U.S., Federal Funds rate (FF), Producer Price Index (PPI) for U.S., Crude Oil Prices (COP) for West Texas Intermediate, and Global Price Index of all commodities (GPI). We use first difference of logarithm except FF. The data are obtained from Federal Reserve Bank of St. Louis. Each data consist of 294 (1Q 1947–2Q 2020), 478 (November 1980–August 2020), 794 (July 1954–August 2020), 1291 (January 1913–July 2020), 416 (January 1986–August 2020), and 343 (January 1992–July 2020) observations, respectively. The lag lengths of ARCH tests are selected by the Akaike information criterion (AIC).

Empirical results are presented in Table 7, which tabulates nominal p values of each test. The results support simulation evidence demonstrated in Section 3. For M2 and FF, the p values of all the tests are small. It is possible that the results are influenced by size distortions because Monte Carlo simulation in Section 3 demonstrates that when time series processes have nonlinear mean properties, $AR(2)$ has overrejections of the null hypothesis, and tends to lead spurious ARCH effects. $T2(2)$, $NP_{pl}(2)$, and $NP_{cv}(2)$ also have overrejections depending on DGP. However, as shown in Section 3, $T3(2)$ has reasonable size properties regardless of DGP. The p values of $T3(2)$ are sufficient small. This indicates that M2 and FF have ARCH effects. While $AR(2)$, $T2(2)$, $T3(2)$, and $NP_{cv}(2)$ have small p values for GDP and PPI similar to those for M2 and FF, $NP_{pl}(2)$ has large p values. The results may be due to poor power performance in the presence of small ARCH effects as shown in Table 5.

For COP and GPI, p values of $AR(2)$ are small, whereas those of $T3(2)$ and $NP_{pl}(2)$ are large. It is possible that this is because $AR(2)$ has overrejection when DGP have nonlinear mean properties. $T3(2)$ has superior size properties for DGP with nonlinear mean properties. The results indicate that COP and GPI have nonlinear mean properties without ARCH effects. The empirical applications support our simulation results and show that $T3(2)$ can appropriately find ARCH effects, whereas there are cases in which other tests lead to unreliable results.

5. Summary and conclusion

This study compares the size and power of the ARCH tests that are robust to misspecified conditional mean models. ARCH tests are important for statistical modeling because the presence of ARCH affects the statistical inference of the conditional mean regression model and the analysis of volatility. However, it is difficult to determine the correct specified conditional mean model and possible to employ a misspecified conditional mean model. This may lead to unreliable results. Therefore, it is necessary to compare robust ARCH tests to various unknown conditional mean model and clarify their size and power properties. The approaches employed in this study are based on two nonparametric regressions: an ARCH test using the Nadaraya-Watson kernel regression and an ARCH test with the polynomial approximation. The two approaches can adapt to various nonlinear models. Since a true model is generally unknown a priori, they are robust to misspecified models. The Monte Carlo simulations evidence that the ARCH tests based on the polynomial regression approach have a better size and power properties than those using the Nadaraya-Watson kernel regression approach for various nonlinear conditional mean models. In particular,

Table 7. Empirical results

	$AR(2)$	$T2(2)$	$T3(2)$	$NP_{pl}(2)$	$NP_{cv}(2)$
GDP	0.012	0.016	0.027	0.906	0.015
M2	0.000	0.000	0.000	0.037	0.003
FF	0.059	0.045	0.011	0.000	0.002
PPI	0.000	0.000	0.000	0.934	0.000
COP	0.000	0.000	0.211	0.609	0.527
GPI	0.000	0.478	0.619	0.594	0.013

the test using the regression approach based on the third-order Taylor approximation has a reasonable and acceptable size and sufficient power for any time series models. The simulation and empirical results further show that the ARCH test using the polynomial approximation approach is useful when testing if DGPs have an ARCH effect and for ARCH without model specifications when the conditional mean model is unknown a priori. Of course, if researchers know DGP a priori or the purpose of researchers is the specification of the conditional mean, nonparametric regressions employed in this study are not useful and parametric approaches are appropriate. However, when it is difficult to specify the parametric model a priori and the purpose of researchers is the test for heteroskedasticity, our analysis is more useful. Robust univariate and multivariate ARCH tests that do not depend on the model specification of the conditional variance in addition to the conditional mean are left for further study.

Funding

This research was supported by KAKENHI [Grant number: 16K03604].

Author details

Daiki Maki¹

E-mail: dmaki@mail.doshisha.ac.jp

Yasushi Ota²

E-mail: yota@mgt.ous.ac.jp

ORCID ID: <http://orcid.org/0000-0003-0534-0031>

¹ Faculty of Commerce, Doshisha University, Kyoto, 602-8580, Japan.

² Faculty of Management, Okayama University of Science, Okayama, Japan.

Citation information

Cite this article as: Robust tests for ARCH in the presence of a misspecified conditional mean: A comparison of nonparametric approaches, Daiki Maki & Yasushi Ota, *Cogent Economics & Finance* (2021), 9: 1862445.

Notes

1. Bühlmann and McNeil (2002) and Meister and Kreiß (2016) introduce nonparametric GARCH models. In addition, studies of nonlinear GARCH models include those of González-Rivera (1998), Pan et al. (2008), and Bauwens et al. (2010).
2. Catani and Ahlgren (2017) propose an LM test for ARCH using high-dimensional vector autoregressive models. In addition, Gel and Chen (2012) introduce bootstrap ARCH tests.
3. Other kernel functions include uniform, Epanechnikov, biweight, and triweight kernel functions. In general, while the type of kernel functions does not have a large impact on the estimation results, the selection of bandwidth significantly influences the estimation results.
4. Sheather and Jones (1991) propose another bandwidth selection that is based on the plug-in method. There are other bandwidth selection methods including those of Hart and Vieu (1990), Ziegler (2006), Li et al. (2013), Cheng et al. (2019), Jones et al. (1996), and Köhler et al. (2014) provide survey of bandwidth selection. The performance of the Nadaraya-Watson kernel regression strongly depends on the bandwidth selection. Since we only employ two bandwidth selections, the results in this study are somewhat limited. The comparison of robust ARCH tests using the Nadaraya-Watson kernel regression using various bandwidth is left for further study.
5. Shimizu (2014) introduces the estimation of nonparametric AR(1)-ARCH(1) using wild bootstrap. Shin and Hwang (2015) apply stationary bootstrap to estimate nonparametric AR(1)-ARCH(1).
6. If you are interested in simulation results of other levels of significance, these results are available upon

request. The results are similar to those of 5% level of significance.

7. Zambom and Kim (2017) propose lag selection in the nonparametric conditional heteroskedastic models. Compared to conventional methods, this method more appropriately selects lag length for various nonlinear models. We fix lag length in this study to investigate the statistical performance of the nonparametric regression models.

References

- Balke, A. P., & Kapetanios, G. (2007). Testing for ARCH in the presence of nonlinearity of unknown form in the conditional mean. *Journal of Econometrics*, 137(2), 472–488. <https://doi.org/10.1016/j.jeconom.2005.08.007>
- Bauwens, L., Preminger, A., & Rombouts, J. V. (2010). Theory and inference for a Markov switching GARCH models. *Econometrics Journal*, 13(2), 218–244. <https://doi.org/10.1111/j.1368-423X.2009.00307.x>
- Becker, R., & Hurn, S. (2009). Testing for nonlinearity in mean in the presence of heteroskedasticity. *Economic Analysis and Policy*, 39(2), 313–326. [https://doi.org/10.1016/S0313-5926\(09\)50023-9](https://doi.org/10.1016/S0313-5926(09)50023-9)
- Bühlmann, P., & McNeil, A. J. (2002). An algorithm for nonparametric GARCH modelling. *Computational Statistics & Data Analysis*, 40(4), 665–683. [https://doi.org/10.1016/S0167-9473\(02\)00080-4](https://doi.org/10.1016/S0167-9473(02)00080-4)
- Catani, P. S., & Ahlgren, N. J. C. (2017). Combined Lagrange multiplier test for ARCH in vector autoregressive models. *Econometrics and Statistics*, 1, 62–84. <https://doi.org/10.1016/j.ecosta.2016.10.006>
- Cheng, T., Gao, J., & Zhang, X. (2019). Nonparametric localized bandwidth selection for Kernel density estimation. *Econometric Reviews*, 38(7), 733–762. <https://doi.org/10.1080/07474938.2017.1397835>
- Engle, R. F. (1982). Autoregressive conditional heteroscedasticity with estimates of the variance of United Kingdom inflation. *Econometrica*, 50(4), 987–1007. <https://doi.org/10.2307/1912773>
- Exterkate, P., Groenen, P. J. K., Heij, C., & van Dijk, D. (2016). Nonlinear forecasting with many predictors using kernel ridge regression. *International Journal of Forecasting*, 32(3), 736–753. <https://doi.org/10.1016/j.ijforecast.2015.11.017>
- Gel, Y. R., & Chen, B. (2012). Robust Lagrange multiplier test for detecting ARCH/GARCH effect using permutation and bootstrap. *The Canadian Journal of Statistics*, 40(3), 405–426. <https://doi.org/10.1002/cjs.11149>
- González-Rivera, G. (1998). Smooth-transition GARCH models. *Studies in Nonlinear Dynamics & Econometrics*, 3(2), 61–78. <https://doi.org/10.2202/1558-3708.1041>
- Hart, J. D., & Vieu, P. (1990). Data-driven bandwidth choice for density estimation based on dependent data. *The Annals of Statistics*, 18(2), 873–890. <https://doi.org/10.1214/aos/1176347630>

- Jones, M. C., Marron, J. S., & Sheather, S. J. (1996). A brief survey of bandwidth selection for density estimation. *Journal of the American Statistical Association*, 91(433), 401–407. <https://doi.org/10.1080/01621459.1996.10476701>
- Katkovnik, V. Y. (1979). Linear and nonlinear methods of nonparametric regression analysis. *Soviet Automatic Control*, 5, 35–46.
- Köhler, M., Schindler, A., & Sperlich, S. (2014). A review and comparison of bandwidth selection methods for kernel regression. *International Statistical Review*, 82(2), 243–274. <https://doi.org/10.1111/insr.12039>
- Lee, J. H. H. (1991). A Lagrange multiplier test for GARCH models. *Economics Letters*, 37(3), 265–271. [https://doi.org/10.1016/0165-1765\(91\)90221-6](https://doi.org/10.1016/0165-1765(91)90221-6)
- Li, Q., Lin, J., & Racine, J. S. (2013). Optimal bandwidth selection for nonparametric conditional distribution and quantile functions. *Journal of Business and Economic Statistics*, 31(1), 57–65. <https://doi.org/10.1080/07350015.2012.738955>
- Lumsdaine, R. L., & Ng, S. (1999). Testing for ARCH in the presence of a possibly misspecified conditional mean. *Journal of Econometrics*, 93(2), 257–279. [https://doi.org/10.1016/S0304-4076\(99\)00011-1](https://doi.org/10.1016/S0304-4076(99)00011-1)
- McMillan, D. G. (2001). Nonlinear predictability of stock market returns: Evidence from nonparametric and threshold models. *International Review of Economics and Finance*, 10(4), 353–368. [https://doi.org/10.1016/S1059-0560\(01\)00093-4](https://doi.org/10.1016/S1059-0560(01)00093-4)
- Meister, A., & Kreiß, J. P. (2016). Statistical inference for nonparametric GARCH models. *Stochastic Processes and Their Applications*, 126(10), 3009–3040. <https://doi.org/10.1016/j.spa.2016.03.010>
- Nadaraya, E. A. (1964). On estimating regression. *Theory of Probability and Its Application*, 9(1), 141–142. <https://doi.org/10.1137/1109020>
- Pan, J., Wan, H., & Tong, H. (2008). Estimation and tests for power-transformed and threshold GARCH models. *Journal of Econometrics*, 142(1), 352–378. <https://doi.org/10.1016/j.jeconom.2007.06.004>
- Pavlidis, E., Paya, I., & Peel, D. (2010). Specifying smooth transition regression models in the presence of conditional heteroskedasticity of unknown form. *Studies in Nonlinear Dynamics and Econometrics*, 14(3), 1–40. <https://doi.org/10.2202/1558-3708.1702>
- Pavlidis, E., Paya, I., & Peel, D. (2013). Nonlinear causality tests and multivariate conditional heteroskedasticity: A simulation study. *Studies in Nonlinear Dynamics and Econometrics*, 17(3), 297–312. <https://doi.org/10.1515/snde-2012-0067>
- Péguin-Feissolle, A., Strikholm, B., & Teräsvirta, T. (2013). Testing the Granger noncausality hypothesis in stationary nonlinear models of unknown functional form. *Communications in Statistics-Simulation and Computation*, 42(5), 1063–1087. <https://doi.org/10.1080/03610918.2012.661500>
- Rudemo, M. (1982). Empirical choice of histograms and kernel density estimators. *Scandinavian Journal of Statistics*, 9(2), 65–78. <https://www.jstor.org/stable/4615859>
- Sheather, S. J., & Jones, M. C. (1991). A reliable data-based bandwidth selection method for kernel density estimation. *Journal of the Royal Statistical Society, Series B*, 53(3), 683–690. <https://doi.org/10.1111/j.2517-6161.1991.tb01857.x>
- Shimizu, K. (2014). Bootstrapping the nonparametric ARCH regression model. *Statistics & Probability Letters*, 87, 61–69. <https://doi.org/10.1016/j.spl.2014.01.002>
- Shin, D. W., & Hwang, E. (2015). Stationary bootstrapping for the nonparametric AR-ARCH model. *Communications for Statistical Applications and Methods*, 22(5), 463–473. <https://doi.org/10.5351/CSAM.2015.22.5.463>
- Silverman, B. W. (1986). *Density estimation for statistics and data analysis*. Chapman & Hall/CRC Monographs on Statistics & Applied Probability, Springer.
- Stone, C. J. (1977). Consistent nonparametric regression. *Annals of Statistics*, 5(4), 595–645. <https://doi.org/10.1214/aos/1176343886>
- Stone, C. J. (1984). An asymptotically optimal window selection rule for kernel density estimates. *Annals of Statistics*, 12(4), 1285–1297. <https://doi.org/10.1214/aos/1176346792>
- Van Dijk, D., Franses, P. H., & Lucas, A. (1999). Testing for ARCH in the presence of additive outliers. *Journal of Applied Econometrics*, 14(5), 539–562. [https://doi.org/10.1002/\(SICI\)1099-1255\(199909/10\)14:5<539::AID-JAE526>3.0.CO;2-W](https://doi.org/10.1002/(SICI)1099-1255(199909/10)14:5<539::AID-JAE526>3.0.CO;2-W)
- Watson, G. S. (1964). Smooth regression analysis. *Sankhya, Series A*, 26, 359–372. <https://www.jstor.org/stable/25049340>
- Zambom, A. Z., & Kim, S. (2017). Lag selection and model selection specification testing in nonparametric autoregressive conditional heteroscedastic models. *Journal of Statistical Planning and Inference*, 186, 13–27. <https://doi.org/10.1016/j.jspi.2017.02.004>
- Ziegler, K. (2006). On local bootstrap bandwidth choice in kernel density estimation. *Statistics and Decisions-International Journal Stochastic Methods and Models*, 24, 291–302. <https://doi.org/10.1524/stdn.2006.24.2.291>



© 2021 The Author(s). This open access article is distributed under a Creative Commons Attribution (CC-BY) 4.0 license.



You are free to:

Share — copy and redistribute the material in any medium or format.

Adapt — remix, transform, and build upon the material for any purpose, even commercially.

The licensor cannot revoke these freedoms as long as you follow the license terms.

Under the following terms:

Attribution — You must give appropriate credit, provide a link to the license, and indicate if changes were made.

You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.

No additional restrictions

You may not apply legal terms or technological measures that legally restrict others from doing anything the license permits.

***Cogent Economics & Finance* (ISSN: 2332-2039) is published by Cogent OA, part of Taylor & Francis Group.**

Publishing with Cogent OA ensures:

- Immediate, universal access to your article on publication
- High visibility and discoverability via the Cogent OA website as well as Taylor & Francis Online
- Download and citation statistics for your article
- Rapid online publication
- Input from, and dialog with, expert editors and editorial boards
- Retention of full copyright of your article
- Guaranteed legacy preservation of your article
- Discounts and waivers for authors in developing regions

Submit your manuscript to a Cogent OA journal at www.CogentOA.com

