

Eibich, Peter; Zai, Xianhua

**Working Paper**

## Are the Grandparents Alright?: The Health Consequences of Grandparental Childcare Provision

CINCH Series, No. 2022/03

**Provided in Cooperation with:**

CINCH – Health Economics Research Center, Universität Duisburg-Essen (UDE)

*Suggested Citation:* Eibich, Peter; Zai, Xianhua (2022) : Are the Grandparents Alright?: The Health Consequences of Grandparental Childcare Provision, CINCH Series, No. 2022/03, University of Duisburg-Essen, CINCH - Health Economics Research Center, Essen, <https://doi.org/10.17185/dupublico/77108>

This Version is available at:

<https://hdl.handle.net/10419/270601>

**Standard-Nutzungsbedingungen:**

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.


Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

**Terms of use:**

*Documents in EconStor may be saved and copied for your personal and scholarly purposes.*

*You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.*

*If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.*



CINCH+

competence in competition + health

series

#2022/03

Peter Eibich and Xianhua Zai

**Are the Grandparents Alright? The Health  
Consequences of Grandparental Childcare Provision**



# Imprint

## EDITOR-IN-CHIEF

Martin Karlsson, Essen

## MANAGING EDITOR

Katharina Blankart, Essen

## EDITORIAL BOARD

Boris Augurzky, Essen

Daniel Avdic, Melbourne (AUS)

Jeanette Brosig-Koch, Magdeburg

Stefan Felder, Basel

Annika Herr, Hannover

Nadja Kairies-Schwarz, Düsseldorf

Hendrik Schmitz, Paderborn

Harald Tauchmann, Erlangen-Nürnberg

Jürgen Wasem, Essen

## CINCH SERIES

CINCH – Health Economics Research Center  
Weststadttürme, Berliner Platz 6-8  
45127 Essen

[www.cinch.uni-due.de](http://www.cinch.uni-due.de)

[cinchseries@cinch-essen.de](mailto:cinchseries@cinch-essen.de)

Phone +49 (0) 201 183 - 3679

Fax +49 (0) 201 183 - 3716

All rights reserved. Essen, Germany, 2014

ISSN 2199-8744 (online)

The working papers published in the Series constitute work in progress circulated to stimulate discussion and critical comments. Views expressed represent exclusively the authors' own opinions and do not necessarily reflect those of the editors.

#2022/03

Peter Eibich and Xianhua Zai

## Are the Grandparents Alright? The Health Consequences of Grandparental Childcare Provision



Peter Eibich<sup>a</sup> and Xianhua Zai<sup>b</sup>

## Are the Grandparents Alright? The Health Consequences of Grandparental Childcare Provision

### Abstract

*This paper examines the causal effect of childcare provision on grandparents' health in the US. We propose the sex ratio among older adults' children as a novel instrument for grandparental childcare provision. Our instrument is rooted in the demographic literature on grandparenthood and exploits that parents of daughters transition to grandparenthood earlier and invest more in their grandchildren than parents of sons. We estimate 2SLS regressions using data from the Health and Retirement Study. The results suggest that childcare provision is not beneficial for grandparents' health and may even be detrimental for physical functioning and subjective health. Specifically, grandparenting worsens functional limitations of IADL by 26 percent and subjective health by 9.5 percent.*

**Keywords:** Grandparents, Childcare provision, Instrumental variables, Health.

**JEL classification:** I10, J13, J14, C26.

---

<sup>a</sup> Max Planck Institute for Demographic Research. Email: [eibich@demogr.mpg.de](mailto:eibich@demogr.mpg.de).

<sup>b</sup> Max Planck Institute for Demographic Research. Email: [zai@demogr.mpg.de](mailto:zai@demogr.mpg.de).

# 1 Introduction

Grandparents across the globe play an important role in raising their grandchildren. For example, the US Census Bureau estimates that in 2011 4.8 million children under 5 ( 24%) received care from their grandparents (Laughlin 2013). In the UK, around 40% of grandparents provide regular care for their grandchildren, and 89% of these provide care at least once a week (Age UK 2017). In the EU, 21% of children under 3 received some childcare from sources other than their parents or formal childcare in 2020 with substantial variation across countries (Eurostat 2022).<sup>1</sup> Grandparental childcare provision can reduce the cost of childrearing for young parents by substituting for formal care or own childcare provision, in particular in contexts with strong family ties (Battistin et al. 2014; Chen et al. 2000). Even in contexts where formal childcare is both available and affordable, grandparents often make important contributions by offering a flexible alternative source of childcare, e.g., in case of illness or during school holidays.

While grandparenting is highly beneficial to parents (Compton 2015), the consequences for grandparents themselves are less clear. Looking after grandchildren might provide grandparents with physical and mental stimulation, thereby helping to maintain their health in old age. This would imply that grandparenting can be considered as “active ageing”, i.e., an activity that benefits both older individuals and the wider society. Yet, keeping up with young children can also be physically strenuous and stressful. The negative health effects of informal care provision by older parents or spouses have been documented extensively in the literature (Bom et al. 2018; Bom and Stöckel 2021; Heger 2017; Schmitz and Westphal 2015). It seems possible that grandparents find caring for young children similarly demanding. Therefore, the overall effect of grandchild care provision on health of grandparents is ambiguous. In this paper, we empirically estimate the health effects of grandparenting for grandparents.

Previous studies on grandchild care provision and grandparents’ health report

---

<sup>1</sup>Formal childcare here includes grandparents, other household members, and professional child carers.

contradictory findings. Several studies report that grandparents caring for their grandchildren are in better health, have fewer mobility limitations and fewer depressive symptoms (Danielsbacka et al. 2019; Di Gessa et al. 2016; Ku et al. 2013; Wang et al. 2020; Zeng et al. 2021). Yet, interpreting these estimates as causal is challenging. First, the transition to grandparenthood is not random. Lai et al. (2021) report that older adults expecting to become grandparents in the future are healthier than those who do not expect this transition. A possible explanation is that healthier individuals are, *ceteris paribus*, able to have more children than individuals in poor health, which in turn means that they are more likely to have grandchildren. Moreover, healthier parents are more likely to survive until they become grandparents and their lifespan overlaps longer with their grandchildren’s lifespan (Margolis and Verdery 2019). Second, health is an important precondition for all activities in old age such that grandparents in poor health are less capable to provide grandchild care. Taken together, these arguments suggest that grandparents providing childcare are positively selected on health.

A few previous studies address the endogeneity of grandchild care provision. Ates (2017) finds that the positive association between grandparenting and health in Germany disappears when introducing individual-fixed effects. While fixed effects can resolve bias from selection on time-invariant unobservable characteristics (e.g., long-term health conditions or family size), it does not address potential reverse causality introduced by an unexpected health shock that reduces a grandparent’s capacity to provide childcare. Brunello and Rocco (2019) and Ku et al. (2012) use instrumental variables (IVs) to address such endogeneity. Brunello and Rocco (2019) use data on European grandparents from the Survey of Health, Ageing and Retirement in Europe (SHARE). Their IV strategy exploits variation in the propensity of grandchild care provision due to the random timing of the survey and changes in the likelihood of grandchild care provision by the age of grandchildren. They find a sizable increase in depressive symptoms for grandparents providing childcare. Ku et al. (2012) examine Taiwanese grandparents and use marital



status of parents and the number of grandchildren as IVs. Their findings indicate that even after addressing the endogeneity, grandchild care provision is beneficial for Taiwanese grandparents' health.

This study examines the causal effect of grandparental childcare provision on grandparents' health in the US. We propose the sex ratio (defined as the number of daughters relative to the total number of children) as a novel instrument to address the endogeneity of grandchild care provision. Our IV is motivated by two insights from the demographic literature on grandparenthood. First, parents of daughters transition to grandparenthood earlier than parents of sons. Second, grandparents are more likely to provide childcare for grandchildren born to a daughter than for those born to a son. We use data from the Health and Retirement Study (HRS) with detailed health information on the number of functional limitations, self-rated health status, and depressive symptoms. Our results indicate that the effects of grandparenting on health are not significant and, in some instances, negative. We therefore conclude that grandparental childcare provision should not be considered as "active ageing". Similar to informal care provision, grandparents' care for their grandchildren can be detrimental to their own health.

This study contributes to the literature by estimating a credibly identified causal effect of grandchild care provision on grandparents' health. We propose a novel instrument motivated by the demographic literature on grandparenthood, and we conduct a battery of tests and falsification exercises that suggest that the IV assumptions likely hold. Our paper is also the first study to provide causal evidence in the US context. The contradictory findings by [Brunello and Rocco \(2019\)](#) and [Ku et al. \(2012\)](#) suggest that the health effects of grandparenting might be context-dependent. The US is a particularly interesting context characterized by both expensive formal childcare compared to some of the European countries examined by [Brunello and Rocco \(2019\)](#) and weaker family ties compared to East Asian societies ([Ku et al. 2012](#)).

The remainder of the paper is organized as follows. Section 2 describes the data, key

variables, and sample statistics. Section 3 first motivates our novel instrument, then describes our estimation strategy, and discusses the assumptions for our IV model. Section 4 presents our main results as well as several robustness checks. Section 5 discusses our findings and concludes.

## 2 Data

### 2.1 Sample Description

We use data from the HRS, a nationally representative longitudinal study of Americans aged 51 and above. Respondents are surveyed every other year since 1992. The survey includes different birth cohorts who enter the study as they become eligible. The core cohort, the HRS cohort, has been followed and interviewed since 1992. Since 1993, the HRS has included the Study of Assets and Health Dynamics Among the Oldest Old (AHEAD) cohort of individuals born before 1924; the Children of the Depression Age (CODA) cohort of people born between 1924 and 1930; and the War Babies cohort (WB) of individuals born between 1942 and 1947. An additional Early Baby Boomers (EBB) cohort of people born between 1948 and 1953 was added to the sample in 2004, and the Mid-Baby Boomers cohort of individuals born between 1954 and 1959 was added in 2010. [Liu and Zai \(2022\)](#) document the details on how respondents enter each survey wave and the number of unique individuals in each wave.

The HRS asks respondents (including cohabiting spouses) detailed information about their own demographic characteristics, health, employment, financial situation, and intergenerational transfers as well as demographic information about their family members such as children and parents. To explore the effect of grandparenting on grandparents' health, we restrict our working sample to HRS respondents aged between 40 to 70 who have at least one child. On the one hand, we aim to include as many potential grandparents as possible to maximize the sample size. On the other hand, we are concerned about the validity of our instrument if we include individuals older than 70.

This older population is more likely to be frail and dependent and thus not able to provide grandchild care. We check the sensitivity of our results to this age restriction in section 4.2 using a sample without age limits. The distribution of age of respondents in Appendix Figure A1 is almost symmetric around 70.

Our study sample includes 84,483 observations (20,508 unique individuals) and covers the period from 1996 (wave 3) to 2014 (wave 12) in which the HRS asks respondents consistent questions on grandparenting.

## 2.2 Dependent Variables

The HRS includes detailed information on the health outcomes of respondents. We mainly focus on three dimensions of health: self-reported health status, physical health, and mental health.

First, the HRS asks respondents to self-report their general health status. Possible answers range from 1 for “excellent”, 2 for “very good”, 3 for “good”, 4 for “fair”, to 5 for “poor”. We create an indicator for poor self-reported health, which is one if self-reported health is “fair” or “poor”, and is zero otherwise. Similar indicators of fair or poor health are also employed in Dave et al. (2006) using the HRS data, in Eibich (2015) using the German Socio-Economic Panel Study (SOEP), and in Kuka (2020) using the Survey of Income and Program Participation (SIPP). We also consider alternative cut-off points to show robustness in section 4.2. While self-reported health is subjective and might be affected by reporting heterogeneity, it is a good predictor of mortality (Idler and Benyamini 1997; DeSalvo et al. 2006; Kuka 2020).

Second, we use more objective measures about physical health conditions. The HRS provides indices of functional limitations, such as limitations in Activities of Daily Living (ADLs) and limitations in Instrumental Activities of Daily Living (IADLs). The ADLs include items such as bathing, eating, dressing, getting in or out of bed, and walking across a room and the IADLs assess difficulties in using the phone, managing money, taking

medications, shopping for groceries, and preparing hot meals.<sup>2</sup> All these indices range from 0 to 5. An index with a value of 5 means that an individual has difficulties with all activities considered, while a value of zero means that the individual has no limitations. We create dichotomous indicators which equal one if an individual reports limitations for four or more items, and zero otherwise. We consider other operationalizations as robustness checks in section 4.2.

Third, we further use information about respondents' mental health. The HRS asks respondents about their mental health using the Center for Epidemiologic Studies Depression (CES-D) score. The CES-D score captures the number of adverse sentiments a respondent experienced all or most of the time in the past two years, including whether an individual was depressed, felt alone, felt sad, had restless sleep, felt everything was an effort, could not get going, felt unhappy, and did not enjoy life. The CES-D scale has been validated as an instrument to identify major depression in older adults (Irwin et al. 1999). The main results in section 4 reports the effect of grandparenting on a binary indicator constructed using five or more depressive symptoms as the cutoff (8 items in total). Other alternative measures are considered as robustness checks in section 4.2.

## 2.3 Treatment Indicators

Our treatment variable of interest is whether individuals provide grandchild care or not. There are two relevant grandparenting questions in the data. First, the HRS asks respondents whether they and their spouse spent 100 or more hours taking care of their grandchildren or great-grandchildren since the last wave.<sup>3</sup> If the answer is yes, respondents are asked to which child they provided grandchild care. This question was not asked in waves 1 and 2.<sup>4</sup>

---

<sup>2</sup>More details on the construction of these measures can be found in Chien et al. (2015).

<sup>3</sup>The question asked in the HRS is "Did you or your husband/wife/partner or your late husband/late wife/late partner spend 100 or more hours in total in the last two years taking care of great-grandchildren/grandchildren?"

<sup>4</sup>In wave 2 of the HRS, the AHEAD cohort was asked whether grandchild care was provided for a year or longer. This question is no longer asked from wave 3. The question is "Which of your children is the parent of those grandchildren (or great-grandchildren)?"

Second, the HRS asks respondents to estimate their childcare hours provided in the last two years. This question is asked separately for the respondent and the spouse.<sup>5</sup> For those who cannot remember the hours or do not know the exact hours or refuse to give the number of hours on grandparenting, the HRS further asks the minimum and maximum values of hours of grandchild care provided.<sup>6</sup> There are both advantages and disadvantages of using each question to construct our treatment indicator of grandparenting. The first question does not distinguish between grandchild care provided by respondents or their spouses, which would introduce measurement errors since we are interested in estimating the health effects on those who are actually grandparenting. On the other hand, the question only requires respondents to answer “yes” or “no” and might thus be less affected by recall bias than asking for the exact number of grandchild care hours provided over the last two years.

For our main analysis, we use the self-reported number of hours of grandchild care by respondents. Among those who are grandparenting, the majority of grandparents provide less than 1,000 hours over two years. Appendix Figure A2 shows the distribution of grandchild care hours reported by HRS respondents for these grandparents. We construct a binary indicator of grandparenting status, which indicates whether the respondent reported 100 or more hours of grandchild care over the last two years. If the number of hours is missing and the minimum and maximum values are above 100, we assume that the respondent is grandparenting. To examine the potential measurement errors in the treatment variable, we use the first question on childcare provided by the respondent and their partner to construct an alternative treatment indicator, which is 1 if the answer is “yes”, i.e., the respondent and their partner provided at least 100 hours of childcare since the last wave. We also explore other cutoffs for the self-reported number of childcare hours as robustness checks in section 4.2.

---

<sup>5</sup>The grandchild care hour question in the HRS is “Roughly how many hours altogether did you spend since the last wave?” for the respondent and “Roughly how many hours altogether did your husband/wife/partner spend since the last wave?” for respondent’s spouse.

<sup>6</sup>The quote in HRS is “Did it amount to a total of less than MAX BREAKPOINT, more than MIN BREAKPOINT, or what?” The MIN BREAKPOINTS are 0, 200, 201, 500, and 501. The MAX BEAKPOINTS are 199, 200, 499, 500, and 5,000.

## 2.4 Sample Statistics

Table 1 presents the summary statistics of the working sample of HRS respondents who are between 40 to 70 in each survey year. The average age of the sample is around 61. About 60 percent of the sample are female. The average educational attainment of the sample is around 12 years. On average, each respondent has between three to four children. The oldest child is on average about 40 years old and the youngest child is on average about 31 years old. About half of the respondents' children are daughters. The majority of the sample is married or living with a partner and white. Approximately 33 to 39 percent of respondents provide some grandchild care according to the different definitions discussed earlier.

The average self-reported health status of respondents is good and about 30 percent of respondents report fair or poor health status. The average ADL score and IADL score are both close to zero, which indicates that our sample is relatively healthy. The share of respondents with more than three functional limitations in ADL or IADL is about 2 percent. The average CESD depression score is 1.5 out of 8 and 12 percent of individuals have four or more depressive symptoms. Detailed definitions of these variables are provided in Appendix Table A1.

## 3 Methods

In this section, we review findings from the demographic literature on grandparenthood and grandparenting to motivate the sex ratio as our valid IV for providing grandchild care. Then we discuss the estimation strategy and provide evidence for the assumptions required for a causal interpretation in the IV framework.

### 3.1 Sex Ratio as an IV for Grandparenting

The transition to grandparenthood as well as the decision to provide grandchild care are endogenous choices, which depend on many factors that are plausibly related to health.

Table 1: Summary Statistics of the Sample

Variable	Mean	S.D.	Obs.
<b><i>Demographics</i></b>			
Age	60.89	5.98	84,483
Female	0.60	0.49	84,483
Education (years)	12.24	3.13	84,268
Number of children	3.72	2.02	84,483
Age of youngest child	30.54	8.37	84,188
Age of oldest child	39.75	6.99	84,188
Income from earnings (\$1,000)	16.41	33.40	84,483
Income from pensions (\$1,000)	3.91	20.68	84,483
Marital status			
Married/partnered	0.76	0.43	84,403
Separated/divorced	0.13	0.34	84,403
Widowed	0.10	0.29	84,403
Never married	0.02	0.12	84,403
Race/ethnicity			
White	0.75	0.43	84,348
Black/African	0.18	0.39	84,348
Other	0.07	0.25	84,348
<b><i>Instrumental variable</i></b>			
Sex ratio	0.50	0.29	84,354
<b><i>Grandparenting</i></b>			
Grandparenting for at least one child (Q1)	0.39	0.49	84,382
Grandparenting for at least 100 hours (Q2)	0.34	0.47	84,483
<b><i>Health variables</i></b>			
Self-report health	2.82	1.12	84,444
Self-report fair or poor health	0.27	0.45	84,444
ADL limitations	0.26	0.80	84,426
Percentage of having ADL limitations (3+)	0.02	0.13	84,426
IADL limitations	0.19	0.67	84,420
Percentage of having IADL limitations (3+)	0.01	0.11	84,420
CESD score	1.54	2.05	79,477
Percentage of having depressive emotions (4+)	0.12	0.32	79,477

Notes: The data used are from the HRS 1996 to 2014 of individuals who are 40 to 70. The definitions of these variables can be found in Appendix Table A1.

For example, parents with larger families are more likely to become grandparents (Margolis and Verdery 2019), and parents who give birth earlier in life are more likely to become grandparents at younger ages. Family size and age at the first birth have been linked to health and mortality of mothers in particular (Mirowsky 2005; Wu and Li 2012), but they are also related to socioeconomic status (Adserà 2017). Beyond the transition to grandparenthood, grandparents' capacity to provide grandchild care depends, among other factors, on the proximity between grandparents and their adult children (Compton 2015), as well as their health (Eibich and Siedler 2020).


In this study, we address the endogeneity of grandparenting by using the sex ratio, defined as the number of daughters divided by the total number of children of a respondent, as an instrument for grandparenting. Our choice of this instrument is motivated by a few stylized facts derived from the demographic literature on fertility and grandparenthood.

It is well-documented that women tend to give birth earlier than men (Margolis and Verdery 2019). This implies, *ceteris paribus*, that parents of daughters will transition to grandparenthood earlier than parents of sons. The gender of a child can be considered as good as randomly determined, thus the gender of a person's first-born child may serve as a suitable instrument that predicts the transition into grandparenthood (and subsequently grandparenting) (Rupert and Zanella 2018). This is also borne out in our data: Fig. 1 shows the share of individuals who are grandparenting by age for individuals with a first-born daughter and those with a first-born son, respectively. At most younger ages, older adults with a first-born daughter are much more likely to provide grandchild care than those with a first-born son. This gap narrows substantially with age and mostly disappears beyond age 70.

Second, older adults with a higher number of daughters are overall more likely to become grandparents and they transition into grandparenthood at younger ages (see Fig. 2). This is in part because the likelihood of becoming a grandparent increases with the number of children (Margolis and Verdery 2019). Some children may remain childless (by choice or



Figure 1: Sex of the first-born child and grandparenting




Notes: The data is the HRS from 1996 to 2014. The sample is limited to individuals with only one child. This graph draws the share of individuals who provide some grandchild care by age for individuals with a first-born daughter and those with a first-born son, respectively. Grandparenting is defined as an indicator that is 1 if the estimated grandchild care hours reported by respondents are at least 100 hours.

involuntarily). Even if the probability of remaining childless is correlated between siblings, we would expect that the likelihood of all children remaining childless is smaller for an older parent with three children than for an older parent with one child.


Third, women show less variation in age at first birth than men (Margolis and Verdery 2019), thus older parents with a first-born son and a second-born daughter might still on average transition into grandparenthood earlier than older parents with two sons. Fig. 3 shows the likelihood of grandparenthood for older adults with two children. Until about age 60, the likelihood of becoming a grandparent for parents with two daughters is considerably higher than for all other groups. The likelihood is slightly higher for parents with a first-born daughter and a second-born son than for parents with a first-born son and a second-born daughter, although differences between these two groups are relatively small. The likelihood of becoming a grandparent tends to be the lowest for parents of two sons, although the differences between groups largely vanish from age 70 onward.

Figure 2: Grandparenthood and the number of daughters



Notes: The data is the HRS from 1996 to 2014. This graph draws the share of individuals who are grandparents by age for all individuals with at least one child. Grandparenthood is defined as an indicator that is 1 if the number of grandchildren reported by the respondent is at least one.

Figure 3: Grandparenthood for older parents with two children




Notes: The data is the HRS from 1996 to 2014 with the respondents limited to having two children. This graph draws the share of individuals who are grandparents by age for individuals with two children. Grandparenthood is defined as an indicator that is 1 if the number of grandchildren reported by the respondent is at least one.

In addition, the sex of a child not only affects their parents' likelihood to become grandparents, but also the extent of their involvement with the grandchild. Maternal grandparents invest more time into grandchild care than paternal grandparents (Compton and Pollak 2011; Danielsbacka et al. 2011). Several explanations have been proposed for this phenomenon – uncertainty around paternal kinship (Danielsbacka et al. 2011), stronger bonds between mothers and daughters (Somary and Strieker 1998), and a longer shared lifetime between maternal grandparents and their grandchildren (Margolis and Verdery 2019). Somary and Strieker (1998) find few differences in grandparenting behavior across lineage, but note that they control for proximity between grandparents and grandchildren, which plays an important role for grandchild care investments (Compton 2015). We find a similar pattern in our data. While there is considerable variation for younger grandparents, from age 60 onwards until about age 75 we observe that older parents with two daughters are most likely to provide grandchild care, whereas grandparents with two sons tend to be the least likely group to provide grandchild care (Fig. 4).

In summary, the literature suggests that the number of children, the sex of children as well as their birth order are related to the likelihood of becoming a grandparent, the age at the transition into grandparenthood as well as grandparental investment into their children. The number of children itself is endogenous and might be related to health of the older parents (Wu and Li 2012). Similarly, the absolute number of daughters is clearly correlated with family size and therefore not a suitable instrument. On the other hand, the sex of the first-born child (“birth order instrument”) is plausibly exogenous, but exploits very limited variation. We therefore propose to use the sex ratio as an instrument. In contrast to the absolute number of daughters, the sex ratio is not correlated with family size ( $\rho = -0.0029$  in our sample), because we essentially condition on the total number of children. The sex ratio does not draw on variation in the birth order of children, but the instrument accounts for variation stemming from parents of daughters becoming grandparents earlier on average than parents of sons and being more likely to provide grandchild care, regardless of the

Figure 4: Grandparenting for older parents with two children



Notes: The data is the HRS from 1996 to 2014 with respondents limited to having two children. This graph draws the share of individuals who are grandparents by age for individuals with two children. Grandparenting is defined as an indicator that is 1 if the estimated grandchild care hours reported by respondents are at least 100 hours.

number of children they have. For example, the birth order instrument and the sex ratio instrument exploit the same variation – parents of a daughter as compared to parents of a son – for parents with one child. For parents with two children, the birth order instrument only distinguishes between parents with a first-born daughter and parents with a first-born son, regardless of the sex of the second child. The sex ratio instrument distinguishes parents with two sons from parents with one daughter and those with two daughters, under the assumption that parents with two daughters are most likely to provide grandchild care and parents with two sons are least likely to provide grandchild care. Since 88% of the HRS sample has two or more children, the sex ratio instrument provides much more variation than the birth order instrument. Therefore we expect that the sex ratio instrument should be stronger than the birth order instrument.

## 3.2 Model Specification

Using the sex ratio as an instrument for grandparenting, we estimate the first stage of our IV model as follows:

$$Grandparent_{it} = \delta Sexratio_{it} + X'_{it}\beta + \epsilon_{it} \quad (1)$$

where  $Grandparent_{it}$  is the grandparenting status of an individual  $i$  in year  $t$ .  $Sexratio_{it}$  is the ratio of the number of daughters to the number of children of an individual  $i$  in year  $t$ .  $X_{it}$  is a vector of covariates. In our preferred specification, we control for individual demographic characteristics such as age (quadratic polynomial), race/ethnicity, religion, gender, birth place and census region fixed effects as well as year-fixed effects, fixed effects for the year of birth of the first-born child, age of the youngest child, cohort fixed effects of the individual, family size and control variables for the economic situation such as the annual income from earnings<sup>7</sup> and the annual income from all pensions and annuities.  $\epsilon_{it}$  are the standard errors clustered at the individual level which allow for correlation within individuals across waves.

We estimate the effect of grandparenting on health in the second stage of the model as follows:

$$Y_{it} = \alpha Grandparent_{it} + X'_{it}\eta + \mu_{it} \quad (2)$$

where  $Y_{it}$  is an indicator of the health status of individual  $i$  in year  $t$ . The other controls  $X_{it}$  are the same as in equation 1. We estimate our IV model using linear two-stage least squares estimation (2SLS).

## 3.3 IV Assumptions

The interpretation of our IV estimates as causal effects requires three assumptions: (i) reliability, i.e., the sex ratio should be correlated with grandparenting, (ii) validity, i.e., the sex ratio should be as good as randomly assigned and should not affect health through any

---

<sup>7</sup>This includes salaries, bonuses, overtime pay, commissions or tips, second job earnings, and professional practice or trade income.

other mechanisms than through grandchild care provision, and (iii) monotonicity, i.e., the sex ratio should affect the likelihood of grandparenting in the same direction (non-negative in this case) for all observations in our sample. In this section, we will discuss the plausibility of these assumptions in details.

### 3.3.1 Reliability

Table 2 shows estimates of the first-stage using equation 1. In column 1, we regress our indicator of grandparenting only on the sex ratio instrument. The estimate suggests that *ceteris paribus* older adults that only have daughters (i.e., a sex ratio of 1) are 6.4 percentage points more likely to provide grandchild care than older adults with only sons (i.e., a sex ratio of 0). For parents with two children, this would imply that every daughter increases the likelihood of grandparenting by 3.2 percentage points. In columns 2-6, we successively add control variables to account for standard demographic characteristics as well as socioeconomic differences of grandparents. The point estimate of the sex ratio instrument is barely affected by the introduction of these controls. The Kleibergen-Paap F-statistic on the strength of the excluded instrument is larger than 53 in all models, which exceeds thresholds that have traditionally been used as a rule-of-thumb. This suggests that the sex ratio is indeed a sufficiently strong predictor of grandparenting, i.e., the reliability assumption holds.

### 3.3.2 Validity

We conduct two falsification exercises to detect potential violations of the validity assumption of our instrument. First, we check for covariate balancing. If the sex ratio is as good as randomly assigned, we would expect that the distribution of covariates that are not affected by the treatment should be similar across the different values of the instrument, or put differently, there is no significant correlation between such covariates and our instrument. If we would find a significant correlation, this would imply that the instrument might be affected

Table 2: First stage estimates

<i>Dependent variable: Grandparenting</i>						
Model	1	2	3	4	5	6
Sex ratio	0.064*** (0.009)	0.064*** (0.009)	0.064*** (0.009)	0.063*** (0.009)	0.063*** (0.009)	0.063*** (0.009)
Year FE + Birth year FE of first-born		Y	Y	Y	Y	Y
Cohort FE + Birth year FE of youngest-born			Y	Y	Y	Y
Demographics				Y	Y	Y
Family size					Y	Y
Socioeconomic controls						Y
Mean of dependent variable	0.34	0.34	0.34	0.34	0.34	0.34
Observations	84,354	84,276	84,187	83,709	83,709	83,709
1st stage Kleibergen-Paap F-stat.istic	53.4	53.6	54.3	53.9	53.9	53.9


Notes: The data used are from the HRS 1996 to 2014 of individuals who are 40 to 70. Each cell reports estimates from a separate specification using equation 1. The sex ratio is defined as the number of daughters divided by the total number of children of an individual. Grandparenting is defined as an indicator that is 1 if the estimated grandchild care hours reported by respondents are at least 100 hours. Column 1 reports estimates without any controls. Column 2 adds year-fixed effects and fixed effects for the year of birth of the first-born child of an individual. Column 3 adds age of the youngest child and cohort fixed effects of the individual. Column 4 includes individual demographics such as age (quadratic polynomial), race, religion, gender, birth place and census region fixed effects. Column 5 further controls for the number of children of individuals. Column 6 adds socio-economic controls such as the annual earnings from wages or salary, bonuses, second job income, and professional practice income as well as the annual sum of pensions and annuities. Standard errors are clustered at the individual level. \*\*\*  $p < 0.01$ , \*\*  $p < 0.05$ , \*  $p < 0.10$ .

by selection. We regress a battery of covariates as dependent variables on our sex ratio IV controlling for (i) year and first-born fixed effects (Model 2 of Table 2), (ii) additional cohort fixed effects for the individual, age fixed effects of the youngest child as well as individual demographic characteristics (Model 4 of Table 2), and (iii) a full specification including also annual pension income and earnings (Model 6 of Table 2).

Figure 5 shows the point estimates and confidence intervals of the regression coefficients on the sex ratio for each dependent variable listed on the vertical axis in three specifications. For all covariates, estimates remain similar across models and none of these estimates is significant at the 95% confidence level, i.e., there is no evidence for a potential violation of the validity assumption of our IV.

We conduct another falsification exercise following Angrist et al. (2010). We estimate the reduced form regression for a subsample of HRS respondents without grandchildren. In

Figure 5: Covariate balance for the sex ratio instrument



Notes: The data used are from the HRS 1996 to 2014 of individuals who are 40 to 70. The graph shows the point estimates and confidence intervals of the regression coefficients on the sex ratio for each covariate as the dependent variable listed on the vertical axis in models 2, 4, 6 of Table 2. The sex ratio is defined as the number of daughters divided by the total number of children of an individual. Grandparenting is defined as an indicator that is 1 if the estimated grandchild care hours reported by respondents are at least 100 hours.

this subsample, the sex ratio instrument is not supposed to predict grandparenting (i.e., the treatment) since none of the respondents have grandchildren. The exclusion restriction requires that the instrument is associated with the outcome only through its effect on the treatment. This implies that there should be no significant relationships between the sex



ratio instrument and the health outcomes in the reduced form regression for individuals without grandchildren, because there is no valid first stage in this subsample. A significant reduced form estimate signals a violation of the IV validity because it would suggest that the sex ratio is related to health through pathways other than grandparenting.

Table 3 reports the reduced form estimates from our preferred model specification for the subsample of respondents who do not have grandchildren. None of the estimates is statistically significant, and the point estimates are very small and close to zero (the reduced form estimates for our working sample are shown in Appendix Table A2 for comparison).

Table 3: Falsification exercise: Reduced form regression

Dependent variable	ADL	IADL	Poor subjective health	Depressed
Sex ratio	0.002 (0.006)	-0.004 (0.005)	0.017 (0.019)	-0.009 (0.015)
Mean of dependent variable	0.021	0.014	0.302	0.121
Number of clusters	5,904	5,902	5,907	5,460
Observations	6,393	6,391	6,395	5,915

Notes: The data used are from the HRS 1996 to 2014 of the subsample of respondents who do not have grandchildren. All models control for year-fixed effects, fixed effects for the year of birth of the first-born child of an individual, age of the youngest child, cohort fixed effects of the individual, individual demographics such as age (quadratic polynomial), race, religion, gender, birth place and census region fixed effects, the number of children of individuals, and socio-economic controls such as the annual earnings from wages or salary, bonuses, second job income, and professional practice income as well as the annual sum of pensions and annuities. Standard errors are clustered at the individual level. \*\*\*  $p < 0.01$ , \*\*  $p < 0.05$ , \*  $p < 0.10$ .

Neither of these two falsification exercises provides evidence that the validity assumption is violated. While we cannot rule out such violations, the results improve our confidence that the sex ratio is a valid instrument for grandchild care provision.

### 3.3.3 Monotonicity Assumption

We test the monotonicity assumption by re-estimating the first stage regression of equation 1 for different subgroups within our working sample. The monotonicity assumption is satisfied if our instrument affects treatment in the same direction for all observations in the sample.

This implies that the estimated effect of the sex ratio on grandparenting should be positive or zero for any arbitrarily defined subsample within our working sample. A significant negative effect in the first stage would imply a violation of monotonicity.<sup>8</sup>

Table 4 shows estimates of the first-stage regression for ten different subsamples defined by demographic characteristics. We note that the size of the first-stage estimate varies considerably – between 3.3 percentage points for men and 8.9 percentage points for older adults with more than four siblings. This suggests that the compliers differ from the overall population, but it does not suggest any violation of monotonicity. For all subsamples, the sex ratio instrument increases the likelihood of grandparenting and estimates are strongly significant.

In summary, these tests and falsification exercises suggest that the three assumptions required for a causal interpretation of our IV estimates are likely to hold.

## 4 Results

### 4.1 The effect of grandparenting on health

Before estimating the causal effect of grandparenting on health using the sex ratio instrument, we examine the relationship between grandparenting and health using ordinary least squares regression (OLS). Comparing OLS and 2SLS estimates will provide an indication of the size and direction of the bias caused by the endogeneity of grandparenting. Table 5 shows the results for our preferred model specification for all four health outcomes. Note that all health indicators are binary measures where a value of 1 represents worse health outcomes.

Overall, the results of Table 5 show that grandparenting is associated with better health across all dimensions of health considered here. The estimates suggest that grandparents caring for their grandchildren have fewer limitations in ADL and IADL, they

---

<sup>8</sup>A negative but insignificant point estimate might either reflect a true zero effect (which does not violate monotonicity) or a violation of monotonicity.

Table 4: Falsification exercise: Monotonicity

Subsample	<i>Dependent variable: Grandparenting</i>									
	Male	Female	$\leq 4$ Children	$> 4$ Children	$\leq 4$ Siblings	$> 4$ Siblings	Married	Not Married	White	Not White
Sex ratio	0.033*** (0.013)	0.082*** (0.011)	0.063*** (0.009)	0.058** (0.023)	0.054*** (0.010)	0.089*** (0.016)	0.058*** (0.010)	0.084*** (0.014)	0.055*** (0.010)	0.087*** (0.016)
Mean of dependent variable	0.285	0.377	0.349	0.318	0.341	0.339	0.349	0.313	0.343	0.332
Number of clusters	8,664	11,600	15,361	6,014	15,494	6,020	16,176	6,570	14,633	5,631
Observations	33,570	50,139	60,830	22,879	61,675	21,538	62,751	20,879	62,832	20,877
Kleibergen-Paap F-stat.istic	6.71	52.89	47.45	6.44	29.56	29.49	33.60	33.92	30.70	28.85

Notes: The data used are from the HRS 1996 to 2014 of individuals who are 40 to 70. Each cell shows first stage estimates from our preferred model specification in equation 1 in each subsample. All models control for year-fixed effects, fixed effects for the year of birth of the first-born child of an individual, age of the youngest child, cohort fixed effects of the individual, individual demographics such as age (quadratic polynomial), race, religion, gender, birth place and census region fixed effects, the number of children of individuals, and socio-economic controls such as the annual earnings from wages or salary, bonuses, second job income, and professional practice income as well as the annual sum of pensions and annuities. Standard errors are clustered at the individual level. \*\*\*  $p < 0.01$ , \*\*  $p < 0.05$ , \*  $p < 0.10$ .

Table 5: OLS estimates

Dependent variable	ADL	IADL	Poor subjective health	Depressed
Grandparenting	-0.011*** (0.001)	-0.009*** (0.001)	-0.039*** (0.004)	-0.019*** (0.003)
Mean of dependent variable	0.018	0.011	0.274	0.115
Number of clusters	20,255	20,255	20,261	19,575
Observations	83,655	83,650	83,673	78,783

Notes: The data used are from the HRS 1996 to 2014 of individuals who are 40 to 70. Each cell shows OLS estimates from our preferred model specification with full controls for each dependent variable. ADL and IADL are dichotomous indicators which equal one if an individual reports limitations for more than three items, and zero otherwise. Poor subjective health is an indicator for poor self-reported health, which is one if self-reported health is “fair” or “poor”, and is zero otherwise. Depressed is a binary indicator constructed using 4 items of depressive symptoms as the cutoff (8 items in total). All models control for year-fixed effects, fixed effects for the year of birth of the first-born child of an individual, age of the youngest child, cohort fixed effects of the individual, individual demographics such as age (quadratic polynomial), race, religion, gender, birth place and census region fixed effects, the number of children of individuals, and socio-economic controls such as the annual earnings from wages or salary, bonuses, second job income, and professional practice income as well as the annual sum of pensions and annuities. Standard errors are clustered at the individual level. \*\*\*  $p < 0.01$ , \*\*  $p < 0.05$ , \*  $p < 0.10$ .

rate their subjective health as better, and they show fewer depressive symptoms. Although these results seemingly support the notion that active ageing is beneficial for older adults’ health, they should not be interpreted as causal effects. It is plausible that grandparents in good health are more likely to provide childcare than grandparents with poor health, which implies that these estimates may reflect reverse causality rather than the causal effect of grandparenting on health.

The results from our 2SLS regressions in Table 6 confirm the presence of such reverse causality. The point estimates for all four health outcomes are positive, suggesting that grandparenting leads to worse health in the form of more functional limitations, more depressive symptoms, and worse self-reported health. However, only the effects on IADLs and subjective health are statistically significant at the 10% confidence level. Tables A3-A6 in the online appendix show that these results are overall robust to the inclusion or exclusion of covariates. The increase in functional limitations of IADL corresponds to a 26% percent change over the mean or 0.076 standard deviations, whereas the change in

subjective health corresponds to a 9.5% increase over the mean or a change of 0.24 standard deviations. While these are rather modest effect sizes, they suggest that childcare provision can be a strenuous activity for grandparents.

Table 6: 2SLS estimates

Dependent variable	ADL	IADL	Poor subjective health	Depressed
Grandparenting	0.032 (0.039)	0.052* (0.028)	0.269* (0.147)	0.097 (0.091)
Mean of dependent variable	0.018	0.011	0.274	0.115
Number of clusters	20,255	20,255	20,261	19,575
Observations	83,655	83,650	83,673	78,783
Kleibergen-Paap F-stat.	53.66	53.61	54.12	54.56
AR F statistic	0.667	3.945	3.674	1.16
AR p-value	0.414	0.047	0.055	0.281

Notes: The data used are from the HRS 1996 to 2014 of individuals who are 40 to 70. Each cell shows 2SLS estimates from our preferred model specification with full controls in equation 2 for each dependent variable. All models control for year-fixed effects, fixed effects for the year of birth of the first-born child of an individual, age of the youngest child, cohort fixed effects of the individual, individual demographics such as age (quadratic polynomial), race, religion, gender, birth place and census region fixed effects, the number of children of individuals, and socio-economic controls such as the annual earnings from wages or salary, bonuses, second job income, and professional practice income as well as the annual sum of pensions and annuities. Standard errors are clustered at the individual level. \*\*\*  $p < 0.01$ , \*\*  $p < 0.05$ , \*  $p < 0.10$ .

## 4.2 Robustness

We assess the sensitivity of our results to the restrictions imposed on our sample and the definition of grandparenting in a series of robustness check. First, we re-estimate our models using the full sample of HRS respondents regardless of age. The results (Panel A in Table 7) are qualitatively similar to those from our main specification in Table 6. Second, we exclude respondents without grandchildren from the working sample. Our sex ratio instrument identifies variation in both the timing of grandparenthood and the likelihood to provide grandchild care for existing grandchildren. This implies that the control group in our main specification consists of both older adults that are not (yet) grandparents as well

as grandparents that do not provide childcare for their grandchildren. Excluding older adults without grandchildren from the sample means that both treatment and control group consist exclusively of grandparents and our instrument identifies variation in the likelihood to provide grandchild care. The results in Panel B of Table 7 are both quantitatively and qualitatively similar to our main results in Table 6.

Table 8 reports the estimates using several alternative definitions of grandparenting to assess whether our results might be affected by measurement errors in the treatment indicator. The definition used in our main specification is a binary indicator whether respondents report providing at least 100 hours of grandchild care over the past two years. The corresponding estimates (Table 6) are repeated in the first row of Table 8 to facilitate comparisons. We consider five alternative definitions: *(i)* an alternative indicator for whether the respondent and their spouse or partner provided at least 100 hours of grandchild care over the past two year as discussed in section 2.3, *(ii)* a binary indicator that is based on the same information as our main specification but excludes observations who do not know the number of care hours and reported a maximum number of hours of childcare provision,<sup>9</sup> *(iii)* a binary indicator for whether respondents provided any grandchild care over the past two years, *(iv)* a binary indicator for whether respondents provided at least 50 hours of grandchild care over the past two years, and *(v)* a binary indicator for whether respondents provided at least 200 hours of grandchild care over the past two years. The results in Table 8 confirm that our results are robust to these changes across different treatment definitions. Both the magnitude as well as the significance of the estimated effects on IADL and subjective health are very similar across specifications.

In addition, we check the sensitivity of our estimates to different outcome definitions. Table 9 reports the results for the health indicators defined with different cutoffs. The baseline column shows our main estimates in Table 6. The sign of the estimates on

---

<sup>9</sup>As discussed in section 2.3, for respondents who report missing grandchild hours, the HRS further asks the minimum and maximum of the bracket range of care hours. We exclude 3,644 respondents who reported 0 minimum care hours and 199 maximum care hours (3,339) and 200 maximum care hours (305) since the actual care hours for this sample are ambiguous.

Table 7: Robustness of estimates to sample change

Dependent variable	ADL	IADL	Poor subjective health	Depressed
<i>A. Sample without age restrictions</i>				
Grandparenting	0.067 (0.062)	0.150** (0.064)	0.487** (0.195)	0.052 (0.101)
Mean of dependent variable	0.037	0.037	0.306	0.107
Number of clusters	27,565	27,566	27,574	26,227
Observations	143,392	143,381	143,386	131,643
Kleibergen-Paap F-stat.istic	47.87	47.79	48.42	53.32
AR F statistic	1.244	6.617	7.582	0.272
AR p-value	0.265	0.010	0.006	0.602
<i>B. Sample restricted to grandparents</i>				
Grandparenting	0.033 (0.042)	0.065** (0.031)	0.274* (0.159)	0.124 (0.100)
Mean of dependent variable	0.018	0.011	0.272	0.114
Number of clusters	19,991	19,991	19,998	19,301
Observations	77,262	77,259	77,278	72,868
Kleibergen-Paap F-stat.	45.23	45.2	45.62	45.69
AR F statistic	0.613	5.132	3.249	1.599
AR p-value	0.434	0.024	0.072	0.206

Notes: Each cell shows 2SLS estimates from our preferred model specification with full controls in equation 2 for each sub-sample. The sample in Panel A uses data from the HRS 1996 to 2014 of all individuals without age limits. The sample in Panel B uses data from the HRS 1996 to 2014 of individuals who are 40 to 70 and excludes those who report no grandchildren. All models control for year-fixed effects, fixed effects for the year of birth of the first-born child of an individual, age of the youngest child, cohort fixed effects of the individual, individual demographics such as age (quadratic polynomial), race, religion, gender, birth place and census region fixed effects, the number of children of individuals, and socio-economic controls such as the annual earnings from wages or salary, bonuses, second job income, and professional practice income as well as the annual sum of pensions and annuities. Standard errors are clustered at the individual level. \*\*\* p<0.01, \*\* p<0.05, \* p<0.10.

different health indicators is consistent with the baseline model while the magnitude and significance varies across health outcome definitions. For ADLs, grandparenting significantly increases the likelihood of reporting at least one or two limitations and the

Table 8: Robustness of estimates to treatment definitions

Dependent variable	ADL	IADL	Poor subjective health	Depressed
<b>Main specification</b>	0.032 (0.039)	0.052* (0.028)	0.269* (0.147)	0.097 (0.091)
Mean of dependent variable	0.018	0.011	0.274	0.115
Kleibergen-Paap F-stat.	53.66	53.61	54.12	54.56
AR F statistic	0.667	3.945	3.674	1.160
AR p-value	0.414	0.047	0.055	0.281
Couples' hours	0.030 (0.038)	0.051* (0.027)	0.260* (0.143)	0.096 (0.091)
Mean of dependent variable	0.018	0.011	0.274	0.115
Kleibergen-Paap F-stat.	52.15	52.11	52.46	50.7
AR F statistic	0.667	3.945	3.674	1.160
AR p-value	0.414	0.047	0.055	0.281
Excluding reported maximum hours	0.042 (0.043)	0.062** (0.031)	0.318** (0.162)	0.100 (0.098)
Mean of dependent variable	0.019	0.012	0.275	0.115
Kleibergen-Paap F-stat.	47.03	46.98	47.35	48.41
AR F statistic	0.958	4.383	4.385	1.065
AR p-value	0.328	0.036	0.036	0.302
> 0 hours	0.031 (0.039)	0.052* (0.028)	0.266* (0.146)	0.096 (0.091)
Mean of dependent variable	0.018	0.011	0.274	0.115
Kleibergen-Paap F-stat.	49.73	49.69	50.05	49.95
AR F statistic	0.667	3.945	3.674	1.160
AR p-value	0.414	0.047	0.055	0.281
> 50 hours	0.032 (0.04)	0.054* (0.029)	0.277* (0.153)	0.099 (0.094)
Mean of dependent variable	0.018	0.011	0.274	0.115
Kleibergen-Paap F-stat.	47.99	47.95	48.3	48.81
AR F statistic	0.667	3.945	3.674	1.160
AR p-value	0.414	0.047	0.055	0.281
> 200 hours	0.045 (0.048)	0.068** (0.034)	0.337* (0.177)	0.111 (0.111)
Mean of dependent variable	0.019	0.012	0.274	0.115
Kleibergen-Paap F-stat.	45.82	45.85	45.99	45.13
AR F statistic	0.914	4.375	4.027	1.033
AR p-value	0.339	0.037	0.045	0.309

Notes: The data used are from the HRS 1996 to 2014 of individuals who are 40 to 70. Each cell shows 2SLS estimates from our preferred model specification with full controls in equation 2 for each dependent variable. All models control for year-fixed effects, fixed effects for the year of birth of the first-born child of an individual, age of the youngest child, cohort fixed effects of the individual, individual demographics such as age (quadratic polynomial), race, religion, gender, birth place and census region fixed effects, the number of children of individuals, and socio-economic controls such as the annual earnings from wages or salary, bonuses, second job income, and professional practice income as well as the annual sum of pensions and annuities. Standard errors are clustered at the individual level. \*\*\*  $p < 0.01$ , \*\*  $p < 0.05$ , \*  $p < 0.10$ .



Table 9: Robustness of estimate to outcome definitions

<i>A. ADL</i>					
<i>Definition</i>	Baseline (4 limitations)	<i>1</i>	<i>2</i>	<i>3</i>	<i>5</i>
Grandparenting	0.032 (0.039)	0.217** (0.110)	0.127* (0.077)	0.059 (0.054)	0.007 (0.024)
<i>B. IADL</i>					
<i>Definition</i>	Baseline (4 limitations)	<i>1</i>	<i>2</i>	<i>3</i>	<i>5</i>
Grandparenting	0.052* (0.028)	0.156 (0.096)	0.056 (0.062)	0.079* (0.043)	0.017 (0.016)
<i>C. Poor subjective health</i>					
<i>Definition</i>	Baseline (“Poor” or “Fair”)	“Poor”			
Grandparenting	0.269* (0.147)	0.152* (0.085)			
<i>D. Depressed</i>					
<i>Definition</i>	Baseline (5 symptoms)	<i>2</i>	<i>3</i>	<i>4</i>	<i>6</i>
Grandparenting	0.097 (0.091)	0.068 (0.139)	0.081 (0.123)	0.135 (0.108)	0.048 (0.074)

Notes: The data used are from the HRS 1996 to 2014 of individuals who are 40 to 70. Each cell shows 2SLS estimates from our preferred model specification with full controls in equation 2 for each dependent variable. The baseline is the outcome definition used in the main tables of health. The other cutoffs are used as alternative definitions to create the health indicators. All models control for year-fixed effects, fixed effects for the year of birth of the first-born child of an individual, age of the youngest child, cohort fixed effects of the individual, individual demographics such as age (quadratic polynomial), race, religion, gender, birth place and census region fixed effects, the number of children of individuals, and socio-economic controls such as the annual earnings from wages or salary, bonuses, second job income, and professional practice income as well as the annual sum of pensions and annuities. Standard errors are clustered at the individual level. \*\*\* p<0.01, \*\* p<0.05, \* p<0.10.

magnitude of these coefficients is much higher (22 percentage points for at least one limitation and 13 percentage points for at least two limitations) than that of the baseline estimate. However, the estimates on at least three and five ADL limitations are much smaller and not statistically significant. For IADLs, grandparenting is consistently harmful across different definitions. The magnitude of the coefficients with at least two or three IADL limitations is similar to that of the baseline estimate. Similar to the pattern of ADL outcomes, the magnitude of the coefficient with at least one IADL limitation is much higher (16 percentage points) and the magnitude shrinks to two percentage points for the IADL measure with five limitations. For subjective health status, grandparenting significantly increases individuals' likelihood to self-report their health as poor by 15 percentage points (compared to 27 percentage points in our baseline specification). For mental health, none of the estimates is statistically significant, although the sign indicates an increase in depressive symptoms across specifications. Overall, our estimates of the effects of grandchild care provision on health are robust across different outcome definitions.

## 5 Discussion

This study examines the effect of childcare provision on grandparents' health in the U.S.. We propose a novel instrument, the sex ratio, for grandparental childcare provision, drawing on insights from the demographic literature on grandparenthood. Our sex ratio instrument measures the share of daughters among all children born to a person, which captures that parents of daughters transition on average earlier into grandparenthood and grandparents are more likely to provide care for grandchildren born to their daughters than to grandchildren born to their sons. We conduct several tests and falsification exercises that support the sex ratio to be a valid instrument for grandparenting.

Our OLS results are in line with earlier studies suggesting that grandchild care provision is indeed positively associated with grandparents' health, but this association is likely driven

by reverse causality. Once we address such endogeneity using the sex ratio as an IV, we find that effects of grandparenting on health are insignificant or even negative. These negative effects are not always precisely estimated, and thus a more conservative conclusion is that – contrary to most previous studies (Choi and Zhang 2021; Di Gessa et al. 2016; Ku et al. 2013; Wang et al. 2020; Zeng et al. 2021) – grandparenting does not improve grandparents’ health.

Our findings suggest that previously reported positive associations between grandparenting and health are biased, likely due to reverse causality. Yet, it is possible that the context of the study also matters and that findings may differ based on, e.g., the role of the family and the strength of family ties. The U.S. is an interesting setting with neither particularly strong family ties (compared to, e.g., East Asia) nor with extensive subsidized formal childcare places (e.g., as in Northern Europe). It seems possible that health effects in this setting are very different from those reported, e.g., for China (Choi and Zhang 2021; Wang et al. 2020).

We also acknowledge a few limitations of our study. For example, it seems plausible that the health effects of grandparenting will differ based on its intensity (Zeng et al. 2021). Unfortunately, information in the HRS on the number of grandparenting hours is only reported retrospectively and the distribution suggests that it is subject to recall bias. Therefore, here we only consider effects of grandparenting at the extensive margin, but future work should examine differences in intensity, e.g., based on time use diaries. Similarly, it seems plausible that the effects of grandparenting may differ based on the tasks taken over by grandparents. For example, taking care of infants during the day or when parents are close-by may be much less demanding than looking after these children overnight. Unfortunately, such data is not available in the HRS. Last, a larger sample (or stronger instruments) are needed to examine treatment effect heterogeneity across relevant subgroups.

In summary, our results show that grandparenting does not improve the health of

grandparents, rather it may even be detrimental to grandparents' health. Instead, it seems that good health is an important precondition for grandparents to provide childcare and this reverse causality causes the frequently observed positive associations documented in the literature. This implies that grandparenting should not be considered as "active ageing" – a socially desirable activity that preserves or improves older adults' health. Instead, similar to informal care provision grandparenting appears to be an activity that older adults engage in to help their family even though it may be detrimental to their own health or well-being.

## References

- Adserà, A. (2017). Education and fertility in the context of rising inequality. *Vienna Yearbook of Population Research*, 15:63–92.
- Age UK (2017). 5 million grandparents take on childcare responsibilities.
- Angrist, J., Lavy, V., and Schlosser, A. (2010). Multiple experiments for the causal link between the quantity and quality of children. *Journal of Labor Economics*, 28(4):773–824.
- Ates, M. (2017). Does grandchild care influence grandparents’ self-rated health? Evidence from a fixed effects approach. *Social Science & Medicine*, 190(C):67–74. Publisher: Elsevier.
- Battistin, E., De Nadai, M., and Padula, M. (2014). Roadblocks on the road to grandma’s house: Fertility consequences of delayed retirement. IZA Discussion Paper 8071, Institute for the Study of Labor (IZA).
- Bom, J., Bakx, P., Schut, F., and van Doorslaer, E. (2018). The impact of informal caregiving for older adults on the health of various types of caregivers: A systematic review. *The Gerontologist*, 59(5):e629–e642.
- Bom, J. and Stöckel, J. (2021). Is the grass greener on the other side? The health impact of providing informal care in the UK and the Netherlands. *Social Science & Medicine*, 269:113562.
- Brunello, G. and Rocco, L. (2019). Grandparents in the blues. The effect of childcare on grandparents’ depression. *Review of Economics of the Household*, 17(2):587–613. Publisher: Springer.
- Chen, F., Short, S. E., and Entwisle, B. (2000). The impact of grandparental proximity on maternal childcare in China. *Population Research and Policy Review*, 19(6):571–590.


- Chien, S., Campbell, N., Hayden, O., Hurd, M., Main, R., Mallett, J., Martin, C., Meijer, E., Miu, A., Moldoff, M., et al. (2015). RAND HRS data documentation, version N. *RAND Corporation*.
- Choi, S.-w. E. and Zhang, Z. (2021). Caring as curing: Grandparenting and depressive symptoms in China. *Social Science & Medicine*, 289:114452.
- Compton, J. (2015). Family proximity and the labor force status of women in Canada. *Review of Economics of the Household*, 13(2):323–358.
- Compton, J. and Pollak, R. A. (2011). Family Proximity, Childcare, and Women’s Labor Force Attachment. *National Bureau of Economic Research Working Paper Series*, No. 17678. featured in NBER digest on 2012-03-30.
- Danielsbacka, M., Tanskanen, A. O., Coall, D. A., and Jokela, M. (2019). Grandparental childcare, health and well-being in Europe: A within-individual investigation of longitudinal data. *Social Science & Medicine*.
- Danielsbacka, M., Tanskanen, A. O., Jokela, M., and Rotkirch, A. (2011). Grandparental Child Care in Europe: Evidence for Preferential Investment in More Certain Kin. *Evolutionary Psychology*, 9(1):147470491100900102. Publisher: SAGE Publications Inc.
- Dave, D., Rashad, I., and Spasojevic, J. (2006). The effects of retirement on physical and mental health outcomes. Technical report, National Bureau of Economic Research.
- DeSalvo, K. B., Bloser, N., Reynolds, K., He, J., and Muntner, P. (2006). Mortality prediction with a single general self-rated health question. *Journal of General Internal Medicine*, 21(3):267–275.
- Di Gessa, G., Glaser, K., and Tinker, A. (2016). The health impact of intensive and nonintensive grandchild care in Europe: New evidence from SHARE. *The Journals of Gerontology: Series B*, 71(5):867–879.

- Eibich, P. (2015). Understanding the effect of retirement on health: Mechanisms and heterogeneity. *Journal of Health Economics*, 43:1–12.
- Eibich, P. and Siedler, T. (2020). Retirement, intergenerational time transfers, and fertility. *European Economic Review*, 124:103392.
- Eurostat (2022). Living conditions in Europe - childcare arrangements.
- Heger, D. (2017). The mental health of children providing care to their elderly parent. *Health Economics*, 26(12):1617–1629. Publisher: John Wiley & Sons, Ltd.
- Idler, E. L. and Benyamini, Y. (1997). Self-rated health and mortality: a review of twenty-seven community studies. *Journal of Health and Social Behavior*, pages 21–37.
- Irwin, M., Artin, K. H., and Oxman, M. N. (1999). Screening for depression in the older adult: Criterion validity of the 10-item Center for Epidemiological Studies Depression Scale (CESD). *Archives of Internal Medicine*, 159(15):1701–1704.
- Ku, L.-J., Stearns, S., Houtven, C., and Holmes, G. (2012). The health effects of caregiving by grandparents in Taiwan: An instrumental variable estimation. *Review of Economics of the Household*, 10(4):521–540. Publisher: Springer.
- Ku, L.-J. E., Stearns, S. C., Van Houtven, C. H., Lee, S.-Y. D., Dilworth-Anderson, P., and Konrad, T. R. (2013). Impact of Caring for Grandchildren on the Health of Grandparents in Taiwan. *The Journals of Gerontology: Series B*, 68(6):1009–1021.
- Kuka, E. (2020). Quantifying the benefits of social insurance: unemployment insurance and health. *Review of Economics and Statistics*, 102(3):490–505.
- Lai, D. W. L., Li, J., and Bai, X. (2021). To be or not to be: relationship between grandparent status and health and wellbeing. *BMC Geriatrics*, 21(1):204.
- Laughlin, L. (2013). Who’s minding the kids? child care arrangements: Spring 2011. Technical Report P70-135, US Census Bureau.

- Liu, Y. and Zai, X. (2022). The unintended effect of Medicaid aging waivers on informal caregiving. *Working Paper*.
- Margolis, R. and Verdery, A. M. (2019). A Cohort Perspective on the Demography of Grandparenthood: Past, Present, and Future Changes in Race and Sex Disparities in the United States. *Demography*, 56(4):1495–1518.
- Mirowsky, J. (2005). Age at First Birth, Health, and Mortality. *Journal of Health and Social Behavior*, 46(1):32–50. Publisher: SAGE Publications Inc.
- Rupert, P. and Zanella, G. (2018). Grandchildren and their grandparents' labor supply. *Journal of Public Economics*, 159:89–103.
- Schmitz, H. and Westphal, M. (2015). Short- and medium-term effects of informal care provision on female caregivers' health. *Journal of Health Economics*, 42:174–185.
- Somary, K. and Strieker, G. (1998). Becoming a Grandparent: A Longitudinal Study of Expectations and Early Experiences as a Function of Sex and Lineage<sup>1</sup>. *The Gerontologist*, 38(1):53–61.
- Wang, H., Fidrmuc, J., and Luo, Q. (2020). Grandparenting and well-being of the elderly in China. Technical report, Bank of Finland, Institute for Economies in Transition. Issue: 18/2020.
- Wu, X. and Li, L. (2012). Family size and maternal health: evidence from the One-Child policy in China. *Journal of Population Economics*, 25(4):1341–1364.
- Zeng, Y., Chen, Y.-C., and Lum, T. Y. S. (2021). Longitudinal impacts of grandparent caregiving on cognitive, mental, and physical health in China. *Aging & Mental Health*, 25(11):2053–2060. Publisher: Routledge.




Figure A1: Distribution of Age of HRS Respondents



Notes: The data is the HRS from 1992 to 2014. This graph draws the distribution of age of HRS respondents. The vertical axis shows the density of age. The age eligibility of respondents is above 50 and the age of their spouses can be any age as shown in the plot.

Figure A2: Distribution of Age of HRS Respondents



Notes: The data used are from the HRS 1996 to 2014 of individuals who are 40 to 70. This graphs draws the distribution of grandchild care hours for respondents who are grandparenting and provide less than 1,000 hours over the last two years.

Table A1: Definitions of Variables

Variable	Definition
<b><i>Demographics</i></b>	
Age	Age of respondents in years
Female	Dichotomous indicator of respondents being female (female=1, 0=male)
Education	Years in school of respondents
Marital status	
Married/partnered	Dichotomous indicator of respondents being married or having a partner living together
Separated/divorced	Dichotomous indicator of respondents being divorced or separated from marriage
Widowed	Dichotomous indicator of respondents having spouses or partners dead
Never married	Dichotomous indicator of respondents never getting married
Race/ethnicity	
White	Dichotomous indicator of respondents being white
Black/African	Dichotomous indicator of respondents being black or Hispanic
Other	Dichotomous indicator of respondents' race other than white or black or African
<b><i>Instrumental variable</i></b>	
Sex ratio	The ratio between the number of daughters to all children
<b><i>Grandparenting</i></b>	
Grandparenting for at least one child (Q1)	Q1: whether the respondent and spouse spent 100 or more hours taking care of their grandchildren or great-grandchildren since the last wave
Grandparenting for at least 100 hours (Q2)	Q2: how many estimate childcare hours provided in the last two years for the respondent and spouse, separately
<b><i>Heath variables</i></b>	
Subjective health	Binary indicator for poor subjective health based on respondent's self-reported general health status, 1 for "fair" or "poor", and 0 for "excellent", "very good" or "good".
ADL	Binary indicator for respondents that report 3 or more difficulties (out of 5 possible) on the index of difficulties in Activities of Daily Living (ADL), indicating respondents having any problem in bathing, eating, getting dressed, getting in/out of bed, and walking across a room
IADL	Binary indicator for respondents that report 3 or more difficulties (out of 5 possible) on the index of difficulties in Instrumental Activities of Daily Living (IADL), indicating respondents having any problem in using the phone, managing money, taking medications, shopping for groceries, and preparing hot meals
Mental health	Binary indicator for poor mental health based on scoring 4 or more points on the Center for Epidemiologic Studies Depression (CESD) scale: sum of five negative indicators minus two positive indicators. The negative indicators measure sentiments all or most of the time: depression, everything is an effort, restless sleep, felt alone, sad, and could not get going. The positive indicators measure whether respondents felt happy and enjoyed life.

Table A2: Reduced form regression of the working sample

Dependent variable	ADL	IADL	Poor subjective health	Depressed
Sex ratio	0.002 (0.002)	0.003** (0.002)	0.017* (0.009)	0.006 (0.006)
Mean of dependent variable	0.018	0.011	0.274	0.115
Number of clusters	20,255	20,255	20,261	19,575
Observations	83,655	83,650	83,673	78,783

Notes: The data used are from the HRS 1996 to 2014 of individuals who are 40 to 70. Each cell shows reduced form estimates from our preferred model specification with full controls. All models control for year-fixed effects, fixed effects for the year of birth of the first-born child of an individual, age of the youngest child, cohort fixed effects of the individual, individual demographics such as age (quadratic polynomial), race, religion, gender, birth place and census region fixed effects, the number of children of individuals, and socio-economic controls such as the annual earnings from wages or salary, bonuses, second job income, and professional practice income as well as the annual sum of pensions and annuities. Standard errors are clustered at the individual level. \*\*\*  $p < 0.01$ , \*\*  $p < 0.05$ , \*  $p < 0.10$ .

Table A3: Robustness of 2SLS estimates on ADL

<i>Dependent variable: ADL</i>						
Model	1	2	3	4	5	6
Grandparenting	0.034 (0.038)	0.029 (0.039)	0.030 (0.039)	0.032 (0.039)	0.032 (0.039)	0.032 (0.039)
Year FE + Birth year FE of first born		Y	Y	Y	Y	Y
Cohort FE + Birth year FE of youngest born			Y	Y	Y	Y
Demographics of R				Y	Y	Y
Family size					Y	Y
Socioeconomic controls						Y
Mean of dependent variable	0.019	0.019	0.019	0.018	0.018	0.018
Number of clusters	20,473	20,416	20,391	20,255	20,255	20,255
Observations	84,297	84,220	84,131	83,655	83,655	83,655
Kleibergen-Paap F-stat.istic	53.29	53.41	54.17	53.73	53.74	53.66
AR F statistic	0.809	0.567	0.607	0.691	0.693	0.667
AR p-value	0.368	0.451	0.436	0.406	0.405	0.414

Notes: The data used are from the HRS 1996 to 2014 of individuals who are 40 to 70. Each cell reports estimates from a separate specification using equation 2 for the dependent variable. Column 1 reports estimates without any controls. Column 2 adds year-fixed effects and fixed effects for the year of birth of the first-born child of an individual. Column 3 adds age of the youngest child and cohort fixed effects of the individual. Column 4 includes individual demographics such as age (quadratic polynomial), race, religion, gender, birth place and census region fixed effects. Column 5 further controls for the number of children of individuals. Column 6 adds into socio-economic controls such as the annual earnings from wages or salary, bonuses, second job income, and professional practice income as well as the annual sum of pensions and annuities. Standard errors are clustered at the individual level. \*\*\*  $p < 0.01$ , \*\*  $p < 0.05$ , \*  $p < 0.10$ .

Table A4: Robustness of 2SLS estimates on IADL

<i>Dependent variable: IADL</i>						
Model	1	2	3	4	5	6
Grandparenting	0.049*	0.046*	0.046*	0.053*	0.053*	0.052*
	(0.028)	(0.028)	(0.028)	(0.028)	(0.028)	(0.028)
Year FE + Birth year FE of first born		Y	Y	Y	Y	Y
Cohort FE + Birth year FE of youngest born			Y	Y	Y	Y
Demographics of R				Y	Y	Y
Family size					Y	Y
Socioeconomic controls						Y
Mean of dependent variable	0.011	0.011	0.011	0.011	0.011	0.011
Number of clusters	20,472	20,415	20,390	20,255	20,255	20,255
Observations	84,291	84,214	84,125	83,650	83,650	83,650
Kleibergen-Paap F-stat.istic	53.22	53.34	54.08	53.67	53.68	53.61
AR F statistic	3.375	2.960	3.065	3.971	3.989	3.945
AR p-value	0.066	0.085	0.080	0.046	0.046	0.047

Notes: The data used are from the HRS 1996 to 2014 of individuals who are 40 to 70. Each cell reports estimates from a separate specification using equation 2 for the dependent variable. Column 1 reports estimates without any controls. Column 2 adds year-fixed effects and fixed effects for the year of birth of the first-born child of an individual. Column 3 adds age of the youngest child and cohort fixed effects of the individual. Column 4 includes individual demographics such as age (quadratic polynomial), race, religion, gender, birth place and census region fixed effects. Column 5 further controls for the number of children of individuals. Column 6 adds into socio-economic controls such as the annual earnings from wages or salary, bonuses, second job income, and professional practice income as well as the annual sum of pensions and annuities. Standard errors are clustered at the individual level. \*\*\*  $p < 0.01$ , \*\*  $p < 0.05$ , \*  $p < 0.10$ .

Table A5: Robustness of 2SLS estimates on poor subjective health

<i>Dependent variable: Poor subjective health</i>						
Model	1	2	3	4	5	6
Grandparenting	0.309** (0.153)	0.311** (0.154)	0.303** (0.153)	0.273* (0.151)	0.275* (0.151)	0.269* (0.147)
Year FE + Birth year FE of first born		Y	Y	Y	Y	Y
Cohort FE + Birth year FE of youngest born			Y	Y	Y	Y
Demographics of R				Y	Y	Y
Family size					Y	Y
Socioeconomic controls						Y
Mean of dependent variable	0.275	0.275	0.275	0.274	0.274	0.274
Number of clusters	20,480	20,422	20,397	20,261	20,261	20,261
Observations	84,316	84,238	84,149	83,673	83,673	83,673
Kleibergen-Paap F-stat.istic	53.65	53.84	54.59	54.19	54.20	54.12
AR F statistic	4.596	4.574	4.375	3.577	3.624	3.674
AR p-value	0.032	0.033	0.037	0.059	0.057	0.055

Notes: The data used are from the HRS 1996 to 2014 of individuals who are 40 to 70. Each cell reports estimates from a separate specification using equation 2 for the dependent variable. Column 1 reports estimates without any controls. Column 2 adds year-fixed effects and fixed effects for the year of birth of the first-born child of an individual. Column 3 adds age of the youngest child and cohort fixed effects of the individual. Column 4 includes individual demographics such as age (quadratic polynomial), race, religion, gender, birth place and census region fixed effects. Column 5 further controls for the number of children of individuals. Column 6 adds into socio-economic controls such as the annual earnings from wages or salary, bonuses, second job income, and professional practice income as well as the annual sum of pensions and annuities. Standard errors are clustered at the individual level. \*\*\*  $p < 0.01$ , \*\*  $p < 0.05$ , \*  $p < 0.10$ .

Table A6: Robustness of 2SLS estimates on Depressed

<i>Dependent variable: Depressed</i>						
Model	1	2	3	4	5	6
Grandparenting	0.157*	0.127	0.118	0.101	0.101	0.097
	(0.094)	(0.093)	(0.093)	(0.092)	(0.092)	(0.091)
Year FE + Birth year FE of first born		Y	Y	Y	Y	Y
Cohort FE + Birth year FE of youngest born			Y	Y	Y	Y
Demographics of R				Y	Y	Y
Family size					Y	Y
Socioeconomic controls						Y
Mean of dependent variable	0.115	0.115	0.115	0.115	0.115	0.115
Number of clusters	19,779	19,726	19,698	19,575	19,575	19,575
Observations	79,356	79,284	79,198	78,783	78,783	78,783
Kleibergen-Paap F-stat.istic	53.51	54.43	55.22	54.68	54.68	54.56
AR F statistic	2.979	1.956	1.710	1.230	1.233	1.160
AR p-value	0.084	0.162	0.191	0.267	0.267	0.281

Notes: The data used are from the HRS 1996 to 2014 of individuals who are 40 to 70. Each cell reports estimates from a separate specification using equation 2 for the dependent variable. Column 1 reports estimates without any controls. Column 2 adds year-fixed effects and fixed effects for the year of birth of the first-born child of an individual. Column 3 adds age of the youngest child and cohort fixed effects of the individual. Column 4 includes individual demographics such as age (quadratic polynomial), race, religion, gender, birth place and census region fixed effects. Column 5 further controls for the number of children of individuals. Column 6 adds into socio-economic controls such as the annual earnings from wages or salary, bonuses, second job income, and professional practice income as well as the annual sum of pensions and annuities. Standard errors are clustered at the individual level. \*\*\*  $p < 0.01$ , \*\*  $p < 0.05$ , \*  $p < 0.10$ .





## CINCH working paper series

- 1 Halla, Martin and Martina Zweimüller. **Parental Responses to Early Human Capital Shocks:** Evidence from the Chernobyl Accident. CINCH 2014.
- 2 Aparicio, Ainhoa and Libertad González. **Newborn Health and the Business Cycle:** Is it Good to be born in Bad Times? CINCH 2014.
- 3 Robinson, Joshua J. **Sound Body, Sound Mind?:** Asymmetric and Symmetric Fetal Growth Restriction and Human Capital Development. CINCH 2014.
- 4 Bhalotra, Sonia, Martin Karlsson and Therese Nilsson. **Life Expectancy and Mother-Baby Interventions:** Evidence from A Historical Trial. CINCH 2014.
- 5 Goebel, Jan, Christian Krekel, Tim Tiefenbach and Nicolas R. Ziebarth. **Natural Disaster, Environmental Concerns, Well-Being and Policy Action:** The Case of Fukushima. CINCH 2014.
- 6 Avdic, Daniel, **A matter of life and death? Hospital Distance and Quality of Care:** Evidence from Emergency Hospital Closures and Myocardial Infarctions. CINCH 2015.
- 7 Costa-Font, Joan, Martin Karlsson and Henning Øien. **Informal Care and the Great Recession.** CINCH 2015.
- 8 Titus J. Galama and Hans van Kippersluis. **A Theory of Education and Health.** CINCH 2015.
- 9 Dahmann, Sarah. **How Does Education Improve Cognitive Skills?:** Instructional Time versus Timing of Instruction. CINCH 2015.
- 10 Dahmann, Sarah and Silke Anger. **The Impact of Education on Personality:** Evidence from a German High School Reform. CINCH 2015.
- 11 Carbone, Jared C. and Snorre Kverndokk. **Individual Investments in Education and Health.** CINCH 2015.
- 12 Zilic, Ivan. **Effect of forced displacement on health.** CINCH 2015.
- 13 De la Mata, Dolores and Carlos Felipe Gaviria. **Losing Health Insurance When Young:** Impacts on Usage of Medical Services and Health. CINCH 2015.


- 14 Tequame, Miron and Nyasha Tirivayi. **Higher education and fertility:** Evidence from a natural experiment in Ethiopia. CINCH 2015.
- 15 Aoki, Yu and Lualhati Santiago. **Fertility, Health and Education of UK Immigrants:** The Role of English Language Skills. CINCH 2015.
- 16 Rawlings, Samantha B., **Parental education and child health:** Evidence from an education reform in China. CINCH 2015.
- 17 Kamhöfer, Daniel A., Hendrik Schmitz and Matthias Westphal. **Heterogeneity in Marginal Non-monetary Returns to Higher Education.** CINCH 2015.
- 18 Ardila Brenøe, Anne and Ramona Molitor. **Birth Order and Health of Newborns:** What Can We Learn from Danish Registry Data? CINCH 2015.
- 19 Rossi, Pauline. **Strategic Choices in Polygamous Households:** Theory and Evidence from Senegal. CINCH 2016.
- 20 Clarke, Damian and Hanna Mühlrad. **The Impact of Abortion Legalization on Fertility and Maternal Mortality:** New Evidence from Mexico. CINCH 2016.
- 21 Jones, Lauren E. and Nicolas R. Ziebarth. **US Child Safety Seat Laws:** Are they Effective, and Who Complies? CINCH 2016.
- 22 Koppensteiner, Martin Foureaux and Jesse Matheson. **Access to Education and Teenage Pregnancy.** CINCH 2016.
- 23 Hofmann, Sarah M. and Andrea M. Mühlenweg. **Gatekeeping in German Primary Health Care – Impacts on Coordination of Care, Quality Indicators and Ambulatory Costs.** CINCH 2016.
- 24 Sandner, Malte. **Effects of Early Childhood Intervention on Fertility and Maternal Employment:** Evidence from a Randomized Controlled Trial. CINCH 2016.
- 25 Baird, Matthew, Lindsay Daugherty, and Krishna Kumar. **Improving Estimation of Labor Market Disequilibrium through Inclusion of Shortage Indicators.** CINCH 2017.
- 26 Bertoni, Marco, Giorgio Brunello and Gianluca Mazzarella. **Does postponing minimum retirement age improve healthy behaviors before retirement?** Evidence from middle-aged Italian workers. CINCH 2017.
- 27 Berniell, Inés and Jan Bietenbeck. **The Effect of Working Hours on Health.** CINCH 2017.
- 28 Cronin, Christopher, Matthew Forsstrom, and Nicholas Papageorge. **Mental Health, Human Capital and Labor Market Outcomes.** CINCH 2017.

- 29 Kamhöfer, Daniel and Matthias Westphal. **Fertility Effects of College Education:** Evidence from the German Educational Expansion. CINCH 2017.
- 30 Jones, John Bailey and Yue Li. **The Effects of Collecting Income Taxes on Social Security Benefits.** CINCH 2017.
- 31 Hofmann, Sarah and Andrea Mühlenweg. **Learning Intensity Effects in Students' Mental and Physical Health –** Evidence from a Large Scale Natural Experiment in Germany. CINCH 2017.
- 32 Vollmer, Sebastian and Juditha Wójcik. **The Long-term Consequences of the Global 1918 Influenza Pandemic:** A Systematic Analysis of 117 IPUMS International Census Data Sets. CINCH 2017.
- 33 Thamarapani, Dhanushka, Rockmore, Marc, and Willa Friedman. **The Educational and Fertility Effects of Sibling Deaths.** CINCH 2018.
- 34 Lemmon, Elizabeth. **Utilisation of personal care services in Scotland:** the influence of unpaid carers. CINCH 2018.
- 35 Avdic, Daniel, Büyükdurmus, Tugba, Moscelli, Giuseppe, Pilny, Adam, and Ieva Sriubaite. **Subjective and objective quality reporting and choice of hospital:** Evidence from maternal care services in Germany. CINCH 2018.
- 36 Hentschker, Corinna and Ansgar Wübker. **Quasi-experimental evidence on the effectiveness of heart attack treatment in Germany.** CINCH 2018.
- 37 Pasha, Mochamad, Rockmore, Marc, and Chih Ming Tan. **Early Life Exposure to Above Average Rainfall and Adult Mental Health.** CINCH 2018.
- 38 Elsner, Benjamin and Florian Wozny. **The Human Capital Cost of Radiation:** Long-run Evidence from Exposure outside the Womb. CINCH 2019.
- 39 de la Mata, Dolores and Carlos Felipe Gaviria Garcés. **Exposure to Pollution and Infant Health:** Evidence from Colombia. CINCH 2019
- 40 Besstremyannaya, Galina and Sergei Golovan. **Physicians' altruism in incentives contracts:** Medicare's quality race. CINCH 2019.
- 41 Hayen, Arthur P., Klein, Tobias J., and Martin Salm. **Does the framing of patient cost-sharing incentives matter?** The effects of deductibles vs. no-claim refunds. CINCH 2019.
- 42 Molina, Teresa. **Pollution, Ability, and Gender-Specific Responses to Shocks.** CINCH 2019.

- 43 Fischer, Martin, Gerdtham, Ulf-G, Heckley, Gawain, Karlsson, Martin, Kjellsson, Gustav, and Therese Nilsson. **Education and Health: Long-run Effects of Peers, Tracking and Years.** CINCH 2019.
- 44 Bannenberg, Norman, Førland, Oddvar, Iversen, Tor, Karlsson, Martin, and Henning Øien. **Preventive Home Visits.** CINCH 2019.
- 45 Pestel, Nico and Florian Wozny. **Low Emission Zones for Better Health: Evidence from German Hospitals.** CINCH 2019.
- 46 Bartoli, Paola, Grembi, Veronica, and The Linh Bao Nguyen. **Ethnic Density and Health at Birth.** CINCH 2019.
- 47 Atal, Juan Pablo, Fang, Hanming, Karlsson, Martin, and Nicolas R. Ziebarth. **Long-Term Health Insurance: Theory Meets Evidence.** CINCH 2020.
- 48 Briody, Jonathan. **Parental Unemployment During the Great Recession and Childhood Adiposity.** CINCH 2020.
- 49 Miller, Grant, Paula, Aureo de, and Christine Valente. **Subjective Expectations and Demand for Contraception.** CINCH 2020.
- 50 Kronenberg, Christoph. **New(spaper) Evidence of a Reduction in Suicide Mentions during the 19<sup>th</sup>-century Gold Rush.** CINCH 2020.
- 51 Avdic, Daniel, Ivets, Maryna, Lagerqvist, Bo, and Ieva Sriubaite. **Providers, Peers and Patients: How do Physicians' Practice Environments Affect Patient Outcomes?** CINCH 2021.
- 52 Alexander Ahammer, Dominik Grübl and Rudolf Winter-Ebmer. **The Health Externalities of Downsizing.** CINCH 2021.
- 53 Emilia Barili, Paola Bertoli, Veronica Grembi, and Veronica Rattini. **COVID Angels Fighting Daily Demons? Mental Health of Healthcare Workers and Religion.** CINCH 2021.
- 54 Signe A. Abrahamsen, Rita Ginja, and Julie Riise. **School Health Programs: Education, Health, and Welfare Dependency of Young Adults.** CINCH 2021.
- 55 Martin Fischer, Martin Karlsson, and Nikolaos Prodromidis. **The Long-Term Effects of Hospital Deliveries.** CINCH 2021.
- 56 Cristina Borra, Libertad González, and David Patiño. **Maternal Age and Infant Health.** CINCH 2021.
- 57 Nicola Barban, Elisabetta De Cao, and Marco Francesconi. **Gene-Environment Effects on Female Fertility.** CINCH 2021.
- 58 Daniel Avdic and Katharina E. Blankart. **A Hard Look at "Soft" Cost-control Measures in Healthcare Organizations: Evidence from Preferred Drug Policies in Germany.** CINCH 2021.

- 59 Sarah Hofmann. **Disease Perception and Preventive Behavior:** The Vaccination Response to Local Measles Outbreaks. CINCH 2022.
- 60 Apostolos Davillas, Victor Hugo de Oliveira, and Andrew M Jones. **A Model of Errors in BMI Based on Self-reported and Measured Anthropometrics with Evidence from Brazilian Data.** CINCH 2022.
- 61 Peter Eibich and Xianhua Zai. **Are the Grandparents Alright? The Health Consequences of Grandparental Childcare Provision.** CINCH 2022.





# DuEPublico

Duisburg-Essen Publications online

UNIVERSITÄT  
DUISBURG  
ESSEN

*Offen im Denken*

ub | universitäts  
bibliothek

Dieser Text wird via DuEPublico, dem Dokumenten- und Publikationsserver der Universität Duisburg-Essen, zur Verfügung gestellt. Die hier veröffentlichte Version der E-Publikation kann von einer eventuell ebenfalls veröffentlichten Verlagsversion abweichen.

**DOI:** 10.17185/duepublico/77108

**URN:** urn:nbn:de:hbz:465-20221107-160716-6

Alle Rechte vorbehalten.