

A Service of

ZBW

Leibniz-Informationszentrum Wirtschaft Leibniz Information Centre for Economics

Kumar, Santosh; Nahlen, Bernard

Working Paper Intergenerational Persistence of Health: Evidence from India

IZA Discussion Papers, No. 15938

Provided in Cooperation with:

IZA – Institute of Labor Economics

Suggested Citation: Kumar, Santosh; Nahlen, Bernard (2023) : Intergenerational Persistence of Health: Evidence from India, IZA Discussion Papers, No. 15938, Institute of Labor Economics (IZA), Bonn

This Version is available at: https://hdl.handle.net/10419/272565

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.



WWW.ECONSTOR.EU



Initiated by Deutsche Post Foundation

DISCUSSION PAPER SERIES

IZA DP No. 15938

Intergenerational Persistence of Health: Evidence from India

Santosh Kumar Bernard Nahlen

FEBRUARY 2023



Initiated by Deutsche Post Foundation

DISCUSSION PAPER SERIES

IZA DP No. 15938

Intergenerational Persistence of Health: Evidence from India

Santosh Kumar University of Notre Dame, Eck Institute for Global Health and IZA

Bernard Nahlen University of Notre Dame and Eck Institute for Global Health

FEBRUARY 2023

Any opinions expressed in this paper are those of the author(s) and not those of IZA. Research published in this series may include views on policy, but IZA takes no institutional policy positions. The IZA research network is committed to the IZA Guiding Principles of Research Integrity.

The IZA Institute of Labor Economics is an independent economic research institute that conducts research in labor economics and offers evidence-based policy advice on labor market issues. Supported by the Deutsche Post Foundation, IZA runs the world's largest network of economists, whose research aims to provide answers to the global labor market challenges of our time. Our key objective is to build bridges between academic research, policymakers and society.

IZA Discussion Papers often represent preliminary work and are circulated to encourage discussion. Citation of such a paper should account for its provisional character. A revised version may be available directly from the author.

ISSN: 2365-9793

IZA – Institute of Labor Economics

Schaumburg-Lippe-Straße 5–9	Phone: +49-228-3894-0	
53113 Bonn, Germany	Email: publications@iza.org	www.iza.org

ABSTRACT

Intergenerational Persistence of Health: Evidence from India

Using nationally representative data, we estimate intergenerational persistence in health in India. Results from the instrumental variable method show that children of anemic mothers are more likely to be anemic, with an intergenerational health correlation of 0.26. Results are robust to the inclusion of confounding factors including the mother's height. We find that the correlation between mothers' anemic status and children's anemic status differs by wealth quintile, indicating that economic status may play a role in the persistence of poor health across generations in developing countries.

JEL Classification:	110, 114, O15
Keywords:	intergenerational mobility, health, anemia, India

Corresponding author: Santosh Kumar Keough School of Global Affairs University of Notre Dame Notre Dame, Indiana USA 46556 E-mail: skumar23@nd.edu

1. Introduction

One of the most topical and urgent public health challenges in low- and middle-income countries (LMICs) is micronutrient deficiencies (MND) among children and women. High levels of MND among women and children may signify the existence of intergenerational transmission of poor health from mother to children. Existing work on intergenerational transmission has focused on the labor market (Azam, 2015) and educational outcomes; however, intergenerational transmission of health is understudied and little is known about the independent effect of maternal health on child health after accounting for genetic factors and unobserved omitted factors in a causal framework (Akbulut-Yuksel and Kugler, 2016; Onyeneho et al., 2019; Halliday, Mazumder, and Wong, 2021).

This study examines the intergenerational transmission of anemia in India and explores the role of economic status in moderating the intergenerational cycle of anemia from mother to child. Prior research has examined intergenerational transmission of health in the context of birthweight, height, weight, etc. (Currie and Moretti, 2007; Akbulut-Yuksel and Kugler, 2016), but the evidence on transmission of anemia from mother to child is limited. Identifying the factors that generate intergenerational transmission of health is important because of the high incidence of anemia globally—about 1.6 billion people are affected by anemia, predominantly in Asia and Africa. Two-thirds of LMICs' anemia cases are in India—an anemia prevalence of 67% among under-five children and 57% among 15-49 years old women (IIPS and ICF, 2021). Furthermore, previous studies related to fetal programming and early-life literature have shown that micronutrient deficiencies in the pre- and post-natal phase of life have adverse effects on health, human capital, labor productivity, and socioeconomic outcomes later in life (Currie and Vogl, 2013; Almond, Currie, and Duque, 2018). A recent modeling study estimates that a scaling-up of antenatal iron-folic acid supplementations could lead to gains of 2.28 million school years and US\$8.25 billion in lifetime income (Perumal et al., 2021). Thus, addressing the anemia problem in LMICs would help spur human capital formation and would lead to substantial economic benefits.

Intergenerational health association (IHA) could be due to genetic factors or due to causal effects of parental behaviors on children's outcomes (Lochner, 2008). Intergenerational transmission of health behaviors and outcomes could be social in nature

and is likely to be determined by socioeconomic environments and is not always purely genetic (Lochner, 2008). Identifying the role of genetic, parental, and socioeconomic factors in the IHA is important to understand the effectiveness of policies or factors in breaking the cycle of transmission of poor health from mother to child. Halliday et al. (2021) estimate a health mobility coefficient in the USA with a range of 0.20-0.25. The intergenerational persistence in birth weight ranged from 0.17-0.20 (Currie and Moretti, 2007), and Akbulut-Yuksel and Kugler (2016) show intergenerational association in BMI and height among natives as well as immigrants in the USA. Most of the studies on intergenerational persistence in health are from high-income countries; thus, this study extends intergenerational persistence in micronutrient deficiency in an LMICs setting.

The objective of this study is twofold: to estimate the intergenerational correlation in anemia between mother and child in India, and to explore heterogeneity in this correlation by the household's economic status after controlling for genetic factors. We accomplish these objectives by using the instrumental variable method to estimate the causal association between the mother's anemic status and the children's anemic status. We employ a leave-out-one instrument at the district level and include the mother's height to further control for the genetic factor. We estimate an intergenerational correlation of 0.40 and 0.26 between mothers' and children's Hemoglobin (Hb) levels and anemic status, respectively. This implies that children of anemic mothers have a 26-percentage point higher probability of being anemic. The heterogeneity analyses by wealth quintiles show that economically disadvantaged children display lower upward mobility compared to children from better-off households.

2. Data and empirical methodology

We use micro-level data from the fourth round of the National Family Health Survey (NFHS) conducted in 2015-16 (IIPS & ICF, 2017). The NFHS is a nationally representative health survey gathering information on population, health, and nutrition from a sample covering 601,509 households and 699,686 women.

The NFHS survey collected hemoglobin (Hb) information for 6-59 months old children and for 15-49 years old pregnant as well as non-pregnant women. The main dependent variable is the Hb level and anemic status of children born in the last five years before the survey. Children are categorized as anemic if the Hb < 11.0 g/dL while the cutoff for pregnant and non-pregnant women is 11.0 g/dL and 12.0 g/dL, respectively. The anemic category includes severely, moderately, and mildly anemic observations. The control variables include gender, age, and birth order of the child and household-level controls include the mother's education, mother's height, religion, social group of the household (caste), and rural dummy. We trim 1 percent of observations at the bottom and top tail of the Hb distribution to exclude outliers. The analytical sample has 198,303 observations.

Table 1 presents the summary statistics for the variables used in the analysis. The average Hb level is 10.59 g/dL and about 58% of the sampled children are anemic. More than half of the women are anemic (56%). The average age of the children is 2 years. Mothers' mean years of schooling is 6 years and about one-half of the households belong to the bottom two wealth groups. More than two-thirds of the children reside in rural areas (76%).

Following the standard method in this literature (Halliday et al., 2021), we estimate the following linear regression model to estimate the intergenerational health association between mother and children:

$$H_{ci} = \alpha_1 + \beta_1 * H_{mi} + \beta_2 * X_{ci} + \mu_d + \epsilon_{ci} \qquad (1)$$

Where H_{ci} denotes the health of the child *c* in the family *i*; H_{mi} denotes the health of the mother *m*; X_{ci} denotes a vector of mothers' and children's level controls; μ_d is district fixed effects to control for time-invariant characteristics of the district. Standard errors are clustered by the district. We weight the regression models with national women's sample weight.

Since the main independent variable, the mother's anemic status, is likely to be endogenous, we employ the Two-Stage Least Square (2SLS) regression model to address endogeneity. For example, unobserved heterogeneity such as the mother's health, dietary behavior, and genetic factors may affect the mothers' as well as children's anemic status, and therefore, the OLS estimates would be biased and inconsistent. Furthermore, mothers and their children share a similar socioeconomic environment and that may affect mothers' and children's health. To address this endogeneity problem, we instrument mothers' anemic status by the leave-out average anemia rate among rural women at the district level. A similar instrument i.e., endogenous variable aggregated at the district or state level has been used in prior studies (Fruehwirth et al., 2019).

The first-stage regression equation is:

$$H_{mi} = \alpha_2 + \beta_3 * average H_{md} + \beta_4 X_{ci} + \eta_d + \epsilon_{m_i}$$
(2)

And the predicted value of the mother's anemia from equation (2) is used in the second stage:

$$H_{ci} = \alpha_3 + \beta_5 * \widehat{H_{mi}} + \beta_6 * X_{ci} + \theta_d + \epsilon_{ci}$$
(3)

The parameter β_5 is the measure of intergenerational persistence of health between mothers and children. $\widehat{H_{ml}}$ is the predicted value of mother's health from equation (2). The instrument relevance condition is satisfied when the district-level average anemia rate is strongly correlated with the mother's anemic status. The exclusion restriction is not testable, but it is reasonable to argue that the exclusion restriction is likely to be satisfied after including a robust set of controls and district-fixed effects.

3. Findings

Table 2 shows that after adjusting for the mother's height, there is a statistically significant association between the mother's anemic status and the anemic status of the children. If a mother is anemic, the anemia probability for children increases by 10.4 percentage points (pp) (col 4). The intergenerational correlation in Hb is 0.14.²

Our preferred 2SLS estimates are presented in Table 3. The first-stage results show that the instrument is statistically significantly correlated with the anemic status of the mother. The *F*-statistics on the excluded instrument is more than 10, indicating that the instrument is strongly correlated with the endogenous variable and that the instrument does not suffer from weak instrument problems. Results further show that the instrument passes the weak-identification test and is robust to weak instrument inference. The 2SLS estimates are

¹Similar findings were reported in Onyeneho et al., (2019). The authors find a Hb correlation of 0.1 and 0.2 for father-child and mother-child samples, respectively.

positive and statistically significant. Col (2) shows that when a mother's Hb level increases by one unit (1 g/dL), children's Hb level increases by 0.40 g/dL. Similarly, children of the anemic mother are 26 pp more likely to be anemic (col 4). The 2SLS estimates are almost 2.5 times larger than the OLS estimates, suggesting that the OLS estimates underestimate the true intergenerational correlation between mothers' and children's health. The OLS and 2SLS estimates cannot be compared directly because the two estimates refer to different population groups. The 2SLS estimate gives the local average treatment effect (LATE) for a subgroup of the population, while OLS estimates the average treatment effect (ATE) for the whole population. LATE can be greater than ATE if there are heterogeneous treatment effects. For the results presented here, the IHA correlation is larger for the LATE subgroup than the whole population.

Previous studies have shown that maternal height captures the genetic factors and affects the nutrition-related intergenerational transmission of health, therefore, it is important to control for maternal height in the empirical model. The coefficients for the mother's height are positive and statistically significant, but the effect size is small. The results in Table 2 show that the inclusion of the mother's height does not change the IHA coefficient much, indicating a muted role of genetic factors in the transmission of health across generations. Our results allude to a greater role of socio-economic environments in shaping the correlation between mother and child anemia since mothers and children share the same socio-economic environment.

Furthermore, the subgroup analysis by wealth index shows that the IHA correlation in Hb levels and any form of anemia is substantially larger for the bottom two wealth quintiles compared with the top two wealth quintiles (Fig. 1). The detailed regression results are reported in Table A1. The pattern of wealth gradient in Fig. 1 indicates that children are more likely to be anemic if they are born in households with lower economic resources. This suggests that improvement in the economic status of the household will limit the extent to which child health is tied to mother health. Government programs designed to improve the economic conditions of populations may break the intergenerational transmission of poor health across generations. Given the confidence interval overlap, our findings related to the wealth quintile, while interesting, should be interpreted with caution.³

4. Conclusion

This paper is a handful of studies to investigate the IHA between mother and child health in a causal framework in India, except the study by Onyeneho et al., (2019). Analyzing large and nationally representative data, we find a high degree of intergenerational persistence in anemia. The study contributes to limited evidence on IHA in developing countries and in the context of anemia which is not self-reported and, therefore, is less likely to suffer from measurement errors. There could be some concerns about the exclusion restriction, for example, micronutrients in the crops grown in the region may affect the instrument as well as the outcome variable; therefore, our 2SLS estimates should be interpreted with caution. These are important findings that highlight the need to address anemia and other health issues in a comprehensive and targeted manner, particularly among disadvantaged populations from lower wealth groups. Future studies should consider other potential factors that may contribute to the persistence of anemia and other health problems across generations, including access to quality health care, nutrition, religion, gender, and other social and economic determinants of health.

Data availability

The authors do not have permission to share data.

Acknowledgment

The authors thank the editor, Prof. Audra Bowlus, an anonymous referee, and Timothy Halliday for valuable suggestions.

 $^{^2}$ It should be noted that overlapping confidence intervals is not an appropriate test of equality because this method does not account for correlation between the estimators, nor does it account for the multiple tests performed. (Schenker and Gentleman, 2001). It is quite possible that coefficients in Fig. 1 are statistically different despite the overlap in the confidence intervals.

References

Almond, D., Currie, J., Duque, V., 2018. Childhood Circumstances and Adult Outcomes: Act II. Journal of Economic Literature. 56 (4): 1360-1446.

Azam, M., 2015. Intergenerational Occupational Mobility among Men in India. The Journal of Development Studies. 51(10), 1389-1408.

Akbulut-Yuksel, M., Kugler, A.D., 2016. Intergenerational persistence of health: do immigrants get healthier as they remain in the U.S. for more generations? Econ. Hum. Biol. 23, 136-148.

Currie, J., Moretti, E., 2007. Biology as destiny? short- and long-run determinants of intergenerational transmission of birth weight. J. Lab. Econ. 25 (2), 231-264.

Currie, J., Vogl, T., 2013. Early-life health and adult circumstance in developing countries. Annual Review of Economics. 5:1-36.

Fruehwirth, J.C., Iyer, S., Zhang, A., 2019. Religion and Depression in Adolescence Journal of Political Economy. 127:3, 1178-1209.

Halliday T.J., Mazumder, B., Wong, A., 2021. Intergenerational mobility in self-reported health status in the US. Journal of Public Economics. 193, 104307.

International Institute for Population Sciences (IIPS) and ICF. 2021. National Family Health Survey (NFHS-5), India, 2019-21: Andhra Pradesh. Mumbai: IIPS.

Lochner, L., 2008. Intergenerational Transmission, in L. Blume and S. Durlauf (eds.), *New Palgrave Dictionary of Economics*, 2nd Edition, Palgrave Macmillan.

Onyeneho, N.G., Corsi, D.J., Kurpad, A., Subramanian, S.V., 2019. Intergenerational influences on childhood anaemia. Matern Child Nutr. 15(2):e12673.

Perumal, N., Blakstad, M.M., Fink, G., Lambiris, M., Bliznashka, L., Danaei, G., Sudfeld, C.R., 2021. Impact of scaling up prenatal nutrition interventions on human capital outcomes in low- and middle-income countries: a modeling analysis. The American Journal of Clinical Nutrition. 114 (5), 1708–1718.

Schenker, N., Gentleman, J.F. 2001. On Judging the Significance of Differences by Examining the Overlap Between Confidence Intervals, The American Statistician, 55:3, 182-186, DOI: <u>10.1198/000313001317097960</u>

Variables	Mean	St. Dev.		
Outcomes				
Child Hb level	10.59	1.38		
Mother Hb level	11.52	1.47		
Child is anemic (%)	0.58	0.50		
Mother is anemic (%)	0.56	0.50		
District anemia rate (%)	0.56	0.12		
Child demographics				
Female (%)	0.48	0.50		
Birth order	2.26	1.40		
Age	2.23	1.30		
Households' demographics				
Mother's education	6.07	5.11		
Religion	0.73	0.45		
Access to improved toilet	0.51	0.50		
Social group (SC-ST)	0.39	0.49		
Wealth groups				
Poorest	0.27	0.44		
Poor	0.24	0.24		
Middle	0.20	0.39		
Rich	0.16	0.37		
Richest	0.13	0.34		
Rural (%)	0.76	0.42		
Number of districts	628			
Number of states	35			

Table 1Descriptive statistics for analytical sample (N= 198,203)

Notes: SC-ST: Scheduled caste and scheduled tribe; mother's education is in years of schooling

0	Child's Hb level		Child i	s anemic
	(1)	(2)	(3)	(4)
Mother's Hb level	0.138*** (0.003)	0.143*** (0.003)		
Mother is anemic			0.104*** (0.004)	0.104*** (0.004)
Mother's height	No	0.003*** (0.0008)	No	-0.00009*** (0.00003)
Child controls	Yes	Yes	Yes	Yes
Household controls	Yes	Yes	Yes	Yes
District fixed effects	Yes	Yes	Yes	Yes
Observations	198.203	198.203	198.203	198.203

Table 2: Intergenerational correlation in Hb and anemia, OLS results

Notes: Hb- hemoglobin level in grams per deciliter (gm/dL). Robust standard errors, clustered at the district level, are reported in parentheses. Child controls include gender, birth order, and age, while household controls include the mother's education, religion, caste, wealth index, and rural dummy.

* Indicates significance at 10% level.

** Significance at 5% level.

*** Significant at 1% level of confidence.

	Child's l	Hb level	Child is anemic		
	(1)	(2)	(3)	(4)	
First-stage results					
District anemia rate	-2.43***	-2.43***	0.967***	0.968***	
	(0.131)	(0.131)	(0.020)	(0.020)	
F-test	344.56	345.89	2289.0	5.67	
Weak identification test					
Cragg-Donald Wald F statistic	767.58	769.90	1221.41	1222.81	
Kleibergen-Paap Wald rk F statistic	344.56	345.89	2289.0	2282.39	
Weak-instrument-robust inference					
Anderson-Rubin Wald test (p-value)	< 0.001	< 0.001	< 0.001	< 0.001	
Stock-Wright LM S statistic	< 0.001	< 0.001	< 0.001	< 0.001	
Two-stage least square (2SLS) results			_		
Mother's Hb level	0.397***	0.395***			
	(0.065)	(0.065)			
Mother is anemic			0.263***	0.262***	
			(0.046)	(0.046)	
Mother's height	No	0.0003*** (0.00008)	No	-0.00009*** (0.00003)	
Child controls	Yes	Yes	Yes	Yes	
Household controls	Yes	Yes	Yes	Yes	
District fixed effects	Yes	Yes	Yes	Yes	
Observations	198,203	198,203	198,203	198,203	

Table 3: Intergenerational correlation in Hb and anemia, 2SLS results

Notes: Hb- hemoglobin level in grams per deciliter (gm/dL). Robust standard errors, clustered at the district level, are reported in parentheses. Child controls include gender, birth order, and age, while household controls include the mother's education, religion, caste, wealth index, and rural dummy. * p < 0.10, ** p < 0.05, *** p < 0.01



Notes: Child controls include gender, birth order, and age, while household controls include the mother's education, mother's height, religion, caste, mother's height, and rural dummy. Using principal component analysis, the wealth index is calculated based on a household's ownership of selected assets, such as televisions and bicycles; materials used for housing construction; and types of water access and sanitation facilities. Households are divided into quintiles from the poorest to the richest.

Appendix See Table A1

Table A.1 Heterogeneous effects by wealth index, 2SLS estimates

	Child's Hb level				Child is anemic					
	Poorest	Poor	Middle	Rich	Richest	Poorest	Poor	Middle	Rich	Richest
	(1)	(2)	(3)	(4)	(5)	(1)	(2)	(3)	(4)	(5)
Mother's Hb level	0.72***	0.41***	0.342**	0.284**	0.509***					
	(0.228)	(0.112)	(0.146)	(0.116)	(0.179)					
Mother is anemic						0.544***	0.343***	0.237**	0.110	0.192
						(0.172)	(0.098)	(0.099)	(0.090)	(0.119)
Child controls	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Household controls	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
District fixed effects	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Observations	52,898	46,891	39,477	32,594	26,343	52,898	46,891	39,477	32,594	26,343

Note: Hb- hemoglobin level in grams per deciliter (gm/dL). Robust standard errors, clustered at the district level, are reported in parentheses. Child controls include gender, birth order, and age, while household controls include the mother's education, mother's height, religion, caste, wealth index, and rural dummy. * p < 0.10, ** p < 0.05, *** p < 0.01