

Ankel-Peters, Jörg et al.

Working Paper

A Protocol for Structured Robustness Reproductions and Replicability Assessments

I4R Discussion Paper Series, No. 143

Provided in Cooperation with:

The Institute for Replication (I4R)

Suggested Citation: Ankel-Peters, Jörg et al. (2024) : A Protocol for Structured Robustness Reproductions and Replicability Assessments, I4R Discussion Paper Series, No. 143, Institute for Replication (I4R), s.l.

This Version is available at:

<https://hdl.handle.net/10419/301917>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.

No. 143

I4R DISCUSSION PAPER SERIES

A Protocol for Structured Robustness Reproductions and Replicability Assessments

Jörg Ankel-Peters

Anna Dreber

Florian Neubauer

Abel Brodeur

Magnus Johannesson

Julian Rose

August 2024

I4R DISCUSSION PAPER SERIES

I4R DP No. 143

A Protocol for Structured Robustness Reproductions and Replicability Assessments

**Jörg Ankel-Peters^{1,2}, Abel Brodeur³, Anna Dreber⁴, Magnus Johannesson⁴,
Florian Neubauer¹, Julian Rose^{1,2,5}**

¹*RWI – Leibniz Institute for Economic Research, Essen/Germany*

²*University of Passau/Germany*

³*University of Ottawa/Germany*

⁴*Stockholm School of Economics, Stockholm/Sweden*

⁵*Ludwig-Maximilians-University Munich/Germany*

AUGUST 2024

Any opinions in this paper are those of the author(s) and not those of the Institute for Replication (I4R). Research published in this series may include views on policy, but I4R takes no institutional policy positions.

I4R Discussion Papers are research papers of the Institute for Replication which are widely circulated to promote replications and meta-scientific work in the social sciences. Provided in cooperation with EconStor, a service of the [ZBW – Leibniz Information Centre for Economics](#), and [RWI – Leibniz Institute for Economic Research](#), I4R Discussion Papers are among others listed in RePEc (see IDEAS, EconPapers). Complete list of all I4R DPs - downloadable for free at the I4R website.

I4R Discussion Papers often represent preliminary work and are circulated to encourage discussion. Citation of such a paper should account for its provisional character. A revised version may be available directly from the author.

Editors

Abel Brodeur
University of Ottawa

Anna Dreber
Stockholm School of Economics

Jörg Ankel-Peters
RWI – Leibniz Institute for Economic Research

A Protocol for Structured Robustness Reproductions and Replicability Assessments

*JÖRG ANKEL-PETERS, ABEL BRODEUR, ANNA DREBER, MAGNUS JOHANNESSON, FLORIAN NEUBAUER, AND JULIAN ROSE**

August 2024

Robustness reproductions and replicability discussions are on the rise in response to concerns about a potential credibility crisis in economics. This paper proposes a protocol to structure reproducibility and replicability assessments, with a focus on robustness. Starting with a computational reproduction upon data availability, the protocol encourages replicators to prespecify robustness tests, prior to implementing them. The protocol contains three different reporting tools to streamline the presentation of results. Beyond reproductions, our protocol assesses adherence to the pre-analysis plans in the replicated papers as well as external and construct validity. Our ambition is to put often controversial debates between replicators and replicated authors on a solid basis and contribute to an improved replication culture in economics.

JEL: A11, C18

Keywords: replication, reproducibility, robustness, research transparency, meta-science.

* Ankel-Peters (corresponding author): RWI – Leibniz Institute for Economic Research and University of Passau (e-mail: peters@rwi-essen.de); Brodeur: University of Ottawa (e-mail: abrodeur@uottawa.ca); Anna Dreber: Stockholm School of Economics (e-mail: anna.dreber@hhs.se); Magnus Johannesson: Stockholm School of Economics (e-mail: magnus.johannesson@hhs.se); Neubauer: RWI – Leibniz Institute for Economic Research (e-mail: neubauer@rwi-essen.de); Rose: RWI – Leibniz Institute for Economic Research, University of Passau, LMU – Department of Sociology (rose@rwi-essen.de). Funding: This work was supported by the German Research Foundation (Deutsche Forschungsgemeinschaft, DFG), Grant No. 3473/1-1 within the DFG Priority Program META-REP (SPP 2317).

1. Introduction

In recent years, the replication crisis has become a significant concern in various scientific fields, including economics. Several studies highlight the gravity of the issue, suggesting that many published findings are not reproducible. For economics, Askarov et al. (2023), Brodeur et al. (2020, 2023), Ferraro & Shukla (2020, 2022), and Ioannidis et al. (2017) have provided compelling evidence for the prevalence of biases in publication processes, underpowered study designs and, more generally, questionable research practices. Low rates of replicability in direct replications (Camerer et al., 2018) resonate with concerns about limited generalizability of empirical findings (Holzmeister et al., 2024; Masselus et al., 2024; Peters et al., 2018; Vivalt, 2020). Replications as a potential remedy remain rare in economics (Ankel-Peters et al., 2023).

Since replications do not seem to occur naturally, systematic and coordinated replication and reproduction programs have emerged (Brodeur et al., 2024a). Following several large-scale replicability and computational reproducibility projects conducted over the past decade (Camerer et al., 2016, 2018; Chang & Li, 2015; Open Science Collaboration, 2015), the boundary has recently been pushed to meta-robustness reproductions (Brodeur et al., 2024b; Campbell et al., 2024). Such projects have a higher explanatory power and visibility than stand-alone replications, and reduce adverse effects on the careers of replicators (Brodeur et al., 2024a).¹ This paper proposes a protocol to structure the post-publication assessment in empirical economics, focusing on computational and robustness reproductions.

A significant challenge in the reproducibility discourse is determining the failure or success of robustness reproductions where results are often subject to intense debates between replicated authors and replicators (Ankel-Peters et al., 2024; Ozier, 2021). While our protocol cannot resolve these debates – for example because perspectives on different analytical choices and robustness results are often subjective (Simonsohn et al., 2020) – it can help structure the exchange of arguments. Most notably, we draw on three tools to present the results of the robustness checks: Reproducibility indicators following Dreber and Johannesson (2024), a visual dashboard that assesses the alignment of original and robustness results in terms of

¹ Terminologically, we follow Dreber & Johannesson (2024) and distinguish between reproductions that redo the original analysis using the same data (computational reproduction) and those that conduct additional analyses (robustness reproduction) as well as replications that collect new data to implement the same study design (direct replication) and a similar study design (conceptual replication).

statistical significance, sign of the effect, and effect size (Bensch, 2024), and specification curves (Simonsohn et al., 2020).

The protocol encourages replicators to pre-register all robustness checks before implementation, thereby countering concerns about so-called null hacking by selective reporting in reproductions (Bryan et al., 2019). Beyond reproduction, the protocol also considers critical issues for assessing the generalizability and replicability of research findings that often receive insufficient attention in the economics literature, that is, external validity, construct validity, and pre-specification diligence. All these dimensions require subjective judgements in which we recommend replicators to be conservative in their assessment and follow an '*in dubio pro reo*' principle. The remainder of the paper is written from the perspective of replicators who adopt the protocol, and it follows the sequential workflow of our protocol with six key elements: Computational reproduction, code inspection, robustness reproduction, testing the paper's supportive analyses, pre-specification, as well as construct and external validity.

2. The Replicability Assessment Protocol

2.1. Data Availability and Computational Reproduction

Computational reproducibility refers to the ability to reproduce the results of a study using the original data and code, following the exact analytical procedures. Therefore, replicators begin by assessing whether the necessary material to reproduce the original paper is available and the ease of use. The precondition to conduct reproductions is the availability of data and code. Some journals have data editors and a priori provide the data (which still is often hampered by restrictions of proprietary data, see Vilhuber (2023)), but for many journals obtaining the data implies contacting the authors (Askarov et al., 2022; Brodeur et al., 2024c; Christensen & Miguel, 2018; Pérignon et al., 2019). Most leading journals have data policies that oblige the authors to share the replication package upon request, but still replicators are dependent on the authors' willingness to cooperate. Again, proprietary data can be an additional barrier. In their assessment of the reproduction package, replicators distinguish between the availability of raw data and analysis data.

As a next step, replicators conduct the computational reproduction. Any issues with the replication package encountered during this process, such as missing data or code, or the need to manually install specific packages, should be documented. When reporting the results, replicators categorize findings based on whether they are fully, partially, or non-reproducible, from both raw data and from analysis data. The distinction between raw and analysis data is crucial because, unlike analysis data, the availability of raw data allows for a comprehensive vetting of the analysis, including the construction of treatment and outcome variables, restriction of the sample, and other decisions that otherwise remain hidden. The decisive criterion for a study to be computationally reproducible is whether the results obtained via re-running the original code match those presented in the paper. If they do not, we consider the paper to be not computationally reproducible. Replicators should explicitly state whether deviations are considered minor or consequential. Similarly, we consider a paper not computationally reproducible if code is missing, making computational reproducibility not feasible using the provided package.

2.2. Code Inspection

When doing the computational reproduction, the replicators carefully inspect the code to identify *objective* (i.e. unambiguous) coding errors. Such errors may include mis-specifying a variable, accidentally coding a variable differently than claimed in the paper, handling missing data erroneously, or making mistakes in sensitive commands such as merging, appending, reshaping, or dropping data, and others. These objective errors contrast with subjective analytical decisions affecting the robustness of results that are informed by methodological or economic appraisals of the respective researcher – and that are often controversial (see below). In case errors are found, the replicator corrects them, and the corrected specification is then used as the basis for the robustness reproduction. It is important to document whether these objective errors are consequential for the original results or not.

2.3. Robustness Reproduction

A crucial step at the beginning of the robustness reproduction is to identify the main outcomes of the original paper. This is done by dissecting the abstract into claims and tracking down the

corresponding coefficient for each claim within the paper. The robustness reproduction then involves testing the stability of these main outcomes to different assumptions, model specifications, estimation techniques, samples, and alterations in the data cleaning and processing. The goal is to assess the sensitivity of the main outcomes to various analytical choices.

To avoid null hacking and selective reporting, the replicators pre-register robustness checks after inspecting the data and code but before implementing them. That is, robustness checks should be pre-specified after coding errors and other flaws have been removed. Robustness checks can also address crucial coding decisions to process the raw data towards the analysis data that are not discussed in the article. Any robustness checks added *after* pre-registration should be clearly labelled when reporting the results.

The protocol provides guidelines for selecting and implementing these robustness checks. The protocol suggests several dimensions for robustness checks: Definition of the sample, outcome and treatment variables, inclusion and definition of control variables, outlier management, handling of missing observations, and model selection. Replicators are free to choose additional robustness checks (e.g., adding or using other data), but the protocol emphasizes that each robustness check and its purpose should be clearly described, including a justification in terms of their economic and methodological legitimacy vis a vis the specification choices in the replicated paper.

Based on these robustness specifications, replicators assess the reproducibility of a study by combining three approaches: First, reproducibility indicators proposed by Dreber and Johannesson (2024). The reproducibility indicators check the degree of reproducibility for single outcomes in a study, or a group of studies, and they are directly comparable across studies. They measure the share of robustness tests supporting the original finding, changes in the average estimated effect and the variation across robustness tests. The second approach is a reproducibility dashboard that visualizes for each outcome variable whether original and robustness results align in terms of statistical significance, sign of the effect, and effect size. In addition, it provides information on relative effect sizes and variation indicators. The reproducibility indicators and the reproducibility dashboard can be created using the Stata command *repframe* (Bensch, 2024). Third, specification curves show the results of all implemented robustness checks in a multiverse. While the reproducibility indicators and the

reproducibility dashboard summarize the overall reproduction results, specification curves are more detailed and also reveal the consequences of individual robustness checks on effect size and significance (Simonsohn et al., 2020).

2.4. Testing the Paper's Supportive Analyses

The previous section of the protocol focused on the robustness of the main outcomes of the replicated paper and how they are estimated. Most papers offer additional statistical analyses to support the findings or address caveats. Such supportive analyses can be tested in this section of the protocol. Prevalent examples are identification strategy checks, weak first stage tests, statistical power analysis as well as attrition or multiple hypotheses testing corrections.

Our protocol recommends replicators identifying those statistical analyses in the paper that are important for the established argument – and test those. For example, they should critically reflect on the identification strategies by applying placebo tests or checking the parallel trends assumption for difference-in-differences studies, and other relevant checks. Testing other supportive analyses could include introducing multiple hypothesis testing, attrition corrections, and others to check if they lead to different results.

2.5. Pre-Analysis Plan

Although pre-registration and pre-analysis plans (PAPs) have increased significantly in recent years, they are most prevalent in experimental studies and hardly existent for work based on secondary data (Ferguson et al., 2023; Ofose & Posner, 2021). Even among experimental studies, many are pre-registered, but not pre-specified, that is, they lack a distinct pre-analysis plan (Brodeur et al., 2024d). A PAP outlines the analytical steps that researchers plan to take before examining the data and, thereby, helps reducing the risk of data mining and p-hacking. By adhering to a pre-specified analysis plan, researchers can ensure that their findings are not driven by post-hoc adjustments or selective reporting. Other critical elements hence are how vigorously the PAP determines the analysis and whether the paper adheres to the PAP (Ofose and Posner 2021). The assessment of the original study's PAP is an essential component of our protocol.

Our protocol prescribes to check whether the original study includes a PAP and whether the analysis was conducted according to the pre-specified plan. If a PAP is available, replicators shall compare the PAP to the analyses that are conducted in the published paper. This step is critical for understanding whether authors have used the PAP to tie their hands and reduce researcher degrees of freedom on the results.

2.6. External and Construct Validity

External validity refers to the extent to which the results of a study can be generalized to other settings, populations, or time periods (Peters et al., 2018). Construct validity is concerned with whether results of a study are generalizable to policy-relevant scaled versions of the treatment or theoretically relevant replicated versions of it (Esterling et al., 2023; Masselus et al., 2024). Our protocol checks whether sufficient information is provided about the treatment under evaluation for the reader to assess a paper's construct and external validity. Specific information about the operationalized treatment is also relevant if other researchers want to implement a sufficiently similar treatment design in a direct or conceptual replication. Both external and construct validity are crucial for assessing the broader applicability and relevance of a study's findings (Esterling et al., 2023). Our protocol includes a structured checklist-like assessment of external and construct validity, guiding replicators through a series of questions to evaluate these aspects. This step examines whether the assessed paper is not only internally consistent but whether it also provides the necessary information for the reader to evaluate to which broader contexts it is generalizable.

3. Conclusion

The reproducibility assessment protocol presented in this paper provides a structured and standardized approach to reproducing and evaluating the robustness of published economics studies. By focusing on key dimensions such as computational reproducibility, robustness checks, pre-analysis plans, and external and construct validity, the protocol offers a comprehensive framework for assessing the reproducibility of research findings. This methodological focus can be complemented on a case-by-case basis, for example if replicators sense that issues so far underrepresented in our protocol are critical to a replicated paper.

Statistical power, multiple hypothesis testing or ethical concerns in RCTs are natural candidates. Our protocol can also be extended into these directions so that ultimately a modular approach can be applied: replicators pick the protocolled dimensions that are most suitable for a replicated paper.

References

- Ankel-Peters, J., Fiala, N., & Neubauer, F. (2023). Do Economists Replicate? *Journal of Economic Behavior & Organization*, 212, 219–232. <https://doi.org/10.1016/j.jebo.2023.05.009>
- Ankel-Peters, J., Fiala, N., & Neubauer, F. (2024). Is economics self-correcting? Replications in the American Economic Review. *Economic Inquiry*, ecin.13222. <https://doi.org/10.1111/ecin.13222>
- Askarov, Z., Doucouliagos, A., Doucouliagos, H., & Stanley, T. D. (2022). The Significance of Data-Sharing Policy. *Journal of the European Economic Association*, jvac053. <https://doi.org/10.1093/jeea/jvac053>
- Askarov, Z., Doucouliagos, A., Doucouliagos, H., & Stanley, T. D. (2023). Selective and (mis)leading economics journals: Meta-research evidence. *Journal of Economic Surveys*, joes.12598. <https://doi.org/10.1111/joes.12598>
- Bensch, G. (2024). *Repframe. A Stata package to calculate, tabulate and visualize Reproducibility and Replicability Indicators based on multiverse analyses*. [Computer software]. <https://github.com/guntherbensch/repframe>
- Brodeur, A., Cook, N., & Heyes, A. (2020). Methods Matter: P-Hacking and Publication Bias in Causal Analysis in Economics. *American Economic Review*, 110(11), 3634–3660. <https://doi.org/10.1257/aer.20190687>
- Brodeur, A., Carrell, S., Figlio, D., & Lusher, L. (2023). Unpacking p-Hacking and Publication Bias. *American Economic Review*, 113(11). <https://doi.org/10.1257/aer.20210795>
- Brodeur, A., Esterling, K., Ankel-Peters, J., Bueno, N. S., Desposato, S., Dreber, A., Genovese, F., Green, D. P., Hepplewhite, M., Hoces De La Guardia, F., Johannesson, M., Kotsadam, A., Miguel, E., Velez, Y. R., & Young, L. (2024a). Promoting Reproducibility and Replicability in Political Science. *Research & Politics*, 11(1), 20531680241233439. <https://doi.org/10.1177/20531680241233439>
- Brodeur, A., Mikola, D., & Cook, N. (2024b). Mass Reproducibility and Replicability: A New Hope. *I4R Discussion Paper Series, No. 107*. <https://doi.org/10.2139/ssrn.4790780>
- Brodeur, A., Cook, N., & Neisser, C. (2024c). P-Hacking, Data type and Data-Sharing Policy. *The Economic Journal*, 134(659), 985–1018. <https://doi.org/10.1093/ej/uead104>
- Brodeur, A., Cook, N. M., Hartley, J. S., & Heyes, A. (2024d). Do Preregistration and Preanalysis Plans Reduce p-Hacking and Publication Bias? Evidence from 15,992 Test Statistics and Suggestions for Improvement. *Journal of Political Economy Microeconomics*, 2(3), 527–561. <https://doi.org/10.1086/730455>
- Bryan, C. J., Yeager, D. S., & O'Brien, J. M. (2019). Replicator Degrees of Freedom Allow Publication of Misleading Failures to Replicate. *Proceedings of the National Academy of Sciences*, 116(51), 25535–25545. <https://doi.org/10.1073/pnas.1910951116>
- Camerer, C. F., Dreber, A., Forsell, E., Ho, T.-H., Huber, J., Johannesson, M., Kirchler, M., Almenberg, J., Altmejd, A., Chan, T., Heikensten, E., Holzmeister, F., Imai, T., Isaksson, S., Nave, G., Pfeiffer, T., Razen, M., & Wu, H. (2016). Evaluating Replicability of Laboratory Experiments in Economics. *Science*, 351(6280), 1433–1436. <https://doi.org/10.1126/science.aaf0918>
- Camerer, C. F., Dreber, A., Holzmeister, F., Ho, T.-H., Huber, J., Johannesson, M., Kirchler, M., Nave, G., Nosek, B. A., Pfeiffer, T., Altmejd, A., Buttrick, N., Chan, T., Chen, Y., Forsell, E., Gampa, A., Heikensten, E., Hummer, L., Imai, T., ... Wu, H. (2018). Evaluating the replicability of social science experiments in Nature and Science between 2010 and 2015. *Nature Human Behaviour*, 2(9), 637–644. <https://doi.org/10.1038/s41562-018-0399-z>

- Campbell, D., Brodeur, A., Dreber, A., Johannesson, M., Kopecky, J., Lusher, L., & Tsoy, N. (2024). The Robustness Reproducibility of the American Economic Review. *I4R Discussion Paper Series*, No. 124.
- Chang, A. C., & Li, P. (2015). Is Economics Research Replicable? Sixty Published Papers from Thirteen Journals Say “Usually Not.” *Finance and Economics Discussion Series*, 2015(83), 1–26. <https://doi.org/10.17016/FEDS.2015.083>
- Christensen, G., & Miguel, E. (2018). Transparency, Reproducibility, and the Credibility of Economics Research. *Journal of Economic Literature*, 56(3), 920–980. <https://doi.org/10.1257/jel.20171350>
- Dreber, A., & Johannesson, M. (2024). A framework for evaluating reproducibility and replicability in economics. *Economic Inquiry*, ecin.13244. <https://doi.org/10.1111/ecin.13244>
- Esterling, K. M., Brady, D., & Schwitzgebel, E. (2023). The Necessity of Construct and External Validity for Generalized Causal Claims. *I4R Discussion Paper Series* No. 18, 18.
- Ferguson, J., Littman, R., Christensen, G., Paluck, E. L., Swanson, N., Wang, Z., Miguel, E., Birke, D., & Pezzuto, J.-H. (2023). Survey of open science practices and attitudes in the social sciences. *Nature Communications*, 14(1), 5401. <https://doi.org/10.1038/s41467-023-41111-1>
- Ferraro, P. J., & Shukla, P. (2020). Feature—Is a Replicability Crisis on the Horizon for Environmental and Resource Economics? *Review of Environmental Economics and Policy*, 14(2), 339–351. <https://doi.org/10.1093/reep/reaa011>
- Ferraro, P. J., & Shukla, P. (2022). Credibility Crisis in Agricultural Economics. *Applied Economic Perspectives and Policy*, 1–17. <https://doi.org/10.1002/aepp.13323>
- Holzmeister, F., Johannesson, M., Böhm, R., Dreber, A., Huber, J., & Kirchler, M. (2024). Heterogeneity in effect size estimates. *Proceedings of the National Academy of Sciences*, 121(32), e2403490121. <https://doi.org/10.1073/pnas.2403490121>
- Ioannidis, J. P. A., Stanley, T. D., & Doucouliagos, H. (2017). The Power of Bias in Economics Research. *The Economic Journal*, 127(605), F236–F265. <https://doi.org/10.1111/eoj.12461>
- Masselus, L., Petrik, C., & Ankel-Peters, J. (2024). Lost in the Design Space? Construct Validity in the Microfinance Literature. <https://doi.org/10.31219/osf.io/nwp8k>
- Ofosu, G. K., & Posner, D. N. (2021). Pre-Analysis Plans: An Early Stocktaking. *Perspectives on Politics*, 1–17. <https://doi.org/10.1017/S1537592721000931>
- Open Science Collaboration. (2015). Estimating the reproducibility of psychological science. *Science*, 349(6251), aac4716. <https://doi.org/10.1126/science.aac4716>
- Ozier, O. (2021). Replication Redux: The Reproducibility Crisis and the Case of Deworming. *The World Bank Research Observer*, 36(1), 101–130. <https://doi.org/10.1093/wbro/lkaa005>
- Pérignon, C., Gadouche, K., Hurlin, C., Silberman, R., & Debonnel, E. (2019). Certify reproducibility with confidential data. *Science*, 365(6449), 127–128. <https://doi.org/10.1126/science.aaw2825>
- Peters, J., Langbein, J., & Roberts, G. (2018). Generalization in the Tropics – Development Policy, Randomized Controlled Trials, and External Validity. *The World Bank Research Observer*, 33(1), 34–64. <https://doi.org/10.1093/wbro/lkx005>
- Simonsohn, U., Simmons, J. P., & Nelson, L. D. (2020). Specification Curve Analysis. *Nature Human Behaviour*, 4(11), 1208–1214. <https://doi.org/10.1038/s41562-020-0912-z>
- Vilhuber, L. (2023). Reproducibility and Transparency versus Privacy and Confidentiality: Reflections from a Data Editor. *Journal of Econometrics*, 235(2), 2285–2294. <https://doi.org/10.1016/j.jeconom.2023.05.001>

Vivalt, E. (2020). How Much Can We Generalize From Impact Evaluations? *Journal of the European Economic Association*, 18(6), 3045–3089. <https://doi.org/10.1093/jeea/jvaa019>

Robustness and Replicability Assessment Protocol

This protocol serves as a comprehensive guide to conduct structured reproducibility assessments of empirical papers in economics. The focus of the protocol is on computational and robustness reproducibility, complemented by a checklist on pre-specification as well as construct and external validity.

1. Data Availability and Computational Reproduction

The replicator should assess whether all necessary material is available to reproduce the results of the original paper. To find reproduction packages, check the journal's website, the authors' websites and as a last resort reach out to the original authors.

Use the provided data and scripts of the original paper and run the code without any corrections/changes. This is to check whether the results match the ones in the paper. Besides the change of working directory, only install missing packages and record them but do not do more to try and run the code. For all other errors due to missing data or incorrect code, document these problems if you were able to identify them, but do not yet attempt to solve them. Note: In case you are not able to computationally reproduce the results of the original paper, but you are able to locate and solve the errors, then you should do so in Section 2 "Code Inspection". Similarly, any coding errors that do not prevent computational reproduction, but you think need to be corrected should be part of the code inspection (see Section 2).

If essential code or data is missing, or if you encounter challenges in computationally reproducing the paper for other reasons, consider reaching out to the original authors for clarification.

Use the following exemplifying table to present the results of the computational reproduction:

Table 1: Results of the computational reproduction

| | Fully | Partially | No |
|---------------------------------|-------|-----------|----|
| Raw data provided | | x | |
| Cleaning code provided | x | | |
| Analysis data provided | x | | |
| Analysis code provided | x | | |
| Reproducible from raw data | | x | |
| Reproducible from analysis data | x | | |

Reproducible from Analysis Data:

- Mark as "Fully" if the analysis data and analysis code are provided, and the reproduced results are identical to the paper's results.
- Mark as "Partially" if the analysis data and analysis code are provided, but there is slight deviation in the reproduced results from the paper.

Reproducible from Raw Data:

- Mark as "Fully" if both raw data and cleaning code are provided, and utilizing these inputs leads to identical results as in the paper.
- Mark as "Partially" if raw data and cleaning code are provided, but utilizing these inputs leads to slight deviations in the results.
- Mark as "No" if no raw data is provided, if raw data is provided without cleaning code, or the code fails to run.

2. Code Inspection

In this section, carefully check the code to identify any objective coding errors. Objective errors refer to errors such as mis-specifying a variable or including missing observations in the generation of variables. This contrasts with analytical decisions, which can be challenged in Section 4. Correct any coding errors and check whether the key results are affected. Document



the changes. Considering the volume of code involved, the process can be efficiently guided by focusing on critical parts, such as:

1. **Main Outcome(s):** Carefully review all code sections that are directly related to generating the main outcomes of the original study. This includes any calculations, transformations, or statistical procedures used to produce the primary results. Additionally, check whether the description in the paper matches the code.
2. **Treatment Variable(s):** Carefully review all code that handles the treatment variable, including any data manipulations, assignments, or computations related to the treatment group or conditions. Additionally, check whether the description in the paper matches the code.
3. **Control Variables:** Carefully review all code relevant to the control variables used in the pertinent estimations. This involves scrutinizing how these variables are handled and integrated into the analysis. Additionally, check whether the description in the paper matches the code.
4. **Estimation Process:** Carefully review all code sections that implement the statistical models or methods in the estimation process itself. This encompasses code used to derive the main outcomes. Additionally, check whether the description of the estimation in the paper matches the code.

Throughout the process of checking these four items, be vigilant in searching for coding errors such as:

- variable misspecification
- coding switcheroo (e.g., variable accidentally differently coded as claimed)
- outlier management
- missing data handling
- sensitive commands such as merging, appending, reshaping, and dropping data.

In case errors are detected, report the corrected results for all main outcomes. If the original analysis path (combination of analytical decisions) is considered a reasonable analysis path, this analysis path with the corrected results will also be included as one analysis path in the multiverse robustness tests as well as in the dashboard and specification curve (see below).

Table 2: Example for recording details on main outcomes

| | (1) Consumption | (2) Food Security |
|--------------------------------------|-------------------------------|-------------------------------|
| Name of display item | Table 1 | Table 1 |
| Column | Column 2 - Panel D | Column 3 - Panel D |
| Estimate | 0.579 | 0.431 |
| Standard Error | 0.175 | 0.062 |
| t/z-value | 3.309 | 6.952 |
| p-value | 0.001 | 0.04 |
| Level of analysis | Household | Household |
| Type of variable | Standardized index | Standardized index |
| Units | Deviation from baseline value | Deviation from baseline value |
| Number of observations | 880 | 885 |
| Sample | Full sample | Full sample |
| Estimation method | OLS | OLS |
| Fixed Effects | Hamlet fixed effect | Hamlet fixed effect |
| Standard Error type | Clustered at household level | Clustered at household level |
| Control variables | Baseline value of outcome | Baseline value of outcome |
| Outcome pre-specified | No | No |
| Outcome construction pre-specified | No | No |
| Minimum detectable effect size (MDE) | 0.49 | 0.17 |

Source: <https://bitss.github.io/ACRE/scoping.html#read-sum>

3. Scoping

Replicators should carefully read the original paper. The focus should be on identifying the main research question, one or several main outcomes, their presentation (Tables, Figures), and the methodology. The main outcomes are those that are mentioned most prominently in the abstract (note: both statistically significant and null-results qualify)¹ and will be subjected to robustness tests to assess the robustness reproducibility of these results. For selection of the main outcomes, the replicator may dissect the abstract into claims and subsequently track the corresponding coefficient(s) in the paper. In case several coefficients correspond to one

¹ Note that for a null result the calculation of the reproducibility indicators is slightly different, see Section 5 for details.



claim, the replicator may choose the one most prominently discussed in the paper (e.g., effect size discussed in the introduction). If no distinction can be made, the replicator can pick one of the qualifying coefficients randomly in a reproducible way.

To provide an overview of the main results, Table 2 can be used. Replicators can also compile a brief summary of the original paper and include it in the introduction of the reproduction.

4. Robustness reproduction

In the following, the protocol provides a concise guide outlining **possible** robustness checks along several dimensions (see Table 3 and Table 4). The specific implementation of these checks may vary depending on the paper. Replicators should carefully assess each dimension and incorporate all relevant ones in their robustness checks. Robustness checks are not limited to those listed here. The robustness tests should be carried out in a multiverse fashion (see Steegen et al., 2016 and Simonsohn et al., 2020), so that all reasonable combinations of the analytical decisions that replicators decide to vary in the robustness tests are estimated. Each such reasonable combination of analytical decisions in the multiverse is referred to as an “analysis path” below. Before proceeding with the implementation of these checks, the protocol recommends to first compile a comprehensive list of the specific checks the replicators intend to conduct for each of the selected main outcomes and formally register them, as described in step 4.1.

4.1. Registering robustness checks

At this stage, replicators have engaged intensively with the original paper, data, and code. They should compile a list of all robustness checks that they came up with during the first steps of the reproduction protocol alongside a short explanation for each robustness check. Replicators should, in a pre-analysis plan, preregister all analytical decisions they want to vary (e.g., outcomes, covariates, outlier management, etc.) alongside all choices within each analytical decision (e.g., which specific covariates to include in which specification). The original choice within each analytical decision should also be included if that is defined as a reasonable choice (note that if the replicators think that the original choice within one analytical decision is not



a reasonable choice and they only defined one reasonable choice for this decision, they should include this choice as one analytical decision with only one option in the multiverse robustness checks). An option is to upload the pre-specification document to OSF folder before implementing any robustness checks.

When implementing the pre-registered robustness checks, replicators should include the full multiverse of all reasonable combinations of analytical decisions and choices they have made. Reasonable refers to all choices and combinations that are sensible and can be backed by logical argumentation (and that are non-redundant). Note that one of the reasonable combinations of analytical decisions (one analysis path) will be the original analysis if replicators interpret this as one of the reasonable analysis paths.²

Should replicators realize during the implementation of the pre-registered robustness checks that they would like to apply more robustness tests than preregistered, they have the flexibility to introduce additional robustness checks if they find them to be pertinent during the reproduction process. However, it is crucial that any such additional checks are clearly labelled as “non-pre-registered” for transparency. Also these additional robustness checks should be conducted in a multiverse fashion so that repliactors add analytical decisions or added choices within each analytical decision and thereafter implement the multiverse analysis of all the reasonable combinations of analytical decisions (all the analysis paths).

4.2. Role of control variables


The protocol recommends discussing the inclusion of each control variable in the context of omitted variables and bad controls. Replicators compile a list of all control variables and assess for each variable whether they are a collider (opening a path to a collider) or a necessary control to avoid omitted variable bias. This exercise is repeated to justify the omission/inclusion of control variables in all robustness checks. Some stylized “Directed Acyclic Graphs” (DAGs) are shown below.

² In case the result for this analysis path was not computationally reproduced in step 2, it should still be included as an analysis path if it is interpreted as a reasonable analysis path but it would give different results from the original analysis. Similarly, if the replicators discover coding errors in step 4, the original analysis path after correcting for these coding errors should still be included as an analysis path if it is interpreted as a reasonable analysis path.


A. Bad controls

The following examples are from Cinelli et al. (2022):

Model 1:



Model 2:




Model 1: Collider bias, controlling for Z opens the path $X \rightarrow Z \leftarrow Y$, but also the colliding path due to the latent parents of Y . (Example: Assume one wants to estimate the effect of attending lectures (X) on good grades (Y). One might then want to control for a poll about attending lectures and good grades (Z), but those who attend and have good grades are more likely to fill out the poll. This therefore introduces a collider bias.)


Model 2: Controlling for Z induces “selection bias” since it opens the colliding path $X \rightarrow Z \leftarrow U \rightarrow Y$. (Example: Assume that job satisfaction has no relationship with income, but education has. At the same time, there is an unobserved variable “work-life balance” that influences income and job satisfaction but has no relationship with education. Controlling in this scenario for income opens up a pathway between education and job satisfaction.)

B. Good controls (leading to omitted variable bias if not controlled for)

Model 3:



Model 4:



Model 1: Z is a common cause of both X and Y (confounder) → controlling closes the back-door path from X to Y.

Model 2: Z is not a common cause of both X and Y, but controlling for Z blocks the back-door path from X to Y due to unobserved confounder U.

4.3. Dimensions and Robustness Checks

Below, the protocol provides a list of dimensions that are relevant to the data processing and analytic decision of researchers (think of each dimension as one analytic decision that can be potentially varied in the robustness tests; but it is also fine to define and include more than one analytic decision within a dimension that is varied in the robustness tests). Researchers take hundreds of decisions when preparing data for analysis. While many of these decisions are taken subconsciously, they can have a large impact on the results. In this part of the reproduction, the protocol aims to show to what extent the results of the original paper depend on these decisions. Along several dimensions, replicators assess the reasonable analytical decisions and choices in each decision for each of the main outcomes. In doing so, they consider the following dimensions (and any other dimensions and analytical decisions that might be relevant for the specific paper):

Table 3: Suggested dimensions for robustness checks

| Dimension |
|---|
| Definition of the analysis sample |
| Definition of the outcome variable(s) |
| Definition of treatment variable(s) |
| Inclusion and definition of control variables |
| Outlier management |
| Missing observations |
| Model selection |

Table 4 provides an example to present the robustness checks by showing for each main outcome the change to the original data manipulation alongside an explanation. It includes each analytical decision that is varied or deviates from the original analysis as one row in the table. Note that if the original analysis is considered as one of the reasonable choices for an analytical decision that is varied in the robustness checks, then this original analysis choice

should be included among the alternative options in Table 4. Note further that if replicators do not think that the original choice is a reasonable choice and only define one reasonable choice for this analytical decision, they should include this analytical decision as one row in the Table with only one choice in the column “Choice(s) in robustness checks”. When presenting their analyses, replicators should sufficiently describe and motivate their implemented robustness checks.

Table 4: Example of robustness checks

| Nr. | Main outcome in Table 1 | Analytical decision | Rationale & Challenge | Choice(s) in Robustness Checks (include also the original analysis choice if that is considered one of the reasonable analytical choices) |
|-----|-------------------------|---------------------|---|---|
| 1. | (1) | Outcome Variable | The authors use indices for all outcomes. This reduces the problem of MHT and allows to include several dimensions into one measure. Yet, indices can be created in many different ways. | 1. Original choice 2. Alter definition of outcome index by including a more comprehensive set of input variables |
| 2. | (1) | Control Variables | Timing of EL4. The baseline took place over 5 months. In particular for income, food security, and consumption, the timing of the survey could matter. | 1. Original choice 2. Add controls for the month in which the interview took place |
| 3. | (1) | Control Variables | The authors do not use any baseline controls. At baseline, slight imbalance for income. | 1. Original choice 2. Add baseline controls for all outcome variables. |

Note: Column “Main outcome in Table 1” should include a reference to the main claims (main outcomes) from Table 1 that the numbered item refers to. Column “Analytical decision” captures the part of the analysis the robustness check refers to. Column “Rational & Challenge” captures the reason for the robustness check and which problems of the original paper it tries to address. Column “Choice(s) in Robustness Checks” includes the analytical choices included in the multiverse robustness checks (note that this can be one choice if the original analysis choice is not considered reasonable and only one alternative reasonable choice is included in the robustness checks).

As mentioned above, the robustness tests of each main outcome are estimated as a multiverse analysis for all the reasonable combinations of the analytical decisions (the combinations of the rows in Table 4).

5. Reporting of reproducibility indicator results

Replicators should report a table including all the results for each of the main outcomes for the reproducibility indicators described below alongside a short interpretation of the results. As described below, different indicators need to be calculated for those original main outcomes reported as statistically significant results and those reported as null results.

The indicators are calculated based on the data extracted from the original study and all robustness checks. In extracting that data adhere to the following general principles:

- A. The correct sign of the effect size and the t/z-value of each analysis path in the multiverse robustness tests need to be recorded, as this is important for the calculation of the indicators.
- B. Some “main outcomes” may involve more than one regression coefficient, such as, for example, the coefficient of a variable and the coefficient of the squared variable for studies testing non-linear relationships. In such cases where the test of a hypothesis of a “main result” depends on the statistical significance of two coefficients both coefficients should be recorded.
- C. For studies using instrumental variables only the 2nd stage results should be included to calculate the reproducibility indicators (i.e., the p-value, effect size and t/z-value of the 2nd stage regression coefficient is used for estimating the reproducibility indicators).
- D. In case the original paper already includes one or more (plausible) robustness tests, replicators should include these analysis paths (robustness tests) in the calculation of the reproducibility indicators in the same way as the analyses paths were added.

To calculate the indicators, the Stata program “repframe” can be used, which is explained in more detail and can be downloaded here: <https://github.com/guntherbenssch/repframe>. See also Dreber and Johannesson (2023) on the calculation of these reproducibility indicators.

Important: If the command “repframe” is used, note that the following procedures (except the specification curve) are all performed by the command so that no further data manipulations are required. Only note that in the two cases where (i) different effect size units are used (e.g. log and non-log) or (ii) the original analysis is included as one analysis path in the multiverse robustness, the help file of the command with instructions on the options “sameunits()” and “orig_in_multiverse()” needs to be consulted.

Reproducibility Indicators for original results reported as statistically significant:

Important: Replicators should estimate the following five indicators only if an original main result is reported as statistically significant in the original paper. An original main result reported as statistically significant in the original paper is defined as: a two-sided p -value < 0.05 or an original result with a two-sided p -value > 0.05 reported as significant by, for instance, using a star for $p < 0.10$ in reporting results or referring to this result as significant at the 10% level, marginally significant or some similar term. If a main result is not defined as an “original main result reported as statistically significant” according to the above definition, it is defined as a “null result”; for main outcomes reported as null results, the three indicators further down are calculated.

1. **Statistical significance indicator:** Share of robustness tests analysis paths in the same direction as the original effect and a two-sided p -value < 0.05 .

Note:

- The original analysis path **should not be included** among the analysis paths in the estimation of the statistical significance indicator.
- For “main outcomes” involving two coefficients, both these coefficients need to have an effect in the same direction as the original result and have a two-

sided p -value < 0.05 to count as a statistically significant robustness test analysis path for the statistical significance indicator.

2. **Relative effect size indicator:** Mean effect size of all the robustness tests analysis paths divided by the original effect size.

Notes:

- The original analysis path **should not be included** among the analysis paths in the estimation of the mean effect size of all the robustness tests analysis paths.
- Only the robustness tests analysis paths that use the same effect size units as the original result should be included. This excludes, for example, robustness tests analysis paths with effect sizes in logs, while the original results are in levels.
- The “relative effect size indicator” should not be estimated for “main outcomes” involving two coefficients.

3. **Relative t/z-value indicator:** Mean t/z-value of all the robustness tests analysis paths divided by the t/z-value of the original effect.³

Notes:

- The original analysis path **should not be included** among the analysis paths in the estimation of the mean t/z-value of all the robustness tests analysis paths.
- For “main outcomes” involving two coefficients, the “relative t/z-value indicator” should be based on the average t/z-value of these two original coefficients and the t/z-value of a robustness test analysis path should be based on the average t/z-value of these two coefficients in the robustness test analysis path.⁴

4. **Variation indicator: effect sizes:** The standard deviation in effect sizes of all the robustness tests analysis paths divided by the standard error of the original effect size.

³ If replicators encounter a different test statistic, please convert the p -value of the test to the equivalent z -value. This can apply both to the original t/z-value if the original test is based on some other test statistic, and/or the robustness checks if some or all robustness checks use some other test statistic.

⁴ In case the command “repframe” is not used: if the main result involves two coefficients and one has a positive and one a negative coefficient, the two original t/z-values must both be assigned positive signs in estimating the average t/z-value of the original result; and for the average t/z-value of a robustness test analysis path the t/z-value of a coefficient must be assigned a positive sign if the coefficient is in the same direction as the original coefficient and a negative sign if the coefficient is in the opposite direction of the original coefficient.



Notes:

- This indicator should not be estimated for “main outcomes” involving two coefficients.
- If the original analysis is included as one analysis path in the multiverse robustness test that analysis path **should be included** in the estimation of the standard deviation.
- Only the robustness tests analysis paths that use the same effect size units as the original result **should be included** in the estimation of the standard deviation. This excludes, for example, robustness tests analysis paths with effect sizes in logs, while the original results are in levels.

5. **Variation indicator: t/z-values:** Standard deviation of t/z-value of all the robustness tests analysis paths.⁵

Notes:

- If the original analysis is included as one analysis path in the multiverse robustness test, that analysis path **should be included** in the estimation of the standard deviation.
- For “main outcomes” involving two coefficients, the “variation indicator: t/z-values” should be based on the average t/z-value of these two coefficients in the robustness test analysis path.⁶

Reproducibility Indicators for original results reported as null results:

Important: The following three indicators are only estimated if an original main result is reported as a null result. For main outcomes reported as statistically significant in the original paper, the five indicators above apply.

⁵ If a different test statistic is encountered, the p-value of the test needs to be converted to the equivalent z-value. This can apply both to the original t/z-value if the original test is based on some other test statistic, and/or the robustness checks if some or all robustness checks use some other test statistic.

⁶ In case replicators are not using the command “repframe” if the main result involves two coefficients and one has a positive and one a negative coefficient, the two original t/z-values must both be assigned positive signs in estimating the average t/z-value of the original result; and for the average t/z-value of a robustness test analysis path the t/z-value of a coefficient must be assigned a positive sign if the coefficient is in the same direction as the original coefficient and a negative sign if the coefficient is in the opposite direction of the original coefficient.



1. **Statistical significance indicator:** Share of robustness tests analysis paths with a two-sided p-value >0.05

Notes:

- The original analysis path **should not be included** among the analysis paths in the estimation of the statistical significance indicator.
 - For “main outcomes” involving two coefficients, such a “main result” will count as a null result ($P > 0.05$) in the statistical significance indicator if at least one of these two coefficients have a p-value > 0.05 .
2. **Variation indicator: effect sizes:** The standard deviation in effect sizes of all the robustness tests analysis paths divided by the standard error of the original effect size.

Notes:

- This indicator should not be estimated for “main outcomes” involving two coefficients.
 - If the original analysis is included as one analysis path in the multiverse robustness test, that analysis path **should be included** in the estimation of the standard deviation.
 - Only the robustness tests analysis paths that use the same effect size units as the original result should be included in the estimation of the standard deviation. This excludes, for example, robustness tests analysis paths with effect sizes in logs, while the original results are in levels.
3. **Variation indicator: t/z-values:** Standard deviation of t/z-value of all the robustness tests analysis paths.⁷

Notes:

- If the original analysis is included as one analysis path in the multiverse robustness test, that analysis path **should be included** in the estimation of the standard deviation.

⁷ If repliactors encounter a different test statistic, they need to convert the p-value of the test to the equivalent z-value. This can apply both to the original t/z-value if the original test is based on some other test statistic, and/or the robustness checks if some or all robustness checks use some other test statistic.



- For “main outcomes” involving two coefficients, the “variation indicator: t/z-values” should be based on the average t/z-value of these two coefficients in the robustness test analysis path.⁸


Aggregated reproducibility indicators on the paper level:

For papers with more than one main outcome included in the robustness tests, the Stata command “repframe” will automatically calculate the robustness indicators on the paper level as the average of the indicators across the main outcomes (e.g., if there are three main outcomes reported as statistically significant in the original paper, the paper level of the “statistical significance indicator” is estimated as the average of the “statistical significance indicator” for these three main outcomes). This estimation is done separately for main outcomes reported as statistically significant in the original paper and main outcomes reported as null results in the original paper (i.e., if an original paper includes at least one main outcome reported as statistically significant and at least one main outcome reported as a null result, there will be two aggregated values of the “statistical significance indicator” for this paper).

Reproducibility Dashboard

The Stata command “repframe” also visualizes a so-called *Reproducibility Dashboard* (see Figure 1). The dashboard highlights confirmatory and non-confirmatory robustness check results and provides indicators tailored to subsets of the analysis paths for which the indicators are most informative. As can be taken from the figure, they are tailored to whether original and robustness results are statistically significant or not. The Reproducibility Dashboard thereby provides a complementary set of statistical significance indicators, relative effective size indicators and variation indicators. Details on the indicators can be found at <https://github.com/guntherbenssch/repframe>.

⁸ In case replicators are not using the command “repframe”: if the main result involves two coefficients and one has a positive and one a negative coefficient, the two original t/z-values must both be assigned positive signs in estimating the average t/z-value of the original result; and for the average t/z-value of a robustness test analysis path the t/z-value of a coefficient must be assigned a positive sign if the coefficient is in the same direction as the original coefficient and a negative sign if the coefficient is in the opposite direction of the original coefficient.

Figure 1: Sample Reproducibility Dashboard

Notes: α = significance level; β = estimate; CI = confidence interval; Δ = mean absolute deviation; o = original estimate; bars over certain values refer to mean values, tildes to median values. A version of the dashboard is applied in Ankel-Peters et al. (2023).


Specification Curve

To complement the previous two reporting tools, specification curve for each main outcome can be used. This curve is based on the multiverse robustness tests that were already implemented above (i.e. the same analytical decisions and analysis paths as above). In case the original paper already includes one or more (plausible) robustness tests, they should be included among the analysis paths in the multiverse robustness tests if they are considered plausible analysis paths. In contrast to the previous reporting tools, the specification curves allow to assess the influence of individual robustness checks on the sensitivity of the results.

A specification curve as suggested by Simonsohn et al. (2020) and presented in Figure 2 should be generated. In case the replicators use Stata (version 16 or newer), the command “speccurve” produces a specification curve. In R the package “spear” can be used. In case an

analysis only includes few analysis paths, a specification curve may not be useful but the results can be shown in a table instead.

Figure 2: Specification Curve



Source: Simonsohn, U., Simmons, J. P., & Nelson, L. D. (2020). Specification curve analysis. *Nature Human Behaviour*, 4(11), 1208-1214.

6. Testing the Paper's Supportive Analyses

In this section, replicators are encouraged to assess additional statistical analyses provided by the original paper to support its findings. Prevalent examples are identification strategy checks, weak first stage tests, statistical power analysis as well as attrition or multiple hypotheses testing corrections. For example, replicators should critically reflect on the identification strategies by applying placebo tests or checking the parallel trends assumption for difference-in-differences studies, and other relevant checks.

Note that these checks are not included in the reproducibility indicator or dashboard.

7. Pre-Analysis Plan

Replicators should check whether the authors registered the study and published a pre-analysis plan (PAP). If the PAP is not publicly available, consider reaching out to the original authors. Please note, which of the following cases applies:

- a. No preregistration
- b. Preregistration without a PAP
- c. Preregistration with a PAP

If the PAP is available, it should be compared to the original paper and deviations highlighted.

- Does the PAP specify the exact design of the study, the exact data collection and exactly how all analyses and tests will be conducted?
- Does the paper follow the pre-registered design and data collection?
- Does the paper carry out all analyses and tests reported in the paper exactly as specified in the PAP?
- Does the paper report all analyses and tests as specified in the PAP?
- Does the paper report additional analyses and tests not specified in the PAP?

8. External and construct validity

Replicators should assess the extent to which the paper provides information regarding external validity and construct validity. Following Peters et al. (2018), Masselus et al. (2024), and Esterling et al. (2023) this checklist can be used:

Construct validity:

- Does the paper provide comprehensive information about the details of the intervention, that would allow other researchers/implementers to implement an intervention that is sufficiently similar to the intervention under evaluation?
- Does the paper describe whether the intervention is an existing intervention implemented by a real-world policy agency (as opposed to an intervention designed by

Leibniz Institute for
Economic Research
UConn
UNIVERSITY OF CONNECTICUT
INSTITUTE for
REPLICATION


researchers for academic purposes)? If not, does the paper describe how the intervention deviates from what is implemented in the real world and whether these deviations are costly?

Researcher involvement and special care:

- Does the paper describe to what extent the authors (or other researchers) were involved in designing the intervention (as opposed to, for example, an NGO or a governmental agency designing the intervention)?
- Does the paper describe who implemented the intervention, and to what extent the researchers were involved in the implementation?

Hawthorne & John Henry effects:

- Does the paper describe whether participants were aware of the randomization/being part of a study?

General Equilibrium Effects:

- Does the paper discuss general equilibrium effects (GEE)?

Generalizability:

- Does the paper determine the scope of generalization regarding the intervention under study?
- Does the paper determine the scope of generalization regarding the population and sample?

9. Documentation of reproduction

To ensure reproducibility of your reproduction, provide all code, data, and a read-me file in a reproduction package. Upload the reproduction package to our server before publishing the reproduction report and/or a working paper.



10. For guidelines on ensuring reproducibility, see:

- [How to Make Sure Your Research Paper is Reproducible? Evidence from 55 Papers. \[Guest Blog Post\] \(worldbank.org\)](#)
- [Reproducible Research - Dimewiki \(worldbank.org\)](#)

11. References

- Ankel-Peters, J., Bensch, G., & Vance, C. (2023). Spotlight on researcher decisions: Infrastructure evaluation, instrumental variables, and specification screening (No. 991). Ruhr Economic Papers.
- Cinelli, C., Forney, A., & Pearl, J. (2022). A crash course in good and bad controls. *Sociological Methods & Research*, 00491241221099552.
- Dreber, A., & Johannesson, M. (2024). A framework for evaluating reproducibility and replicability in economics. *Economic Inquiry*, ecin.13244. <https://doi.org/10.1111/ecin.13244>
- Esterling, K. M., Brady, D., & Schwitzgelb, E. (2023). The Necessity of Construct and External Validity for Generalized Causal Claims. I4R Discussion Paper Series No. 18.
- Masselus, L., Petrik, C., & Ankel-Peters, J. (2024). Lost in the Design Space? Construct Validity in the Microfinance Literature. <https://doi.org/10.31219/osf.io/nwp8k>
- Peters, J., Langbein, J., & Roberts, G. (2018). Generalization in the tropics—development policy, randomized controlled trials, and external validity. *The World Bank Research Observer*, 33(1), 34-64.
- Simonsohn, U., Simmons, J. P., & Nelson, L. D. (2020). Specification curve analysis. *Nature Human Behaviour*, 4(11), 1208-1214.
- Steege, S., Tuerlinckx, F., Gelman, A., & Vanpaemel, W. (2016). Increasing Transparency Through a Multiverse Analysis. *Perspectives on Psychological Science*, 11(5), 702–12.