# **ECONSTOR** Make Your Publications Visible.

A Service of

ZBW

Leibniz-Informationszentrum Wirtschaft Leibniz Information Centre for Economics

Smetana, Sergiy; Tamásy, Christine; Mathys, Alexander; Heinz, Volker

#### Article

# Sustainability Assessment of Agribusiness Clusters: A Case Study Based on Regional Sustainability Assessment Methodology

German Journal of Agricultural Economics (GJAE)

#### Provided in Cooperation with:

Gesellschaft für Wirtschafts- und Sozialwissenschaften des Landbaues e.V. (GEWISOLA)

*Suggested Citation:* Smetana, Sergiy; Tamásy, Christine; Mathys, Alexander; Heinz, Volker (2017) : Sustainability Assessment of Agribusiness Clusters: A Case Study Based on Regional Sustainability Assessment Methodology, German Journal of Agricultural Economics (GJAE), ISSN 2191-4028, Deutscher Fachverlag, Frankfurt a. M., Vol. 66, Iss. 3, pp. 159-171, https://doi.org/10.52825/gjae.v66i3.2066

This Version is available at: https://hdl.handle.net/10419/305012

#### Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

#### Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.



# WWW.ECONSTOR.EU

# Sustainability Assessment of Agribusiness Clusters: A Case Study Based on Regional Sustainability Assessment Methodology

Sergiy Smetana German Institute of Food Technologies, Quakenbrück, Germany

Christine Tamásy University of Vechta, Germany

Alexander Mathys ETH Zürich, Switzerland

Volker Heinz German Institute of Food Technologies, Quakenbrück, Germany

#### Abstract

Socioeconomic and environmental sustainability is becoming critical for the evaluation of regional development. Regional planning and policy development depend on accurate sustainability assessment for the estimation of genesis and evolution of agribusiness clusters. The lack of a universal methodology creates challenges for the analysis of regional issues and the role of separate industries in the sustainability of regional development. This research suggests and reviews the application of Regional Sustainability Assessment Methodology (RSAM) by using a case study approach with a special attention to agribusiness clusters. The empirical analyses reveal the importance of resource cycling in the regions, their dependency on external resources and self-sufficiency as indicators of regional sustainability (social, economic and environmental resource use efficiency) in a dynamic perspective, applicable to diverse regions and separate aspects of sustainability. The study areas include three sub-national level regions in Germany: two forming an agribusiness cluster, specialized in livestock-meat production (Vechta Landkreis and Cloppenburg Landkreis) and one control region (Hochsauerlandkreis). The application of RSAM revealed varying rates of resource use efficiency, selfsufficiency and dependency on external resources for both static and dynamic perspectives. The assessment indicated the potential of RSAM application for the static and dynamic assessment of regions with agribusiness clusters. At the same time, static analysis of separate aspects of sustainability (economic, social and environmental) was not informative. RSAM application to the separate types of resource analysis allowed identification of sustainability hotspots (unbalanced development, "weak sustainability" and increased dependency on separate resources).

#### **Key Words**

regional sustainability assessment; agribusiness cluster; case study approach; input-output analysis; life cycle analysis

## **1** Introduction

Modern rural agribusiness clusters are the drivers and the response to the globalization changes of food trade. VAN WITTELOOSTUIJN highlighted this paradox as one of the most important propositions for future research, indicating that: "globalisation will trigger regionalisation, and regionalisation will boost globalisation" (VAN WITTELOOSTUIJN, 2009). Despite the extensive literature about industrial development analysis in Germany (STERNBERG and LITZENBERGER, 2004; ROCHA and STERNBERG, 2005; BRENNER, 2006; ALECKE and UNTIEDT, 2008; TITZE, BRACHERT and KUBIS, 2011; BRACHERT et al., 2011; KAHL and HUNDT, 2015; WROBEL, 2015) the identification and classification of agribusiness clusters is a challenging task (HOFF-MANN, HIRSCH and SIMONS, 2015). It is connected with high data aggregation at the national level of analysis, when overall industrial production is taken into account without a proper detail level. Most of the areas of food production in such case could not be accounted as regional industrial clusters (TITZE et al., 2011). There are quite a few examples of using scientific methods to identify clusters linked to the food industry (STEINER et al., 2007; WANDEL, 2010; DIEZ-VIAL, 2011; GELLYNCK et al., 2011). They illustrate

the hurdles of agribusiness clusters identification and characterization as different methodological approaches result in sometimes quite opposite results. Such a great variety is caused by the differences in regional clusters definitions and assessment techniques.

Mentioned studies demonstrate a wide range of approaches towards the identification and assessment of rural agribusiness clusters. It is logical that the analysis of their development is a challenging task, especially in terms of their role in the network of global interconnected value chains, driving forces and evolution of agribusiness clusters, sustainable consequences of agribusiness clusters globalization. Industrial clusters development has a direct impact on the regional development, both in positive and negative perspective, depending on the state of cluster life cycle (BRENNER and GILDNER, 2006; MENZEL and FORNAHL, 2007; MENZEL and FORNAHL, 2010; BROEKEL et al., 2015). The sustainability assessment of rural agribusiness clusters from regional perspective is needed. It would indicate the endogenous and exogenous interconnection properties of the systems, relevant "hotspots" and identify the evolutionary dynamic paths of their development.

Despite multiple hurdles of agribusiness cluster identification for the regional scale, some authors (HOFFMANN et al., 2015) were able not only to identify clusters but also point towards their specialization (e.g. Cloppenburg and Vechta Lankreises for meat processing). They were able to perform analysis of the clusters with high level of disaggregation (up to the county level). Following the publications (MARTIN and SUNLEY, 2003; MARTIN and SUNLEY, 2007; HOFFMANN et al., 2015), our study concentrated on the comparative assessment of development sustainability of the identified agribusiness clusters.

The selection of agribusiness clusters, as a point of interest was dictated by their controversial nature. They are triggering the economic development of regions from one side and are responsible for more than 50% of all greenhouse gas emissions and more than 70% of water footprint globally (PACHAURI et al., 2014; SCHMIDT and MERCIAI, 2014; HOEKSTRA et al., 2012; PFISTER and BAYER, 2014). The highest share of the environmental impact of food is associated with meat production (SCHMIDT and MERCIAI, 2014; STEINFELD et al., 2006). It sets a lot of questions about the sustainability of development of the regions, which form agribusiness clusters relying on meat production. Regions of Vechta and Cloppenburg are known as an agribusiness cluster of intensive livestock production and meat processing (TAMÁSY, 2013; HOFFMANN et al., 2015; SMETANA et al., 2016). That is why these counties were selected as the main study area.

The aim of this paper is to empirically indicate sustainability issues of sub-national regions development influenced by cluster formation by using RSAM as an assessment tool. Following the previous research (SMETANA et al., 2015; SMETANA et al., 2016), the paper concentrates on the dynamic application of RSAM. For the dynamic sustainability assessment the study includes socioeconomic activities of the regions, which form agribusiness clusters. The analysis will focus on two similar rural regions, which compose the Oldenburger Münsterland and one rural-urban region, which will serve as a control area (Hochsauerlandkreis). All three regions of the same level (NUTS3) are located in the west of Northern Germany. The Oldenburger Münsterland region (Vechta Landkreis and Cloppenburg Landkreis) has a common historical development, economic structure and dynamics. The economic development of the region is based on intensive agricultural production, mainly livestock farming, and related food processing industry. The economic success of the Oldenburger Münsterland is connected with the placement of many globally operating agricultural and food businesses, which clearly indicate the region as an agribusiness cluster (HOFF-MANN et al., 2015). At the same time a high number of livestock and their concentration in the region cause over-supply of manure, air pollution, groundwater deterioration and related health issues (TAMÁSY, 2013). The third region (Hochsauerlandkreis), used for comparison, is characterized with economic development which relies on manufacturing and service industries. It is not known as an agribusiness cluster. Moreover, large areas (almost 40,000 ha) are preserved as natural reserves mainly due to middle mountain landscape. The economic structure of the region is based, in a big degree, on the service sector, development of the metal industry, electrical engineering, engineering and plastics industry. The region is also holding a central residual landfill site for waste treatment.

Therefore, this paper commences with some conceptual considerations and methodological background of sustainability assessment with application of RSAM for the identification of regional development paths. This is followed by the results of the regional sustainability comparison in terms of resource use efficiency and resource transferability. Using the three rural and semi-urban sub-national regions as a case study, static and dynamic properties of agribusiness clusters are analyzed and presented. The generalized regional analysis is followed by the identification of the critical aspects in regional development, affecting the sustainability. Finally, the paper concludes with a brief discussion of the results and application possibilities of RSAM.

# 2 **Conceptual Considerations**

Currently sustainability is a guiding approach for the development of humanity (ROBÈRT et al., 2013; CAMPBELL and GARMESTANI, 2012; O'RIORDAN and VOISEY, 1998). Even though, it might have certain conflicts of interpretation and understanding (VOINOV, 2008; GATTO, 1995), its application is proven to provide a sufficient guideline for regional planning. The idea of sustainability is in equilibrium of multiple dimensions included in three main pillars (social, economic and environmental). Such multi-criteria balance, which should also preserve resources in non-degraded state for future generations, demands for multi-criteria assessment techniques (MUNDA and SAISANA, 2011). The multidimensional social development triggered the advancement of corresponding sustainability assessment techniques. They covered a wide number of social (GUINÉE et al., 2011; ZAMAGNI, 2012; COOKE et al., 2005; FLORIDA et al., 2008; ABEL et al., 2010), economic (WECKMAN et al., 2008; BOSKER, 2009) and environmental aspects (BOSKER, 2009; RODRÍGUEZ and ARIAS, 2008; PHILLIPS, 2011). However, most of the assessment methods do not provide an answer to the integrated estimation of "a sustaining degree" for different regions (products, technologies etc.) with weighted interlinked value estimation for various assessment categories. Neither can they assess the developmental properties of industrial clusters. Moreover, advanced integrated or multi-criteria approaches, developed in the last decade (KURKA, 2013; CABELLO et al., 2014; AGOSTINHO and ORTEGA, 2012; SINGH et al., 2012; VAN PASSEL and MEUL, 2012; KOURTIT et al., 2014; TENERELLI and CARVER, 2012), revealed sustainability of an object either with a number of indices (not interlinked with each other) or concentrated only on a few indicators, important for the development. Therefore, at a given time, published and available to the public review methodologies do not provide an integrated sustainability assessment of multiple aspects of regional development.

Among the variety of sustainability assessment indices, life-cycle and input-output methods have a great potential for sustainability assessment of a regional development. Input-Output Analysis (IOA), well-known in economic studies, was successfully combined with environmental aspects (including life cycle assessment) and applied to the assessment of production economies at national level (LEONTIEF, 1951; HENDRICKSON et al., 1998; DANIELS et al., 2011). IOA techniques were also applied for the regional industrial cluster identification (TITZE et al., 2011). Providing its potential combination possibilities with Social Accounting Matrices (ALLAN et al., 2010; MORILLA et al., 2007; LEHMANN et al., 2013) it might become an integrated technique for sustainability assessment at national or regional levels. It has not been completed due to the hurdles of a single accounting unit selection, overlapping of social and economic aspects and regional variations.

Regional Sustainability Assessment Methodology (RSAM) is an original methodology developed to assess relative regional sustainability, based on adapted and combined methods of input-output table analysis, life cycle assessment and resource cycling indices (SMETANA et al., 2015; SMETANA et al., 2016). It is aimed to provide an assessment of relative sustainability levels for regions and indicate the "hotspot" aspects of regional development (SMETANA et al., 2016). RSAM, however, should identify the dynamic paths and opportunities for the development of the regions, based on the main economic activities, which have not been previously tested with real regional data. As Vechta and Cloppenburg counties are characterized as regions, which origin the agribusiness cluster of livestock-meat production (HOFFMANN et al., 2015; BRACHERT et al., 2011; TITZE et al., 2011) their dynamic analysis was selected as an appropriate testing ground for RSAM. We argue that amount, quality and changes of the natural (environmental), social and economic resources and their availability determine regional (local) cluster development paths. Such approach is well-supported in the literature (DEVINE-WRIGHT, FLEMING and CHADWICK, 2001; ADGER, 2003; JORDAN et al., 2010; BIRCH et al., 2010; BEN LETAIFA and RABEAU, 2013; DELGADO et al., 2016).

Despite multiple indicators of sustainability aiming at simplification of sustainability measurements (SINGH et al., 2012; SICHE et al., 2008; BÖHRINGER and JOCHEM, 2007; MAYER, 2008), RSAM is placed as a holistic assessment system, capable of multiple issues analysis and further adaptation. That is why it is based on a combination of Input-Output Analysis and efficiency estimations of resource flows, which account for the true cost of activities. It differentiates RSAM from common evaluations of sustainability, which are based on operational and professional judgement rather than on quantitative approaches (SHAKER, 2015; PHILLIPS, 2015). The applicability of RSAM for the identification of development paths of agribusiness clusters is an empirical task to solve further in the article.

# 3 Methods

Input-Output (IO) tables' analysis is a well-known methodology used for the indication of purchasing interdependencies between sectors of the economy (LEONTIEF, 1951; MILLER and BLAIR, 2009). It was improved with further developed IO techniques, which included environmental impacts (HENDRICKSON et al., 1998; MATTHEWS and SMALL, 2000) and social accounting (MORILLA et al., 2007; ALLAN et al., 2010). IO tables' analysis consists of linear balance models, which account for multiple connections of system elements. IO tables' analysis was a basis for RSAM data compilation, selected due to comparison abilities inside sectors and with other matrix systems and indices. The relation of IO tables based models and Life Cycle Assessment (HERTWICH, 2005) indicated the potential of RSAM for a combination of social, environmental and economic data in a single assessment system. The basis of RSAM on IO tables' analysis allowed data transparency and multiple combination and recombination abilities. They assured the flexibility of RSAM and its independence from specific data.

Traditional IO analysis perceives sectors of economy as producers and consumers at the same time. It, therefore, accounts for the transfer of resources from one industry (a producer) to all the others in the forms of a value or a price. The accounting of such transfers is usually presented in a form of matrix, which is then applicable for the matrix analysis. Moreover, some authors introduced the application of IO analysis to the matrices formed with physical units (mass, energy, number) (MILLER and BLAIR, 2009). Therefore, IO matrices are the representation of a certain amount of resources stored and the transfers of the resources between sectors. Balanced and non-balanced models of IO matrices are a viable presentation of the complex system character of regional systems and their economies.

RSAM followed a simple algorithm for the assessment of resource use in the regions. It required data collection for the comparable regions, inputoutput tables' construction, estimation of resources cycling in the regions, comparison and analysis of results (SMETANA et al., 2016). Further, the data were compiled into the corresponding initial resource matrices, which were the basis for more complex IO matrices construction. The data for the assessment was collected from official statistical sources (LWK. 2015; NLWKN, 2015; RDG, 2015). The data sources contain publicly available information on sub-national regions (NUTS2 and NUTS3 levels), classified according to the Nomenclature of Units for Territorial Statistics. The agricultural association of Lower Saxony (LWK, 2015) contains updated information (including prices) on biotic resources. Lower Saxony data bank for water data of State Enterprise for Water Management, Coastal and Nature Conservation (NLWKN, 2015) is a reliable source of information for water supply and wastewater management. The biggest amount of data was mined from The Regional Database Germany (RDG, 2015).

There are a few benefits of using the data from the indicated sources. First of all, they represent the official sources of data collected directly from the regional statistics offices which accumulate the annual data on a constant basis. It assures the consistency and comparability of the data. Second, the datasets represent a wide variety of economic, environmental and social indicators, which can be used to identify the interactions between different components of the regional system. While economic data represent the interactions between production and consumption components, environmental data are presented with amounts of natural resources (biotic, mineral, water, energy, land resources and wastes generated and treated). The social component of sustainability in RSAM model is including information on the characterization of different components of the society (working population, direct consumers, service suppliers, social beneficiaries and regulators). This way the RSAM model tried to cover the whole population, which performs different functions from social benefits suppliers and consumers to the regulators, which act as transformers of the social goods distribution.

The analysis required initial resource IO matrices construction and their further transformation into single unit IO tables (based on the actual price index). The prices of the resources were acquired from multiple sources, which include regional statistical offices (Landesbetrieb für Statistik und Kommunikationstechnologie Niedersachsen), regional minister's offices (Niedersächsisches Ministerium für Wirtschaft, Arbeit und Verkehr) and other available sources. The integrated monetized matrix represents the estimation of separate resource development at given time frame (usually annual). Then the same calculation procedure was used to estimate the efficiency of resource use for the whole region. The detailed methodology of static RSAM application is presented in published papers (SMETANA et al., 2015; SMETANA et al., 2016).

Methodologically, except IO table analysis RSAM included flows cycling analyses (FINN, 1976;

ALLESINA and ULANO-WICZ, 2004), which assess resource transfer data within an outlined area. The efficiency of resource use in the regions was indicated as a relation of the resources amount cycled in the system to the amount of resources, leaving the system, according to the cycling estimations based on adapted Han (HCI), Finn (FCI) and Comprehensive cycling indices (CCI) (FINN, 1976; ALLE-SINA and ULANOWICZ, 2004). Cycling indices represent the complexity of the network in the system. The higher amount of alternative routes for resource transfer would result in higher cycling indices. The self-sufficiency index (sum of minors of included matrices) demonstrated the strength of a region to satisfy the needs for a specific resource with own resources. Further analysis included the interpretation of the results and repetition of the application procedure for the time series (dynamic assessment) and separate production impact assessment. The last two stages could

indicate the development of clusters based on specific industries. It was performed as a case study on two selected economically similar "rural districts", which formed a single agribusiness cluster, and one "ruralurban district" of level NUTS3 (Nomenclature of Units for Territorial Statistics) for comparison. Selected regions, compared in this case study, were situated in the west part of Northern Germany (Figure 1).

RSAM application for the regional characterization allows assessment of separate productions available in the region. It was performed with the use of multiplicators for IO tables. Once the RSAM is applied for the regional analysis, then the identification



Figure 1. The location of case study regions on the map of Germany

Source: authors

of specific production share for the regional sustainability is possible. This article estimates the relative importance of biotic resources (as an indicator of the agri-food based regional development) and their potential impact on the development of regions. Production properties as a driver of the economy were also separately evaluated for the efficiency of resource use. Dynamic trends were analyzed as a series of RSAM applied to the regions at different annual time frames. Trend analysis of time series was used to estimate possible paths for regional development. CCI) in Region (V) and Region (H). Region (C) was different due to the higher FCI and CCI in social aspects, whereas economic aspects had lower values. Environmental resources had the lowest cycling indices and self-sufficiency scores for all the regions. The analysis of separate aspects of sustainability and even separate production sectors was more informative with sectoral IO tables' analysis of RSAM presented further (Table 3).

## 4 **Empirical Results**

#### 4.1 Static Comparison

Static sustainability comparison of regional development of the three regions was performed for 2010 as the most detailed statistic data were well presented for the year from a single source for all the studied regions (it decreased the influence of data inconsistency). Regional comparison with RSAM showed that regions with similar economic and social sectors have similar indicator results (Table 1). Rural-urban region (H) is characterized by higher rates of resources cycling in the system and lower dependency on the external flows. The rate of self-sufficiency of Region (H) was higher than in the other two regions.

Presented analysis of the studied regions reflected the overall regional assessment and the differences between regions caused by the variations in organization and economic activities. However, it did not provide information on separate pillars of sustainability or drivers for agribusiness cluster development. That is why we performed an analysis of RSAM applicability for the assessment of social, environmental and economic aspects (Table 2). The results indicated higher rates of economic resources cycling (FCI and

Table 1.Integrated regional system comparison results (2010)

Regions	НСІ	FCI	CCI	I,O/Total	Self- sufficiency
(V)	0.15	0.93	1.06	0.39	23.56
(C)	0.14	0.99	1.13	0.4	17.72
<b>(H)</b>	0.27	1.14	1.31	0.35	28.93

Note: HCI – adapted Han cycling index; FCI – Finn cycling index; CCI – Comprehensive cycling index; I,O/Total – relation of inputs or outputs to total flows; (V) – Vechta Landkreis; (C) – Cloppenburg Landkreis; (H) – Hochsauerlandkreis Source: authors

Table 2.	Integrated regional system comparison results for
	separate aspects of sustainability (2010)

Regions	Aspects	НСІ	FCI	CCI	I,O/Total	Self- sufficiency
( <b>V</b> )	Soc	0.74	3.94	4.50	0.15	22.29
	Eco	8.73	7.29	8.32	0.05	19.45
	Env	0.09	1.20	1.37	0.38	7.05
( <b>C</b> )	Soc	1.33	7.21	8.23	0.07	47.52
	Eco	28.07	1.38	1.57	0.02	58.90
	Env	0.10	1.20	1.37	0.38	7.12
( <b>H</b> )	Soc	0.70	3.31	3.78	0.18	19.24
	Eco	7.17	6.21	7.10	0.06	16.32
	Env	0.10	1.20	1.38	0.38	7.43

Note: HCI – adapted Han cycling index; FCI – Finn cycling index; CCI – Comprehensive cycling index; I,O/Total – relation of inputs or outputs to total flows; (V) – Vechta Landkreis; (C) – Cloppenburg Landkreis; (H) – Hochsauerlandkreis; Soc – social; Eco – economic; Env – environmental Source: authors

 Table 3.
 Producer resources comparison of the regions in 2010

Regions	HCI	FCI	CCI	I/O	I/Total	O/Total	Self- sufficiency		
Absolute (monetized) values based comparison									
( <b>V</b> )	1.37	3.21	3.66	0.21	0.12	0.57	647		
( <b>C</b> )	1.18	3.67	4.19	0.24	0.14	0.59	279		
<b>(H</b> )	0.78	2.96	3.38	0.32	0.18	0.54	99		

Note: (V) – Vechta ; (C) – Cloppenburg ; (H) – Hochsauerlandkreis; HCI – adapted Han cycling index; FCI – Finn cycling index; CCI – Comprehensive cycling index; I/O – relation of inputs to outputs; I/Total – relation of inputs to total flows; O/Total – relation of outputs to total flows

Source: authors

#### 4.2 Dynamic Trends Comparison

The comparison of resource cycling rates was based on five-year period, which represented the ability of RSAM to reflect the short-term changes and indicate tendencies for further regional development. The identified trends demonstrated insignificant annual changes in regional development for the economically and socially similar regions (V and C) (Figure 2). They had similar stable results throughout the five year period with variations in the range of 0.2-9% for adapted Han Cycling Index (HCI) with the only exception of a 27.5% drop in 2012 for Region (V). Region (H) had overall higher indices of HCI (Figure 2). It also had a higher level of the index annual fluctuations (4-23%) which indicated higher regional dependence on annual changes of resource structure.

Trend analysis of adapted HCI results indicated that the amount of inner resources cycling in regions will be decreased in all three regions. Region (H) inner resources cycling will drop 4% by within the next two years comparing to the value of 2008, while the same measure will drop by 12.5% and 15% for Region (C) and Region (V) accordingly.

The analysis of the data with Finn Cycling Index (FCI) revealed similar results (Figure 3). Region (H) had a higher index of inner resources cycling (FCI = 1.2-1.5) and higher annual fluctuations (2.3-24.6%). The other regions had lower resource cy-

Figure 2. Results of adapted Han cycling index (HCI) applied to the regions



Note: V – Vechta; C – Cloppenburg; H – Hochsauerlandkreis; HCI – Han cycling index Source: authors

Figure 3. Results of adapted Finn cycling index (FCI) applied to the regions



Note: V – Vechta; C – Cloppenburg; H – Hochsauerlandkreis; FCI – Finn cycling index Source: authors

Figure 4. Results of external flows relation to the total amount of flows in the region



Note: V – Vechta; C – Cloppenburg; H – Hochsauerlandkreis; E/Total – relation of external flows to the total flow amount Source: authors

cling indices: ~0.9 for Region (V) and ~0.95 for Region (C). Annual changes of the cycling indices achieved 0.5-9.3% for Region (V) and 1.4-5.0% for Region (C).

Trend analysis of FCI application results revealed similar results of decreasing dependency on resources cycling in the regions. All three regions forecasted to have the FCI decreased within the range of 5.4-8%.

 Table 4.
 Biotic resources comparison of the study regions in 2010

Regions	HCI	FCI	CCI	I/O	I/Total	O/Total	Self-sufficiency	
Absolute values comparison								
( <b>V</b> )	4.47	3.52	4.02	0.37	0.17	0.46	719	
( <b>C</b> )	5.85	5.56	6.35	0.25	0.12	0.48	62741	
( <b>H</b> )	8.08	36.10	41.23	0.05	0.02	0.42	352280	
Monetized values comparison								
( <b>V</b> )	5.86	1.30	1.48	0.60	0.30	0.50	871	
( <b>C</b> )	6.48	2.02	2.31	0.42	0.24	0.57	20412	
<b>(H</b> )	20.05	10.95	12.50	0.16	0.07	0.46	366345	

Note: (V) – Vechta; (C) – Cloppenburg; (H) – Hochsauerlandkreis; HCI – Han cycling index; FCI – Finn cycling index; CCI – Comprehensive cycling index; I/O – relation of inputs to outputs; I/Total – relation of inputs to total flows; O/Total – relation of outputs to total flows Source: authors

The dependence on external flows was indicated as a relation of external flows to the total amount of resource flows in a region. The higher representation of external resources was highlighted for Region (V) and Region (C). Their dependency on external resources was around 40% (Figure 4). Region (H) had a lower overall dependency on external resources (0.25-0.35%). Annual fluctuation of results was 0.4-0.8% for Region (V) and 1.1-2% Region (C). The dependence on external resources in Region (H) fluctuated in a higher degree at 3.3-31.2%.

Trend analysis of dependence relations on external resources indicated minor possible changes in the future for the regions. Region (H) is unlikely to change the overall score as it was predicted to lower the dependency on 1.5% within the next two years comparing to the value in 2008. Region (V) should lower the dependency on 3.9%, while Region (C) should probably increase the use of external resources at 2%.

#### 4.1 Sustainability Assessment of Driving Forces of Regional Development

The sustainability analysis of separate aspects of regional development was performed with the use of the same methods of RSAM (see Methods). In order to reflect on the drivers of agribusiness cluster development this paper presents the analyses of overall producer interactions (Table 3) and biotic resources (Table 4). The lowest rates of producer resources cycling were outlined for Region (H). Producers in Region (V) and Region (C) had higher cycling indices of resources and therefore higher self-sufficiency. At the same time, the dependency of producers on external resource flows was similar between the three regions. High dependency on the external resource flows could result in negative changes in case of scarcity of external resources. Self-sufficiency of production activities was much higher in regions, which form the agribusiness cluster (V and C).

Similarly, separate aspects of regional development could be analyzed with RSAM for any important issue of sustainable development. In this article aggregated IO tables included assessment of lands, energy, water, biotic resources, minerals, producers, consumers, service suppliers, correctors (regulators), social beneficiaries and wastes. But it should be considered that the analysis on separate issues does not reflect their relative contribution towards the overall level of regional sustainability, but rather indicates the comparative value.

Another example, illustrating the application of RSAM for the drivers of agribusiness characterization is presented for biotic resource flows in 2010 (Table 4). On contrary to producer IO matrix it was possible to perform an analysis of absolute values and weighted monetized indicators for biotic resources.

IO table analysis for biotic resources (absolute values) identified the lowest levels of resource cycling for Region (V). Region (V) was also characterized with the highest levels of input flows. Monetized weighted IO matrix analysis indicated similar results for Region (V). Region (H) had the highest cycling rates of biotic resources in the region and the lowest dependency on external flows (both for absolute and monetized weighted values).

### **5** Discussion

Sustainability assessment at regional level has its specifics. We defined regional sustainability as the ability of a regional system (outlined with administrative, cultural, social, economic or other criteria) to sustain current socioeconomic and environmental conditions with possibilities for future development or current state preservation (SMETANA et al., 2015). Such regional definition as well refers to industrial cluster definitions (CRUZ and TEIXEIRA, 2010; BRENNER, 2004; BRENNER, 2006). Regional (sub-national) scale of analysis allows local data aggregation and integration, but keeps the possibilities for the detailed analysis of the key influencing factors of regional development. Identification of links between diverse elements, which influence the sustainability of regional development (LEIN, 2014; GRAYMORE et al., 2010; GRAYMORE et al., 2008) is a useful characteristic for the analysis of endogenous and exogenous drivers of cluster development. For example, the agribusiness cluster (the Oldenburger Münsterland) was dependent on external biotic resource flows. Together with higher than in the control region (Hochsauerlandkreis) inner cycling rates of economic production resources it indicated the active role of food production chains in the cluster region. Despite relatively high inner cycling rates of resources in the agribusiness cluster no considerable dynamic changes were observed, which pointed to the mature stage of cluster development. At the same time, the dependency of the region on the external resources is "preserving" deep shocking effects in the future. Therefore, a system vision of regional factors influencing the development of a region could identify the factors and solutions to the problems of sustainable development for regions.

We argue that the level of resources cycling in the system (region or cluster) and regional dependency on external resource flows indicated the sustainability of the system performance. In this case, the comparison of the flow rates with other systems allowed indication of problem areas, while a few-year comparison allowed evolutionary trend analysis in a dynamic perspective.

The application of RSAM for the static comparison of the regions revealed the possibilities of relative efficiency estimation of resources cycling in the system (region, cluster). RSAM application was indicative to distinguish the regions, which were characterized with different socioeconomic and environmental structure, which corresponds well to the previous study (SMETANA et al., 2016). At the same time, the differentiation between similar regions (V) and (C) was not indicative.

RSAM analysis of separate pillars of sustainability revealed differences in economic and social factors, whereas the efficiency of resource use in environmental sector was similarly low for all the regions. RSAM analysis of separate issues of sustainability allowed determination of differences and similarities between regions, thus, allowing for the selection of regional aspects aimed for further deeper analysis and hotspot identification. At the same time, at a given level of data aggregation and analysis, it was not possible to provide a substantial basis for sustainable hotspots allocation of the regions in 2010.

In order to exclude the potential impact of the annual changes on the resource use (and potential mistakes of sustainability levels indication) we performed a dynamic regional assessment for the fiveyear period. Further dynamics of a regional system were evaluated with trend analysis for the series of time periods. The analysis was reflective and confirmed the conclusions on the overall higher sustainability rates of Region (H), which was reflected in higher rates of resources cycling and lower dependency on external resources over the five-year period. It was also characterized with the prediction of decreased dependency on external resources.

Development and application of RSAM is foreseen to provide information about the current state of sustainability of regional development, to identify the "problem" fields and to indicate more sustainable alternatives for the development of a region. As compared regions had a high level of dependency on the use of biotic resources due to the cluster formation (Regions V and C are known as regions of intensive agricultural production), it was necessary to analyze the relative performance of biotic resources (environmental pillar) and production industries (economic pillar). The analysis indicated that despite relatively efficient use of economic resources in Region (H), the efficiency of resource use by producers was lower than in the other two regions, while dependency on external resources was higher and self-sufficiency was therefore, lower. Biotic resources were used with the highest efficiency in Region (H), which was explained with the lower economic dependency on the biotic resources (compared to the other regions). As Region (H) had a comparatively low share of external resource flows in the economy, the self-sufficiency measure was very high. The application of RSAM for the analysis of only biotic resources should not be performed separately from general regional assessment, as it would emphasize the value of biotic resources and neglect the role of other components of regional development. RSAM application for biotic resource efficiency assessment also revealed minor differentiation between Region (V) and Region (C), which indicated the potential sensitivity of RSAM application.

The use of RSAM, presented in this article, indicated its abilities for sustainability analysis of diverse regions and their dynamics. It is also applicable for the analysis of separate sustainability aspects of regional development with a high level of details according to the type of resources. RSAM provides the results in the same relative form, which can be compared and analyzed. Such a diverse apparatus for data analysis makes the RSAM a tool, which can provide a support for the identification of agribusiness cluster development in rural areas. Therefore, a further testing and application of RSAM is foreseen for diverse regions in various countries, potentially forming industrial clusters. It was indicated that RSAM is applicable for the analysis of specific sustainability components and their internal and external performance according to the integrated RSAM model. It can be used for the "hotspot analysis" of sustainability issues in relation to inter and intra-dynamics. The answer to such issues could be interesting for the actors involved in regional research and planning (e.g. government and scholars). Moreover, the interest in the research from community demonstrated the relevance of the research application for public acknowledgement with regional sustainability problems.

# 6 Conclusions

The study presented Regional Sustainability Assessment Methodology (RSAM) application to the analysis of static regional sustainability conditions and dynamic trends of regional development. A case study approach of RSAM use for regions of agribusiness cluster analysis in comparison to a control region was performed.

Similarities of socioeconomic development of regions were reflected in comparable indices of RSAM. Thus, Vechta and Cloppenburg regions (V and C) had comparable endogenous recourse cycling rates, dependencies on external resources and self-sufficiency rates. The control region – Hochsauerlandkreis (H) had different results. Such results could potentially indicate the applicability of RSAM for the identification of industrial clusters through the comparison of similarities between neighboring regions. However, in order to claim the significance of such conclusions, further testing of the methodology with more diverse industrial clusters and regions should be performed.

Agribusiness cluster formation in rural areas requires the increased magnitude of external resource flows as the development of an industrial center depends on the external demand and supply. Similar to the other studies (WILK and FENSTERSEIFER, 2003; FENSTERSEIFER, 2007) it was highlighted in the case of RSAM application for Regions (V) and (C). On the other hand, regions which depend on inner resources for the development (H) do not form industrial clusters. The conclusion should be further investigated with a wider range of industrial clusters in order to confirm the applicability of RSAM for cluster identification.

The characterization of agribusiness clusters was performed with RSAM for the resources reflecting producer and biotic sectors of the economy. Producer activities analysis with RSAM revealed almost a double rate of economic resources cycling in agribusiness cluster, comparing to the control Region (H). The high level of resources cycling in cluster regions triggered also higher rates of self-sufficiency. At the same time high dependency on the external resource flows holds a risk of resources use efficiency leveling to the rates of the control region, which might add another point to the conclusions of other authors about the selfdestructing forces in clusters (BROEKEL et al., 2015; MENZEL and FORNAHL, 2010).

Application of RSAM for biotic resource analysis reflected regions of the agribusiness cluster as dependent on the external resource flows and having low cycling rates of inner resources compared to the control region (H). RSAM analysis also revealed the relative position of the regions in terms of biotic resources use efficiency and self-sufficiency. Despite producer and biotic resource characterization, the results of separate sustainability pillars with RSAM analysis did not indicate the specific drivers of agribusiness cluster formation at the regional scale. Neither were they illustrative for the hotspot characterization.

Dynamic analysis based on the results of RSAM application for the five-year period, indicated the stable nature of the results to characterize the differences of sustainability condition of regional development. Forecasting trend analysis allowed indication of the potential changes of regional sustainability. The overall resource efficiency use (as an indicator of sustainability) was reflective on the annual changes for all the indices, but demonstrated the overall stable position between different socioeconomic regions.

# References

- ABEL, J.R., I. DEY and T.M. GABE (2010): Productivity and the density of human capital. Staff Report. Federal Reserve Bank of New York, No. 440.
- ADGER, W.N. (2003): Social Capital, Collective Action, and Adaptation to Climate Change. In: Economic Geography 79 (4): 387-404.
- AGOSTINHO, F. and E. ORTEGA (2012): Integrated food, energy and environmental services production as an alternative for small rural properties in Brazil. In: Energy 37 (1): 103-114.
- ALECKE, B. and G. UNTIEDT (2008): Die räumliche Konzentration von Industrie und Dienstleistungen in Deutschland. In: Review of Regional Research 28 (1): 61-92.
- ALLAN, G., P. MCGREGOR and K. SWALES (2010): The Importance of Revenue Sharing for the Local Economic Impacts of a Renewable Energy Project: A Social Accounting Matrix Approach. In: Regional Studies 45 (9): 1171-1186.
- ALLESINA, S. and R.E. ULANOWICZ (2004): Cycling in ecological networks: Finn's index revisited. In: Computational biology and chemistry 28 (3): 227-33.
- BEN LETAIFA, S. and Y. RABEAU (2013): Too close to collaborate? How geographic proximity could impede entrepreneurship and innovation. In: Journal of Business Research 66 (10): 2071-2078. Available at: http://linking hub.elsevier.com/retrieve/pii/S014829631300057X.
- BIRCH, K., L. LEVIDOW and T. PAPAIOANNOU (2010): Sustainable Capital? The Neoliberalization of Nature and Knowledge in the European "Knowledge-based Bioeconomy." In: Sustainability 2 (9): 2898-2918.
- BÖHRINGER, C. and P.E.P. JOCHEM (2007): Measuring the immeasurable - A survey of sustainability indices. In: Ecological Economics 63 (1): 1-8. Available at: http://www.sciencedirect.com/science/article/pii/S09218 00907002029, accessed July 13, 2014.
- BOSKER, M. (2009): The spatial evolution of regional GDP disparities in the 'old' and the 'new' Europe. In: Papers in Regional Science 88 (1): 3-27.
- BRACHERT, M., M. TITZE and A. KUBIS (2011): Identifying industrial clusters from a multidimensional perspective: Methodical aspects with an application to Germany. In: Papers in Regional Science 90 (2): 419-439. Available at: http://dx.doi.org/10.1111/j.1435-5957.2011.00356.x.
- BRENNER, T. (2004): Local industrial clusters: existence, emergence and evolution. Routledge, London.
- (2006): Identification of Local Industrial Clusters in Germany. In: Regional Studies 40 (9): 991-1004. Available at: http://dx.doi.org/10.1080/00343400601047408.
- BRENNER, T. and A. GILDNER (2006): The long-term Implications of Local Industrial Clusters. In: European Planning Studies 14 (9): 1315-1328. Available at: http://www.tandfonline.com/doi/abs/10.1080/09654310 600933371.
- BROEKEL, T., T. BRENNER and M. BUERGER (2015): An Investigation of the Relation between Cooperation Intensity and the Innovative Success of German Regions. In: Spatial Economic Analysis 10 (1): 52-78. Available at: http://www.tandfonline.com/doi/abs/10.10 80/17421772.2014.992359.

- CABELLO, J.M., E. NAVARRO, F. PRIETO, B. RODRÍGUEZ and F. RUIZ (2014): Multicriteria development of synthetic indicators of the environmental profile of the Spanish regions. In: Ecological Indicators 39: 10-23.
- CAMPBELL, D.E. and A.S. GARMESTANI (2012): An energy systems view of sustainability: emergy evaluation of the San Luis Basin, Colorado. In: Journal of environmental management 95 (1): 72-97.
- COOKE, P., N. CLIFTON and M. OLEAGA (2005): Social capital, firm embeddedness and regional development. In: Regional Studies 39 (8): 1065-1077.
- CRUZ, S.C.S. and A.A.C. TEIXEIRA (2010): The Evolution of the Cluster Literature: Shedding Light on the Regional Studies-Regional Science Debate. In: Regional Studies 44 (9): 1263-1288. Available at: http://www.tand fonline.com/doi/abs/10.1080/00343400903234670, accessed May 28, 2014.
- DANIELS, P.L., M. LENZEN and S.J. KENWAY (2011): The ins and outs of water use-a review of multi-region inputoutput analysis and water footprints for regional sustainability analysis and policy. In: Economic Systems Research 23 (4): 353-370. Available at: http://dx.doi.org/ 10.1080/09535314.2011.633500.
- DELGADO, M., M.E. PORTER and S. STERN (2016): Defining clusters of related industries. In: Journal of Economic Geography 16 (1): 1-38. Available at: http://joeg.oxford journals.org/lookup/doi/10.1093/jeg/lbv017.
- DEVINE-WRIGHT, P., P.D. FLEMING and H. CHADWICK (2001): Role of social capital in advancing regional sustainable development. In: Impact Assessment and Project Appraisal 19 (2): 161-167.
- DIEZ-VIAL, I. (2011): Geographical cluster and performance: The case of Iberian ham. In: Food Policy 36 (4): 517-525. Available at: http://www.sciencedirect.com/science/ article/pii/S0306919211000613, accessed February 10, 2016.
- FENSTERSEIFER, J.E. (2007): The emerging Brazilian wine industry: Challenges and prospects for the Serra Gaúcha wine cluster. In: International Journal of Wine Business Research 19 (3): 187-206. Available at: http://dx.doi.org/ 10.1108/17511060710817221.
- FINN, J. (1976): Measures of ecosystem structure and function derived from analysis of flows. In: Journal of theoretical Biology (23): 363-380.
- FLORIDA, R., C. MELLANDER and K. STOLARICK (2008): Inside the black box of regional development - human capital, the creative class and tolerance. In: Journal of Economic Geography 8 (5): 615-649.
- GATTO, M. (1995): Sustainability: Is it a Well Defined Concept? In: Ecological Applications 5 (4): 1181-1183. Available at: http://www.jstor.org/stable/2269365.
- GELLYNCK, X., B. KÜHNE and R.D. WEAVER (2011): Relationship quality and innovation capacity of chains: the case of the traditional food sector in the EU. In: International Journal on Food System Dynamics 2 (1): 1-22.
- GRAYMORE, M.L.M., N.G. SIPE and R.E. RICKSON (2008): Regional sustainability: How useful are current tools of sustainability assessment at the regional scale? In: Ecological Economics 67 (3): 362-372.
- (2010): Sustaining Human Carrying Capacity: A tool for regional sustainability assessment. In: Ecological Economics 69 (3): 459-468.

- GUINÉE, J.B., R. HEIJUNGS, G. HUPPES, A. ZAMAGNI, P. MASONI, R. BUONAMICI, T. EKVALL and T. RYDBERG (2011): Life cycle assessment: past, present, and future. In: Environmental science & technology 45 (1): 90-6.
- HENDRICKSON, C., A. HORVATH, S. JOSHI and L. LAVE (1998): Economic Input-Output Models for Environmental Life-Cycle Assessment. In: Environmental science & technology 32 (7): 184A-191A.
- HERTWICH, E.G. (2005): Life cycle approaches to sustainable consumption: a critical review. In: Environmental science & technology 39 (13): 4673-4684.
- HOEKSTRA, A.Y., M.M. MEKONNEN, A.K. CHAPAGAIN, R.E. MATHEWS and B.D. RICHTER (2012): Global monthly water scarcity: blue water footprints versus blue water availability. In: PloS one 7 (2): e32688. Available at: http://www.pubmedcentral.nih.gov/article render.fcgi?artid=3290560&tool=pmcentrez&rendertyp e=abstract, accessed October 31, 2013.
- HOFFMANN, J., S. HIRSCH and J. SIMONS (2015): Identification of spatial agglomerations in the German food processing industry. In: Papers in Regional Science 96:139-162. Available at: http://dx.doi.org/10.1111/ pirs.12171.
- JORDAN, S.J., S.E. HAYES, D. YOSKOWITZ, L.M. SMITH, J.K. SUMMERS, M. RUSSELL and W.H. BENSON (2010): Accounting for natural resources and environmental sustainability: linking ecosystem services to human well-being. In: Environmental science & technology 44 (5): 1530-1536.
- KAHL, J. and C. HUNDT (2015): Employment performance in times of crisis: A multilevel analysis of economic resilience in the German biotechnology industry. In: Competitiveness Review 25 (4): 371-391. Available at: http://dx.doi.org/10.1108/CR-12-2014-0038.
- KOURTIT, K., C. MACHARIS and P. NIJKAMP (2014): A multi-actor multi-criteria analysis of the performance of global cities. In: Applied Geography 49: 24-36. Available at: http://www.sciencedirect.com/science/article/pii/S01 43622813002117, accessed January 18, 2015.
- KURKA, T. (2013): Application of the analytic hierarchy process to evaluate the regional sustainability of bioenergy developments. In: Energy 62: 393-402. Available at: http://linkinghub.elsevier.com/retrieve/pii/S0360544 213008190, accessed October 23, 2013.
- LEHMANN, A., E. ZSCHIESCHANG, M. TRAVERSO, M. FINKBEINER and L. SCHEBEK (2013): Social aspects for sustainability assessment of technologies – challenges for social life cycle assessment (SLCA). In: The International Journal of Life Cycle Assessment 18: 1581-1592. Available at: http://link.springer.com/10.1007/ s11367-013-0594-0, accessed August 9, 2013.
- LEIN, J. (2014): Toward a Remote Sensing Solution for Regional Sustainability Assessment and Monitoring. In: Sustainability 6 (4): 2067-2086.
- LEONTIEF, W. (1951): Input-Output Economics. In: Scientific American 185 (4): 15-21.
- LWK (2015): Die Landwirtschaftskammer Niedersachsen. Available at: http://www.lwk-niedersachsen.de/index. cfm/portal/landwirtschaftskammer.html.
- MARTIN, R. and P. SUNLEY (2003): Deconstructing clusters: chaotic concept or policy panacea? In: Journal of Economic Geography 3 (1): 5-35. Available at: http://joeg.ox fordjournals.org/content/3/1/5.abstract.

- MARTIN, R. and P. SUNLEY (2007): Complexity thinking and evolutionary economic geography. In: Journal of Economic Geography 7 (5): 573-601.
- MATTHEWS, H. and M. SMALL (2000): Extending the boundaries of life-cycle assessment through environmental economic input-output models. In: Journal of Industrial Ecology 4 (3): 7-10. Available at: http://gdi.ce. cmu.edu/docs/extending-the.pdf, accessed March 24, 2014.
- MAYER, A.L. (2008): Strengths and weaknesses of common sustainability indices for multidimensional systems. In: Environment international 34 (2): 277-91.
- MENZEL, M.-P. and D. FORNAHL (2007): Cluster Life Cycles - Dimensions and Rationales of Cluster Development. In: SSRN Electronic Journal (2007-076). Available at: http://www.ssrn.com/abstract=1025970.
- (2010): Cluster life cycles dimensions and rationales of cluster evolution. In: Industrial and Corporate Change 19 (1): 205-238. Available at: http://icc.oxfordjournals.org/ cgi/doi/10.1093/icc/dtp036.
- MILLER, R.E. and P.D. BLAIR (2009): Input-output analysis: foundations and extensions. Cambridge University Press, Cambridge.
- MORILLA, C., G. DIAZ-SALAZAR and M. CARDENETE (2007): Economic and environmental efficiency using a social accounting matrix. In: Ecological economics 60 (4): 774-786. Available at: http://www.sciencedirect.com/ science/article/pii/S0921800906000735, accessed January 5, 2015.
- MUNDA, G. and M. SAISANA (2011): Methodological Considerations on Regional Sustainability Assessment Based on Multicriteria and Sensitivity Analysis. In: Regional Studies 45 (2): 261-276.
- NLWKN (2015): Niedersächsischen Landesbetrieb für Wasserwirtschaft, Küsten- und Naturschutz (NLWKN). Available at: http://www.wasserdaten.niedersachsen.de/.
- O'RIORDAN, T. and H. VOISEY (1998): The transition to sustainability: the politics of Agenda 21 in Europe. Earthscan, London.
- PACHAURI, R.K., M.R. ALLEN, V.R. BARROS, J. BROOME, W. CRAMER, R. CHRIST, J.A. CHURCH, L. CLARKE, Q. DAHE and P. DASGUPTA (2014): Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. IPCC, Geneva, Switzerland.
- PFISTER, S. and P. BAYER (2014): Monthly water stress: Spatially and temporally explicit consumptive water footprint of global crop production. In: Journal of Cleaner Production 73: 52-62.
- PHILLIPS, J. (2011): The conceptual development of a geocybernetic relationship between sustainable development and Environmental Impact Assessment. In: Applied Geography 31 (3): 969-979.
- (2015): A quantitative approach to determine and evaluate the indicated level and nature of global ecological sustainability. In: Area 47 (2): 197-210. Available at: http://dx.doi.org/10.1111/area.12174.
- RDG (2015): The Regional Database Germany ("Regionaldatenbank Deutschland"). Available at: www.regional statistik.de.
- ROBÈRT, K., G. BROMAN and G. BASILE (2013): Analyzing the concept of planetary boundaries from a strategic

sustainability perspective: How does humanity avoid tipping the planet? In: Ecology and Society 18 (2): 5.

- ROCHA, H.O. and R. STERNBERG (2005): Entrepreneurship: The role of clusters theoretical perspectives and empirical evidence from Germany. In: Small Business Economics 24 (3): 267-292.
- RODRÍGUEZ, X.A. and C. ARIAS (2008): The effects of resource depletion on coal mining productivity. In: Energy Economics 30 (2): 397-408.
- SCHMIDT, J.H. and S. MERCIAI (2014): Life cycle assessment of the global food consumption. In 9<sup>th</sup> International Conference LCA of Food, San Francisco, USA.
- SHAKER, R.R. (2015): The spatial distribution of development in Europe and its underlying sustainability correlations. In: Applied Geography 63: 304-314. Available at: http://www.sciencedirect.com/science/article/pii/S01436 22815001745, accessed August 3, 2015.
- SICHE, J.R., F. AGOSTINHO, E. ORTEGA and A. ROMEIRO (2008): Sustainability of nations by indices: Comparative study between environmental sustainability index, ecological footprint and the emergy performance indices. In: Ecological Economics 66 (4): 628-637. Available at: http://www.sciencedirect.com/science/article/pii/S09218 0090700523X, accessed September 30, 2015.
- SINGH, R.K., H.R. MURTY, S.K. GUPTA and A.K. DIKSHIT (2012): An overview of sustainability assessment methodologies. In: Ecological Indicators 15 (1): 281-299.
- SMETANA, S., C. TAMÁSY, A. MATHYS and V. HEINZ (2015): Sustainability and regions: sustainability assessment in regional perspective. In: Regional Science Policy & Practice 7 (4): 163-186. Available at: http://doi.wiley.com/10.1111/rsp3.12068.
- (2016): Measuring Relative Sustainability of Regions Using Regional Sustainability Assessment Methodology. In: Geographical Analysis 48 (4): 391-410.
- STEINER, B., M. FRITZ, U. RICKERT and G. SCHIEFER (2007): Clustering and innovation in Canada: evidence from the Alberta food processing sector. In: Innovation and system dynamics in food networks 2007. Proceedings of the 1<sup>st</sup> European Forum on Innovation and System Dynamics in Food Networks, Innsbruck-Igis, Austria, 15-17 February, 2007. Universität Bonn-ILB: 111-128.
- STEINFELD, H., P. GERBER, T. WASSENAAR, V. CASTEL, M. ROSALES and C. DE HAAN (2006): Livestock's long shadow. Environmental issues and options. Rome: Food and Agriculture Organization of the United Nations (FAO). Available at: ftp://ftp.fao.org/docrep/fao/010/ a0701e/a0701e.pdf.
- STERNBERG, R. and T. LITZENBERGER (2004): Regional clusters in Germany - their geography and their relevance for entrepreneurial activities. In: European Planning Studies 12 (6): 767-791. Available at: http://dx.doi.org/ 10.1080/0965431042000251855.
- TAMÁSY, C. (2013): Areas of intensive livestock agriculture as emerging alternative economic spaces? In: Applied Geography 45: 385-391.

- TENERELLI, P. and S. CARVER (2012): Multi-criteria, multiobjective and uncertainty analysis for agro-energy spatial modelling. In: Applied Geography 32 (2): 724-736. Available at: http://www.sciencedirect.com/science/ article/pii/S0143622811001664, accessed May 21, 2015.
- TITZE, M., M. BRACHERT and A. KUBIS (2011): The Identification of Regional Industrial Clusters Using Qualitative Input-Output Analysis (QIOA). In: Regional Studies 45 (1): 89-102.
- VAN PASSEL, S. and M. MEUL (2012): Multilevel and multiuser sustainability assessment of farming systems. In: Environmental Impact Assessment Review 32 (1): 170-180.
- VAN WITTELOOSTUIJN, A. (2009): Globalisation in the food industry: the impact on market structures and firm postures. In: Bunte, F. and H. Dagevos (eds.): The Food Economy. Global issues and challenges. Wageningen Academic Publishers, The Netherlands: 23-42.
- VOINOV, A. (2008): Understanding and communicating sustainability: global versus regional perspectives. In: Environment, Development and Sustainability 10 (4): 487-501. Available at: http://link.springer.com/10.1007/ s10668-006-9076-x.
- WANDEL, J. (2010): The cluster-based development strategy in Kazakhstan's agro-food sector: A critical assessment from an" Austrian" perspective. IAMO Discussion Paper 128. IAMO, Halle (Saale).
- WECKMAN, G., S. LAKSHMINARAYANAN, J.H. MARVEL and A. SNOW (2008): An integrated stock market forecasting model using neural networks. In: International Journal of Business Forecasting and Marketing Intelligence 1 (1): 30-49.
- WILK, E. DE O. and J.E. FENSTERSEIFER (2003): Use of resource based view in industrial cluster strategic analysis. In: International Journal of Operations & Production Management 23 (9): 995-1009. Available at: http://dx.doi.org/10.1108/01443570310491747.
- WROBEL, M. (2015): 'One for all and all for one': Cluster, employment, and the global economic crisis. Evidence from the German mechanical engineering industry. In: Papers in Regional Science 94 (2): 273-294. Available at: http://dx.doi.org/10.1111/pirs.12065.
- ZAMAGNI, A. (2012): Life cycle sustainability assessment. In: The International Journal of Life Cycle Assessment 17 (4): 373-376.

Contact author:

#### **DR. SERGIY SMETANA**

German Institute of Food Technologies (DIL e.V.) Prof.-von-Klitzing-Str. 7, 49610 Quakenbrück, Germany e-mail: s.smetana@dil-ev.de