

Beran, Jan; Droullier, Frieder

Article — Published Version

On strongly dependent zero-inflated INAR(1) processes

Statistical Papers

Provided in Cooperation with:

Springer Nature

Suggested Citation: Beran, Jan; Droullier, Frieder (2023) : On strongly dependent zero-inflated INAR(1) processes, Statistical Papers, ISSN 1613-9798, Springer, Berlin, Heidelberg, Vol. 65, Iss. 4, pp. 2527-2553,
<https://doi.org/10.1007/s00362-023-01496-z>

This Version is available at:

<https://hdl.handle.net/10419/307038>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.



<https://creativecommons.org/licenses/by/4.0/>



On strongly dependent zero-inflated INAR(1) processes

Jan Beran¹ · Frieder Droullier¹

Received: 8 November 2022 / Revised: 15 May 2023 / Published online: 29 September 2023

© The Author(s) 2023

Abstract

We consider INAR(1) processes modulated by an unobserved strongly dependent $0 - 1$ process. The observed process exhibits zero inflation and long memory. A simple method is proposed for estimating the INAR-parameters without modelling the unobserved modulating process. Asymptotic results for the estimators are derived, and a zero-inflation test is introduced. Asymptotic rejection regions and asymptotic power under long-memory alternatives are derived. A small simulation study illustrates the asymptotic results.

Keywords INAR process · Zero-inflation · Long-range dependence · Modulation

1 Introduction

Time series that consist of counts often exhibit zero inflation (see e.g. Young et al. 2022 and references therein). For instance, instrumental detection limits can lead to an increased number of zeroes. Also ecological data sets tend to contain a large proportion of zeroes that are classified by ecologists as "false zero counts" (Martin et al. 2005). Examples can also be found in other areas such as medicine, psychology, traffic or insurance, among others (Young et al. 2022). In this paper we consider zero inflation generated by a long-memory process. This is in contrast to the statistical literature where zero inflation processes are usually assumed to be iid or at most weakly dependent. The issue of zeroes occurring in long batches has been discussed in the recent applied literature. In many applications, zero inflation is related to missing values due to faulty measurement devices, instrumental detection limits or biological reasons. Also, the occurrence of batches of zeroes may be caused by other unobserved time series (see e.g. Che et al. 2018). A natural approach to handling such data is to use models that exhibit long-range dependence with respect to zero inflation. Note that, in a related paper, Möller et al. (2018) discuss models that tend to produce relatively long

✉ Jan Beran
jan.beran@uni-konstanz.de

¹ Department of Mathematics and Statistics, University of Konstanz, Konstanz, Germany

runs of zeroes (e.g. ZT-Bar(1) processes). However, asymptotically, the autocorrelation function of their models decays exponentially and is thus summable. In contrast, the models considered here have long memory in the sense that autocorrelations decay hyperbolically and are not summable. Note also that a long-memory INAR model, a so-called fractional INARFIMA, is suggested in Quoreshi (2014). However, the INARFIMA model, as specified by Quoreshi, is not well defined, because it relies on a fractional thinning operator that yields infinity when applied to non-negative sequences.

More specifically, we consider INAR(1) processes modulated in a multiplicative way by an unobserved strongly dependent 0 – 1 process. Thus, the observed process is assumed to be of the form

$$Y_j = W(j) X_j \quad (j = 1, 2, \dots) \tag{1}$$

where the stationary 0–1-process $W(j)$ ($j \in \mathbb{N}$) is independent of the INAR(1) process X_j ($j \in \mathbb{N}$), and the autocovariances $\gamma_W(k) = cov(W(j), W(j+k))$ are such that

$$\gamma_W(k) \underset{k \rightarrow \infty}{\sim} c_{\gamma,W} k^{2d-1} \tag{2}$$

for some constants $0 < c_{\gamma,W} < \infty$, $d \in (0, \frac{1}{2})$. Note that Eq. (2) implies that the spectral density

$$f_W(\lambda) = \frac{1}{2\pi} \sum_{k=-\infty}^{\infty} \gamma_W(|k|) e^{-ik\lambda} \tag{3}$$

has a pole at the origin, of the form

$$f_W(\lambda) \underset{\lambda \rightarrow 0}{\sim} c_{f,W} |\lambda|^{-2d} \tag{4}$$

where $c_{f,W} = c_{\gamma,W} \pi^{-1} \Gamma(2d) \sin(\pi/2 - \pi d)$ (Zygmund 1968; Beran et al. 2013, Theorem 1.3). We will use the notation

$$p_{0,W} = P(W(j) = 0) = 1 - P(W(j) = 1) \tag{5}$$

More specifically, we will assume X_j to be a Poisson INAR(1) process (McKenzie 1985; Al-Osh and Alzaid 1987) defined by

$$X_j = \alpha \circ X_{j-1} + \varepsilon_j$$

where $\alpha \in (0, 1)$, and ε_j are iid Poisson variables with intensity λ . The thinning operator "o" (Steutel and Van Harn 1979; Gauthier and Latour 1994) means that, conditionally on X_{j-1} , $\alpha \circ X_{j-1}$ is a binomial random variable generated by X_{j-1} Bernoulli trials with success probability α . The unknown parameters are α and λ . Alternatively, one may also use the parameterization α and $\mu_X = E(X_j) = \lambda/(1-\alpha)$.

Two questions regarding model Eq. (1) are addressed in this paper: a) estimation of the INAR(1) parameters α and μ_X without modelling the unobservable process W ; b) testing for zero-inflation as specified by model Eq. (1).

Since the introduction of INAR models by McKenzie (1985) and Al-Osh and Alzaid (1987), the literature on integer-valued time series based on binomial thinning (Steutel and Van Harn 1979) has grown steadily. Some references are Du and Li (1991); Gauthier and Latour (1994); Da Silva and Oliveira (2004); Freeland and McCabe (2004); Freeland and McCabe (2005); Gourieroux and Jasiak (2004); Jung et al. (2005); Puig and Valero (2007); Drost et al. (2008); Park and Kim (2012); Pedeli and Karlis (2013); Schweer and Weiss (2014); Pedeli et al. (2015) and Jentsch and Weiss (2019). Excellent overviews on integer-valued processes and references can be found for instance in Weiss (2018) and Davis et al. (2021). Testing for zero-inflation in INAR(1) models is discussed in a recent paper by Weiss et al. (2019). Also see Pavlopoulos and Karlis (2008); Barreto-Souza (2015); Weiss (2013), and Bourguignon and Weiss (2017) for over- and underdispersed INAR processes. References to long-memory processes can be found for instance in Beran (1994); Giraitis et al. (2012); Beran et al. (2013) and Pipiras and Taqqu (2017).

The paper is organized as follows. Basic results are established in Sect. 2. Question a) is addressed in Sect. 3. In Sect. 4, a test is developed for the null hypothesis of a non-inflated INAR(1) process against the alternative of a zero-inflated process as defined in Eq. (1). Asymptotic rejection regions are derived, as well as an asymptotic lower and upper bound for the power of the test. A small simulation study in Sect. 5 illustrates the results. Final remarks in Sect. 6 conclude the paper. Proofs are given in the appendix.

2 Basic results

2.1 Expected value and autocovariance function

Let Y_j be generated by model Eq. (1) where X_j is a Poisson INAR(1) process. We will use the notation $\mathbf{1}\{A\}$ for the indicator function of a set (or event) A . Moreover, we define

$$\begin{aligned} p_{0,W} &= P(W(j) = 0), p_{0,Y} = P(Y_j = 0), p_{0,X} = P(X_j = 0), \\ \mu_X &= E(X_j), \mu_W = E[W(j)], \mu_Y = E(Y_j), \\ \gamma_X(k) &= cov(X_j, X_{j+k}), \gamma_W(k) \\ &= cov(W(j), W(j+k)), \gamma_Y(k) = cov(Y_j, Y_{j+k}) \quad (j, k \geq 0). \end{aligned}$$

Also, for $k < 0$, we set $\gamma_X(k) = \gamma_X(-k)$, $\gamma_W(k) = \gamma_W(-k)$, $\gamma_Y(k) = \gamma_Y(-k)$, and, for $\lambda \in [-\pi, \pi]$,

$$f_X(\lambda) = \frac{1}{2\pi} \sum_{k=-\infty}^{\infty} \gamma_X(k) e^{-ik\lambda}$$

$$\begin{aligned}
 f_W(\lambda) &= \frac{1}{2\pi} \sum_{k=-\infty}^{\infty} \gamma_W(k) e^{-ik\lambda} \\
 f_{WX}(\lambda) &= \frac{1}{2\pi} \sum_{k=-\infty}^{\infty} \gamma_W(k) \gamma_X(k) e^{-ik\lambda} \\
 f_Y(\lambda) &= \frac{1}{2\pi} \sum_{k=-\infty}^{\infty} \gamma_Y(k) e^{-ik\lambda}
 \end{aligned}$$

Due to mutual independence and stationarity of the two processes X_j and $W(j)$ we have

$$\mu_Y = \mu_W \mu_X$$

and

$$\begin{aligned}
 \gamma_Y(k) &= E[W(0)W(k)]E[X_0X_k] - \mu_W^2\mu_X^2 \\
 &= \mu_X^2\gamma_W(k) + \mu_W^2\gamma_X(k) + \gamma_W(k)\gamma_X(k)
 \end{aligned} \tag{6}$$

Noting that

$$\gamma_X(k) = \alpha^k \mu_X(k \geq 0)$$

where $\mu_X = \lambda/(1 - \alpha)$, assumption Eq. (2) implies

$$\gamma_Y(k) \underset{k \rightarrow \infty}{\sim} \mu_X^2 c_{\gamma,W} k^{2d-1} \tag{7}$$

and

$$f_Y(\lambda) = \mu_X^2 f_W(\lambda) + \mu_W^2 f_X(\lambda) + f_{WX}(\lambda) \underset{\lambda \rightarrow 0}{\sim} \mu_X^2 c_{f,W} |\lambda|^{-2d} \tag{8}$$

Thus, although the original process X_j has exponentially decaying autocovariances, the observed process Y_j inherits long memory from the modulating process $W(j)$.

Example 1 Let Z_j ($j \in \mathbb{Z}$) be a stationary Gaussian process with zero mean, variance one and spectral density function f_Z such that

$$f_Z(\lambda) \underset{\lambda \rightarrow 0}{\sim} c_{f,Z} |\lambda|^{-2d}$$

for some $0 < c_{f,Z} < \infty$, $d \in (0, \frac{1}{2})$. Also, let $\kappa \neq 0$, and denote by Φ and ϕ the standard normal distribution and density function respectively. Then the process

$$W(j) = \mathbf{1}\{Z_j \leq \kappa\} \quad (j \in \mathbb{Z})$$

is stationary with expected value $\mu_W = \Phi(\kappa)$ and spectral density

$$f_W(\lambda) \underset{\lambda \rightarrow 0}{\sim} c_{f,W} |\lambda|^{-2d}$$

where $c_{f,W} = c_{f,Z\varphi(\kappa)}$.

Example 2 Another example of a modulating process with long memory can be obtained from a renewal reward process defined by

$$\tilde{W}(t) = \xi_0 \mathbf{1}\{0 \leq t < \tau_0\} + \sum_{j=-1}^{\infty} \xi_j \mathbf{1}\{\tau_{j-1} \leq t < \tau_{j-1} + T_j\} \quad (t > 0) \quad (9)$$

where τ_j ($j = 1, 2, \dots$) is a renewal process with intensity ν , and $\xi_j \in \{0, 1\}$ are iid random variables with $p_\xi = P(\xi_j = 1) \in (0, 1)$, independent of the process τ_j . Moreover, the interarrival times $T_j = \tau_j - \tau_{j-1}$ are assumed to have a finite expected value $E(T) = \mu$ and a marginal distribution function F_T such that

$$1 - F_T(u) \sim c_{F,T} u^{-\alpha} \quad (u \rightarrow \infty)$$

for some constants $0 < c_{F,T} < \infty$, $1 < \alpha < 2$. Finally, to achieve stationarity, the distribution function of τ_0 is assumed to be given by $F_{\tau_0}(x) = \mu^{-1} \int_0^x (1 - F_T(u)) du$. Then \tilde{W} is a continuous time stationary process with expected value $\mu_W = p_\xi$ and autocovariance function

$$\gamma_{\tilde{W}}(u) = cov(\tilde{W}(0), \tilde{W}(u)) = p_\xi (1 - p_\xi) P(\tau_0 > u) \underset{u \rightarrow \infty}{\sim} c_{\gamma,W} u^{2d-1}$$

where

$$d = 1 - \frac{\alpha}{2} \in \left(\frac{1}{2}, 1\right)$$

Setting $W(j) = \tilde{W}(j)$ ($j = 1, 2, \dots$), we obtain a discrete time stationary zero–one process with long memory as defined by Eq. (4) (see e.g. Beran et al. 2013, and references therein).

2.2 Sample mean and sample autocovariances

Standard methods for estimating the parameters of a Poisson INAR(1) process include moment and conditional least squares estimation. For both methods, estimators are functions of the sample mean and the lag-one sample autocorrelation. It is therefore of interest to study the effect of modulation on these statistics.

Consider first the sample mean $\bar{y}_n = n^{-1} \sum_{j=1}^n Y_j$ as an estimator of μ_Y . With respect to the expected value we have $E(\bar{y}_n) = \mu_W \mu_X$. Thus, \bar{y}_n is biased, unless $P(W(t) = 1) = 1$. The asymptotic distribution of \bar{y}_n follows from the asymptotic

distribution of $\bar{w}_n = n^{-1} \sum_{j=1}^n W(j)$, as stated in the next theorem. We will use the notation

$$v(d) = \frac{2 \sin \pi d}{d(2d + 1)} \Gamma(1 - 2d)$$

Also, " \rightarrow_d " and " \rightarrow_p " will denote convergence in distribution and in probability respectively.

Theorem 1 *Let the process W be such that*

$$Z_{n,W} = \frac{\bar{w}_n - \mu_W}{\sqrt{\text{var}(\bar{w}_n)}} \xrightarrow{d} Z_W \quad (n \rightarrow \infty)$$

where Z_W is a standard normal random variable. Then, under the assumptions given above,

$$n^{\frac{1}{2}-d} \frac{\bar{y}_n - \mu_W \mu_X}{\sqrt{c_{f,W} v(d)}} \xrightarrow{d} \mu_X Z_W$$

Remark 1 In Theorem 1, the assumption that X_t is an INAR(1) process is not needed. More generally, the result holds whenever the modulating process W has the properties given above, the modulated process Y_t is defined by Eq. (1), and X_t is a weakly stationary process with expected value μ_X and summable autocovariances $\gamma_X(k)$.

An analogous result can be obtained for sample autocovariances. In particular,

$$\hat{\gamma}_Y(k) = n^{-1} \sum_{j=1}^{n-k} (Y_j - \bar{y}_n)(Y_{j+k} - \bar{y}_n) \quad (k \geq 0)$$

converges in probability to

$$\gamma_Y(k) = \mu_X^2 \gamma_W(k) + (\mu_W^2 + \gamma_W(k)) \gamma_X(k)$$

Thus, the asymptotic bias of the unadjusted sample autocovariance is equal to

$$B_Y(k) = \gamma_Y(k) - \gamma_X(k) = \mu_X^2 \gamma_W(k) + (\mu_W^2 + \gamma_W(k) - 1) \gamma_X(k) \quad (10)$$

3 Conditional estimators

The modulating process W is usually not observable. Therefore the question arises whether consistent estimation of the INAR(1) parameters characterizing X_j is possible, without having to model the unobserved process W . A simple approach is to make use of the fact that $P(Y_j = X_j | Y_j > 0) = 1$. For instance, we may consider conditional moment estimators.

More specifically, for a Poisson INAR(1) process with intensity λ , we have $\mu_X = E(X) = \lambda/(1 - \alpha)$ and

$$E(X_j | X_j > 0) = \frac{\mu_X}{1 - \exp(-\mu_X)} =: g(\mu_X)$$

Moreover,

$$\begin{aligned} E(\mathbf{1}\{Y_j > 0\} Y_j) &= E(\mathbf{1}\{W(j) = 1\} \mathbf{1}\{X_j > 0\} X_j) \\ &= (1 - p_{0,W}) \mu_X \end{aligned}$$

$$E(\mathbf{1}\{Y_j > 0\}) = (1 - p_{0,W}) (1 - \exp(-\mu_X))$$

and hence

$$R = \frac{E(\mathbf{1}\{Y_j > 0\} Y_j)}{E(\mathbf{1}\{Y_j > 0\})} = E(X_j | X_j > 0)$$

Therefore,

$$R_n = \frac{n^{-1} \sum_{j=1}^n \mathbf{1}\{Y_j > 0\} Y_j}{n^{-1} \sum_{j=1}^n \mathbf{1}\{Y_j > 0\}}$$

provides a consistent nonparametric estimator of $E(X_j | X_j > 0)$. This motivates the definition of a conditional moment estimator of μ_X as the solution $\hat{\mu}_X$ of

$$g(\hat{\mu}_X) = R_n \tag{11}$$

It is straightforward to prove that Eq. (11) provides a consistent estimator:

Theorem 2 *Let $\hat{\mu}_X$ be defined by Eq. (11). Then*

$$\hat{\mu}_X \xrightarrow{p} \mu_X.$$

Remark 2 In Theorem 2, the assumption that X_t is an INAR(1) process is not needed. Suppose that X_t is a nonnegative integer valued weakly stationary process with a summable autocovariance function γ_X . Then, calculating $E(X_t | X_t > 0)$, the function g and the nonparametric estimator R_n can be adapted accordingly to obtain a consistent estimator $\hat{\mu}_X$ via (??).

While consistency of $\hat{\mu}_X$ is not influenced by the modulating process, this is not true for the asymptotic distribution of $\hat{\mu}_X$. This is discussed in Theorem 4 (Sect. 4).

The usual moment estimator of the INAR(1)-parameter α is obtained by the lag-one sample autocorrelation. However, as we saw above, for the modulated

process Y_j sample autocovariances are asymptotically biased estimators of γ_X . We may therefore consider an estimator based on the conditional autocovariance $cov(Y_j, Y_{j+1}|Y_j, Y_{j+1} > 0)$. By analogous arguments as above it is easy to see that

$$cov(Y_j, Y_{j+1}|Y_j, Y_{j+1} > 0) = cov(X_j, X_{j+1}|X_j, X_{j+1} > 0). \tag{12}$$

However, in terms of the INAR(1) parameters α and μ_X , the right hand side of Eq. (12) is a rather complicated nonlinear function. As it turns out, a much simpler expression can be obtained for the noncentered conditional moment $E(X_j X_{j+1}|X_j, X_{j+1} > 0)$. Specifically, for a Poisson INAR(1) process the following formulas hold:

Lemma 1 *Let X_j be a Poisson INAR(1) process with parameters μ_X and α . Then*

$$P(X_j X_{j+1} > 0) = 1 - 2e^{-\mu_X} + e^{-2\mu_X} e^{\mu_X \alpha}$$

and

$$E(X_j X_{j+1}|X_j, X_{j+1} > 0) = \frac{\alpha \mu_X + \mu_X^2}{1 - 2 \exp(-\mu_X) + \exp(-2\mu_X) \exp(\mu_X \alpha)}$$

A consistent nonparametric estimator of $E(X_j X_{j+1}|X_j, X_{j+1} > 0)$ is given by

$$E_n = \frac{n^{-1} \sum_{j=1}^{n-1} \mathbf{1}\{Y_j Y_{j+1} > 0\} Y_j Y_{j+1}}{n^{-1} \sum_{j=1}^{n-1} \mathbf{1}\{Y_j Y_{j+1} > 0\}}$$

Setting

$$h(\alpha, \mu_X) = \frac{\alpha \mu_X + \mu_X^2}{1 - 2 \exp(-\mu_X) + \exp(-2\mu_X) \exp(\mu_X \alpha)}$$

we define $\hat{\alpha}$ by

$$h(\hat{\alpha}, \hat{\mu}_X) = E_n \tag{13}$$

where $\hat{\mu}_X$ is obtained from (11). Equation (13) provides a consistent estimator:

Theorem 3 *Let $\hat{\alpha}$ be defined by Eq. (13). Then*

$$\hat{\alpha} \xrightarrow{p} \alpha$$

4 Testing for zero inflation

4.1 Definition of the test statistic

In this section, we consider the question how to test for zero-inflation due to modulation as specified by Eqs. (1), (5), (2) and (3). As before, we assume that the modulating

process W is not observable. Using the notation

$$p_{0,X} = P(X_j = 0) = \exp(-\mu_X), p_{0,Y} = P(Y_j = 0), p_{0,W} = P(W(t) = 0)$$

we may formulate the null hypothesis H_0 and the alternative H_1 as

$$H_0 : p_{0,Y} = p_{0,X}, H_1 : p_{0,Y} > p_{0,X}$$

For some general results on zero-inflation tests for INAR processes, see e.g. the recent paper by Weiss et al. (2019). Note that in our situation, we are testing explicitly for zero-inflation in the *marginal* distribution. We may therefore use a very simple statistic that compares the empirical relative frequency of zeroes with the frequency modelled by an INAR(1) process. Since $P(Y_j = X_j | Y_j > 0) = 1$, we suggest to use the consistent conditional estimator of μ_X defined in Eq. (11). Thus, let $\hat{\mu}_X$ be defined by Eq. (11), and let

$$\hat{p}_{0,X} = \exp(-\hat{\mu}_X)$$

and

$$\tilde{p}_{0,Y} = \frac{1}{n} \sum_{t=1}^n \mathbf{1}\{Y_t = 0\}$$

We define the test statistic

$$T_n = \tilde{p}_{0,Y} - \hat{p}_{0,X}$$

Then, under H_0 , T_n converges to zero in probability, whereas under the alternative H_1 , it converges to a positive constant. Asymptotic rejection regions are derived in the following section.

4.2 Asymptotic rejection regions

Before deriving the asymptotic distribution of T_n , we need to consider the joint asymptotic distribution of $\hat{\mu}_X$ and $\tilde{p}_{0,X}$. The following notation is adopted from Weiss et al. (2019):

$$\sigma_{11} = \mu_X \frac{1 + \alpha}{1 - \alpha} \tag{14}$$

$$\sigma_{22} = e^{-\mu_X} \left[1 - e^{-\mu_X} + 2e^{-\mu_X} \sum_{j=1}^{\infty} \frac{\mu_X^j}{j!} \frac{\alpha^j}{1 - \alpha^j} \right] \tag{15}$$

and

$$\sigma_{12} = -\mu_X e^{-\mu_X} \frac{1 + \alpha}{1 - \alpha} \tag{16}$$

Also, denote by

$$r_{0,X} = \frac{p_{0,X}}{1 - p_{0,X}}$$

the odds ratio based on $p_{0,X}$.

Theorem 4 *Let*

$$\zeta_n = \sqrt{n} (\hat{\mu}_X - \mu_X, \tilde{p}_{0,Y} - p_{0,X})$$

and suppose that H_0 holds. Then

$$\zeta_n \xrightarrow{d} \zeta$$

where $\zeta = (\zeta_1, \zeta_2)$ is a zero mean bivariate normal random variable with covariance matrix $\Sigma_\zeta = (c_{ij})_{i,j=1,2}$. The entries in Σ_ζ are defined by

$$c_{11} = \frac{\sigma_{11} + 2\mu_X r_{0,X} p_{0,X}^{-1} \sigma_{12} + \mu_X^2 r_{0,X}^2 p_{0,X}^{-2} \sigma_{22}}{(1 - \mu_X r_{0,X})^2} \tag{17}$$

$$c_{22} = \sigma_{22} \tag{18}$$

$$c_{12} = \frac{\sigma_{12} + \mu_X r_{0,X} p_{0,X}^{-1} \sigma_{22}}{1 - \mu_X r_{0,X}} \tag{19}$$

The asymptotic distribution of T_n under H_0 is given by the following Theorem:

Theorem 5 *Under H_0 ,*

$$\sqrt{n} T_n \xrightarrow{d} \sigma_T Z_T$$

where Z_T is a standard normal variable,

$$\sigma_T^2 = \frac{e^{-2\mu_X} \sigma_{11} + 2e^{-\mu_X} \sigma_{12} + \sigma_{22}}{(1 - \mu_X r_{0,X})^2}$$

and σ_{ij} are given by Eqs. (14), (16), (15).

Based on Theorem 5, asymptotic rejection regions at a level of significance $\alpha \in (0, 1)$ can be defined by

$$T_n > z_{1-\alpha} \frac{\sigma_T}{\sqrt{n}} \tag{20}$$

where $z_{1-\alpha}$ denotes the $(1 - \alpha)$ -quantile of a standard normal variable.

4.3 Asymptotic power

We consider the asymptotic distribution of T_n under the alternative as specified by Eqs. (1), (2), (4) and (5) with $0 < p_{0,W} < 1$. Note first that

$$\tilde{p}_{0,Y} = \frac{1}{n} \sum_{j=1}^n \mathbf{1}\{Y_j = 1\} = \frac{1}{n} \sum_{j=1}^n \mathbf{1}\{X_j = 0\} + \frac{1}{n} \sum_{j=1}^n \mathbf{1}\{Y_j = 0, X_j > 0\}$$

converges in probability to

$$\begin{aligned} P(Y_t = 0) &= p_{0,X} + P(W(t) = 0, X_t > 0) \\ &= p_{0,X} + p_{0,W} (1 - p_{0,X}) \end{aligned}$$

On the other hand, $\hat{p}_{0,X}$ converges in probability to $p_{0,X}$. Therefore, under H_1 ,

$$T_n \xrightarrow{p} p_{0,W} (1 - p_{0,X}) > 0$$

and

$$\lim_{n \rightarrow \infty} P\left(T_n > z_{1-\alpha} \frac{\sigma_T}{\sqrt{n}}\right) = 1$$

A more detailed result is given by the following Theorem. We will use the notation

$$\begin{aligned} S_{n,1,W} &= \sum_{j=1}^n (\mathbf{1}\{W(j) = 0\} - p_{0,W}) \\ \varphi(x) &= \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}x^2} \\ \nu(d) &= \frac{2 \sin \pi d}{d(2d + 1)} \Gamma(1 - 2d) \end{aligned}$$

and

$$q = \frac{p_{0,X}}{\sqrt{c_{f,W} \nu(d)}}$$

Theorem 6 Suppose that H_1 holds, as specified by Eqs. (1), (2), (4) and (5) with $0 < p_{0,W} < 1$. Moreover, assume that

$$\frac{S_{n,1,W}}{\sqrt{\text{var}(S_{n,1,W})}} \xrightarrow{d} Z_{1,W} \tag{21}$$

Table 1 Simulated mean (first number) and variance (second number) of unconditional and conditional estimates of μ_X and α under H_0 . For each combination of parameters and sample size, the results are based on 10,000 simulations

n	$\mu_X = 0.5$ with $\lambda = 0.4, \alpha = 0.2$		$\hat{\alpha}_{\text{uncond}}$	$\hat{\alpha}_{\text{cond}}$
	\bar{y}	$\hat{\mu}_X$		
100	0.50, 0.8	0.49, 2.6	0.18, 1.2	0.26, 7.6
200	0.50, 0.8	0.50, 2.6	0.19, 1.2	0.26, 13.2
400	0.50, 0.8	0.50, 2.4	0.20, 1.2	0.26, 20.8
800	0.50, 0.8	0.50, 2.4	0.20, 1.6	0.24, 27.2
1000	0.50, 1.0	0.50, 3.0	0.20, 1.0	0.23, 28.0
$\mu_X = 1$ with $\lambda = 0.4, \alpha = 0.6$				
100	1.00, 3.9	0.98, 6.5	0.56, 0.8	0.51, 7.0
200	1.00, 3.8	0.99, 6.6	0.58, 0.8	0.56, 11.0
400	1.00, 4.0	0.99, 6.8	0.59, 0.8	0.59, 16.4
800	1.00, 4.0	1.00, 6.4	0.60, 0.8	0.61, 25.6
1000	1.00, 4.0	1.00, 7.0	0.60, 1.0	0.62, 29.0

where $Z_{1,W}$ is a standard normal random variable. Then, for n large enough,

$$n^{d-\frac{1}{2}} \frac{1}{q + q^{-1}n^{2d-1}} \varphi \left(qn^{\frac{1}{2}-d} \right) \leq 1 - P \left(\sqrt{n}T_n > z_{1-\alpha} \right) \leq n^{d-\frac{1}{2}} \frac{1}{q} \varphi \left(qn^{\frac{1}{2}-d} \right) \tag{22}$$

Remark 3 Note that the rate at which the two bounds in Eq. (22) tend to zero, as $n \rightarrow \infty$, is slower for larger values of d . Thus, the power of the test becomes weaker with increasing long memory in the modulating process W . Also note that, with increasing d , the bounds become less sharp.

5 Simulations

We consider the model $Y_j = W(j)X_j$ where $W(j) = \tilde{W}(j)$ and \tilde{W} is the renewal reward process defined in Eq. (9). The following INAR(1) parameters are used: a) $\mu_W = 0.5$ with $\lambda = 0.4$ and $\alpha = 0.2$; b) $\mu_W = 1$ with $\lambda = 0.4$ and $\alpha = 0.6$. For W we consider $\mu_W = p_{0,W} = 0.5, 0.9, 1$, and $d = 0.1, 0.4$. In the simulations of the zero-inflation test we also add a case of extreme long memory, with $d = 0.49$. For each parameter combination and sample sizes $n = 100, 200, 400, 800$ and 1000 , ten thousand simulations were carried out. Simulated means and variances of the conditional moment estimates $\hat{\mu}_X$, as defined in Eq. (11) and $\hat{\alpha}_{\text{cond}}$, as defined in Eq. (13), are shown in Tables 1 ($\mu_W = 1$), 2 and 3 ($\mu_W = 0.5, 0.9$). For comparison, the unconditional estimates \bar{y} and $\hat{\alpha}_{\text{uncond}}$ are also given. Under H_0 , we observe an undisturbed INAR(1) process. As expected, the conditional estimates have a larger variance than the unconditional ones (Table 1). Under H_1 , we have $\mu_W < 1$, so that \bar{y} underestimates μ_X . Due to zero inflation, the absolute value of the bias increases as μ_W decreases. In contrast, no relevant finite sample bias is observed for the conditional estimator $\hat{\mu}_X$. For the unconditional estimator

Table 2 Simulated mean (first number) and variance (second number) of unconditional and conditional estimates of μ_X and α under H_1 with $\mu_X = 0.5$ ($\alpha=0.2, \lambda = 0.4$). For each combination of parameters and sample size, the results are based on 10,000 simulations

	$d = 0.1$		$d = 0.4$		$\hat{\mu}_X$	$\hat{\alpha}_{uncond}$	$\hat{\alpha}_{cond}$	$\hat{\mu}_X$	$\hat{\alpha}_{uncond}$	$\hat{\alpha}_{cond}$
	\bar{y}	$\hat{\mu}_X$	\bar{y}	$\hat{\mu}_X$						
$\mu_W = 0.5$										
$n = 100$	0.25, 0.9	0.49, 5.7	0.20, 1.8	0.24, 8.9	0.25, 1.7	0.20, 1.8	0.24, 8.9	0.48, 6.8	0.24, 2.1	0.24, 8.5
200	0.25, 1.0	0.50, 5.2	0.22, 1.8	0.26, 16.4	0.25, 2.4	0.22, 1.8	0.26, 16.4	0.49, 6.6	0.27, 2.2	0.25, 15.8
400	0.25, 1.2	0.50, 5.2	0.23, 2.0	0.26, 29.6	0.25, 3.6	0.23, 2.0	0.26, 29.6	0.50, 6.8	0.29, 2.8	0.25, 27.6
800	0.25, 1.6	0.50, 4.8	0.23, 2.4	0.27, 50.4	0.25, 5.6	0.23, 2.4	0.27, 50.4	0.50, 6.4	0.30, 3.2	0.26, 45.6
1000	0.25, 1.0	0.50, 5.0	0.24, 2.0	0.26, 57.0	0.25, 7.0	0.24, 2.0	0.26, 57.0	0.50, 6.0	0.30, 3.0	0.25, 50.0
$\mu_W = 0.9$										
$n = 100$	$d = 0.1$		$d = 0.4$		$d = 0.4$		$d = 0.4$		$d = 0.4$	
200	0.45, 0.9	0.49, 2.8	0.19, 1.3	0.26, 7.9	0.45, 1.1	0.19, 1.3	0.26, 7.9	0.49, 3.0	0.20, 1.4	0.25, 7.7
400	0.45, 1.0	0.50, 2.8	0.20, 1.4	0.26, 14.2	0.45, 1.4	0.20, 1.4	0.26, 14.2	0.50, 3.0	0.21, 1.6	0.26, 13.4
800	0.45, 0.8	0.50, 2.8	0.29, 1.2	0.25, 21.6	0.45, 2.0	0.29, 1.2	0.25, 21.6	0.50, 2.8	0.22, 1.6	0.25, 21.2
1000	0.45, 1.0	0.50, 3.0	0.21, 1.6	0.24, 30.4	0.45, 3.2	0.21, 1.6	0.24, 30.4	0.50, 3.2	0.22, 2.4	0.24, 28.8
			0.21, 1.0	0.24, 32.0	0.45, 3.0	0.21, 1.0	0.24, 32.0	0.50, 3.0	0.22, 2.0	0.24, 31.0

Table 3 Simulated mean (first number) and variance (second number) of unconditional and conditional estimates of μ_X and α under H_1 with $\mu_X = 1$ ($\alpha=0.6, \lambda = 0.4$). For each combination of parameters and sample size, the results are based on 10,000 simulations

	$d = 0.1$		$d = 0.4$		$\hat{\mu}_X$	$\hat{\alpha}_{\text{uncond}}$	$\hat{\alpha}_{\text{cond}}$	$\hat{\mu}_X$	$\hat{\alpha}_{\text{uncond}}$	$\hat{\alpha}_{\text{cond}}$
	\bar{y}	$\hat{\mu}_X$	\bar{y}	$\hat{\mu}_X$						
$\mu_W = 0.5$										
$n = 100$	0.50, 3.7	0.97 11.2	0.45, 1.9	0.43, 11.3	0.50, 6.9	0.97, 14.8	0.53, 2.2	0.43, 10.7		
200	0.50, 4.0	0.99 11.4	0.47, 2.2	0.49, 18.6	0.50, 10.0	0.98, 15.8	0.57, 2.4	0.48, 17.6		
400	0.50, 4.8	0.99 10.8	0.48, 2.4	0.54, 28.8	0.50, 14.8	0.99, 14.4	0.60, 2.4	0.54, 26.8		
800	0.50, 5.6	1.00 11.2	0.49, 2.4	0.58, 43.2	0.50, 24.0	1.00, 16.0	0.61, 2.4	0.58, 40.0		
1000	0.50, 6.0	1.00 11.0	0.50, 3.0	0.59, 50.0	0.50, 27.0	1.00, 16.0	0.62, 2.0	0.59, 47.0		
$\mu_W = 0.9$										
$n = 100$	0.90, 4.1	0.98, 7.0	0.53, 1.0	0.49, 7.7	0.90, 5.3	0.97, 7.3	0.56, 1.0	0.49, 7.5		
200	0.90, 4.2	0.99, 6.8	0.55, 1.0	0.55, 11.6	0.90, 6.4	0.99, 7.6	0.58, 1.0	0.55, 11.6		
400	0.90, 4.4	1.00, 6.8	0.56, 1.2	0.59, 18.0	0.90, 8.0	0.99, 7.2	0.59, 1.2	0.59, 17.6		
800	0.90, 4.8	1.00, 7.2	0.57, 0.8	0.61, 28.0	0.90, 11.2	1.00, 8.0	0.60, 1.6	0.61, 27.2		
1000	0.90, 5.0	1.00, 7.0	0.57, 1.0	0.62, 32.0	0.90, 12.0	1.00, 8.0	0.60, 1.0	0.61, 31.0		

Table 4 Simulated relative rejection frequencies based on asymptotic rejection regions defined by Eq. (20) with a level of significance of 0.05. For each combination of parameters and sample size, the results are based on 10,000 simulations

H_0 $\mu_X = 0.5$ with $\lambda = 0.4, \alpha = 0.2$					$\mu_X = 1$ with $\lambda = 0.4, \alpha = 0.6$				
$n = 100$	200	400	800	1000	$n = 100$	200	400	800	1000
0.04	0.04	0.04	0.04	0.05	0.03	0.02	0.02	0.02	0.02
H_1 with $\mu_W = 0.5, d = 0.1$									
0.43	0.64	0.86	0.97	0.98	0.70	0.92	0.99	0.99	1.00
H_1 with $\mu_W = 0.5, d = 0.4$									
0.39	0.59	0.80	0.93	0.95	0.64	0.84	0.94	0.97	0.98
H_1 with $\mu_W = 0.5, d = 0.49$									
0.38	0.57	0.77	0.89	0.91	0.60	0.80	0.90	0.95	0.96
H_1 with $\mu_W = 0.9, d = 0.1$									
0.09	0.12	0.17	0.26	0.25	0.10	0.13	0.26	0.38	0.47
H_1 with $\mu_W = 0.9, d = 0.4$									
0.12	0.14	0.19	0.24	0.30	0.12	0.17	0.22	0.30	0.39
H_1 with $\mu_W = 0.9, d = 0.49$									
0.10	0.13	0.17	0.25	0.28	0.12	0.17	0.23	0.33	0.37

of α , the effect of zero-inflation is much less clear. A noticeable bias of the unconditional estimator can only be observed for $p_{0,W} = 0.5$ and $d = 0.1$. A possible reason for the rather minor effect is that the potential bias under zero-inflation may be compensated by a modified lag-one autocorrelation of the observed process due to dependence in W .

Tables 4 show results for the zero-inflation test based on Eq. (20). Under H_0 , the simulated levels are reasonably close to the nominal value of 0.05. As expected, for smaller values of $p_{0,W}$, the test has higher power. The effect of long memory is less obvious for mild zero-inflation ($p_{0,W} = 0.9$). For $p_{0,W} = 0.5$, the simulation results show a clearer picture, with a tendency for slightly lower power when d is large.

Finally, Table 5 shows results for the case where X_j is generated by an INAR(2) process (Alzaid and Al-Osh 1990), but is misspecified as an INAR(1) model. Specifically, we simulate $X_j = \alpha_1 \circ X_{j-1} + \alpha_2 \circ X_{j-2} + \varepsilon_j$ where $(\alpha_1, \alpha_2) = (0.2, 0.2)$ and $(0.6, 0.2)$ respectively, and ε_j are iid Poisson variables with $\lambda = 0.4$. For $(\alpha_1, \alpha_2) = (0.6, 0.2)$, the simulated rejection frequencies under H_0 are very close to the nominal level of 0.05. For $(\alpha_1, \alpha_2) = (0.2, 0.2)$, rejection frequencies tend to be slightly too high. A possible explanation is that for $\alpha_1 = 0.6$ the relative influence of X_{j-1} on X_j is much stronger than for $\alpha_1 = 0.2$. In this sense, the INAR(2) process is closer to an INAR(1) process for larger values of the ratio α_1/α_2 , and the misspecification is less noticeable. On the other hand, under H_1 , rejection probabilities tend to be higher for $\alpha_1 = 0.2$. This may be explained, at least partially, by the fact that for this parameter setting the level of significance is higher than the nominal level of

Table 5 Simulated relative rejection frequencies under misspecification. The results are based on 10,000 simulations of an INAR(2) process with $(\alpha_1, \alpha_2) = (0.2, 0.2)$ and $(0.6, 0.2)$ respectively, and ε_j iid Poisson variables with $\lambda = 0.4$. The alternative considered here is $\mu_W = 0.5$, with $d = 0.1$ and $d = 0.4$ respectively. For each simulated series, a (misspecified) INAR(1) model was fitted using the methods in Sect. 3 and the test defined in Sect. 4 was applied, at a nominal level of significance of 0.05

H_0 $(\alpha_1, \alpha_2) = (0.2, 0.2)$					$(\alpha_1, \alpha_2) = (0.6, 0.2)$				
$n = 100$	200	400	800	1000	$n = 100$	200	400	800	1000
0.04	0.05	0.06	0.09	0.10	0.08	0.08	0.07	0.05	0.04
H_1 with $d = 0.1$									
0.59	0.82	0.96	0.99	0.99	0.62	0.56	0.45	0.46	0.53
H_1 with $d = 0.4$									
0.53	0.76	0.91	0.97	0.98	0.63	0.56	0.49	0.48	0.53

0.05. Generally speaking, one may conjecture that the test is likely to be fairly robust to mild deviations from an INAR(1) model. If serious deviations from the ideal model are expected, a modification of the test may be preferable (see Remarks 1 and 2). An interesting task for future research is to develop adaptive tests for zero-inflation that are applicable without the need of specifying which member of a certain family of "ideal models" (e.g. the family of all INAR(p) models with $p = 1, 2, \dots$) generates the unmodulated process X_j .

6 Final remarks

In this paper, we introduced a zero-inflated INAR(1) model where zero-inflation is caused by a long-memory process. Parameter estimation and testing for zero-inflation was considered. Various extensions of this model are of interest. For instance, modulation may occur at the level of the innovations of the INAR(1) process. The structure of this type of modulated INAR(1) processes is much more complex, and will be considered elsewhere. Another obvious extension is to consider model (1), however replacing X_j by a general INAR(p) process. It is expected that the methods considered here can be extended to this case using analogous arguments as for $p = 1$.

Finally note that instead using the test statistic discussed in Sect. 4, one may consider the lengths of zero runs (for related results see e.g. Jazi et al. 2012, Qi et al. 2019). An advantage of the test considered here is its simplicity. In particular, under the alternative, the distribution of runs can be quite complicated. On the other hand, the estimator of μ_X used here, omits parts of the information that may be useful for detecting deviations from the null hypothesis. The development of more efficient tests, for instance based on runs, is an interesting task for future research.

Acknowledgements We would like to thank the referees for insightful comments that helped to improve the presentation of the results.

Funding Open Access funding enabled and organized by Projekt DEAL.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

7 Appendix—Proofs, Tables

7.1 Proofs

Proof of Theorem 1 First of all, $E(\bar{y}_n) = \mu_W \mu_X = p_{0,W} \mu_X$, because the two processes X and W are independent of each other. Moreover,

$$\begin{aligned} n^{-1} \sum_{j=1}^n Y_j &= n^{-1} \sum_{j=1}^n W(j) (X_j - \mu_X) + \mu_X n^{-1} \sum_{j=1}^n W(j) \\ &= \mu_W \mu_X + n^{-1} \sum_{j=1}^n W(j) (X_j - \mu_X) + \mu_X n^{-1} \sum_{j=1}^n (W(j) - \mu_W) \\ &= \mu_W \mu_X + n^{-1} S_{n,1} + \mu_X n^{-1} S_{n,2}. \end{aligned}$$

Now,

$$E [W(j) (X_j - \mu_X)] = 0$$

and

$$\begin{aligned} |cov (W(j) (X_j - \mu_X), W(j+k) (X_{j+k} - \mu_X))| &= |\gamma_W(k) + \mu_W^2| |\gamma_X(k)| \\ &\leq c |\gamma_X(k)| \end{aligned}$$

for a suitable constant $0 < c < \infty$. Therefore

$$n^{-1} \sum_{j=1}^n Y_j - \mu_W \mu_X = O_p \left(n^{-\frac{1}{2}} \right)$$

Furthermore,

$$S_{n,2} = \bar{w}_n - \mu_W = \sqrt{var(\bar{w}_n)} Z_{n,W}$$

Now Eqs. (7) and (8) imply

$$var(\bar{w}_n) \underset{n \rightarrow \infty}{\sim} c_{f,W} v(d) n^{2d-1}$$

(see e.g. Beran et al. 2013, Corollary 1.2). Since $d > 0$, $n^{-1}S_{n,1}$ is asymptotically negligible and the result follows. \square

Proof of Theorem 2 The function g is monotonically increasing, with $\lim_{\mu \rightarrow 0} g(\mu) = 1$ and $\lim_{\mu \rightarrow \infty} g(\mu) = \infty$. Since $R_n \geq 1$, Eq. (11) has a unique solution. Moreover, W is a stationary 0 – 1–process, and

$$\begin{aligned} cov(\mathbf{1}\{Y_j > 0\} Y_j, \mathbf{1}\{Y_{j+k} > 0\} Y_{j+k}) &= cov(Y_j, Y_{j+k}) = \gamma_Y(k) \\ &= \mu_X^2 \gamma_W(k) + \mu_W^2 \gamma_X(k) + \gamma_W(k) \gamma_X(k) \end{aligned}$$

Tschebyshev’s inequality then implies

$$n^{-1} \sum_{j=1}^n \mathbf{1}\{Y_j > 0\} Y_j \rightarrow_p E(\mathbf{1}\{Y > 0\} Y) = (1 - p_{0,W}) \mu_X$$

and

$$n^{-1} \sum_{j=1}^n \mathbf{1}\{Y_j > 0\} \rightarrow_p (1 - p_{0,W}) (1 - \exp(-\mu_X))$$

so that

$$R_n \xrightarrow{p} \frac{\mu_X}{1 - \exp(-\mu_X)}$$

\square

Proof of Lemma 1 First of all,

$$\begin{aligned} P(X_j X_{j+1} > 0) &= \sum_{k=1}^{\infty} \frac{\mu_X^k}{k!} e^{-\mu_X} \left[1 - (1 - \alpha)^k + (1 - e^{-\lambda}) (1 - \alpha)^k \right] \\ &= \sum_{k=1}^{\infty} \frac{\mu_X^k}{k!} e^{-\mu_X} \left[1 - e^{-\lambda} (1 - \alpha)^k \right] \\ &= (1 - e^{-\mu_X}) - e^{-(\lambda + \mu_X) + \mu_X(1 - \alpha)} \sum_{k=1}^{\infty} \frac{[\mu_X (1 - \alpha)]^k}{k!} e^{-\mu_X(1 - \alpha)} \\ &= (1 - e^{-\mu_X}) - e^{-\mu_X} \left(1 - e^{-\mu_X(1 - \alpha)} \right) \\ &= 1 - 2e^{-\mu_X} + e^{-2\mu_X} e^{\mu_X \alpha} \end{aligned}$$

Then

$$E(X_j X_{j+1} | X_j, X_{j+1} > 0) = \frac{E(X_j X_{j+1})}{P(X_j X_{j+1} > 0)} = \frac{\gamma_X(1) + \mu_X^2}{P(X_j X_{j+1} > 0)}$$

$$= \frac{\alpha\mu_X + \mu_X^2}{1 - 2 \exp(-\mu_X) + \exp(-2\mu_X) \exp(\mu_X \alpha)}$$

□

Proof of Theorem 3 First note that

$$\begin{aligned} E(\mathbf{1}\{Y_j Y_{j+1} > 0\} Y_j Y_{j+1}) &= E(\mathbf{1}\{W(j) W(j+k) > 0\}) \\ &\quad E(\mathbf{1}\{X_j X_{j+1} > 0\} X_j X_{j+k}), \\ E(\mathbf{1}\{Y_j Y_{j+1} > 0\}) &= E(\mathbf{1}\{W(j) W(j+k) > 0\}) E(\mathbf{1}\{X_j X_{j+1} > 0\}) \end{aligned}$$

so that

$$\begin{aligned} \frac{E(\mathbf{1}\{Y_j Y_{j+1} > 0\} Y_j Y_{j+1})}{E(\mathbf{1}\{Y_j Y_{j+1} > 0\})} &= \frac{E(\mathbf{1}\{X_j X_{j+1} > 0\} X_j X_{j+k})}{E(\mathbf{1}\{X_j X_{j+1} > 0\})} \\ &= E(X_j X_{j+1} | X_j, X_{j+1} > 0) \end{aligned}$$

Then,

$$n^{-1} \sum_{j=1}^{n-1} \mathbf{1}\{Y_j Y_{j+1} > 0\} Y_j Y_{j+1} \xrightarrow{p} E(\mathbf{1}\{Y_j Y_{j+1} > 0\} Y_j Y_{j+1})$$

and

$$n^{-1} \sum_{j=1}^{n-1} \mathbf{1}\{Y_j Y_{j+1} > 0\} \xrightarrow{p} E(\mathbf{1}\{Y_j Y_{j+1} > 0\})$$

implies

$$E_n \xrightarrow{p} E(X_j X_{j+1} | X_j, X_{j+1} > 0)$$

Now, for any fixed $\mu_X > 0$, $h(\alpha, \mu_X)$ is monotonically increasing in α with

$$\begin{aligned} \lim_{\alpha \rightarrow 0} h(\alpha, \mu_X) &= \frac{\mu_X^2}{(1 - \exp(-\mu_X))^2} = E^2(X|X > 0) \\ \lim_{\alpha \rightarrow 1} h(\alpha, \mu_X) &= \frac{\mu_X(\mu_X + 1)}{1 - \exp(-\mu_X)} = E(X^2|X > 0) \end{aligned}$$

Hence, together with Theorem 2, we may conclude that, for n large enough, (13) has a unique solution and $\hat{\alpha} \rightarrow_p \alpha$. □

Proof of Theorem 4 Recall that

$$\tilde{p}_{0,Y} = \frac{1}{n} \sum_{j=1}^n \mathbf{1} \{Y_j = 0\}$$

and

$$g(\hat{\mu}_X) = \frac{n^{-1} S_n}{1 - \tilde{p}_{0,Y}}$$

where

$$g(\mu) = \frac{\mu_X}{1 - p_{0,X}} = \frac{\mu_X}{1 - \exp(-\mu_X)} = E(X_j | X_j > 0)$$

and

$$S_n = \sum_{j=1}^n \mathbf{1} \{Y_j > 0\} Y_j = \sum_{j=1}^n Y_j = n\bar{y}$$

Under H_0 , we have

$$n^{-1} S_n = \bar{y} = \bar{x}$$

so that

$$g(\hat{\mu}_X) = \frac{\bar{x}}{1 - \tilde{p}_{0,Y}} \quad (23)$$

Since $\hat{\mu}_X$ is consistent, we may use a Taylor expansion

$$g(\hat{\mu}_X) = g(\mu_X) + g'(\mu_X)(\hat{\mu}_X - \mu_X) + o_p(\hat{\mu}_X - \mu_X)$$

to obtain

$$\hat{\mu}_X - \mu_X = \frac{1}{g'(\mu_X)} \left[\frac{\bar{x}}{1 - \tilde{p}_{0,Y}} - g(\mu_X) \right] + o_p(\hat{\mu}_X - \mu_X) \quad (24)$$

Moreover, under H_0 , we also have $p_{0,Y} = p_{0,X}$ and

$$\frac{1}{1 - \tilde{p}_{0,Y}} = \frac{1}{1 - p_{0,X}} \left[1 + \frac{\tilde{p}_{0,Y} - p_{0,X}}{1 - p_{0,X}} + o_p(\tilde{p}_{0,Y} - p_{0,X}) \right]$$

Now

$$\frac{\bar{x}}{1 - \tilde{p}_{0,Y}} - g(\mu_X) = \frac{\bar{x} - \mu_X}{1 - \tilde{p}_{0,Y}} + \left(\frac{\mu_X}{1 - \tilde{p}_{0,Y}} - g(\mu_X) \right)$$

$$= A_n + B_n$$

The second term can be written as

$$\begin{aligned} B_n &= \frac{\mu_X}{1 - \tilde{p}_{0,Y}} - g(\mu_X) \\ &= \frac{\mu_X}{1 - p_{0,X}} \left[1 + \frac{\tilde{p}_{0,Y} - p_{0,X}}{1 - p_{0,X}} + o_p(\tilde{p}_{0,Y} - p_{0,X}) \right] - g(\mu_X) \\ &= g(\mu_X) \frac{\tilde{p}_{0,Y} - p_{0,X}}{1 - p_{0,X}} + o_p(\tilde{p}_{0,Y} - p_{0,X}) \end{aligned}$$

The first term can be simplified to

$$A_n = \frac{\bar{x} - \mu_X}{1 - p_{0,X}} + o_p(\tilde{p}_{0,Y} - p_{0,X})$$

Thus,

$$\begin{aligned} \hat{\mu}_X - \mu_X &= \frac{1}{g'(\mu_X)} (A_n + B_n) + o_p(\hat{\mu}_X - \mu_X) \\ &= \frac{(\bar{x} - \mu_X) + g(\mu_X)(\tilde{p}_{0,Y} - p_{0,X})}{g'(\mu_X)(1 - p_{0,X})} + o_p(\hat{\mu}_X - \mu_X) + o_p(\tilde{p}_{0,Y} - p_{0,X}) \end{aligned}$$

Applying Theorem 3.1.1 in Weiss et al. (2019), this can be simplified to

$$\hat{\mu}_X - \mu_X = \frac{Z_1 + g(\mu_X) Z_2}{(1 - p_{0,X}) g'(\mu_X)} n^{-\frac{1}{2}} + o_p(n^{-\frac{1}{2}})$$

where $Z = (Z_1, Z_2)$ is a zero mean bivariate random variable with covariance matrix $\Sigma = (\sigma_{ij})_{i,j=1,2}$, as defined by Eqs. (14), (16), (15). Note furthermore that $p_{0,X} = \exp(-\mu_X)$,

$$g'(\mu) = \frac{1 - (\mu + 1) \exp(-\mu)}{(1 - \exp(-\mu))^2}$$

so that

$$\begin{aligned} \frac{1}{g'(\mu_X)(1 - p_{0,X})} &= \frac{1 - \exp(-\mu_X)}{1 - (\mu_X + 1) \exp(-\mu_X)} \\ &= \frac{1}{1 - \mu_X r_{0,X}} \end{aligned}$$

and

$$\frac{g(\mu_X)}{g'(\mu_X)(1 - p_{0,X})} = \frac{\mu_X}{1 - (\mu_X + 1) \exp(-\mu_X)}$$

$$= \frac{\mu_X}{1 - p_{0,X}} \frac{1}{1 - \mu_X r_{0,X}}$$

Thus, we obtain

$$\hat{\mu}_X - \mu_X = \frac{1}{1 - \mu_X r_{0,X}} \left(Z_1 + \frac{\mu_X}{1 - p_{0,X}} Z_2 \right) n^{-\frac{1}{2}} + o_p \left(n^{-\frac{1}{2}} \right)$$

Moreover, by definition,

$$\tilde{p}_{0,Y} - p_{0,X} = n^{-\frac{1}{2}} Z_2 + o_p \left(n^{-\frac{1}{2}} \right)$$

Therefore

$$\zeta_n = \sqrt{n} (\hat{\mu}_X - \mu_X, \tilde{p}_{0,Y} - p_{0,X}) \xrightarrow{d} \zeta$$

where ζ has the covariance matrix Σ_ζ defined by Eqs. (17), (18) and (19). □

Proof of Theorem 5 Note first that

$$\begin{aligned} \hat{p}_{0,X} &= \exp(-\hat{\mu}_X) = p_{0,X} \exp(-(\hat{\mu}_X - \mu_X)) \\ &= p_{0,X} (1 - (\hat{\mu}_X - \mu_X) + o_p(\hat{\mu}_X - \mu_X)) \end{aligned}$$

Theorem 4 implies

$$\hat{p}_{0,X} - p_{0,X} = -p_{0,X} \zeta_1 n^{-\frac{1}{2}} + o_p \left(n^{-\frac{1}{2}} \right)$$

and

$$\begin{aligned} \sqrt{n} T_n &= \tilde{p}_{0,Y} - \hat{p}_{0,X} \\ &= \zeta_2 + p_{0,X} \zeta_1 + o_p(1) \end{aligned}$$

where ζ is defined in Theorem 4. Then formulas Eqs. (17),(18),(19) lead to

$$\begin{aligned} \sigma_T^2 &= \lim_{n \rightarrow \infty} n \cdot \text{var} (T_n) \\ &= \text{var} (\zeta_2) + 2p_{0,X} \text{cov} (\zeta_1, \zeta_2) + p_{0,X}^2 \text{var} (\zeta_1) \\ &= \frac{\sigma_{22} + 2p_{0,X} \sigma_{12} + \sigma_{11} p_{0,X}^2}{(1 - \mu_X r_{0,X})^2} \\ &= \frac{e^{-2\mu_X} \sigma_{11} + 2e^{-\mu_X} \sigma_{12} + \sigma_{22}}{(1 - \mu_X r_{0,X})^2} \end{aligned}$$

□

Proof of Theorem 6 First note that

$$\tilde{p}_{0,Y} = A_n + B_n$$

where

$$A_n = \frac{1}{n} \sum_{j=1}^n \mathbf{1}\{X_j = 0\} = p_{0,X} + n^{-\frac{1}{2}} Z_2 + o_p\left(n^{-\frac{1}{2}}\right)$$

and

$$\begin{aligned} B_n &= \frac{1}{n} \sum_{j=1}^n \mathbf{1}\{Y_j = 0, X_j > 0\} \\ &= p_{0,W} (1 - p_{0,X}) + n^{-1} S_{1,n} - n^{-1} S_{2,n} \end{aligned}$$

with

$$\begin{aligned} n^{-1} S_{1,n} &= n^{-1} \sum_{j=1}^n \mathbf{1}\{W(j) = 0\} - p_{0,W} \\ n^{-1} S_{2,n} &= n^{-1} \sum_{j=1}^n \mathbf{1}\{W(j) = 0\} \mathbf{1}\{X_j = 0\} - p_{0,W} p_{0,X} \end{aligned}$$

Also note that

$$\begin{aligned} & cov(\mathbf{1}\{W(j) = 0\}, \mathbf{1}\{W(j+k) = 0\}) \\ &= cov(1 - W(j), 1 - W(j+k)) \\ &= \gamma_W(k) \end{aligned}$$

By assumption,

$$\gamma_W(k) \underset{k \rightarrow \infty}{\sim} c_{\gamma,1,W} k^{2d-1}$$

where

$$c_{\gamma,W} = 2c_{f,W} \Gamma(1 - 2d) \sin \pi d$$

(see e.g. Beran et al. 2013, Theorem 1.3). Moreover,

$$\begin{aligned} & E[\mathbf{1}\{W(j) = 0\} \mathbf{1}\{X_j = 0\} \mathbf{1}\{W(j+k) = 0\} \mathbf{1}\{X_{j+k} = 0\}] \\ &= E[\mathbf{1}\{W(j) = 0\} \mathbf{1}\{W(j+k) = 0\}] E[\mathbf{1}\{X_j = 0\} \mathbf{1}\{X_{j+k} = 0\}] \\ &= (\gamma_{1,W}(k) + p_{0,W}^2) (\gamma_X(k) + p_{0,X}^2) \end{aligned}$$

$$E^2 [\mathbf{1}\{W(j) = 0\} \mathbf{1}\{X_j = 0\}] = p_{0,W}^2 p_{0,X}^2,$$

and hence

$$\begin{aligned} cov(\mathbf{1}\{W(j) = 0\} \mathbf{1}\{X_j = 0\}, \mathbf{1}\{W(j+k) = 0\} \mathbf{1}\{X_{j+k} = 0\}) \\ = p_{0,X}^2 \gamma_W(k) + p_{0,W}^2 \gamma_X(k) + \gamma_W(k) \gamma_X(k) \end{aligned}$$

This implies

$$cov(\mathbf{1}\{W(j) = 0\} \mathbf{1}\{X_j = 0\}, \mathbf{1}\{W(j+k) = 0\} \mathbf{1}\{X_{j+k} = 0\}) \underset{k \rightarrow \infty}{\sim} c_{\gamma,W} k^{2d-1} p_{0,X}^2$$

Also,

$$\begin{aligned} E(\mathbf{1}\{W(j) = 0\} \mathbf{1}\{W(j+k) = 0\} \mathbf{1}\{X_{j+k} = 0\}) \\ = p_{0,X} (\gamma_W(k) + p_{0,W}^2) \end{aligned}$$

and

$$E(\mathbf{1}\{W(j) = 0\}) E(\mathbf{1}\{W(j+k) = 0\} \mathbf{1}\{X_{j+k} = 0\}) = p_{0,X} p_{0,W}^2$$

so that

$$cov(\mathbf{1}\{W(j) = 0\}, \mathbf{1}\{W(j+k) = 0\} \mathbf{1}\{X_{j+k} = 0\}) = p_{0,X} \gamma_W(k)$$

We then obtain

$$var(n^{-1}(S_1 - S_2)) = n^{-2} [var(S_1) + var(S_2) - 2cov(S_1, S_2)]$$

with

$$var(S_1) = \sum_{k=-(n-1)}^{n-1} (n - |k|) \gamma_W(k) \underset{n \rightarrow \infty}{\sim} c_{f,W} \nu(d) n^{2d+1}$$

$$\begin{aligned} var(S_2) &= p_{0,X}^2 \sum_{k=-(n-1)}^{n-1} (n - |k|) \gamma_W(k) + p_{0,W}^2 \sum_{k=-(n-1)}^{n-1} (n - |k|) \gamma_X(k) \\ &\quad + 2 \sum_{k=-(n-1)}^{n-1} (n - |k|) \gamma_W(k) \gamma_X(k) \\ &\underset{n \rightarrow \infty}{\sim} p_{0,X}^2 c_{f,W} \nu(d) n^{2d+1} \end{aligned}$$

and

$$\text{cov}(S_1, S_2) = p_{0,X} \sum_{k=-(n-1)}^{n-1} (n - |k|) \gamma_{1,W}(k) \underset{n \rightarrow \infty}{\sim} p_{0,X} c_{f,W\nu}(d) n^{2d+1}$$

Thus, overall

$$\text{var}\left(n^{-1}(S_1 - S_2)\right) \underset{n \rightarrow \infty}{\sim} c_{f,W\nu}(d) (1 - p_{0,X})^2 n^{2d-1}$$

Moreover, under the given assumptions, $n^{\frac{1}{2}-d}(S_{1,n}, S_{2,n})$ are asymptotically jointly normal, so that

$$\tilde{p}_0 = p_{0,X} + p_{0,X} (1 - p_{0,X}) + n^{d-\frac{1}{2}} \sqrt{c_{f,W\nu}(d)} (1 - p_{0,X}) Z_3 + o_p\left(n^{d-\frac{1}{2}}\right)$$

where $Z_3 \sim N(0, 1)$. Hence, we have

$$\begin{aligned} \sqrt{n}T_n &= \sqrt{n}(\tilde{p}_0 - \hat{p}_0) \\ &= \sqrt{n}p_{0,X} (1 - p_{0,X}) + n^d \sqrt{c_{f,W\nu}(d)} (1 - p_{0,X}) Z_3 + o_p\left(n^{d-\frac{1}{2}}\right) \end{aligned}$$

For

$$\Delta_n = 1 - P\left(\sqrt{n}T_n > z_{1-\alpha}\right)$$

this yields

$$\begin{aligned} \Delta_n &= 1 - P\left(\sqrt{n}p_{0,X} (1 - p_{0,X}) + n^d \sqrt{c_{f,W\nu}(d)} (1 - p_{0,X}) Z_3 > z_{1-\alpha}\right) + o(\Delta_n) \\ &= P\left(Z_3 > n^{\frac{1}{2}-d} \frac{p_{0,X}}{\sqrt{c_{f,W\nu}(d)}}\right) + o(\Delta_n) \end{aligned}$$

Inequality (22) then follows from

$$\frac{\exp\left(-\frac{1}{2}x^2\right)}{\sqrt{2\pi}(x + x^{-1})} < 1 - \Phi(x) < \frac{\exp\left(-\frac{1}{2}x^2\right)}{\sqrt{2\pi}x}$$

(Gordon 1941) by plugging in

$$\begin{aligned} x + x^{-1} &= n^{\frac{1}{2}-d} \frac{p_{0,X}}{\sqrt{c_{f,W\nu}(d)}} + n^{d-\frac{1}{2}} \frac{\sqrt{c_{f,W\nu}(d)}}{p_{0,X}} \\ &= n^{\frac{1}{2}-d} \left(q + q^{-1}n^{2d-1}\right) \end{aligned}$$

□

7.2 Tables

See Tables 1, 2, 3 and 4.

References

- Al-Osh MA, Alzaid AA (1987) First-order integer-valued autoregressive (INAR(1)) process. *J Time Ser Anal* 8(3):261–275
- Alzaid AA, Al-Osh MA (1990) An integer-valued p th-order autoregressive structure (INAR(p)) process. *J Appl Probab* 27(2):314–324
- Barreto-Souza W (2015) Zero-modified geometric INAR(1) process for modelling count time series with deflation or inflation of zeros. *J Time Ser Anal* 36(6):839–852
- Beran J (1994) *Statistics for long-memory processes*. Chapman and Hall, CRC Press, New York
- Beran J, Feng Y, Ghosh S, Kulik R (2013) *Long-memory processes-probabilistic properties and statistical methods*. Springer, New York
- Bourguignon M, Weiss CH (2017) An INAR(1) process for modeling count time series with equidispersion, underdispersion and overdispersion. *Test* 26(4):847–868
- Che Z, Purushotham S, Cho K, Sontag D, Yan Liu Y (2018) Recurrent neural networks for multivariate time series with missing values. *Sci Rep* 8:6085. <https://doi.org/10.1038/s41598-018-24271-9>
- Da Silva EM, Oliveira VL (2004) Difference equations for the higher order moments and cumulants of the INAR (1) model. *J Time Ser Anal* 25(3):317–333
- Davis R, Fokianos K, Holan S, Joe H, Livsey J, Lund R, Pipiras V, Ravishanker N (2021) Count time series: a methodological review. *J Am Stat Assoc* 116:1–50
- Drost FC, Van Den Akker R, Werker BJM (2008) Local asymptotic normality and efficient estimation for INAR (p) models. *J Time Ser Anal* 29(5):783–801
- Du JG, Li Y (1991) The integer-valued autoregressive (INAR (p)) model. *J Time Ser Anal* 12(2):129–142
- Freeland RK, McCabe BPM (2004) Forecasting discrete valued low count time series. *Int J Forecast* 20(3):427–434
- Freeland RK, McCabe BPM (2005) Asymptotic properties of CLS estimators in the Poisson AR(1) model. *Stat Probab Lett* 73(2):147–153
- Gauthier G, Latour A (1994) Convergence forte des estimateurs des paramètres d'un processus GENAR(p). *Ann Sci Math Québec* 18(1):49–71
- Giraitis L, Koul HL, Surgailis D (2012) *Large sample inference for long memory processes*. Imperial College Press, London
- Gordon RD (1941) Values of Mills' ratio of area to bounding ordinate of the normal probability integral for large values of the argument. *Ann Math Stat* 12:364–366
- Gourieroux C, Jasiak J (2004) Heterogeneous INAR(1) model with application to car insurance. *Insur Math Econ* 34(2):177–192
- Jazi MA, Jones G, Lai CD (2012) First-order integer valued AR processes with zero inflated Poisson innovations. *J Time Ser Anal* 33:954–963
- Jentsch C, Weiss CH (2019) Bootstrapping INAR models. *Bernoulli* 25(3):2359–2408
- Jung RC, Ronning G, Tremayne AR (2005) Estimation in conditional first order autoregression with discrete support. *Stat Papers* 46:195–224
- Martin TG, Wintle BA, Rhodes JR, Kuhnert PM, Field SA, Low-Choy SJ, Tyre AJ, Possingham HP (2005) Zero tolerance ecology: improving ecological inference by modelling the source of zero observations. *Ecol Lett* 8:1235–1246
- McKenzie E (1985) Some simple models for discrete variate time series. *Water Resour Bull.* 21(4):645–650
- Möller TA, Weiß CH, Kim HY, Sirchenko A (2018) Modeling zero inflation in count data time series with bounded support. *Methodol Comput Appl Probab* 20:589–609
- Park Y, Kim HY (2012) Diagnostic checks for integer-valued autoregressive models using expected residuals. *Stat Papers* 53(4):951–970
- Pavlopoulos H, Karlis D (2008) INAR (1) modeling of overdispersed count series with an environmental application. *Environmetrics* 19(4):369–393
- Pedeli X, Karlis D (2013) On composite likelihood estimation of a multivariate INAR(1) model. *J Time Ser Anal* 34(2):206–220

- Pedeli X, Davison AC, Fokianos K (2015) Likelihood estimation for the INAR(p) model by saddlepoint approximation. *J Am Stat Assoc* 110(511):1229–1238
- Pipiras V, Taqqu MS (2017) Long-range dependence and self-similarity. Cambridge University Press, Cambridge
- Puig P, Valero J (2007) Characterization of count data distributions involving additivity and binomial subsampling. *Bernoulli* 13(2):544–555
- Qi X, Li Q, Zhu F (2019) Modeling time series of count with excess zeroes and ones based on INAR(1) model with zero-and-one inflated poisson innovations. *J Comput Appl Math* 346:572–590
- Qureshi A (2014) A long-memory integer-valued time series model, INARFIMA, for financial application. *Quantit Financ* 14(12):2225–2235
- Schweer S, Weiss CH (2014) Compound Poisson INAR(1) processes: stochastic properties and testing for overdispersion. *Comput Stat Data Anal* 77:267–284
- Steutel FW, Van Harn K (1979) Discrete analogues of self-decomposability and stability. *Ann Probab* 7(5):893–899
- Weiss CH (2013) Integer-valued autoregressive models for counts showing underdispersion. *J Appl Stat* 40(9):1931–1948
- Weiss CH (2018) An introduction to discrete-valued time series. Wiley, Hoboken
- Weiss CH, Homburg A, Puig P (2019) Testing for zero inflation and overdispersion in INAR(1) models. *Stat Papers* 60:823–848
- Young DS, Roemmele ES, Yeh P (2022) (2022) Zero-inflated modeling part I: traditional zero-inflated count regression models, their applications, and computational tools. *WIREs Comput Stat* 14:e1541. <https://doi.org/10.1002/wics.1541>
- Zygmund A (1968) Trigonometric series, vol 1. Cambridge University Press, Cambridge

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.