

Ebner, Bruno; Henze, Norbert

Article — Published Version

On the eigenvalues associated with the limit null distribution of the Epps-Pulley test of normality

Statistical Papers

Provided in Cooperation with:

Springer Nature

Suggested Citation: Ebner, Bruno; Henze, Norbert (2022) : On the eigenvalues associated with the limit null distribution of the Epps-Pulley test of normality, Statistical Papers, ISSN 1613-9798, Springer, Berlin, Heidelberg, Vol. 64, Iss. 3, pp. 739-752, <https://doi.org/10.1007/s00362-022-01336-6>

This Version is available at:

<https://hdl.handle.net/10419/308594>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.



<https://creativecommons.org/licenses/by/4.0/>



On the eigenvalues associated with the limit null distribution of the Epps-Pulley test of normality

Bruno Ebner¹ · Norbert Henze¹

Received: 20 September 2021 / Revised: 19 April 2022 / Accepted: 16 June 2022 /
Published online: 5 July 2022
© The Author(s) 2022

Abstract

The Shapiro–Wilk test (SW) and the Anderson–Darling test (AD) turned out to be strong procedures for testing for normality. They are joined by a class of tests for normality proposed by Epps and Pulley that, in contrast to SW and AD, have been extended by Baringhaus and Henze to yield easy-to-use affine invariant and universally consistent tests for normality in any dimension. The limit null distribution of the Epps–Pulley test involves a sequences of eigenvalues of a certain integral operator induced by the covariance kernel of a Gaussian process. We solve the associated integral equation and present the corresponding eigenvalues.

Keywords Test for normality · Integral operator · Fredholm determinant

Mathematics Subject Classification Primary 62F03 ; Secondary 65C60 · 65R20

1 Introduction

Let $X, X_1, X_2 \dots$ be a sequence of independent and identically distributed (i.i.d) random variables with unknown distribution. To test the hypothesis H_0 that the distribution of X is some unspecified normal distribution, there is a myriad of testing procedures, among which the tests of Shapiro–Wilk (SW) and Anderson–Darling (AD) deserve special mention, see, e.g., the monographs of D’Agostino and Stephens (1996) and Thode (2002). There is, however, a further test which was proposed by Epps and Pulley (1983). This test, which is based on the empirical characteristic function, comes as a serious competitor to SW and AD, as shown in simulation studies (see, e.g., Bar-

✉ Bruno Ebner
Bruno.Ebner@kit.edu

Norbert Henze
Norbert.Henze@kit.edu

¹ Institute of Stochastics, Karlsruhe Institute of Technology (KIT), Englerstr. 2, 76133 Karlsruhe, Germany

inghaus et al. 1989; Betsch and Ebner 2020). Baringhaus and Henze (1988) extended the approach of Epps and Pulley to test for normality in any dimension. By now, the BHEP-test (an acronym coined by Csörgö (1989) after earlier developers of the idea) is known to be an affine-invariant and universally consistent test of normality in any dimension, and limit distributions of the test statistic have been obtained under H_0 as well as under fixed and contiguous alternatives to normality (see the review article Ebner and Henze 2020). In this paper, we revisit the limit null distribution of the Epps–Pulley test statistic in the univariate case. The test statistic involves a positive tuning parameter β , and, based on $X_1, \dots, X_n$, is denoted by $T_{n,\beta}$. It is given by

$$T_{n,\beta} = n \int_{-\infty}^{\infty} \left| \psi_n(t) - e^{-t^2/2} \right|^2 \varphi_{\beta}(t) dt,$$

where $\psi_n(t) = n^{-1} \sum_{j=1}^n \exp(itY_{n,j})$ is the empirical characteristic function of the scaled residuals $Y_{n,1}, \dots, Y_{n,n}$. Here, $Y_{n,j} = S_n^{-1}(X_j - \bar{X}_n)$, $j = 1, \dots, n$, and $\bar{X}_n = n^{-1} \sum_{j=1}^n X_j$, $S_n^2 = n^{-1} \sum_{j=1}^n (X_j - \bar{X}_n)^2$ denote the sample mean and the sample variance of $X_1, \dots, X_n$, respectively. Moreover,

$$\varphi_{\beta}(t) = \frac{1}{\beta\sqrt{2\pi}} \exp\left(-\frac{t^2}{2\beta^2}\right), \quad t \in \mathbb{R},$$

is the density of the centred normal distribution with variance β^2 . A closed-form expression of $T_{n,\beta}$ that is amenable to computational purposes is

$$\begin{aligned} T_{n,\beta} &= \frac{1}{n} \sum_{j,k=1}^n \exp\left(-\frac{\beta^2}{2}(Y_{n,j} - Y_{n,k})^2\right) \\ &\quad - \frac{2}{\sqrt{1+\beta^2}} \sum_{j=1}^n \exp\left(-\frac{\beta^2 Y_{n,j}^2}{2(1+\beta^2)}\right) + \frac{n}{\sqrt{1+2\beta^2}}. \end{aligned}$$

The limit null distribution of $T_{n,\beta}$, as $n \rightarrow \infty$, is that of

$$T_{\infty} := \int_{-\infty}^{\infty} Z^2(t) \varphi_{\beta}(t) dt.$$

Here, $Z(\cdot)$ is a centred Gaussian element of the Hilbert space $L^2 = L^2(\mathbb{R}, \mathcal{B}, \varphi_{\beta}(t)dt)$ (of equivalence classes) of Borel-measurable real-valued functions that are square-integrable with respect to $\varphi_{\beta}(t)dt$, and the covariance function of $Z(\cdot)$ is given by

$$K(s, t) = \exp\left(-\frac{(s-t)^2}{2}\right) - \left(1 + st + \frac{(st)^2}{2}\right) \exp\left(-\frac{s^2}{2} - \frac{t^2}{2}\right), \quad s, t \in \mathbb{R} \quad (1.1)$$

(see Henze and Wagner 1997). The kernel K is the starting point of this paper. Writing $\sim$ for equality in distribution, it is well-known that

$$T_{\infty} \sim \sum_{j=0}^{\infty} \lambda_j N_j^2,$$

where $\lambda_0, \lambda_1, \dots$ is the sequence of nonzero eigenvalues associated with the integral operator $\mathbb{A} : L^2 \rightarrow L^2$ defined by

$$(\mathbb{A}f)(s) := \int_{-\infty}^{\infty} K(s, t) f(t) \varphi_{\beta}(t) dt,$$

and $N_0, N_1, \dots$ is a sequence of i.i.d. standard normal random variables. In the next section, we obtain the eigenvalues of $\mathbb{A}$ by numerical methods. In Sect. 3 the sum of powers of the largest eigenvalues is compared to normalized cumulants. The difference should be close to 0 if the eigenvalues have been computed correctly. Section 4 demonstrates that the results can be applied to fit a Pearson system of distributions, and that the fit is reasonable to approximate critical values of the Epps-Pulley test. The article ends by some concluding remarks. Finally, Appendix A extends the results to the cases in which no parameters, only the mean and only the variance, are estimated.

2 Solution of a Fredholm integral equation

To obtain the values $\lambda_0, \lambda_1, \lambda_2, \dots$, that determine the distribution of T_{∞} , one has to solve the integral equation

$$\int_{-\infty}^{\infty} K(s, t) f(t) \varphi_{\beta}(t) dt = \lambda f(s), \quad s \in \mathbb{R}.$$

In general, this task is considered a hard problem, and solutions for kernels associated with testing problems involving composite hypotheses are very sparse, see Stephens (1976, 1977) for the classical tests of normality and exponentiality that are based on the empirical distribution function. In what follows, we use a result of Zhu et al. (1997) to obtain the eigenvalues of $\mathbb{A}$ by a stable numerical method. To this end, let

$$K_0(s, t) = \exp\left(-\frac{(s-t)^2}{2}\right), \quad s, t \in \mathbb{R},$$

and

$$\begin{aligned} \phi_1(s) &= \frac{s^2}{\sqrt{2}} \exp\left(-\frac{s^2}{2}\right), \\ \phi_2(s) &= s \exp\left(-\frac{s^2}{2}\right), \end{aligned}$$

$$\phi_3(s) = \exp\left(-\frac{s^2}{2}\right), \quad s \in \mathbb{R}.$$

Notice that

$$K(s, t) = K_0(s, t) - \sum_{j=1}^3 \phi_j(s)\phi_j(t), \quad s, t \in \mathbb{R}. \quad (2.1)$$

The first step is to solve the eigenvalue problem for the covariance kernel K_0 . The associated eigenvalue problem, which leads to the kernel K_0 , was solved in the context of machine learning in Chapter 4 of Zhu et al. (1997). Here, we use the formulation given in Rasmussen and Williams (2008), Sect. 4.3.1. The eigenvalues of K_0 are given by

$$\lambda_k^{(0)} = \sqrt{\frac{2}{\sqrt{1+4\beta^2} + 2\beta^2 + 1}} \left(\frac{2\beta^2}{\sqrt{1+4\beta^2} + 2\beta^2 + 1} \right)^k, \quad k = 0, 1, 2, \dots,$$

with corresponding normalized eigenfunctions

$$\psi_k(x) = h_k \exp\left(-\left(\frac{\sqrt{\beta^{-2}+4}}{4\beta} - (4\beta)^{-1}\right)x^2\right) H_k\left(\left(\beta^{-4} + 4\beta^{-2}\right)^{1/4} x/\sqrt{2}\right),$$

$$k = 0, 1, 2, \dots$$

(see also the errata to Rasmussen and Williams (2008) on the books homepage). Here, $h_k^{-2} = (4\beta^2 + 1)^{-1/4} 2^k k!$, and $H_k(x) = (-1)^k \exp(x^2) \frac{d^k}{dx^k} \exp(-x^2)$ is the k th order Hermite polynomial.

Remark 2.1 Note that for the special case $\beta = 1$ the eigenvalues $\lambda_k^{(0)}$ coincides with the formula given in (6) of Baringhaus (1996). In that article, the limit distribution of a modified statistic $T_{n,\beta}^{(0)}$, which originates from $T_{n,\beta}$ by replacing $\psi_n(t)$ with $\psi_n^{(0)}(t) = n^{-1} \sum_{j=1}^n \exp(itX_j)$, i.e., the problem is to test for standard normality and thus no estimation of parameters is involved, is analyzed, cf. Appendix A. The corresponding covariance kernel is $K_1(s, t) = K_0(s, t) - \phi_3(s)\phi_3(t)$, $s, t \in \mathbb{R}$, and explicit formulae for the eigenvalues and eigenfunctions are given, see Baringhaus (1996), p. 3878.

To solve the eigenvalue problem of $\mathbb{A}$ figuring in (1.1), we adapt the methodology in Stephens (1976). Define

$$a_{j,k} = \int_{-\infty}^{\infty} \psi_j(x)\phi_k(x)\varphi_\beta(x) dx,$$

$$S_k(\lambda) = 1 + \sum_{j=0}^{\infty} \frac{a_{j,k}^2}{1/\lambda - \lambda_j^{(0)}},$$

$$S_{k,\ell}(\lambda) = \sum_{j=0}^{\infty} \frac{a_{j,k}a_{j,\ell}}{1/\lambda - \lambda_j^{(0)}}, \quad \lambda > 0.$$

With this notation, we can formulate our main result.

Theorem 2.2 *The eigenvalues of $\mathbb{A}$ are the reciprocals of the solutions $\lambda > 0$ of the equation*

$$D(\lambda) = d(\lambda)S_2(\lambda) \left(S_1(\lambda)S_3(\lambda) - S_{1,3}^2(\lambda) \right) = 0, \quad (2.2)$$

where $d(\lambda) = \prod_{k=0}^{\infty} (1/\lambda - \lambda_k^{(0)})$ is the Fredholm determinant connected to the eigenvalue problem of K_0 . Moreover, none of the reciprocals of the eigenvalues $\lambda_k^{(0)}$ of K_0 solve equation (2.2).

Proof Since $a_{j,1}a_{j,2} = a_{j,2}a_{j,3} = 0$ holds for all $j = 0, 1, 2, \dots$, we use Theorem 2.2 of Sukhatme (1972) to see that the Fredholm determinant for the eigenvalue problem takes the form

$$D(\lambda) = d(\lambda) \det \begin{pmatrix} S_1(\lambda) & 0 & S_{1,3}(\lambda) \\ 0 & S_2(\lambda) & 0 \\ S_{1,3}(\lambda) & 0 & S_3(\lambda) \end{pmatrix} = d(\lambda)S_2(\lambda) \left(S_1(\lambda)S_3(\lambda) - S_{1,3}^2(\lambda) \right).$$

Hence, the reciprocals of the roots of $D(\lambda)$ are the eigenvalues of $\mathbb{A}$. By direct calculation, it follows that $a_{j,2} = 0$ if j is even, and we have $a_{j,1} = a_{j,3} = 0$ if j is odd. Consequently, none of the reciprocals of the eigenvalues $\lambda_k^{(0)}$ is a root of $D(\lambda)$ and thus a solution of the eigenvalue problem associated with the kernel K . $\square$

According to Theorem 2.2, the eigenvalues of $\mathbb{A}$ are the roots of $S_2(\lambda)$ and of $S_1(\lambda)S_3(\lambda) - S_{1,3}^2(\lambda)$. The reciprocals of the roots have been obtained numerically, and the largest twenty eigenvalues are displayed in Table 1 for different values of β . Note that, since these values tend to be very small, the reciprocal approach used here leads to numerically stable procedures to find the roots of the Fredholm determinant.

3 Accuracy of the numerical solutions

The accuracy of the values presented in Table 1 may be judged by a comparison with results of Henze (1990). That paper gives the first four cumulants of the distribution of T_{∞} in the special case $\beta = 1$. The m -th cumulant of T_{∞} is

$$\kappa_m(\beta) = 2^{m-1}(m-1)! \sum_{j=1}^{\infty} \lambda_j^m = 2^{m-1}(m-1)! \int_{-\infty}^{\infty} K_m(x, x) \varphi_{\beta}(x) dx, \quad m \geq 1,$$

Table 1 Eigenvalues of $\mathbb{A}$ for different tuning parameters β , here $E-j$ stands for 10^{-j}

$\lambda_j \backslash \beta$	0.25	0.5	1	2	3
λ_0	4.07235E-04	1.01443E-02	7.42748E-02	1.54164E-01	1.59960E-01
λ_1	3.96229E-05	2.98027E-03	4.48104E-02	1.29257E-01	1.45877E-01
λ_2	8.87536E-07	2.13968E-04	8.41907E-03	4.99665E-02	7.56703E-02
λ_3	7.41169E-08	5.45396E-05	4.58684E-03	3.98239E-02	6.69664E-02
λ_4	2.36367E-09	5.42325E-06	1.07998E-03	1.70946E-02	3.68745E-02
λ_5	1.81032E-10	1.27337E-06	5.51939E-04	1.31547E-02	3.19372E-02
λ_6	6.72990E-12	1.46554E-07	1.45739E-04	6.00412E-03	1.82413E-02
λ_7	4.87430E-13	3.26023E-08	7.12110E-05	4.49725E-03	1.55292E-02
λ_8	1.97617E-14	4.08130E-09	2.01821E-05	2.14175E-03	9.10854E-03
λ_9	1.37638E-15	8.73898E-10	9.53839E-06	1.56980E-03	7.64324E-03
λ_{10}	5.90134E-17	1.15555E-10	2.83684E-06	7.71666E-04	4.57714E-03
λ_{11}	3.99302E-18	2.40495E-11	1.30684E-06	5.55566E-04	3.79317E-03
λ_{12}	1.78040E-19	3.30498E-12	4.02503E-07	2.79917E-04	2.31011E-03
λ_{13}	1.17813E-20	6.72882E-13	1.81702E-07	1.98509E-04	1.89295E-03
λ_{14}	5.40732E-22	9.51525E-14	5.74665E-08	1.01995E-04	1.16876E-03
λ_{15}	3.51545E-23	1.90367E-14	2.55205E-08	7.13642E-05	9.46891E-04
λ_{16}	1.64986E-24	2.75201E-15	8.24056E-09	3.71939E-05	5.90055E-04
λ_{17}	1.05731E-25	5.42770E-16	3.61045E-09	2.56128E-05	4.70590E-04
λ_{18}	6.60890E-28	1.07411E-17	1.18543E-09	2.86884E-06	8.19769E-05
λ_{19}	7.53533E-29	3.77859E-18	5.13526E-10	3.67008E-06	1.22699E-05

where $K_1(x, y) := K(x, y)$ and

$$K_m(x, y) = \int_{-\infty}^{\infty} K_{m-1}(x, z) K(z, y) \varphi_{\beta}(z) dz$$

for $m \geq 2$ (see e.g., Chapter 5 of Shorack and Wellner (1986)). We have

$$\kappa_1(1) = \mathbb{E}(T_{\infty}) = \sum_{j=1}^{\infty} \lambda_j = 1 - \frac{\sqrt{3}}{2} = 0.133974596 \dots, \quad (3.1)$$

$$\kappa_2(1) = \mathbb{V}(T_{\infty}) = 2 \sum_{j=1}^{\infty} \lambda_j^2 = \frac{2\sqrt{5}}{5} + \frac{5}{6} - \frac{155\sqrt{2}}{128} = 0.015236301 \dots$$

and thus

$$\sum_{j=1}^{\infty} \lambda_j^2 = \frac{1}{\sqrt{5}} + \frac{5}{12} - \frac{155}{128\sqrt{2}} = 0.0076181509 \dots$$

Furthermore,

$$\begin{aligned}\kappa_3(1) &= \mathbb{E}(T_\infty - \kappa_1(1))^3 = 8 \sum_{j=1}^{\infty} \lambda_j^3 = 0.00400343 \dots, \\ \kappa_4(1) &= \mathbb{E}(T_\infty - \kappa_1(1))^4 = 48 \sum_{j=1}^{\infty} \lambda_j^4 = 0.001654655 \dots\end{aligned}\quad (3.2)$$

and thus

$$\sum_{j=1}^{\infty} \lambda_j^3 = 0.0005004285291 \dots \quad \text{and} \quad \sum_{j=1}^{\infty} \lambda_j^4 = 0.00003447197917 \dots$$

From Table 2, we see that the corresponding sums of the first 20 numerical values of the eigenvalues (as well of their squares and cubes) agree approximately with the values figuring in (3.1) and (3.2), respectively, in most cases up to five significant digits.

The results of Henze (1990) have been partially generalized in Henze and Wagner (1997), Theorem 2.3, for the first three cumulants and a fixed tuning parameter β , and they thus lead to general formulae in the univariate case. For the sake of completeness, we restate the formulae of the first two cumulants here. For the first cumulant, we have

$$\kappa_1(\beta) = 1 - (2\beta^2 + 1)^{-1/2} \left[1 + \frac{\beta^2}{2\beta^2 + 1} + \frac{3\beta^4}{(2\beta^2 + 1)^2} \right],$$

and the second cumulant is

$$\begin{aligned}\kappa_2(\beta) &= \frac{2}{\sqrt{1 + 4\beta^2}} + \frac{2}{1 + 2\beta^2} \left[1 + \frac{2\beta^4}{(1 + 2\beta^2)^2} + \frac{9\beta^8}{4(1 + 2\beta^2)^4} \right] \\ &\quad - \frac{4}{\sqrt{1 + 4\beta^2 + 3\beta^4}} \left[1 + \frac{3\beta^4}{2(1 + 4\beta^2 + 3\beta^4)} + \frac{3\beta^8}{2(1 + 4\beta^2 + 3\beta^4)^2} \right].\end{aligned}$$

The formula for the third cumulant is found in Henze and Wagner (1997), Theorem 2.3, for the case $d = 1$. Table 2 exhibits the normalized cumulants, together with the corresponding sums of the first 20 eigenvalues taken from Table 1. We stress that by now no formula for the fourth cumulant is known in the literature for general tuning parameter β .

4 Pearson system fit for approximation of critical values

The first four cumulants can directly be used in packages that implement the Pearson system of distributions (see Sect. 4.1 of Johnson et al. (1994)). In the statistical computing language R (see R Core Team 2021), we use the package `PearsonDS`

Table 2 Sums over different powers of the first 20 eigenvalues and corresponding theoretical cumulants for different values of β . The entry denoted by * could not be computed due to numerical instabilities

β	0.25	0.5	1	2	3
$\sum_{j=0}^{19} \lambda_j$	4.47822E-04	1.34000E-02	1.33975E-01	4.19722E-01	5.83761E-01
$\sum_{j=0}^{19} \lambda_j^2$	1.67411E-07	1.11838E-04	7.61814E-03	4.50863E-02	6.02196E-02
$\sum_{j=0}^{19} \lambda_j^3$	6.75980E-11	1.04392E-06	5.00428E-04	6.01902E-03	8.02464E-03
$\sum_{j=0}^{19} \lambda_j^4$	2.75053E-14	1.06687E-08	3.44719E-05	8.52858E-04	1.16350E-03
$\kappa_1(\beta)$	4.47822E-04	1.34000E-02	1.33975E-01	4.19753E-01	5.84700E-01
$\kappa_2(\beta)/2$	1.66500E-07	1.11838E-04	7.61814E-03	4.50863E-02	6.02202E-02
$\kappa_3(\beta)/8$	*	1.07115E-06	5.00429E-04	6.01903E-03	8.02468E-03

(see Becker and Klößner 2017) to approximate critical values of the Epps–Pulley test statistic. The Epps–Pulley test is implemented in the R-Package `mmt` (see Butsch and Ebner 2020) by using the function `BHEP`. Table 3 shows simulated empirical critical values of the Epps–Pulley statistic for sample sizes $n \in \{10, 25, 50, 100, 200\}$ and levels of significance $\alpha \in \{0.1, 0.05, 0.01\}$. For each combination of n and β , the entries corresponding to different values of α are based on 10^6 replications under the null hypothesis. Each entry in a row named ' ∞ ' is the calculated $(1 - \alpha)$ -quantile of the fitted Pearson system using the cumulants given in Table 2. We conclude that, for larger sample sizes, the simulated critical values are close to the approximated counterparts of the Pearson system. Moreover, we have corroborated the results of Henze (1990) for the special case $\beta = 1$, and we have extended these results for general $\beta > 0$.

5 Conclusions

We have solved the eigenvalue problem of the integral operator associated with the covariance kernel K of the limiting Gaussian process that occurs in the limit null distribution of the Epps–Pulley test statistic. In view of a comparison with the first three known cumulants from the literature, Table 2 shows that the eigenvalues obtained by numerical methods are very close to the corresponding theoretical values. In Sect. 5 of Ebner and Henze (2021), the authors present a Monte Carlo based approximation method to find stochastic approximations of the eigenvalues. A comparison of Table 1 and Table 1 of Ebner and Henze (2021) reveals that there are some significant differences for some values of β , which can be explained by the approximation of the eigenvalues by a Monte Carlo method in Ebner and Henze (2021). This observation is of particular interest, since the largest eigenvalue is used in the derivations of approximate Bahadur efficiencies. Recent results concerning this topic for the Epps–Pulley test are presented in Ebner and Henze (2021) and, for other normality tests based on the empirical distribution function, in Milošević et al. (2021).

We point out the difficulties encountered if one tries to generalize our findings to the multivariate case, i.e. to obtain the eigenvalues associated with the limit null

Table 3 Empirical critical values (simulated with 10^6 replications) and approximated critical values by the Pearson system (denoted by ' ∞ ') for different levels of significance α

α	$n \setminus \beta$	0.25	0.5	1	2	3
0.1	10	7.28E-04	0.0245	0.277	0.817	1.03
	25	9.58E-04	0.0289	0.288	0.814	1.04
	50	1.05E-03	0.0304	0.289	0.811	1.04
	100	1.10E-03	0.0310	0.290	0.812	1.04
	200	1.13E-03	0.0314	0.291	0.811	1.04
	∞	1.14E-03	0.0319	0.292	0.812	1.04
0.05	10	1.06E-03	0.0343	0.355	0.99	1.22
	25	1.39E-03	0.0403	0.371	1.00	1.25
	50	1.51E-03	0.0420	0.374	1.01	1.25
	100	1.57E-03	0.0427	0.376	1.01	1.25
	200	1.60E-03	0.0429	0.378	1.01	1.25
	∞	1.61E-03	0.0429	0.379	1.01	1.25
0.01	10	1.91E-03	0.0589	0.543	1.39	1.65
	25	2.58E-03	0.0696	0.570	1.44	1.72
	50	2.75E-03	0.0711	0.575	1.45	1.74
	100	2.78E-03	0.0720	0.581	1.46	1.75
	200	2.78E-03	0.0717	0.585	1.46	1.75
	∞	2.74E-03	0.0700	0.585	1.46	1.74

distribution of the BHEP test of multivariate normality, see (Baringhaus and Henze 1988; Henze and Wagner 1997; Henze and Zirkler 1990). The d -variate analog to the covariance kernel K in (1.1) is given in Theorem 2.1 of Henze and Wagner (1997), namely, writing $\|\cdot\|$ for the Euclidean norm and $^\top$ for the transpose of vectors, we have

$$K(s, t) = \exp\left(-\frac{\|s - t\|^2}{2}\right) - \left\{1 + s^\top t + \frac{(s^\top t)^2}{2}\right\} \exp\left(-\frac{\|s\|^2 + \|t\|^2}{2}\right),$$

$$s, t \in \mathbb{R}^d. \quad (5.1)$$

The first step is to derive explicit expressions for eigenvalues w.r.t. the kernel

$$K_0(s, t) = \exp\left(-\|s - t\|^2/2\right), \quad s, t \in \mathbb{R}^d$$

(for a starting point, see Baringhaus 1996). The second step is to find the corresponding multivariate representation of (2.1), which seems to be non-standard, since the quadratic summand $(s^\top t)^2$ in (5.1) does not factorize easily. Both problems have to be solved in order to successfully apply the method presented in Sect. 2.

Finally, it is an interesting question whether the results may be extended to other recent tests of normality associated with the empirical characteristic function, such as Ebner (2020), or to other empirical integral transformations, such as the moment generating function, see Henze and Koch (2020), or for multivariate versions see Ebner

et al. (2021) as well as Henze and Visagie (2020). In each of these papers an explicit formula of the covariance kernel under the null hypothesis is derived, but it is again unclear how to find explicit expressions for the eigenvalues of the reduced kernel formula. Hence we leave these problems open for future work.

Acknowledgements The authors thank two anonymous referees for helpful comments that improved the paper.

Funding Open Access funding enabled and organized by Projekt DEAL.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

A Approximation of eigenvalues in case of testing under partially known parameters

In the spirit of the work of Stephens (1976), we provide the approximation of the eigenvalues for the following three related cases:

1. both parameters known: The test statistic is applied to $Y_j = (X_j - \mu)/\sigma$, where μ and σ are the parameters under the null hypothesis. The covariance kernel then reduces to

$$K_1(s, t) = K_0(s, t) - \phi_3(s)\phi_3(t), \quad s, t \in \mathbb{R},$$

and the Fredholm determinant is $D(\lambda) = d(\lambda)S_3(\lambda)$. The zeros of $D(\lambda)$ (note that some zeros of $d(\lambda)$ are not zeros of $D(\lambda)$) provide the eigenvalues in Table 4.

2. the mean is unknown, but the variance is known: The test statistic is applied to $Y_{n,j} = (X_j - \bar{X}_n)/\sigma$ and the covariance kernel reduces to

$$K_2(s, t) = K_0(s, t) - \sum_{j=2}^3 \phi_j(s)\phi_j(t), \quad s, t \in \mathbb{R}.$$

Here, the Fredholm determinant takes the form $D(\lambda) = d(\lambda)S_2(\lambda)S_3(\lambda)$. The zeros of $D(\lambda)$ provide the eigenvalues in Table 5. Note that none of the zeros of $d(\lambda)$ are zeros of $D(\lambda)$.

3. the mean is known, but the variance is unknown: The test statistic is applied to $Y_{n,j} = S_n^{-1}(X_j - \mu)$. The covariance kernel reduces to

$$K_3(s, t) = K_0(s, t) - \phi_1(s)\phi_1(t) - \phi_3(s)\phi_3(t), \quad s, t \in \mathbb{R},$$

Table 4 Eigenvalues of $\mathbb{A}$ applied to kernel K_1 for different tuning parameters β , here $E-j$ stands for 10^{-j}

$\lambda_j \backslash \beta$	0.25	0.5	1	2	3
λ_0	5.26225E-02	1.42136E-01	2.36068E-01	2.37985E-01	2.02640E-01
λ_1	4.39314E-03	3.61356E-02	1.27399E-01	1.87334E-01	1.77471E-01
λ_2	1.63425E-04	4.18408E-03	3.44419E-02	8.84416E-02	1.04356E-01
λ_3	1.13824E-05	8.95872E-04	1.62590E-02	6.45254E-02	8.70835E-02
λ_4	5.07537E-07	1.23168E-04	5.02500E-03	3.28673E-02	5.37419E-02
λ_5	3.29555E-08	2.46420E-05	2.23123E-03	2.29768E-02	4.35288E-02
λ_6	1.57622E-09	3.62572E-06	7.33137E-04	1.22143E-02	2.76763E-02
λ_7	9.07802E-11	6.99678E-07	3.14550E-04	8.30900E-03	2.19590E-02
λ_8	4.89513E-12	1.06731E-07	1.06963E-04	4.53918E-03	1.42528E-02
λ_9	2.72830E-13	2.01376E-08	4.49058E-05	3.03064E-03	1.11393E-02
λ_{10}	8.47202E-16	5.81399E-10	6.45455E-06	1.11118E-03	5.67138E-03
λ_{11}	2.63109E-18	1.62478E-11	9.31517E-07	4.08788E-04	2.89477E-03
λ_{12}	8.17116E-21	4.67570E-13	1.34787E-07	1.50730E-04	1.48022E-03
λ_{13}	2.53765E-23	1.37512E-14	1.95378E-08	5.56673E-05	7.57908E-04
λ_{14}	7.88098E-26	4.04788E-16	2.83563E-09	2.05829E-05	3.88457E-04
$\sum_{j=0}^{14} \lambda_j$	5.71910E-02	1.83503E-01	4.22631E-01	6.63970E-01	7.55041E-01
$\sum_{j=0}^{14} \lambda_j^2$	2.78845E-03	2.15266E-02	7.34401E-02	1.05574E-01	9.74343E-02
$\sum_{j=0}^{14} \lambda_j^3$	1.45803E-04	2.91876E-03	1.52687E-02	2.10636E-02	1.59814E-02
$\sum_{j=0}^{14} \lambda_j^4$	7.66842E-06	4.09848E-04	3.37053E-03	4.51932E-03	2.86707E-03
$\kappa_1(\beta)$	5.71910E-02	1.8350E-01	4.22649E-01	6.66667E-01	7.70584E-01
$\kappa_2(\beta)/2$	2.78845E-03	2.1527E-02	7.34401E-02	1.05577E-01	9.75077E-02
$\kappa_3(\beta)/8$	1.45802E-04	2.9188E-03	1.52687E-02	2.10636E-02	1.59818E-02
$\kappa_4(\beta)/48$	7.66638E-06	4.0985E-04	3.37053E-03	4.51932E-03	2.86707E-03

and the Fredholm determinant is $D(\lambda) = d(\lambda)(S_1(\lambda)S_3(\lambda) - S_{1,3}^2(\lambda))$. The zeros of $D(\lambda)$ provide the eigenvalues in Table 6. Note that some zeros of $d(\lambda)$ are also zeros of $D(\lambda)$.

Note that the sums of powers of the eigenvalues are close to the respective cumulants in all three Tables 4 - 6, which confirms the good approximation of the eigenvalues.

Table 5 Eigenvalues of $\mathbb{A}$ applied to kernel K_2 for different tuning parameters β , here E-j stands for 10^{-j} . The entry denoted by * could not be computed due to numerical instabilities

$\lambda_j \backslash \beta$	0.25	0.5	1	2	3
λ_0	4.39314E-03	3.61356E-02	1.27399E-01	1.87334E-01	1.77471E-01
λ_1	4.07235E-04	1.01443E-02	7.42748E-02	1.54164E-01	1.59960E-01
λ_2	1.13824E-05	8.95872E-04	1.62590E-02	6.45254E-02	8.70835E-02
λ_3	8.87536E-07	2.13968E-04	8.41907E-03	4.99665E-02	7.56703E-02
λ_4	3.29965E-08	2.46420E-05	2.23123E-03	2.29768E-02	4.35288E-02
λ_5	2.36367E-09	5.42325E-06	1.07998E-03	1.70946E-02	3.68745E-02
λ_6	9.88175E-11	6.99683E-07	3.14550E-04	8.30900E-03	2.19590E-02
λ_7	6.72990E-12	1.46554E-07	1.45739E-04	6.00412E-03	1.82413E-02
λ_8	3.00073E-13	2.01411E-08	4.49058E-05	3.03064E-03	1.11393E-02
λ_9	1.97617E-14	4.08130E-09	2.01821E-05	2.14175E-03	9.10854E-03
λ_{10}	9.17796E-16	5.83944E-10	6.45455E-06	1.11118E-03	5.67138E-03
λ_{11}	5.90134E-17	1.15555E-10	2.83684E-06	7.71666E-04	4.57714E-03
λ_{12}	2.81901E-18	1.70012E-11	9.31517E-07	4.08788E-04	2.89477E-03
λ_{13}	1.78040E-19	3.30498E-12	4.02503E-07	2.79917E-04	2.31011E-03
λ_{14}	8.68183E-21	4.96305E-13	1.34787E-07	1.50730E-04	1.48022E-03
$\sum_{j=0}^{14} \lambda_j$	4.81268E-03	4.74207E-02	2.30200E-01	5.18487E-01	6.60957E-01
$\sum_{j=0}^{14} \lambda_j^2$	1.94656E-05	1.40954E-03	2.20888E-02	6.64619E-02	7.47405E-02
$\sum_{j=0}^{14} \lambda_j^3$	8.48536E-08	4.82298E-05	2.48243E-03	1.06496E-02	1.09279E-02
$\sum_{j=0}^{14} \lambda_j^4$	3.72505E-10	1.71565E-06	2.93941E-04	1.82038E-03	1.74279E-03
$\kappa_1(\beta)$	4.81268E-03	4.74207E-02	2.30200E-01	5.18519E-01	6.61914E-01
$\kappa_2(\beta)/2$	1.94653E-05	1.40954E-03	2.20888E-02	6.64619E-02	7.47408E-02
$\kappa_3(\beta)/8$	8.46157E-08	4.82310E-05	2.48243E-03	1.06496E-02	1.09279E-02
$\kappa_4(\beta)/48$	*	1.71569E-06	2.93941E-04	1.82038E-03	1.74279E-03

Table 6 Eigenvalues of $\mathbb{A}$ applied to kernel K_3 for different tuning parameters β , here $E-j$ stands for 10^{-j}

$\lambda_j \backslash \beta$	0.25	0.5	1	2	3
λ_0	5.26225E-02	1.42136E-01	2.36068E-01	2.37985E-01	2.02640E-01
λ_1	1.63425E-04	4.18408E-03	4.48104E-02	1.29257E-01	1.45877E-01
λ_2	3.96229E-05	2.98027E-03	3.44419E-02	8.84416E-02	1.04356E-01
λ_3	5.07537E-07	1.23168E-04	5.02500E-03	3.98239E-02	6.69664E-02
λ_4	7.41169E-08	5.45396E-05	4.58684E-03	3.28673E-02	5.37419E-02
λ_5	1.57622E-09	3.62572E-06	7.33137E-04	1.31547E-02	3.19372E-02
λ_6	1.81032E-10	1.27337E-06	5.51939E-04	1.22143E-02	2.76763E-02
λ_7	4.89513E-12	1.06731E-07	1.06963E-04	4.53918E-03	1.55292E-02
λ_8	4.87430E-13	3.26023E-08	7.12110E-05	4.49725E-03	1.42528E-02
λ_9	1.37638E-15	8.73898E-10	9.53839E-06	1.56980E-03	7.64324E-03
λ_{10}	3.99302E-18	2.40495E-11	1.30684E-06	5.55566E-04	3.79317E-03
λ_{11}	1.17813E-20	6.72882E-13	1.81702E-07	1.98509E-04	1.89295E-03
λ_{12}	3.51545E-23	1.90367E-14	2.55195E-08	7.13642E-05	9.46891E-04
λ_{13}	1.05731E-25	5.42770E-16	3.60561E-09	2.56128E-05	4.70590E-04
λ_{14}	7.53533E-29	3.77859E-18	1.43728E-10	3.67008E-06	1.22699E-04
$\sum_{j=0}^{14} \lambda_j$	5.28261E-02	1.49483E-01	3.26406E-01	5.65205E-01	6.77846E-01
$\sum_{j=0}^{14} \lambda_j^2$	2.76915E-03	2.02289E-02	5.89695E-02	8.41984E-02	8.29135E-02
$\sum_{j=0}^{14} \lambda_j^3$	1.45718E-04	2.87160E-03	1.32867E-02	1.64331E-02	1.30781E-02
$\sum_{j=0}^{14} \lambda_j^4$	7.66805E-06	4.08143E-04	3.11106E-03	3.55180E-03	2.28777E-03
$\kappa_1(\beta)$	5.28261E-02	1.49483E-01	3.26425E-01	5.67901E-01	6.93371E-01
$\kappa_2(\beta)/2$	2.76915E-03	2.02289E-02	5.89695E-02	8.42017E-02	8.29872E-02
$\kappa_3(\beta)/8$	1.45717E-04	2.87160E-03	1.32867E-03	1.64331E-02	1.30786E-02
$\kappa_4(\beta)/48$	7.66704E-06	4.08145E-04	3.11106E-03	3.55180E-03	2.28779E-03

References

- Baringhaus L (1996) Fibonacci numbers, Lucas numbers and integrals of certain Gaussian processes. *Proc Am Math Soc* 124(12):3875–3884
- Baringhaus L, Danschke R, Henze N (1989) Recent and classical tests for normality - a comparative study. *Communications in Statistics - Simulation and Computation* 18:363–379
- Baringhaus L, Henze N (1988) A consistent test for multivariate normality based on the empirical characteristic function. *Metrika* 35(1):339–348
- Becker, M., Klößner, S.: PearsonDS: Pearson Distribution System. R package version 1.1 (2017)
- Betsch S, Ebner B (2020) Testing normality via a distributional fixed point property in the Stein characterization. *TEST* 29(1):105–138
- Butsch L, Ebner B (2020) mnt: Affine Invariant Tests of Multivariate Normality, R package version 1.3
- Csörgő S (1989) Consistency of some tests for multivariate normality. *Metrika* 36:107–116
- D'Agostino RB, Stephens MA (1986) (eds) Goodness-of-fit techniques. *Statistics: textbooks and monographs*, vol 68. Dekker, New York
- Ebner B (2020) On combining the zero bias transform and the empirical characteristic function to test normality. *ALEA* 18:1029–1045
- Ebner B, Henze N (2020) Tests for multivariate normality—a critical review with emphasis on weighted L^2 -statistics. *TEST* 29(4):845–892

- Ebner B, Henze N (2021) Bahadur efficiencies of the Epps-Pulley test for normality. *Zapiski Nauchnykh Semin* 501:302–314
- Ebner B, Henze N, Strieder D (2021) Testing normality in any dimension by Fourier methods in a multivariate Stein equation. *Can J Stat*. <https://doi.org/10.1002/cjs.11670>
- Epps TW, Pulley LB (1983) A test for normality based on the empirical characteristic function. *Biometrika* 70(3):723–726
- Henze N (1990) An approximation to the limit distribution of the Epps-Pulley test statistic for normality. *Metrika* 37(1):7–18
- Henze N, Koch S (2020) On a test of normality based on the empirical moment generating function. *Stat Pap* 61(1):17–29
- Henze N, Visagie J (2020) Testing for normality in any dimension based on a partial differential equation involving the moment generating function. *Ann Inst Stat Math* 72(5):1109–1136
- Henze N, Wagner T (1997) A new approach to the BHEP tests for multivariate normality. *J Multivar Anal* 62(1):1–23
- Henze N, Zirkler B (1990) A class of invariant consistent tests for multivariate normality. *Commun Stat Theory Methods* 19(10):3595–3617
- Johnson NL, Kotz S, Balakrishnan N (1994) Continuous univariate distributions, vol 1, 2nd edn. Wiley, New York
- Milošević B, Nikitin YY, Obradović M (2021) Bahadur efficiency of edf based normality tests when parameters are estimated. *Zapiski Nauchnykh Semin*, vol 501
- R Core Team (2021) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna
- Rasmussen CE, Williams CKI (2008) Gaussian processes for machine learning. In: Adaptive computation and machine learning. MIT Press, Cambridge
- Shorack GR, Wellner JA (1986) Empirical processes with applications to statistics. Wiley, New York
- Stephens MA (1976) Asymptotic results for goodness-of-fit statistics with unknown parameters. *Ann Stat* 4(2):357–369
- Stephens MA (1977) Goodness of fit for the extreme value distribution. *Biometrika* 64(3):583–588
- Sukhatme S (1972) Fredholm determinant of a positive definite kernel of a special type and its application. *Ann Math Stat* 43(6):1914–1926
- Thode HC (2002) Testing for normality, vol 164. Statistics: textbooks and monographs. Dekker, New York
- Zhu H, Williams CK, Rohwer R, Morciniec M (1997) Gaussian regression and optimal finite dimensional linear models. In: Bishop CM (ed) Neural networks and machine learning. Springer, Berlin

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.