

A Service of

ZBU

Leibniz-Informationszentrum Wirtschaft Leibniz Information Centre for Economics

Huber, Johannes; Meyer-Gohde, Alexander

# Working Paper Iterative refinement of the QZ decomposition for solving linear DSGE models

IMFS Working Paper Series, No. 217

**Provided in Cooperation with:** Institute for Monetary and Financial Stability (IMFS), Goethe University Frankfurt am Main

*Suggested Citation:* Huber, Johannes; Meyer-Gohde, Alexander (2025) : Iterative refinement of the QZ decomposition for solving linear DSGE models, IMFS Working Paper Series, No. 217, Goethe University Frankfurt, Institute for Monetary and Financial Stability (IMFS), Frankfurt a. M.

This Version is available at: https://hdl.handle.net/10419/311846

#### Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

#### Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.



# WWW.ECONSTOR.EU

Institute for Monetary and Financial Stability

# Johannes Huber and Alexander Meyer-Gohde

# Iterative Refinement Of The QZ Decomposition For Solving Linear DSGE Models

Institute for Monetary and Financial Stability GOETHE UNIVERSITY FRANKFURT

WORKING PAPER SERIES NO. 217 (2025)

This Working Paper is issued under the auspices of the Institute for Monetary and Financial Stability (IMFS). Any opinions expressed here are those of the author(s) and not those of the IMFS. Research disseminated by the IMFS may include views on policy, but the IMFS itself takes no institutional policy positions.

The IMFS aims at raising public awareness of the importance of monetary and financial stability. Its main objective is the implementation of the "Project Monetary and Financial Stability" that is supported by the Foundation of Monetary and Financial Stability. The foundation was established on January 1, 2002 by federal law. Its endowment funds come from the sale of 1 DM gold coins in 2001 issued at the occasion of the euro cash introduction in memory of the D-Mark.

The IMFS Working Papers often represent preliminary or incomplete work, circulated to encourage discussion and comment. Citation and use of such a paper should take account of its provisional character.

Institute for Monetary and Financial Stability Goethe University Frankfurt House of Finance Theodor-W.-Adorno-Platz 3 D-60629 Frankfurt am Main www.imfs-frankfurt.de | info@imfs-frankfurt.de

# ITERATIVE REFINEMENT OF THE QZ DECOMPOSITION FOR SOLVING LINEAR DSGE MODELS

#### JOHANNES HUBER AND ALEXANDER MEYER-GOHDE

ABSTRACT. The standard approach to solving linear DSGE models is to apply the QZ method. It is a one-shot algorithm that leaves the researcher with little alternative than to seek a different algorithm should the result be numerically unsatisfactory. We develop an iterative implementation of QZ that delivers the standard result as its first iteration and further refinements at each subsequent iteration. We demonstrate that our algorithm successful corrects for accuracy losses identified in particular cases of a macro finance model and does not erroneously attempt to refine sufficiently accurate solutions. *JEL classification codes*: C61, C63, E17

Keywords: Numerical accuracy; DSGE; Solution methods

# 1. INTRODUCTION

The major computational hurdle in the solution of linear dynamic stochastic general equilibrium (DSGE) models is the solution of the associated matrix quadratic equation - the current standard in the literature is to use a generalized Schur or QZ decomposition (Moler and Stewart, 1973; Golub and van Loan, 2013) to solve this equation.<sup>1</sup> Implementations of this algorithm has been provided to the profession as a one-shot tool, requiring researchers to take it or leave it. This is problematic if the accuracy of the QZ solution is perhaps lacking. We extend the methodology of solving linear DSGE models with QZ or generalized Schur decompositions by providing an iterative version that allows users to improve on a QZ solution with QZ itself.

(Huber) UNIVERSITY OF REGENSBURG, UNIVERSITÄTSSTRASSE 31, 93053 REGENSBURG, GERMANY (Meyer-Gohde) GOETHE-UNIVERSITÄT FRANKFURT AND INSTITUTE FOR MONETARY AND FINANCIAL

STABILITY (IMFS), THEODOR-W.-ADORNO-PLATZ 3, 60629 FRANKFURT AM MAIN, GERMANY

*E-mail addresses*: johannes1.huber@ur.de, meyer-gohde@econ.uni-frankfurt.de.

Date: February 19, 2025.

This research was supported by the DFG through grant nr. 465469938 "Numerical diagnostics and improvements for the solution of linear dynamic macroeconomic models" and grant nr. 465135565 "Models of Imperfect Rationality and Redistribution in the context of Retirement". Meyer-Gohde's Website and Github Repository.

<sup>1</sup>See Dynare (Adjemian, Bastani, Juillard, Mihoubi, Perendia, Ratto, and Villemot, 2011) (henceforth Dynare) and especially Villemot (2011) for Dynare's specific implementation of QZ.

#### HUBER AND MEYER-GOHDE

We provide an eigenvalue preserving transformation that reinterprets the companion linearization of the matrix quadratic as a subspace problem in increments of the solution or solvent. Beginning at the zero matrix, the first iteration reproduces the standard QZ approach. Further iterations refine the solution - while theoretically refinements should not be possible, this is numerically not the case: as Tisseur (2000) shows for the quadratic eigenvalue problem, QZ's backward stability is broken by the block structure that is necessary to maintain the mapping to the underlying matrix quadratic problem. We assess our iterative QZ implementation in a sets of experiments using the model of Jermann (1998) that has numerically unstable parameterization nearby to the standard parameterization. We find that our iterative QZ method successfully terminates when the solution is accurate enough, as measured by Meyer-Gohde's (2022) practical forward error bounds, and overcomes numerically inaccuracies by running a few iterations through improvement increments.

## 2. Iterative QZ

The linear, or first-order, approximation of a DSGE model at the steady state can typically be stated as

$$0 = AE_t[y_{t+1}] + By_t + Cy_{t-1} + D\varepsilon_t$$
(1)

where the coefficient matrices A, B and C are  $n_y \times n_y$  and D is  $n_y \times n_e$ , the endogenous variables  $y_t \in \mathbb{R}^{n_y}$  are a vector of size  $n_y$ ; and the vector of  $n_e$  exogenous shocks are contained in  $\varepsilon_t \in \mathbb{R}^{n_e}$ , where  $n_y$  and  $n_e$  are positive integers  $(n_y, n_e \in \mathbb{N})$  and  $\varepsilon_t$  has a known mean zero distribution.

The solution to the linearized model (1) is a linear solution in the form,

$$y_t = P \ y_{t-1} + Q \ \varepsilon_t \tag{2}$$

a recursive solution for  $y_t$  as a function of its own past,  $y_{t-1}$ , and shocks,  $\varepsilon_t$ .

Using our linear solution (2), it yields through substitution into (1) - and recognizing that the expectation  $E_t[\varepsilon_{t+1}] = 0$  is known - the following two equations

$$0 = AP^{2} + BP + C, \quad 0 = (AP + B)Q + D \tag{3}$$

With the former solution being quadratic with potentially multiple solutions, a selection criteria has to be used and generally a unique (semi) stable solution P is sought by the literature, that is a P with all its eigenvalues inside the open unit circle. As Lan and

Meyer-Gohde (2014) prove the latter can be uniquely solved for Q if such a P can be found, our focus will be the former, the unilateral quadratic matrix equations (UQME).

# 2.1. Standard QZ

To enable our interative QZ approach, we will first express the UQME in (3) as a subspace problem by forming the first companion linearization of the matrix quadratic problem (Hammarling, Munro, and Tisseur, 2013)

$$\mathscr{AX} = \mathscr{BXM} \tag{4}$$

with

$$\mathscr{X} = \begin{bmatrix} I \\ P \end{bmatrix} \qquad \qquad \mathscr{A} = \begin{bmatrix} C & B \\ 0 & I \end{bmatrix}, \qquad \qquad \mathscr{B} = \begin{bmatrix} 0 & -A \\ I & 0 \end{bmatrix}, \qquad \qquad \mathscr{M} = P$$

Clearly, any *P* satisfying (3) is a solution of (4). Further note that the eigenvalues of  $\mathcal{M}$  are a subset of the generalized eigenvalues of the matrix pencil  $\mathcal{A} - \lambda \mathcal{B}$ , i.e.,  $\operatorname{eig}(\mathcal{M}) \subset \operatorname{eig}(\mathcal{A}_0, \mathcal{B}_0)$ .

We will assume the conditions for the existence of the unique solvent P are fulfilled, i.e., Blanchard and Kahn's (1980) celebrated rank and order conditions, and hence that there exist a unique solvent P satisfying the UQME in (3) with eigenvalues on or inside the unit circle and all remaining eigenvalues associated with the UQME (or latent roots of the UQME, see Meyer-Gohde (2022)) are outside the unit circle.

The problem in (4) is numerically an eigenvalue problem and can thus be solved using the QZ or generalized Schur decompostion of Moler and Stewart (1973). We will derive the solution by working directly with the linear algebraic problem instead of the dynamic model as is usually done to keep the notation simple. The decomposition provides unitary Q and Z and upper triangular S and T with  $Q^* \mathscr{B}Z = S$  and  $Q^* \mathscr{A}Z = T$ , where \* indicates the complex conjugation of Q that delivers its inverse by virtue of it being a unitary matrix and where the eigenvalues of the matrix pencil  $P_{\mathscr{B}\mathscr{A}}(z) = \mathscr{B}z - \mathscr{A}$ ,  $\rho(P_{\mathscr{B}\mathscr{A}}) = \rho(P_{ST}) = \{t_{ii}/s_{ii}, \text{ if } s_{ii} \neq 0; \infty, \text{ if } s_{ii} = 0; \emptyset, \text{ if } s_{ii} = 1, \dots, 2n_y\},^2$  can be ordered arbitrarily to form

$$\begin{bmatrix} T_{11} & T_{12} \\ 0 & T_{22} \end{bmatrix} \begin{bmatrix} W^s \\ W^u \end{bmatrix} = \begin{bmatrix} S_{11} & S_{12} \\ 0 & S_{22} \end{bmatrix} \begin{bmatrix} W^s \\ W^u \end{bmatrix} P$$
(5)

<sup>&</sup>lt;sup>2</sup>If both  $s_{ii} = t_{ii} = 0$  for some *i*, then any  $z \in \mathbb{C}$  will satisfy  $|\mathscr{B}z - \mathscr{A}| = 0$ . This is the mundane singularity of King and Watson (1998) that reflects a poorly specified model, e.g., a repeated equation, and the definition in the text can be interpreted to ruling out such misspecifications by limiting the analysis to regular pencils.

where  $Z^* \begin{bmatrix} I & P' \end{bmatrix}' = \begin{bmatrix} W^{s'} & W^{u'} \end{bmatrix}'$ . We assume the decomposition is ordered so that the unstable eigenvalues are in the lower right blocks of S and T (hence  $S_{22}$  and  $T_{22}$ ), wherefor the lower block equation gives

$$T_{22}W^{u} = S_{22}W^{u}P \Rightarrow W^{u} = T_{22}^{-1}S_{22}W^{u}P.$$
 (6)

Consequently, the eigenvalues of  $T_{22}^{-1}S_{22}$  are inside the unit circle and P is a (semi) stable solvent, so that we can iterate

$$W^{u} = T_{22}^{-1} S_{22} W^{u} P = \left(T_{22}^{-1} S_{22}\right)^{2} W^{u} P^{2} = \left(T_{22}^{-1} S_{22}\right)^{j} W^{u} P^{j} \xrightarrow{j \to \infty} 0$$
(7)

for a unit root stable *P*. Using the definition of  $W^u$  we receive

$$0 = W^{u} = Z_{21}^{*} + Z_{22}^{*}P \Rightarrow P = -Z_{22}^{*}{}^{-1}Z_{21}^{*} = Z_{21}Z_{11}^{-1}$$
(8)

The equivalence  $Z_{22}^{*}{}^{-1}Z_{21}^{*} = -Z_{21}Z_{11}^{-1}$  follows from the properties of unitary matrices and  $Z_{21}Z_{11}^{-1} = Q_{11}S_{11}^{-1}T_{11}Q_{11}^{-1}$  from the first block rows of  $\mathscr{A}$  and  $\mathscr{B}$  in (4) and the upper triangularity of S and T. From  $Q_{11}S_{11}^{-1}T_{11}Q_{11}^{-1}$ , it follows that the recursion in P is indeed stable from the ordering of the eigenvalues above, i.e. the eigenvalues of the upper left block of the generalized Schur decomposition,  $det(S_{11}\lambda - T_{11}) = 0$ , are inside the unit circle. So the QZ decomposition applied to our matric pencil will recover the unique (semi) stable solvent P if it exists consistent with our assumption its existence.

## 2.2. Iterative QZ

While the QZ decomposition analytically recovers the unique solvent, numerically it is subject to finite precision. Hence, the decomposition delivers  $\hat{P} = P + \mathcal{O}(\epsilon)$  which differs from the true solution P via numerical errors  $\mathcal{O}(\epsilon)$ . If the solution  $\hat{P}$  is unsatisfactory, that is, if the errors,  $\mathcal{O}(\epsilon)$ , are too large, we would like an alternative to abandoning the QZ solution and having to find an alternate solution. That is, we would like to improve on  $\hat{P}$  using QZ. To this end we will derive an interative QZ approach that takes this approximate solution as the initialization or output from the previous iteration  $P_{k-1}$ . Using this  $P_{k-1}$ , we define a matrix to transform the original, one-shot problem

$$U_{k} \equiv \begin{bmatrix} I & 0 \\ P_{k-1} & I \end{bmatrix}, \qquad \qquad U_{k}^{-1} \equiv \begin{bmatrix} I & 0 \\ -P_{k-1} & I \end{bmatrix}$$
(9)

We note that  $U_k$  is an eigenvalue preserving transform<sup>3</sup> with

$$\mathscr{A}_{k} \equiv \mathscr{A}U_{k}, \qquad \mathscr{B}_{k} \equiv \mathscr{B}U_{k} \tag{10}$$

giving

$$\mathscr{AX} = \mathscr{BXM} \Rightarrow \mathscr{A}U_k U_k^{-1} \mathscr{X} = \mathscr{B}U_k U_k^{-1} \mathscr{XM} \Rightarrow \mathscr{A}_k \hat{\mathscr{X}}_k = \mathscr{B}_k \hat{\mathscr{X}}_k \mathscr{M}.$$
(11)

where  $\hat{\mathscr{X}}_k$  defines the potential improvement that is key to the iterative nature of our approach

$$\hat{\mathscr{X}}_{k} \equiv U_{k}^{-1} \mathscr{X} = \begin{bmatrix} I & 0 \\ -P_{k-1} & I \end{bmatrix} \begin{bmatrix} I \\ P \end{bmatrix} = \begin{bmatrix} I \\ P-P_{k-1} \end{bmatrix}$$
(12)

and thus (with  $\Delta P_k \equiv P - P_{k-1}$ )

$$\mathcal{A}_{k}\begin{bmatrix}I\\\Delta P_{k}\end{bmatrix} = \mathcal{B}_{k}\begin{bmatrix}I\\\Delta P_{k}\end{bmatrix}\mathcal{M}$$
(13)

and

$$\mathscr{A}_{k} = \begin{bmatrix} C + BP_{k-1} & B \\ P_{k-1} & I \end{bmatrix}, \qquad \qquad \mathscr{B}_{k} = \begin{bmatrix} -AP_{k-1} & -A \\ I & 0 \end{bmatrix}$$
(14)

As above with the pencil  $\mathscr{A}$ ,  $\mathscr{B}$  we can apply the QZ decomposition to the pencil defined by  $\mathscr{A}_k$ ,  $\mathscr{B}_k$ . The decomposition provides unitary  $Q_k$  and  $Z_k$  and upper triangular  $S_k$  and  $T_k$  with  $Q_k^* \mathscr{B}_k Z_k = S_k$  and  $Q_k^* \mathscr{A}_k Z_k = T_k$ , where the eigenvalues of the matrix pencil  $P_{\mathscr{B}_k \mathscr{A}_k}(z) = \mathscr{B}_k z - \mathscr{A}_k$ ,  $\rho(P_{\mathscr{B}_k \mathscr{A}_k}) = \rho(P_{S_k T_k}) =$  $\{t_{k,ii}/s_{k,ii}, \text{ if } s_{k,ii} \neq 0; \infty, \text{ if } s_{k,ii} = 0; \emptyset, \text{ if } s_{k,ii} = t_{k,ii} = 0; i = 1, ..., 2n_y\}$ . From the eigenvalue preserving transformation behind  $\mathscr{A}_k$  and  $\mathscr{B}_k$ , follows that  $\rho(P_{\mathscr{B}_k \mathscr{A}_k}) = \rho(P_{\mathscr{B} \mathscr{A}})$ . And analogous to (5) we receive

$$\begin{bmatrix} T_{k,11} & T_{k,12} \\ 0 & T_{k,22} \end{bmatrix} \begin{bmatrix} W_k^s \\ W_k^u \end{bmatrix} = \begin{bmatrix} S_{k,11} & S_{k,12} \\ 0 & S_{k,22} \end{bmatrix} \begin{bmatrix} W_k^s \\ W_k^u \end{bmatrix} P$$
(15)

where  $Z_k^* \begin{bmatrix} I & \Delta P'_k \end{bmatrix}' = \begin{bmatrix} W_k^{s'} & W_k^{u'} \end{bmatrix}'$ . We assume the decomposition is ordered so that the unstable eigenvalues are in the lower right blocks of  $S_k$  and  $T_k$  (hence  $S_{k,22}$  and  $T_{k,22}$ ), wherefor the lower block equation gives

$$T_{k,22}W_k^u = S_{k,22}W_k^u P \Rightarrow W_k^u = T_{k,22}^{-1}S_{k,22}W_k^u P.$$
(16)

<sup>&</sup>lt;sup>3</sup>See the appendix for details.

Since the eigenvalues of  $T_{k,22}^{-1}S_{k,22}$  are inside the unit circle and P is the (semi) stable (but unknown) solvent or  $\mathcal{M}$  we can iterate

$$W_{k}^{u} = T_{k,22}^{-1} S_{k,22} W_{k}^{u} P = \left( T_{k,22}^{-1} S_{k,22} \right)^{2} W_{k}^{u} P^{2} = \left( T_{k,22}^{-1} S_{k,22} \right)^{j} W_{k}^{u} P^{j} \xrightarrow{}_{j \to \infty} 0.$$
(17)

Using the definition of  $W_k^u$  we receive

$$0 = W_k^u = Z_{k,21}^* + Z_{k,22}^* \Delta P_k \Rightarrow \Delta P_k = -Z_{k,22}^* {}^{-1}Z_{k,21}^* = Z_{k,21}Z_{k,11}^{-1}.$$
 (18)

Due to finite precision arithmetic, the algorithm will compute  $\Delta \hat{P}_k = \Delta P_k + \mathcal{O}_k(\epsilon)$  which is only an approximation of the true potential improvement  $\Delta P_k$  subject to numerical errors  $\mathcal{O}_k(\epsilon)$ . Therefore, the updated approximation of P yields

$$P_k = P_{k-1} + \Delta \widehat{P}_k, \tag{19}$$

We initialize with  $P_{-1} = 0$ , such that the  $P_0$  recovered above is the standard QZ solution derived by Klein (2000), Sims (2001), and Uhlig (1999) and is the standard method in Dynare.<sup>4</sup> We thusly interpret  $P_k$  for k > 0 as iterative QZ improvements and define the following iterative QZ algorithm

# Algorithm 1: Iterative QZ Algorithm

**Given:** *A*, *B*, *C*, and a convergence criterion  $\epsilon$ Set  $P_{-1} = 0$ **While** *criterion*( $\Delta P_k$ ) >  $\epsilon$  **do** 

Define 
$$\mathcal{A}_{k} = \begin{bmatrix} C + BP_{k-1} & B \\ P_{k-1} & I \end{bmatrix}$$
 and  $\mathcal{B}_{k} = \begin{bmatrix} -AP_{k-1} & -A \\ I & 0 \end{bmatrix}$   
Apply the QZ algorithm  $Q_{k}^{*}\mathcal{B}_{k}Z_{k} = S_{k}$  and  $Q_{k}^{*}\mathcal{A}_{k}Z_{k} = T_{k}$ , arranged with the unstable eigenvalues are in the lower right blocks of  $S_{k}$  and  $T_{k}$   
Set  $\Delta P_{k} = Z_{k,21}Z_{k,11}^{-1}$   
Set  $P_{k} = P_{k-1} + \Delta P_{k}$   
Advance  $k = k + 1$   
end

**Return:**  $P_k$ 

<sup>&</sup>lt;sup>4</sup>See the appendix for the details of implementing the algorithm in Dynare in accordance with its classification of variables.

Ideally we would solve all problems exactly. With finite precision, the best we can hope for is to solve a problem within machine precision,  $\epsilon$ . With each calculation, we invariably inherit potential rounding or floating point errors that limit the possible accuracy of our results. Furthermore, these errors are magnified in ill-conditioned problems. Accordingly, we follow Higham (2002, ch. 3) and calculate a target threshold for the solution of the matrix quadratic based on the number of floating point calculations. Adding C and  $AP^2 + BP$  involves  $ny^2$  operations and adding  $AP^2$  and BP another  $ny^2$ , each with a potential finite precision error  $\varepsilon_{MP}$ . The inner product x'y for vectors of length  $n_y$  can be bounded by  $\gamma(n_y) \equiv \frac{n_y \varepsilon_{MP}}{1 - n_y \varepsilon_{MP}}$  accordingly the product of B and P has floating point errors bounded by  $ny^2\gamma(n_y)$  and that of A, P, and P by  $(ny^2\gamma(n_y))^2$ . This gives in total  $ny^2 \left[ \varepsilon_{MP} + \varepsilon_{MP} \gamma(n_y) \left( \varepsilon_{MP} + \varepsilon_{MP} \gamma(n_y) \right) \right]$  which following Higham (2002, Lemma 3.3) using the rules of running error analysis gives  $n_y^2 (\varepsilon_{MP} + \gamma(n_y + 2) + \gamma(2 * n_y + 2))$ . For ill conditioned problems, these error can be potentiated into errors of the solution via the condition number  $\Psi$  (see Higham (2002) in general and Meyer-Gohde (2022) for the specific DSGE matrix quadratic problem). Combining, the realistic level of accuracy for a matrix quadratic problem with a given conditioning and dimension is

$$\epsilon = \Psi * n_{\nu}^2 \left( \varepsilon_{MP} + \gamma (n_{\nu} + 2) + \gamma (2 * n_{\nu} + 2) \right)$$
(20)

which we take as our convergence criterion  $\epsilon$ .

We expect our iterative QZ algorithm to perform favorably in terms of improving the accuracy of a solution for two reasons. Firstly, the algorithm attempts to set  $\Delta P_k$  to close the gap between the current initialization  $P_{k-1}$  and the solution to the matrix quadratic P. That is, it solves for  $\Delta P_k$  such that  $P_k = P$ . Of course, it is limited like the original QZ (i.e., the first iteration of our method) by finite precision arithmetic. However, the addition of the terms  $P_{k-1}$  in the lower left of  $\mathcal{A}_k$  and  $-AP_{k-1}$  in the upper left of  $\mathcal{B}_k$  removes two of the five zero/identity blocks in the original formulation of  $\mathcal{A}$  and  $\mathcal{B}$ . This is significant as the breakdown of the backward stability of the QZ algorithm for solving the matrix quadratic problem as identified by Tisseur (2000) is caused by the large number of mandatory zeros and ones to maintain the mapping of the matrix quadratic to a generalized eigenvalue problem, which, as Meyer-Gohde (2022) shows, leads to QZ based algorithms in the DSGE literature being more prone to numerical instability than other, less common methods. We will now confirm that with the reduction of these mandatory zeros and ones by 40%, our iterative QZ is able to overcome numerical instability as identified in a specific macro-finance example by Meyer-Gohde (2022).

#### HUBER AND MEYER-GOHDE

### **3.** A MACRO FINANCE: APPLICATION

We run through a set of experiments to assess the iterative QZ in the model of Jermann (1998). We test how our algorithm performs under both conditions when the standard QZ works well and when it does not - namely Meyer-Gohde's (2022) parameterizations and formulations of Jermann's (1998) macro finance model. This model is attractive as it successfully replicates key finance variables, such as the average equity premium, in a production based asset pricing framework - that is, with endogenous production and consumption in a compact parsimonious DSGE model. Furthermore, Heiberger, Klarl, and Maußner (2017) analyze a similar model, with habit formation in labor along with consumption, for the numerical accuracy of linear DSGE solutions.

Jermann's (1998) model of habit formation and adjustment costs gives agents a strong motive to care about volatile consumption streams but inhibits their ability to respond which drive the success in matching the macro and finance moments. The habit stock,  $X_t$ , is internal as households internalize the effect of consumption today on the habit they will face tomorrow, yielding marginal utility from consumption,  $\lambda_t$ , in the pricing kernel  $m_{t+1} \equiv \beta \lambda_{t+1}/\lambda_t$  of

$$\lambda_t \equiv \partial \left( u \left( c_t, X_t \right) + \beta E_t \left[ u \left( c_{t+1}, X_{t+1} \right) \right] \right) / \partial c_t \tag{21}$$

If habit formation is external, this is simply  $u_c(c_t, X_t)$ , here with the habit internalized it is a function of the previous period's consumption,  $X_t \equiv X(c_{t-1})$ , and is given by

$$\lambda_t = u_c(c_t, X(c_{t-1})) + \beta E_t[u_X(c_{t+1}, X(c_t))X_c(c_t)]$$

We assume  $u(\cdot_t) = (\cdot_t)^{1-\tau}/(1-\tau)$  with curvature  $\tau$  and habit  $\cdot_t = c_t - bc_{t-1}$  with the degree of habit formation *b*. Households' ability to reduce the volatility of consumption streams as desired is impeded by adjustment costs in capital accumulation

$$k_t = (1 - \delta/k_{t-1})k_{t-1}$$

where the capital adjustment cost function is  $\phi(i_t/k_{t-1}) = \frac{\tilde{b}}{1-\xi} (i_t/k_{t-1})^{1-\xi} + \tilde{c}$  with  $\tilde{b}$  and  $\tilde{c}$  set such that the steady state matches the case without adjustment costs and  $1/\xi$  is the elasticity of the investment-capital ratio with respect to Tobin's q.

We examine two parameterizations that can be found in table 1. The baseline calibration is the calibration given by Jermann (1998) slightly adjusted to match the his empirical moments, the average risk premium, the risk free rate and the standard deviations of output, investment and consumption growth, in the most accurate linear method exactly.

	τ	b	$\beta^* = \beta \overline{a}^{1-\tau}$	ξ	$\sigma_Z$	
Baseline	7.92	0.74	0.99	3.75	0.999%	
Alternative	29.96	0.66	0.94	0.76	0.997%	
	α	$\overline{a}$	δ	$\rho_Z$	$N_{ss}$	$SS_{adjc}$
Common	0.36	0.005	0.025	0.99	1	1

TABLE 1. Calibrations for Jermann's (1998) DSGE Macro-Finance Model

The alternative parameterization is merely a near-by calibration with an increase in the curvature of the utility kernel - increasing households' unconditional sensitivity to volatile consumption streams - partially attenuated by a reduction in degree of habit formation but an significant increase in the elasticity in the adjustment costs - making them, however, more able to respond to this volatility. Just as different methods that are theoretically identical can have different numerical consequences, so too can different formulations of a model that are theoretically identical have different numerical consequences. Consider the application of the Lucas asset pricing equation to pricing economic capital

$$1 = E_t \left[ e^{\hat{m}_{t+1} + \hat{R}_{t+1}} \right]$$
(22)

or equivalently

$$1 = \beta E_t \left[ e^{-\tau \overline{a} + \hat{R}_{t+1}} \frac{\mu_{t+1}}{\mu_t} \right], \text{ as } e^{\hat{m}_t} = \beta e^{-\tau \overline{a}} \frac{\mu_t}{\mu_{t-1}}$$
(23)

and  $\mu_t$  is deterministically detrended  $\lambda_t$  from (21) above. In the results above, the formulation (23) was used, now we replace (23) with (22). This theoretically makes no difference, but numerically changes the problem significantly (Meyer-Gohde, 2022). To see this intuitively, notice that previously, there were entries in the *B* matrix (associated with contemporaneous variables) as well as in the *A* matrix (associated with future variables) in the row associated with this equation. Now with (22), there are only entries in the *A* matrix (associated with future variables) in the row associated with this equation.

Table 2 contains the results for both parameterizations and the reformulation of the Euler equation. The data row repeats the moments in post-war US data reported by Jermann (1998) in his study. The first set of results reports the baseline calibration and model definition. We see that our iterative QZ algorithm terminates after the initial iteration and produces the same results as Dynare's QZ and the best (in terms of forward error which is Dynare's cyclic reduction (CR) method) performing algorithm for this constellation of Jermann's (1998) model as found by Meyer-Gohde (2022).<sup>5</sup> Thus, the

<sup>&</sup>lt;sup>5</sup>Note that the forward errors for our iterative QZ and Dynare's QZ differ by roughly an order of magnitude. For the experiments with Jermann's (1998) model, we implement the naïve version of our

	E[rp]	$E\left[R_{t}^{f} ight]$	$std(\log y_t)$	$std(\log c_t)$	Cond. Num.	Forward Error
Data	6.18	0.8	0.01	0.51	-	_
Baseline Calibration						
Iterative QZ (1 iteration)	6.18	0.8	0.01	0.51	9.4e+05	1.12e-12
Dynare QZ	6.18	0.8	0.01	0.51	9.4e+05	2.06e-13
In Meyer-Gohde (2022) (Best: Dynare CR)	6.18	0.8	0.01	0.51	9.4e+05	2.23e-15
Alternative Calibration						
Iterative QZ (2 iterations)	6.23	0.824	0.01	0.519	1.06e+05	3.07e-08
Dynare QZ	6.09	0.696	0.01	0.584	1.22e+05	1.83e-01
In Meyer-Gohde (2022) (Best: Binder-Peseran)	6.23	0.824	0.01	0.519	1.06e+05	1.77e-15
Alternative Calibration, Alter	rnate Equ	ations				
Iterative QZ (8 iterations)	6.23	0.824	0.01	0.519	1.05e+05	7.19e-06
Dynare QZ	6.29	-0.326	0.01	0.531	7.65e+04	1.16e+01
In Meyer-Gohde (2022) (Best: AIM)	6.23	0.824	0.01	0.519	1.05e+05	1.72e-15

### TABLE 2. Jermann (1998) Moments and Errors

expected level of accuracy is returned by the standard QZ algorithm and the macro finance results in terms of moments of economic variables are identical in the QZ and the most accurate algorithm. The next set of results is for the alternative calibration with higher utility curvature counterbalanced by a lower degree of habit formation and higher elasticity of adjustment costs. Our iterative QZ algorithm does *not* terminate after the initial iteration and chooses instead to run a second iteration. Compared with the standard one-shot implementation of QZ there is a gain of forward error accuracy of

algorithm as laid out above. Dynare's implementation reduces the dimensionality of the QZ problem by grouping variables and structuring the matrix quadratic according to the classification of "static", "purely forward", "purely backward looking", and "mixed" variables, see Villemot (2011). The analogous dimension reduction for our iterative QZ algorithm is detailed in the appendix. We choose to report the naïve results here, particularly for the baseline calibration, to highlight the importance of the actual numerical implementation of QZ that leads to a one order of magnitude improvement in the forward error. roughly seven degrees of magnitude and our iterative QZ successfully recovers the macro finance moments produced by the most accurate method reported in Meyer-Gohde (2022). The last set of results with both the alternative calibration and formulation of the Euler equations presents a formidable challenge - many methods in Meyer-Gohde (2022) failed to produce a solution at all and Dynare's QZ even predicts a *negative* risk-free rate for this case. With 8 iterations, our iterative QZ has to put in extra effort with this particular problem to achieve the target forward error. Upon convergence, the algorithm again is roughly seven degrees of magnitude more accurate than Dynare's QZ and produces moments that match those of the most accurate method for this constellation. Thus, for the model of Jermann (1998), we conclude that our method successfully identifies situations where the standard QZ method is sufficient and rectifies those that are not.

In sum, our experiments clearly demonstrate the utility and reliability of our algorithm - iterating over the standard QZ results when they are not sufficiently accurate and refraining from iterating when the are. The only additional costs over the standard implementation of QZ are the calculations of conditioning numbers and forward errors for each model, which is standard numerical practice to ensure numerical stability.

# 4. CONCLUSION

We have introduced and developed an iterative implementation of the QZ algorithm for solving linear DSGE models as alternatives to the one-shot QZ methods in the literature. The first iteration initialized on the zero matrix returns the standard QZ output, further iterations provide improvement increments and we continue until the solution achieves the accuracy permitted by Meyer-Gohde's (2022) forward error bounds.

In a set of experiments using Jermann's (1998) macro finance model, we find that our proposed algorithm returns the standard QZ output should it already be accurate enough and otherwise runs a few iterations until this accuracy is achieved. Thus we have provided an entirely QZ based method to improve the accuracy of solutions of linear DSGE models alongside the alternative iterative methods developed in Meyer-Gohde and Saecker (2022), Meyer-Gohde (2023), Binder and Meyer-Gohde (2024), and Huber, Meyer-Gohde, and Saecker (2023). As these studies already demonstrated, there is likely no single one-size-fits-all algorithm and our iterative QZ based algorithm should be considered a complement of the different methods for solving linear DSGE models.

#### REFERENCES

- ADJEMIAN, S., H. BASTANI, M. JUILLARD, F. MIHOUBI, G. PERENDIA, M. RATTO, AND S. VILLEMOT (2011): "Dynare: Reference Manual, Version 4," Dynare Working Papers 1, CEPREMAP.
- BINDER, M., AND A. MEYER-GOHDE (2024): "Revisiting the Fully Recursive Computation of Multivariate Linear Rational Expectations Models," mimeo, Goethe University Frankfurt, Institute for Monetary and Financial Stability (IMFS).
- BLANCHARD, O. J., AND C. M. KAHN (1980): "The Solution of Linear Difference Models under Rational Expectations," *Econometrica*, 48(5), 1305–1311.
- GOLUB, G. H., AND C. F. VAN LOAN (2013): *Matrix Computations*. The Johns Hopkins University Press, fourth edn.
- HAMMARLING, S., C. J. MUNRO, AND F. TISSEUR (2013): "An Algorithm for the Complete Solution of Quadratic Eigenvalue Problems," ACM Transactions On Mathematical Software, 39(3), 18:1–18:19.
- HEIBERGER, C., T. KLARL, AND A. MAUSSNER (2017): "On the numerical accuracy of first-order approximate solutions to DSGE models," *Macroeconomic Dynamics*, 21(7), 1811–1826.
- HIGHAM, N. J. (2002): Accuracy and Stability of Numerical Algorithms. Society for Industrial and Applied Mathematics, second edn.
- HUBER, J., A. MEYER-GOHDE, AND J. SAECKER (2023): "Solving Linear DSGE Models With Structure-Preserving Doubling Methods," IMFS Working Paper Series 195, Goethe University Frankfurt, Institute for Monetary and Financial Stability (IMFS).
- JERMANN, U. J. (1998): "Asset Pricing in Production Economies," Journal of Monetary Economics, 41(2), 257–275.
- KING, R. G., AND M. W. WATSON (1998): "The Solution of Singular Linear Difference Systems under Rational Expectations," *International Economic Review*, 39(4), 1015–26.
- KLEIN, P. (2000): "Using the Generalized Schur Form to Solve a Multivariate Linear Rational Expectations Model," *Journal of Economic Dynamics and Control*, 24(10), 1405–1423.
- LAN, H., AND A. MEYER-GOHDE (2014): "Solvability of Perturbation Solutions in DSGE Models," Journal of Economic Dynamics and Control, 45, 366–388.
- MEYER-GOHDE, A. (2022): "Backward Error and Condition Number Analysis of Linear DSGE Solutions," mimeo, Goethe University Frankfurt, Institute for Monetary and

Financial Stability (IMFS).

- —— (2023): "Solving Linear DSGE Models with Bernoulli Methods," IMFS Working Paper Series 182, Goethe University Frankfurt, Institute for Monetary and Financial Stability (IMFS).
- MEYER-GOHDE, A., AND J. SAECKER (2022): "Solving Linear DSGE Models with Newton Methods," IMFS Working Paper Series 174, Goethe University Frankfurt, Institute for Monetary and Financial Stability (IMFS).
- MOLER, C. B., AND G. W. STEWART (1973): "An Algorithm for Generalized Matrix Eigenvalue Problems," *SIAM Journal on Numerical Analysis*, 10(2), 241–256.
- SIMS, C. A. (2001): "Solving Linear Rational Expectations Models," Computational Economics, 20(1-2), 1–20.
- TISSEUR, F. (2000): "Backward error and condition of polynomial eigenvalue problems," *Linear Algebra and its Applications*, 309(1), 339–361.
- UHLIG, H. (1999): "A Toolkit for Analysing Nonlinear Dynamic Stochastic Models Easily," in Computational Methods for the Study of Dynamic Economies, ed. by R. Marimon, and A. Scott, chap. 3, pp. 30–61. Oxford University Press.
- VILLEMOT, S. (2011): "Solving Rational Expectations Models at First Order: What Dynare Does," Dynare Working Papers 2, CEPREMAP.

#### APPENDIX

#### 4.1. Eigenvalue Preservation

The eigenvalue preservation can be seen by comparison. The eigenvalues of the pencil defined by  $\mathscr{A}$ ,  $\mathscr{B}$  are given by  $\lambda \in \mathbb{C}$  such that

$$0 = \det(\mathscr{A} - \lambda \mathscr{B}) \Rightarrow 0 = \det(A\lambda^2 + B\lambda + C)$$
(A1)

The eigenvalues of the pencil defined by  $\mathcal{A}_k$ ,  $\mathcal{B}_k$  are given by  $\mu \in \mathbb{C}$  such that

$$0 = \det\left(\mathscr{A}_k - \mu \mathscr{B}_k\right) \tag{A2}$$

$$= \det \left( \begin{bmatrix} C + BP_{k-1} & B \\ P_{k-1} & I \end{bmatrix} - \mu \begin{bmatrix} -AP_{k-1} & -A \\ I & 0 \end{bmatrix} \right)$$
(A3)

$$= \det\left( \begin{bmatrix} AP_{k-1}\mu + C + BP_{k-1} & A\mu + B\\ -I\mu + P_{k-1} & I \end{bmatrix} \right)$$
(A4)

(A5)

As  $-I\mu + P_{k-1}$  and I commute

$$0 = \det\left(\mathscr{A}_k - \mu\mathscr{B}_k\right) \tag{A6}$$

$$= \det \left[ \left( AP_{k-1}\mu + C + BP_{k-1} \right) I - \left( A\mu + B \right) \left( P_{k-1} - I\mu \right) \right]$$
(A7)

$$= \det \left[ AP_{k-1}\mu + C + BP_{k-1} - A\mu P_{k-1} - BP_{k-1} + A\mu^2 + B\mu \right]$$
(A8)

$$= \det\left(C + A\mu^2 + B\mu\right) \tag{A9}$$

Inspection shows that  $\lambda$  and  $\mu$  are identical.

#### 4.2. Detailed Dynare Topology

Here we summarize the details in the matrix quadratic that follows from the typology of variables from Dynare as laid out in Villemot (2011). See Meyer-Gohde and Saecker (2022) for additional details.

Subdividing the system of equations in accordance with the QR decomposition yields



where  $n^d$  is the number of dynamic variables, the sum of number of purely backward-looking,  $n^{--}$ , mixed  $n^m$ , and purely forward-looking variables,  $n^{++}$ . The number of forward-looking variables,  $n^+$ , is the sum of the number of mixed,  $n^m$ , and purely forward-looking variables,  $n^{++}$ , and the number of backward-looking variables,  $n^-$ , is the sum of the number of purely backward-looking,  $n^{--}$  and mixed variables  $n^m$ . Hence, the number of endogenous variables is the sum of the number of static,  $n^s$ , and dynamic variables,  $n^d$ , or

the sum of the number of static,  $n^s$ , purely backward-looking,  $n^{--}$ , mixed  $n^m$ , and purely forward-looking variables,  $n^{++}$ . The dimensions satisfy the following

$$n^{d} = n^{--} + n^{m} + n^{++}, \quad n^{+} = n^{m} + n^{++}, \quad n^{-} = n^{--} + n^{m}, \quad n = n^{s} + n^{d} = n^{s} + n^{--} + n^{m} + n^{++}$$

The transition matrix, P, from (2) that solves the matrix equation (3) can be subdivided in accordance to Dynare's typology as

The matrix quadratic can be expressed as

$$\mathbf{M}(\mathbf{P}_{n \times n}) = \mathbf{A}_{n \times n} \mathbf{P}^2 + \mathbf{B}_{n \times n} \mathbf{P} + \mathbf{C}_{n \times n}$$
$$= \underbrace{\left(\mathbf{A}\mathbf{P} + \mathbf{B}\right)}_{\equiv \mathbf{G}} \mathbf{P} + \mathbf{C}$$

For a solvent P of the matrix quadratic, taking the structure of C from the Dynare typology above into account yields

$$\mathbf{M}(\mathbf{P}) = 0 = \mathbf{G}\mathbf{P} + \mathbf{C}$$

$$n^{s} \quad n^{--} \quad n^{m} \quad n^{++}$$

$$= \mathbf{G} \quad n \begin{bmatrix} \mathbf{P}_{\cdot,s} & \mathbf{P}_{\cdot,--} & \mathbf{P}_{\cdot,m} & \mathbf{P}_{\cdot,++} \end{bmatrix} + \frac{n^{--}}{n^{m}} \begin{bmatrix} \mathbf{0} & \mathbf{A}_{-}^{-} & \mathbf{0} \\ n^{s \times n^{--}} & \mathbf{0} \end{bmatrix}$$

$$\mathbf{A}_{-}^{\mathbf{A}_{-}} = \mathbf{O}_{-}$$

Following Meyer-Gohde and Saecker (2022), who apply corollary 4.5 of Lan and Meyer-Gohde (2014), if P is the unique solvent of M(P) stable with respect to the closed unit circle, **G** has full rank and hence the columns of P associated with nonzero columns in C, the static and forward-looking vari-

ables are zero  $\rightarrow \mathbf{P}_{\bullet,s} = \mathbf{0}_{n \times n^{s}}, \ \mathbf{P}_{\bullet,++} = \mathbf{0}_{n \times n^{++}}, \ \text{whence } \mathbf{P} \text{ is } \mathbf{P} = n \begin{bmatrix} \mathbf{0} & | \mathbf{P}_{\bullet,--} & | \mathbf{P}_{\bullet,m} & | \mathbf{0} \end{bmatrix} \text{ and}$  $\mathbf{M}(\mathbf{P}) = \begin{bmatrix} \mathbf{0} & \mathbf{M}(\mathbf{P})^{--} & \mathbf{M}(\mathbf{P})^{m} & \mathbf{0}_{n \times n^{m}} \end{bmatrix} \text{. Consequentially, the first } n^{s} \text{ rows of the matrix quadratic, taking}$  n  $n^{--} & n^{m} \\ n^{m} & \mathbf{P}_{m,\bullet} \\ n^{m} & \mathbf{P}_{m,\bullet} \end{bmatrix} \text{ as given, yield } n^{s} \begin{bmatrix} \mathbf{P}_{s,--} & | & \mathbf{P}_{s,m} \end{bmatrix} \text{ as}$   $n^{++} & \mathbf{P}_{++,\bullet} \end{bmatrix}$   $n^{--} & n^{m} \\ n^{s} \begin{bmatrix} \mathbf{P}_{s,--} & | & \mathbf{P}_{s,m} \end{bmatrix} = -\begin{bmatrix} \mathbf{A}_{n^{s} \times n^{s}} \end{bmatrix}^{-1} \begin{pmatrix} n^{--} & n^{m} & n^{--} & n^{m} \\ \mathbf{A}_{n^{s} \times n^{+}}^{*} n^{n++} & \mathbf{P}_{n++,--} & \mathbf{P}_{n++,m} \end{bmatrix} n^{m} \begin{bmatrix} \mathbf{P}_{-,--} & \mathbf{P}_{-,m} \\ \mathbf{P}_{m,--} & \mathbf{P}_{m,m} \end{bmatrix}$ 

 $n^{s} \quad n^{--} \quad n^{m} \quad n^{++}$ and the first  $n^{s}$  rows of P are  $\mathbf{P}_{s,\bullet} = n^{s} \begin{bmatrix} \mathbf{0} & \mathbf{P}_{s,--} & \mathbf{P}_{s,m} & \mathbf{0} \end{bmatrix}$ . The last  $n^{d}$  columns and rows of P solve the reduced matrix quadratic equation

Recalling that  $\mathbf{P}_{\bullet,++} = \mathbf{0}_{n \times n^{++}}$ ,  $\tilde{\mathbf{P}}$  can be reduced and four submatrices  $\mathbf{P}^{--}$ ,  $\mathbf{P}^{-}$ ,  $\mathbf{P}^{+}$  and  $\mathbf{P}^{++}$  defined via

$$\tilde{\mathbf{P}} = n^{m} \frac{n^{--}}{n^{m}} \frac{n^{++}}{n^{++}} \frac{n^{--}}{n^{m}} \frac{n^{++}}{n^{++}} \frac{n^{--}}{n^{m}} \frac{n^{m}}{n^{++}} \frac{n^{++}}{n^{--}} \frac{n^{m}}{n^{m}} \frac{n^{++}}{n^{++}} \frac{n^{--}}{n^{m}} \frac{n^{m}}{n^{++}} \frac{n^{++}}{n^{m}} \frac{n^{--}}{n^{m}} \frac{n^{m}}{n^{m}} \frac{n^{++}}{n^{m}} \frac{n^{--}}{n^{m}} \frac{n^{m}}{n^{m}} \frac{n^{--}}{n^{m}} \frac{n^{m}}{n^{m}} \frac{n^{--}}{n^{m}} \frac{n^$$

where 
$$\mathbf{P}_{n^{--} \times n^{-}}^{--} \equiv n^{--} \begin{bmatrix} \mathbf{P}_{--,--} & \mathbf{P}_{--,m} \end{bmatrix}, \quad \mathbf{P}_{n^{--} \times n^{-}}^{--} \equiv \frac{n^{m}}{n^{--}} \begin{bmatrix} \mathbf{P}_{m,--} & \mathbf{P}_{m,m} \\ \mathbf{P}_{--,--} & \mathbf{P}_{--,m} \end{bmatrix}, \quad \mathbf{P}_{n^{+} \times n^{-}}^{+-} \equiv \frac{n^{m}}{n^{++}} \begin{bmatrix} \mathbf{P}_{m,--} & \mathbf{P}_{m,m} \\ \mathbf{P}_{++,--} & \mathbf{P}_{++,m} \end{bmatrix},$$

and  $\mathbf{P}^{++}_{n^{++} \times n^{-}} \equiv n^{++} \begin{bmatrix} \mathbf{P}_{++,--} & \mathbf{P}_{++,m} \end{bmatrix}$ .

We can now write the matrix quadratic as  $\frac{1}{2}$ 

$$n^{--} n^{m} n^{++}$$

$$n^{--} \begin{bmatrix} \mathbf{0} \\ \mathbf{A}^{-} \\ n^{d \times n^{-}} \end{bmatrix} = \mathbf{\tilde{M}}(\mathbf{\tilde{P}}) = n^{d} \begin{bmatrix} \mathbf{\tilde{M}}(\mathbf{\tilde{P}})^{--} & \mathbf{\tilde{M}}(\mathbf{\tilde{P}})^{m} \end{bmatrix} = \mathbf{0} \\ n^{d \times n^{d}} \end{bmatrix} = \mathbf{0} \\ n^{d \times n^{d}}$$

Multiplying this out and keeping only the generically nonzero columns gives

$$\tilde{\mathbf{A}}^{+}_{n^{d} \times n^{+} n^{+} \times n^{-} n^{-} \times n^{-}} + \tilde{\mathbf{A}}^{0+}_{n^{d} \times n^{+} n^{+} \times n^{-}} + \tilde{\mathbf{A}}^{0-}_{n^{d} \times n^{-} n^{-} \times n^{-}} + \tilde{\mathbf{A}}^{-}_{n^{d} \times n^{-}} = \mathbf{0}$$
(A10)

where

$$\mathbf{\tilde{A}}_{n^{d} \times n^{d}}^{0} = n^{d} \begin{bmatrix} \mathbf{\tilde{A}}_{n^{d} \times n^{-}}^{0} & \mathbf{0} \end{bmatrix} + n^{d} \begin{bmatrix} \mathbf{0} & \mathbf{\tilde{A}}_{n^{d} \times n^{+}}^{0} \end{bmatrix}$$

As stated by Villemot (2011), the allocation of the middle columns (the  $n^m$  ones associated with "mixed" variables) between  $\tilde{\mathbf{A}}^{0-}$  and  $\tilde{\mathbf{A}}^{0+}$  is not unique.

The companion linearization of (A10) analogous to the main text is

$$n^{-} n^{+} n^{-} n^{-} n^{-} n^{-} n^{+} n^{-} n^{-$$

 $n^{m} n^{++} n^{--} n^{m}$ where  $I^{+}_{n^{m} \times n^{+}} = n^{m} \begin{bmatrix} I \\ n^{m} \times n^{m} \end{bmatrix} n^{m} + 1$  and  $I^{-}_{n^{m} \times n^{-}} = n^{m} \begin{bmatrix} 0 \\ n^{m} \times n^{--} \end{bmatrix} I$  are selection matrices. This is (4), note that  $n^{d} + n^{m} = n^{-} + n^{+}$ 

$$\mathscr{A}_{(n^{d}+n^{m})\times(n^{-}+n^{+})(n^{-}+n^{+})\times n^{-}} = \mathscr{B}_{(n^{d}+n^{m})\times(n^{-}+n^{+})(n^{-}+n^{+})\times n^{-}(n^{-}+n^{+})\times n^{-}}$$
(A12)

now with

$$\mathscr{X} = \frac{n^{-}}{n^{+}} \begin{bmatrix} I \\ n^{-} \times n^{-} \\ \mathbf{P}^{+} \\ n^{+} \times n^{-} \end{bmatrix} \qquad \mathscr{A} = \frac{n^{d}}{n^{m}} \begin{bmatrix} \tilde{\mathbf{A}}^{-} & \tilde{\mathbf{A}}^{0+} \\ n^{d} \times n^{-} & n^{d} \times n^{+} \\ \mathbf{0} \\ n^{m} \times n^{-} & n^{m} \times n^{+} \end{bmatrix}, \qquad \mathscr{B} = \frac{n^{d}}{n^{m}} \begin{bmatrix} -\tilde{\mathbf{A}}^{0-} & -\tilde{\mathbf{A}}^{+} \\ n^{d} \times n^{-} & n^{d} \times n^{+} \\ I^{-} \\ n^{m} \times n^{-} & n^{m} \times n^{+} \end{bmatrix}, \qquad \mathscr{M} = \mathbf{P}^{-}_{n^{-} \times n^{-}}$$

Now we can begin with an approximate solution  $\mathbf{P}^+_{k-1}$  and define the eigenvalue preserving transformation

$$U_{k} \equiv \frac{n^{-} \quad n^{+}}{n^{+} \left[ \begin{array}{ccc} I & \mathbf{0} \\ n^{-} \times n^{-} & n^{-} \\ \mathbf{P}_{k-1}^{+} & I \\ n^{+} \times n^{-} \end{array} \right]}, \qquad \qquad U_{k}^{-1} \equiv \frac{n^{-} \left[ \begin{array}{ccc} I & \mathbf{0} \\ n^{-} \times n^{-} & n^{-} \\ \mathbf{0}_{k}^{-1} \\$$

giving

$$\mathscr{A}_k \hat{\mathscr{X}}_k = \mathscr{B}_k \hat{\mathscr{X}}_k \mathscr{M} \tag{A14}$$

or

as

$$\hat{\mathscr{X}}_{k} \equiv U_{k}^{-1} \mathscr{X}_{k} = \frac{n^{-} \begin{bmatrix} I \\ n^{-} \times n^{-} \\ n^{+} \end{bmatrix} \begin{pmatrix} 0 \\ n^{-} \times n^{+} \\ -\mathbf{P}_{k-1}^{+} \\ n^{+} \times n^{-} \end{bmatrix} \begin{pmatrix} n^{-} \begin{bmatrix} I \\ n^{-} \times n^{-} \\ n^{+} \end{bmatrix} \begin{pmatrix} n^{-} \begin{bmatrix} I \\ n^{-} \times n^{-} \\ \mathbf{P}_{k}^{+} \\ n^{+} \times n^{-} \end{bmatrix} = \frac{n^{-} \begin{bmatrix} I \\ n^{-} \times n^{-} \\ n^{+} \end{bmatrix} \begin{pmatrix} I \\ \mathbf{P}_{k}^{+} \\ n^{+} \times n^{-} \end{pmatrix} \begin{pmatrix} I \\ \mathbf{P}_{k-1}^{+} \\ n^{+} \times n^{-} \end{pmatrix}$$

and

$$\mathcal{A}_{k} \equiv \mathcal{A}U_{k} = \begin{bmatrix} n^{-} & n^{+} & n^{-} & n^{+} \\ \frac{\tilde{\mathbf{A}}_{k}^{-} + \tilde{\mathbf{A}}_{n}^{0+} \mathbf{P}_{k-1}^{+} & \tilde{\mathbf{A}}_{n}^{0+} \\ \frac{n^{d} \times n^{-} & n^{d} \times n^{+} n^{+} \times n^{-}}{n^{d} \times n^{+} n^{+} \times n^{-}} & \frac{\tilde{\mathbf{A}}_{n}^{0+} \\ n^{d} \times n^{+} & n^{d} \times n^{+}}{n^{d} \times n^{+}} \end{bmatrix}, \quad \mathcal{B}_{k} \equiv \mathcal{B}U_{k} = \begin{bmatrix} n^{d} & -\tilde{\mathbf{A}}_{n}^{0-} - \tilde{\mathbf{A}}_{n}^{+} \mathbf{P}_{k-1}^{+} & -\tilde{\mathbf{A}}_{n^{d} \times n^{+}} \\ \frac{1}{n^{d} \times n^{-}} & n^{d} \times n^{+} \\ n^{m} \times n^{+} & n^{+} \times n^{-} & n^{m} \times n^{+} \end{bmatrix}, \quad \mathcal{B}_{k} \equiv \mathcal{B}U_{k} = \begin{bmatrix} n^{d} & -\tilde{\mathbf{A}}_{n}^{0-} - \tilde{\mathbf{A}}_{n}^{+} \mathbf{P}_{k-1}^{+} & -\tilde{\mathbf{A}}_{n^{d} \times n^{+}} \\ \frac{1}{n^{d} \times n^{-}} & n^{d} \times n^{+} \\ n^{m} \times n^{+} & n^{m} \times n^{+} & n^{m} \end{pmatrix}, \quad (A16)$$

As in the main text, we can apply the QZ decomposition to the pencil defined by  $\mathcal{A}_k$ ,  $\mathcal{B}_k$ , delivering unitary  $Q_k$  and  $Z_k$  and upper triangular  $S_k$  and  $T_k$  with  $Q_k^* \mathcal{B}_k Z_k = S_k$  and  $Q_k^* \mathcal{A}_k Z_k = T_k$ 

$$n^{-} n^{+} n^{-} n^{-} n^{+} n^{-} n^{-} n^{+} n^{-}$$

$$n^{-} \left[ \frac{T_{k,11}}{n^{-} \times n^{-}} \frac{T_{k,12}}{n^{-} \times n^{+}} \right] n^{-} \left[ \frac{w_{k}^{s}}{n^{-} \times n^{-}} \right] = n^{-} \left[ \frac{S_{k,11}}{n^{-} \times n^{-}} \frac{S_{k,12}}{n^{-} \times n^{+}} \right] n^{-} \left[ \frac{w_{k}^{s}}{n^{-} \times n^{-}} \right] n^{-} \left[ \frac{w_{k}^{s}}{n^{-} \times n^{-}} \frac{n^{-} \left[ \frac{S_{k,11}}{n^{-} \times n^{-}} \frac{S_{k,22}}{n^{+} \times n^{+}} \right] n^{+} \left[ \frac{w_{k}^{s}}{n^{+} \times n^{-}} \right] n^{-} \left[ \frac{w_{k}^{s}}{n^{-} \times n^{-}} \right] n^{-} \left[ \frac{w_{k}^{s}}{n^{-} \times n^{-}} \frac{n^{-} \left[ \frac{w_{k}^{s}}{n^{-} \times n^{-}} \frac{w_{k}^{s}}{n^{-} \times n^{-}} \right] n^{-} \left[ \frac{w_{k}^{s}}{n^{-} \times n^{-}} \frac{w_{k}^{s}}{n^{-} \times n^{-}} \right] n^{-} \left[ \frac{w_{k}^{s}}{n^{-} \times n^{-}} \frac{w_{k}^{s}}{n^{-} \times n^{-}} \frac{w_{k}^{s}}{n^{-} \times n^{-}} \frac{w_{k}^{s}}{n^{-} \times n^{-}} \right] n^{-} \left[ \frac{w_{k}^{s}}{n^{-} \times n^{-}} \right] n^{-} \left[ \frac{w_{k}^{s}}{n^{-} \times n^{-}} \frac{w_{k}^{$$

where

$$\frac{n^{-}}{Z_{k}^{*}} \frac{n^{-}}{\left(\frac{I}{n^{-} \times n^{-}}\right)} = \frac{n^{-}}{n^{+}} \left(\frac{w_{k}^{s}}{\frac{n^{-} \times n^{-}}{w_{k}^{u}}}\right) \Leftrightarrow \frac{n^{-}}{n^{+}} \left(\frac{I}{\frac{n^{-} \times n^{-}}{\Delta \mathbf{P}_{k}^{+}}}\right) = \frac{n^{-}}{\left(\frac{w_{k}^{s}}{\frac{n^{-} \times n^{-}}{\omega_{k}^{u}}}\right)} = \frac{n^{-}}{\left(\frac{w_{k}^{s}}{\frac{n^{-} \times n^{-}}{\omega_{k}^{u}}}\right)} \tag{A18}$$

and

We assume the decomposition is ordered so that the unstable eigenvalues are in the lower right blocks of  $S_k$  and  $T_k$  (hence  $S_{k,22}$  and  $T_{k,22}$ ), whence the lower block equation gives

$$T_{k,22}w_k^u = S_{k,22}w_k^u \mathbf{P}^- \Rightarrow w_k^u = T_{k,22}^{-1}S_{k,22}w_k^u \mathbf{P}^- = 0$$
(A20)

18

as the eigenvalues of  $T_{k,22}^{-1}S_{k,22}$  are inside the unit circle and  $\mathbf{P}^-$  is the (semi) stable (but unknown) solvent or  $\mathcal{M}$  - see the main text. Using the definitions of  $w_k^s$  and  $w_k^u$  and inserting  $w_k^u = 0$  gives

$$n^{-} \qquad n^{-} \qquad n^{-} \qquad n^{+} \qquad n^{-}$$

$$n^{-} \left[ \frac{I}{n^{-} \times n^{-}} \right]_{n^{+} \times n^{-}} = n^{-} \left[ \frac{Z_{k,11}}{n^{-} \times n^{-}} \qquad \frac{Z_{k,12}}{n^{-} \times n^{+}} \right]_{n^{-} \times n^{+}} n^{-} \left[ \frac{w_{k}^{s}}{n^{-} \times n^{-}} \right]_{n^{+} \times n^{-}} n^{-} \left[ \frac{Z_{k,21}}{n^{+} \times n^{-}} \qquad n^{+} \times n^{+} \right]_{n^{+} \times n^{-}} n^{+} \left[ \frac{w_{k}^{s}}{n^{-} \times n^{-}} \right]_{n^{+} \times n^{-}} n^{+} \left[ \frac{w_{k}^{s}}{n^{+} \times n^{-}} \right]_{n^{+} \times n^{-}} n^{+} n^{+}$$

and

$$n^{-} \begin{bmatrix} Z_{k,11}^{*} & Z_{k,12}^{*} \\ n^{-} \times n^{-} & n^{-} \\ R^{+} \begin{bmatrix} Z_{k,21}^{*} & Z_{k,22}^{*} \\ n^{+} \times n^{-} & n^{+} \times n^{+} \end{bmatrix} n^{-} \begin{bmatrix} I \\ n^{-} \times n^{-} \\ \Delta \mathbf{P}_{k}^{+} \\ n^{+} \times n^{-} \end{bmatrix} = n^{-} \begin{bmatrix} w_{k}^{s} \\ n^{-} \times n^{-} \\ n^{+} \end{bmatrix}$$
(A22)

the second block equations (noting that the first block of the first definition implies  $w_k^s = Z_{k,11}^{-1}$ )

+

$$0 = w_k^u = Z_{k,21}^* + Z_{k,22}^* \Delta \mathbf{P}_k^+ \Rightarrow \Delta \mathbf{P}_k^+ = -Z_{k,22}^{*-1} Z_{k,21}^* = Z_{k,21} Z_{k,11}^{-1}$$
(A23)

Using the definition of  $\Delta \mathbf{P}_k^+,$  this defines an update of  $\mathbf{P}_{k-1}^+$  given by

$$\mathbf{P}_{k}^{+} = \mathbf{P}_{k-1}^{+} + \Delta \mathbf{P}_{k}^{+} \tag{A24}$$

Upon convergence of  $\mathbf{P}_k^+$ ,  $\mathbf{P}^-$  then follows from  $w_k^s = Z_{k,11}^{-1}$  and  $w_k^u = 0$  inserted into the pencil

$$n^{-} n^{+} n^{+} n^{-} n^{-} n^{+} n^{-} n^{-} n^{+} n^{-}$$

$$n^{-} \left[ \frac{T_{k,11}}{n^{-} \times n^{-}} \frac{T_{k,12}}{n^{-} \times n^{+}} \right] n^{-} \left[ \frac{Z_{k,11}^{-1}}{n^{-} \times n^{-}} \right] = n^{-} \left[ \frac{S_{k,11}}{n^{-} \times n^{-}} \frac{S_{k,12}}{n^{-} \times n^{+}} \right] n^{-} \left[ \frac{Z_{k,11}^{-1}}{n^{-} \times n^{-}} \frac{R^{-}}{n^{+} \times n^{+}} \right] n^{+} \left[ \frac{R^{-}}{n^{-} \times n^{-}} \frac{R^{-}}{n^{+} \times n^{-}} \right] n^{+} \left[ \frac{R^{-}}{n^{+} \times n^{-}} \frac{R^{-}}{n^{+} \times n^{-}} \right] n^{+} \left[ \frac{R^{-}}{n^{-} \times n^{-}} \frac{R^{-}}{n^{-} \times n^{-}} \right] n^{+} \left[ \frac{R^{-}}{n^{-} \times n^{-}} \frac{R^{-}}{n^{-} \times n^{-}} \right] n^{-} \left[ \frac{R^{-}}{n^{-} \times n^{-}} \frac{R^{-}}{n^{-} \times n^{-}}$$

with the first block equation being

$$T_{k,11}Z_{k,11}^{-1} = S_{k,11}Z_{k,11}^{-1}\mathbf{P}^{-}$$
(A26)

the eigenvalue assumptions ensure that  $S_{k,11}$  is invertible, hence,

$$\mathbf{P}^{-} = Z_{k,11} S_{k,11}^{-1} T_{k,11} Z_{k,11}^{-1}$$
(A27)

Finally, the (generically) non zero elements of the first  $n^s$  rows of P ( $\mathbf{P}_{s,\bullet} = n^s \quad n^{--} \quad n^m \quad n^{++}$  $n^s \begin{bmatrix} \mathbf{0} & \mathbf{P}_{s,--} & \mathbf{P}_{s,m} & \mathbf{0} \end{bmatrix}$ ) solve

$$n^{--} n^{m} = \left[ \check{\mathbf{A}}_{n^{s} \times n^{s}}^{0s} \right]^{-1} \begin{pmatrix} n^{--} n^{m} \\ n^{--} \left[ \mathbf{P}_{--,--} - \mathbf{P}_{--,m} \right] \\ \check{\mathbf{A}}_{n^{s} \times n^{+}}^{+} \mathbf{P}_{k}^{+} \mathbf{P}^{-} + \check{\mathbf{A}}_{n^{s} \times n^{d}}^{0d} n^{m} \left[ \mathbf{P}_{m,--} - \mathbf{P}_{m,m} \right] \\ n^{++} \left[ \mathbf{P}_{++,--} - \mathbf{P}_{++,m} \right] + \check{\mathbf{A}}_{n^{s} \times n^{-}}^{-} \end{pmatrix}$$

# where from above

# IMFS WORKING PAPER SERIES

# Recent Issues

216/2025	Michael Haliassos, Thomas Jansson, Yigitcan Karabulut	Wealth Inequality: Opportunity for Me or for Others?
215/2024	Michael D. Bauer, Eric Offner, Glenn D. Rudebusch	Green Stocks and Monetary Policy Shocks: Evidence from Europe
214 / 2024	Michael D. Bauer, Daniel Huber, Eric Offner, Marlene Renkel, Ole Wilms	Corporate Green Pledges
213 / 2024	Athanasios Orphanides	The Federal Reserve's Evolving Interpretation and Implementation of Its Mandate
212 / 2024	Matthias Rumpf Michael Haliassos Tetyana Kosyakova Thomas Otter	Do Financial Advisors Have Different Beliefs than Lay People?
211 / 2024	Michael Haliassos	Wealth Accumulation: The Role of Others
210 / 2024	Kamila Duraj Daniela Grunow Michael Haliassos Christine Laudenbach Stephan Siegel	Rethinking the Stock Market Participation Puzzle: A Qualitative Approach
209 / 2024	Balint Tatar Volker Wieland	Policy Rules and the Inflation Surge: The Case of the ECB
208 / 2024	Reimund Mink	Helmut Schlesinger: Wegbereiter und Garant der deutschen Geld- und Stabilitätspolitik wird 100
207 / 2024	Alexander Meyer-Gohde	Solving and analyzing DSGE models in the frequency domain
206 / 2024	Jochen Güntner Magnus Reif Maik Wolters	Sudden Stop: Supply and Demand Shocks in the German Natural Gas Market
205 / 2024	Alina Tänzer	Multivariate Macroeconomic Forecasting: From DSGE and BVAR to Artificial Neural Networks
204 / 2024	Alina Tänzer	The Effectiveness of Central Bank Purchases of long-term Treasury Securities:A Neural Network Approach

203 / 2024	Gerhard Rösl	A present value concept for measuring welfare
202 / 2024	Reimund Mink Karl-Heinz Tödter	Staatsverschuldung und Schuldenbremse
201 / 2024	Balint Tatar Volker Wieland	Taylor Rules and the Inflation Surge: The Case of the Fed
200 / 2024	Athanasios Orphanides	Enhancing resilience with natural growth targeting
199 / 2024	Thomas Jost Reimund Mink	Central Bank Losses and Commercial Bank Profits – Unexpected and Unfair?
198 / 2024	Lion Fischer Marc Steffen Rapp Johannes Zahner	Central banks sowing the seeds for a green financial sector? NGFS membership and market reactions
197 / 2023	Tiziana Assenza Alberto Cardaci Michael Haliassos	Consumption and Account Balances in Crises: Have We Neglected Cognitive Load?
196 / 2023	Tobias Berg Rainer Haselmann Thomas Kick Sebastian Schreiber	Unintended Consequences of QE: Real Estate Prices and Financial Stability
195 / 2023	Johannes Huber Alexander Meyer-Gohde Johanna Saecker	Solving Linear DSGE Models With Structure Preserving Doubling Methods
194 / 2023	Martin Baumgärtner Johannes Zahner	Whatever it takes to understand a central banker – Embedding their words using neural networks
193 / 2023	Alexander Meyer-Gohde	Numerical Stability Analysis of Linear DSGE Models – Backward Errors, Forward Errors and Condition Numbers
192 / 2023	Otmar Issing	On the importance of Central Bank Watchers
191 / 2023	Anh H. Le	Climate Change and Carbon Policy: A Story of Optimal Green Macroprudential and Capital Flow Management
190 / 2023	Athanasios Orphanides	The Forward Guidance Trap
189 / 2023	Alexander Meyer-Gohde Mary Tzaawa-Krenzler	Sticky information and the Taylor principle
188 / 2023	Daniel Stempel Johannes Zahner	Whose Inflation Rates Matter Most? A DSGE Model and Machine Learning Approach to Monetary Policy in the Euro Area

187 / 2023	Alexander Dück Anh H. Le	Transition Risk Uncertainty and Robust Optimal Monetary Policy
186 / 2023	Gerhard Rösl Franz Seitz	Uncertainty, Politics, and Crises: The Case for Cash
185 / 2023	Andrea Gubitz Karl-Heinz Tödter Gerhard Ziebarth	Zum Problem inflationsbedingter Liquiditätsrestriktionen bei der Immobilienfinanzierung