

Wallner, Christian; Wystup, Uwe

Working Paper

Efficient computation of option price sensitivities for options of American style

CPQF Working Paper Series, No. 1

Provided in Cooperation with:

Frankfurt School of Finance and Management

Suggested Citation: Wallner, Christian; Wystup, Uwe (2004) : Efficient computation of option price sensitivities for options of American style, CPQF Working Paper Series, No. 1, HfB - Business School of Finance & Management, Centre for Practical Quantitative Finance (CPQF), Frankfurt a. M., <https://nbn-resolving.de/urn:nbn:de:101:1-2009032302>

This Version is available at:

<https://hdl.handle.net/10419/40183>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.

No. 1

**Efficient Computation of Option Price Sensitivities
for Options of American Style**

**Christian Wallner
Uwe Wystup**

July 2004

Author: *Christian Wallner*
Ostpreußenstraße 6 a
21391 Reppenstedt
Germany
Christian.wallner@gmx.net

Prof. Dr. Uwe Wystup
HfB - Business School of
Finance & Management
Frankfurt/Main
Germany
<http://www.mathfinance.de>

Publisher: HfB - Business School of Finance & Management
Phone: +49 (0) 69 154 008-0 ■ Fax: +49 (0) 69 154 008-728
Sonnemannstr. 9-11 ■ D-60314 Frankfurt/M. ■ Germany

Abstract

No front-office software can survive without providing derivatives of option prices with respect to underlying market or model parameters, the so called Greeks. If a closed form solution for an option exists, Greeks can be computed analytically and they are numerically stable. However, for American style options, there is no closed-form solution. The price is computed by binomial trees, finite difference methods or an analytic approximation. Taking derivatives of these prices leads to instable numerics or misleading results, specially for Greeks of higher order. We compare the computation of the Greeks in various pricing methods and conclude with the recommendation to use *Leisen-Reimer trees*.

Keywords: American Options, Greeks, Leisen-Reimer trees.

JEL classification: C63, F31

Contents

1	Introduction	4
1.1	Option Price Sensitivities	4
1.2	Approximation by Finite Difference Quotients	5
2	Computation Methods	5
2.1	Binomial Trees	5
2.1.1	The Method of Leisen and Reimer	6
2.2	Finite Differences	7
2.3	Analytic Approximations	10
2.3.1	Approximation by Barone-Adesi and Whaley	10
3	Comparison of the Methods	12
3.1	Value Function	12
3.2	Delta and Gamma	13
3.3	Theta	13
3.4	Rho (domestic) and Rho (foreign)	13
3.5	Vega and Volga	13
3.6	Vanna	14
4	Summary	14

1 Introduction

We examine which is a suitable method to compute Greeks for American style call and put options in the Black-Scholes model. We choose an exchange rate for the underlying following a geometric Brownian motion,

$$dS_t = S_t[(r_d - r_f)dt + \sigma dW_t], \quad (1)$$

under the risk-neutral measure. As usual r_d denotes the domestic interest rate, r_f the foreign interest rate, σ the volatility. The analysis we do is also applicable to equity options, but we take the foreign exchange market as an example. For contract parameters maturity in years T , strike K and put/call indicator ϕ , which is +1 for a call and -1 for a put, the payoff of the option is

$$[\phi(S_T - K)]^+ = \max[0, \phi(S_T - K)]. \quad (2)$$

We denote by $V(t, x)$ the value of an American style put or call at time t if the spot S_t takes the value x . It is well known (see e.g. Karatzas and Shreve [14]) that in this model the value at time zero is given by

$$V(0, S_0) = \sup_{\tau \in \mathcal{T}} \mathbb{E}[e^{-r_d \tau} [\phi(S_\tau - K)]^+], \quad (3)$$

where $\mathcal{T}$ is the set of all stopping times taking values in $[0, T]$. A closed-form solution for this optimization problem has not yet been found.

1.1 Option Price Sensitivities

Delta	Δ	V_x
Gamma	Γ	V_{xx}
Theta	Θ	V_t
Rho (domestic)	ρ_d	V_{r_d}
Rho (foreign)	ρ_f	V_{r_f}
Vega		V_σ
Volga		$V_{\sigma\sigma}$
Vanna		$V_{x\sigma}$

Table 1: Commonly used Greeks, t is running time and $x = S_0$

Option price sensitivities, the so-called *Greeks* of option values are derivatives with respect to market variables or model parameters. The most commonly used Greeks are listed in Table 1. Numerous relationships and properties of the Greeks for European style options are presented in Reiss and Wystup [19]. Other relevant publications include the work by Carr [7], Broadie and Glasserman [6] in the case of Monte Carlo simulations, Pelsser and Vorst [17] in the case of binomial trees, the work by Eric Benhamou [3] and [4], who uses Malliavin calculus, and the contribution by Rogers and Stapleton [20] using binomial trees with a random number of steps. Joubert and Rogers [13] use a lookup table for a fast, accurate and inelegant valuation of American options. Formulae for Greeks of many exotic foreign exchange options are published in Hakala and Wystup [10].

Order	Difference Quotient
1	$\frac{f(x+h)-f(x)}{h}$
1	$\frac{f(x)-f(x-h)}{h}$
2	$\frac{f(x+h)-f(x-h)}{2h}$

Table 2: Approximation for $\frac{\partial f}{\partial x}$

Order	Difference Quotient
2	$\frac{f(x-h)-2f(x)+f(x+h)}{h^2}$
2	$\frac{2f(x-2h)-f(x-h)-2f(x)-f(x+h)+2f(x+2h)}{14h^2}$
4	$\frac{-f(x-2h)+16f(x-h)-30f(x)+16f(x+h)-f(x+2h)}{12h^2}$

Table 3: Approximation for $\frac{\partial^2 f}{\partial x^2}$

1.2 Approximation by Finite Difference Quotients

We summarize the common methods of numerical differentiation in Tables 2 and 3. For vanna one can use

$$\frac{f(x_i + h, x_k + h) - f(x_i - h, x_k + h) - f(x_i + h, x_k - h) + f(x_i - h, x_k - h)}{4h^2}. \quad (4)$$

2 Computation Methods

2.1 Binomial Trees

The computation of option values with binomial trees was introduced by Cox, Ross and Rubinstein (CRR) [8], where the assumption is used that the log-returns are binomially distributed. It is known that in the limiting case this converges to the continuous Black-Scholes model. Some of the enhancements include Jarrow and Rudd [12], who developed a moment matching method for the parameters. Tian [23] constructed binomial and trinomial trees and showed how to compute the model parameters to obtain weak convergence to the Black-Scholes model in the Lindeberg sense. Hull and White [11] enhanced

the precision of the binomial model using a control variate technique, as it is common in Monte Carlo simulations. Leisen and Reimer [15] modify the parameters of the binomial tree to minimize the oscillating behavior of the value function. We review this technique in the following section.

2.1.1 The Method of Leisen and Reimer

As the convergence of the binomial tree based value to the limit is not monotone but rather oscillatory (see Figure 1), the goal here is to achieve maximum precision with a minimum number of time steps N . However, one can not expect that decreasing the step size $\Delta T = T/N$ will yield a more precise value when using the methods by Cox-Ross-Rubinstein, Tian or Jarrow-Rudd. Leisen and Reimer [15] developed a method in which the parameters u , d and p of the binomial tree can be altered in order to get better convergence behavior.

Instead of choosing the parameters p , u and d to get convergence to the normal distribution Leisen-Reimer suggest to use inversion formulae reverting the standard method – they use normal approximations to determine the binomial distribution $B(n, p)$. In particular, they suggest the following three inversion formulae to replace p (probability of an up move) by $p(d_-)$.

Camp-Paulson-Inversion formula (for arbitrary n)

$$p(z) = \left(\frac{b}{a}\right)^2 \sqrt[3]{\frac{(9a-1)(9b-1) + 3z\sqrt{a(9b-1)^2 + b(9a-1)^2 - 9abz^2}}{(9b-1)^2 - 9bz^2}} \quad (5)$$

with $a = n - j$, $b = j + 1$ and z as input values for the standard normal distribution one uses in the Black-Scholes formula.

Peizer-Pratt-Inversion formula 1 ($n = 2j + 1$)

$$p(z) = \frac{1}{2} + \text{sign}(z) \frac{1}{2} \sqrt{1 - \exp \left[- \left(\frac{z}{n + \frac{1}{3}} \right)^2 \left(n + \frac{1}{6} \right) \right]} \quad (6)$$

Peizer-Pratt-Inversion formula 2 ($n = 2j + 1$)

$$p(z) = \frac{1}{2} + \text{sign}(z) \frac{1}{2} \sqrt{1 - \exp \left[- \left(\frac{z}{n + \frac{1}{3} + \frac{1}{10 \cdot (n+1)}} \right)^2 \left(n + \frac{1}{6} \right) \right]} \quad (7)$$

Then the model parameters are defined by

$$u = e^{(r_d - r_f)\Delta t} \frac{p(d_+)}{p(d_-)}, \quad (8)$$

$$d = \frac{e^{(r_d - r_f)\Delta t} - p(d_-)u}{1 - p(d_-)}, \quad (9)$$

$$d_{\pm} = \frac{\ln \frac{S_0}{K} + (r_d - r_f \pm \frac{1}{2}\sigma)T}{\sigma\sqrt{T}}. \quad (10)$$

Using this method, Leisen and Reimer observe much better convergence behavior.

To compute the Greeks, one can easily use approximations for delta, gamma and theta directly from the tree if the tree satisfies $u = 1/d$, as for example in the CRR model. Let $\Delta T = T/N$ be the step size of an option with maturity T and

$$V_n^i, \quad i = 0, \dots, n,$$

be the value of the option at time $n\Delta T$, $n \leq N$, if the underlying is $S_n^i = Su^i d^{n-i}$. Then the approximations are given by

$$\Delta \approx \frac{V_1^1 - V_1^0}{S(u - d)} \quad (11)$$

$$\Gamma \approx \frac{\frac{V_2^2 - V_2^1}{S(u^2 - 1)} - \frac{V_2^1 - V_2^0}{S(1 - d^2)}}{S(u^2 - d^2)} \quad (12)$$

$$\Theta \approx \frac{V_0^0 - V_2^1}{2\Delta T}. \quad (13)$$

Vega, Volga and the Rhos can be computed using the difference quotients in Tables 2 and 3, Vanna based on Equation (4). Leisen-Reimer trees do not satisfy $u = 1/d$. Nevertheless, delta and gamma can be computed as described above. Theta needs to be determined numerically since at $2\Delta T$ we have $S_2^1 = Sud \neq S$ for the value.

2.2 Finite Differences

The implementation we use for finite differences is essentially based on the *PREMIA2 project* [18] or Andersen and Brotherton-Ratcliff [1].

The value $u(t, X_t = \log(S_t))$ of a European style option in the Black-Scholes model obeys the PDE

$$\begin{cases} \frac{\partial u}{\partial t}(t, x) + \frac{\sigma^2}{2} \frac{\partial^2 u}{\partial x^2}(t, x) + (r_d - r_f - \frac{\sigma^2}{2}) \frac{\partial u}{\partial x}(t, x) - r_d u(t, x) = 0 & \text{in } [0, T) \times \mathbb{R}, \\ u(T, x) = \psi(\exp(x)), \forall x \in \mathbb{R}. \end{cases},$$

where ψ is the payoff at maturity T .

Let $x = \log(S_0)$. Then we let the log spot range in $D \triangleq [x - l, x + l]$ with a suitably chosen l , usually about 3 to 4 standard deviations. We discretize the range using the grid $\{x_i\}$ defined by

$$x_i \triangleq x - l + \frac{2il}{M}, \quad \text{for } 1 \leq i \leq M - 1.$$

We approximate the differential operator

$$A\phi \triangleq \frac{1}{2}\sigma^2 \frac{\partial^2 \phi}{\partial x^2} + (r_d - r_f - \frac{\sigma^2}{2}) \frac{\partial \phi}{\partial x} - r_d \phi$$

by a discrete operator A_h

$$A_h u_h(t, x_i) = \frac{\sigma^2}{2} \frac{\partial^2 u_h}{\partial x^2}(t, x_i) + (r_d - r_f - \frac{\sigma^2}{2}) \frac{\partial u_h}{\partial x}(t, x_i) - r_d u_h(t, x_i),$$

where the functions $u_h(t, \cdot)$ are defined by

$$\begin{aligned} \frac{\partial^2 u_h}{\partial x^2}(t, x_i) &= \frac{1}{h^2} (u_h(t, x_{i+1}) - 2u_h(t, x_i) + u_h(t, x_{i-1})), \\ \frac{\partial u_h}{\partial x}(t, x_i) &= \frac{1}{2h} (u_h(t, x_{i+1}) - u_h(t, x_{i-1})). \end{aligned}$$

Now we determine $u_h(t, x_i)$, ($0 \leq i \leq M$) such that for $0 \leq t \leq T$, $1 \leq i \leq M-1$ the conditions

$$\begin{cases} \frac{d}{dt} u_h(t, x_i) + A_h u_h(t, x_i) = 0, \\ u_h(T, x_i) = \psi(x_i), \\ u_h(t, x-l) = \psi(x-l), \\ u_h(t, x+l) = \psi(x+l) \end{cases} \quad (14)$$

hold. We let $u_h(t) \triangleq (u_h(t, x_1), \dots, u_h(t, x_{M-1}))^T$ and

$$\begin{aligned} \alpha &\triangleq \frac{\sigma^2}{2h^2} - \frac{1}{2h}(r_d - r_f - \frac{\sigma^2}{2}), \\ \beta &\triangleq -\frac{\sigma^2}{h^2} - r_d, \\ \gamma &\triangleq \frac{\sigma^2}{2h^2} + \frac{1}{2h}(r_d - r_f - \frac{\sigma^2}{2}). \end{aligned}$$

Then we can write the operator A_h applied to $u_h(t, \cdot)$ as $A_h u_h(t, \cdot) = M^h u_h(t) + v^h$, where

$$M^h = \begin{bmatrix} \beta & \gamma & 0 & \cdots & 0 & 0 \\ \alpha & \beta & \gamma & 0 & \cdots & 0 \\ 0 & \alpha & \beta & \gamma & \cdots & 0 \\ 0 & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & \alpha & \beta & \gamma \\ 0 & 0 & 0 & \cdots & \alpha & \beta \end{bmatrix}, \quad v^h = \begin{bmatrix} \psi(x-l)\alpha \\ 0 \\ \vdots \\ 0 \\ \psi(x+l)\gamma \end{bmatrix}. \quad (15)$$

For the time-discretization we use the standard- θ -scheme ($\theta \in [0, 1]$). We choose the step size k such that $T = Nk$ and construct the approximation

$$u_{h,k}(t, x) = \sum_{n=0}^N u_h^n(x) \mathbf{1}_{[nk, (n+1)k]}(t),$$

where $u_h^0, \dots, u_h^N$ satisfy the equations

$$\begin{cases} u_h^N = \psi_h, \\ u_h^n(x-l) = \psi(x-l) \quad \text{for } 0 \leq n \leq N-1, \\ u_h^n(x+l) = \psi(x+l) \quad \text{for } 0 \leq n \leq N-1, \\ \frac{u_h^{n+1} - u_h^n}{k} + A_h(u_h^{n+1} + \theta(u_h^n - u_h^{n+1})) = 0 \quad \text{for } 0 \leq n \leq N-1. \end{cases} \quad (16)$$

For $\theta = 0$ we obtain the fully explicit scheme, for $\theta = 1$ the fully implicit scheme and for $\theta = \frac{1}{2}$ the so-called Crank-Nicholson scheme.

In the explicit case $\theta = 0$ the definition of A_h reduces the approximation scheme (16) to

$$\begin{cases} u_h^N = \psi \\ \text{for } 1 \leq n \leq M-1 : \\ u_h^n(x_i) = p_1 u_h^{n+1}(x_{i-1}) + p_2 u_h^{n+1}(x_i) + p_3 u_h^{n+1}(x_{i+1}), \end{cases}$$

where

$$p_1 = k \left(\frac{\sigma^2}{2h^2} - \frac{b}{2h} \right), \quad p_2 = 1 - k \left(r_d + \frac{\sigma^2}{h^2} \right), \quad p_3 = k \left(\frac{\sigma^2}{2h^2} + \frac{b}{2h} \right), \quad (17)$$

and $b = r_d - r_f - \frac{1}{2}\sigma^2$. The scheme is stable if $k \leq \frac{h^2}{\sigma^2 + (r_d - r_f)h^2}$.

In all other cases $1 \geq \theta > 0$ we need to solve a system of linear equations at each time step

$$\mathbf{M}u_{k,h}(jk, \cdot) = \mathbf{N}u_{k,h}((j+1)k, \cdot)$$

where the tri-diagonal matrices $\mathbf{M}$ and $\mathbf{N}$ take the form

$$\begin{pmatrix} b_1 & c_1 & 0 & \cdots & 0 & 0 \\ a_2 & b_2 & c_2 & 0 & \cdots & 0 \\ 0 & a_3 & b_3 & c_3 & \cdots & 0 \\ 0 & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & a_{M-1} & b_{M-1} & c_{M-1} \\ 0 & 0 & 0 & \cdots & a_M & b_M \end{pmatrix}.$$

$\mathbf{M}$ is given by

$$a_i = \theta k \left(\frac{b}{2h} - \frac{\sigma^2}{2h^2} \right), \quad b_i = 1 + \theta k \left(r + \frac{\sigma^2}{h^2} \right), \quad c_i = -\theta k \left(\frac{b}{2h} + \frac{\sigma^2}{2h^2} \right),$$

and $\mathbf{N}$ is given by

$$a_i = (1 - \theta)k \left(\frac{\sigma^2}{2h^2} - \frac{b}{2h} \right), \quad b_i = 1 - (1 - \theta)k \left(r + \frac{\sigma^2}{h^2} \right), \quad c_i = (1 - \theta)k \left(\frac{b}{2h} + \frac{\sigma^2}{2h^2} \right).$$

Solving a system of equations of the kind $\mathbf{M}u = v$, where u and v are M -dimensional vektors can be carried out with the Gauss-Seidel-Factorisation, which is based on the fact that a regular matrix can be decomposed into the product $\mathbf{M} = \mathbf{L}\mathbf{U}$ with a lower triangular matrix $\mathbf{L}$ and an upper triangular matrix $\mathbf{U}$ whose diagonal entries are all equal to 1. The solution of a system of the form $\mathbf{L}\mathbf{U}z = v$ will be done in two steps $\mathbf{L}y = v$, $\mathbf{U}z = y$.

One realizes that if $\mathbf{M}$ is triangular, then $\mathbf{L}$ and $\mathbf{U}$ are triangular as well and hence we only need to find the upper diagonal of $\mathbf{U}$ and the two diagonals of $\mathbf{L}$. The computation of $\mathbf{L}$, $\mathbf{U}$ and v happens in the same step

$$\left| \begin{array}{l} b'_M \triangleq b_M, \quad y_M \triangleq v_M \\ \text{For } 1 \leq i \leq M-1, \text{ } i \text{ increasing:} \\ \quad b'_i = b_i - c_i a_{i+1} / b'_{i+1}, \\ \quad y_i = v_i - c_i y_{i+1} / b'_{i+1}. \\ \\ \quad u_1 = y_1 / b'_1 \\ \text{For } 2 \leq i \leq M, \text{ } i \text{ decreasing:} \\ \quad z_i = (y_i - a_i u_{i-1}) / b'_i. \end{array} \right.$$

Remark. We require the pivot-elements b_i to be non-zero.

To determine the Greeks it appears advantageous to use the information contained in the grid rather than the formulae in Tables 2 and 3 to compute approximations for delta, gamma and theta through

$$\Delta_h = \frac{u_h^0(\exp(x+h)) - u_h^0(\exp(x-h))}{S(e^h - e^{-h})}, \quad (18)$$

$$\Gamma_h = \frac{\frac{u_h^0(\exp(x+h)) - u_h^0(\exp(x))}{S(e^h - 1)} - \frac{u_h^0(\exp(x)) - u_h^0(\exp(x-h))}{S(1 - e^{-h})}}{S(e^h - e^{-h})}, \quad (19)$$

$$\Theta_h = \frac{u_h^k(e^x) - u_h^0(e^x)}{k}. \quad (20)$$

2.3 Analytic Approximations

Since there is no closed form solution available for American style call or put options and the need for fast computation is eminent, several analytic approximations have been developed. However, one needs to be careful using these for the computation of derivatives, as it is well-known that approximating a function does not necessarily imply that the approximation is also a good approximation of the function's derivatives.

2.3.1 Approximation by Barone-Adesi and Whaley

We outline the method to compute the value function for American style options proposed by MacMillan [16] and Barone-Adesi and Whaley [2].

We introduce the notation $v_S = \frac{\partial v}{\partial S}$, $v_{SS} = \frac{\partial^2 v}{\partial S^2}$ and $v_t = \frac{\partial v}{\partial t}$ and X for the strike. Furthermore, we let $V(S, T)$ be the value of an American style options and $v(S, T)$ be the value of a European style option. The values of calls will be denoted by $C(S, T)$ and $c(S, T)$, the values of puts by $P(S, T)$ and $p(S, T)$ respectively. The key idea for the approximation rests on the fact that since the Black-Scholes PDE holds for both the European and the American style option, the early-exercise-premium

$$\varepsilon_C(S, T) = C(S, T) - c(S, T) \quad (21)$$

must satisfy

$$\frac{1}{2}\sigma^2 S^2 \varepsilon_{SS} - r_d \varepsilon + (r_d - r_f) S \varepsilon_S + \varepsilon_t = 0. \quad (22)$$

Using the abbreviations $\tau \triangleq T - t$, $K(\tau) \triangleq 1 - e^{-r_d \tau}$, $M \triangleq \frac{2r_d}{\sigma^2}$, $N \triangleq \frac{2(r_d - r_f)}{\sigma^2}$ and $\varepsilon_C(S, K) \triangleq K(\tau)f(S, K)$ Equation (22) implies

$$S^2 f_{SS} + N S f_S - \frac{M}{K} f - (1 - K) M f_K = 0. \quad (23)$$

The authors now argue that the term $(1 - K) M f_K = 0$ is negligible for small and large τ ¹. The resulting ordinary differential equation

$$S^2 f_{SS} + N S f_S - \frac{M}{K} f = 0 \quad (24)$$

has the general solution

$$f(S) = a_1 S^{q_1} + a_2 S^{q_2}, \quad (25)$$

where the roots of the characteristic polynomial are given by

¹This hints at a weaker quality of the method for medium length maturities.

$$q_{1,2} = \frac{-(N-1) \mp \sqrt{(N-1)^2 + 4\frac{M}{K}}}{2}$$

with $q_1 < 0$ and $q_2 > 0$ since $\frac{M}{K} > 0$. Since $q_1 < 0$, $a_1 \neq 0$ would imply $\lim_{S \rightarrow 0} f(S) = \infty$, whence we must have $a_1 = 0$.

Using equation

$$C(S, \tau) = c(S, \tau) + K a_2 S^{q_2} \quad (26)$$

we can derive restrictions on a_2 , namely

1. for $S = 0$ Equation (26) implies $C(S, T) = 0$.
2. $C(S, \tau)$ must be increasing in S . Therefore, $a_2 > 0$.
3. the r.h. side of (26) must not intersect the line $S - X$, but only touch it in at the optimal exercise level S^* . For $S \leq S^*$ the value of the American call is given by Equation (26). For $S > S^*$ its value is $S - X$.

In order to find S^* we differentiate

$$S^* - X = c(S^*, \tau) + K a_2 (S^*)^{q_2}. \quad (27)$$

with respect to S^* and obtain

$$1 = e^{(b-r_d)\tau} \mathcal{N}(d_1(S^*)) + K q_2 a_2 (S^*)^{q_2}, \quad (28)$$

where $d_1(S^*) = \frac{\ln \frac{S^*}{X} + (b + \frac{\sigma^2}{2})\tau}{\sigma\sqrt{\tau}}$ and $b = r_d - r_f$.

Then one solves Equation (28) for a_2 and plugs the result into Equation (27) to reach

$$S^* - X = c(S^*, \tau) + \frac{1 - e^{(b-r_d)\tau} \mathcal{N}[d_1(S^*)]}{q_2}, \quad (29)$$

where S^* is the only unknown and can be easily found numerically. As a result we get

$$C(S, \tau) = \begin{cases} c(S, \tau) + A_2 \left(\frac{S}{S^*}\right)^{q_2}, & \text{if } S < S^* \\ S - X, & \text{otherwise,} \end{cases} \quad (30)$$

where $A_2 = \frac{(1 - e^{(b-r_d)\tau} \mathcal{N}[d_1(S^*)])}{q_2}$.

Remark: Note that A_2 is only positive if $b < r_d$, i.e. $r_f > 0$, which is usually satisfied. Similarly for puts Equation (21) holds in the form

$$\varepsilon_P(S, T) = P(S, T) - p(S, T). \quad (31)$$

Here in Equation (25) we need $a_2 = 0$ and hence

$$P(S, \tau) = p(S, \tau) + K a_1 S^{q_1}. \quad (32)$$

To get a_1 we employ the optimal exercise level S^{**} defined by

$$X - S^{**} = p(S^{**}, \tau) - \frac{1 - e^{(b-r_d)\tau} \mathcal{N}[-d_1(S^{**})]}{q_1}. \quad (33)$$

The American style put is then approximated by

$$P(S, \tau) = \begin{cases} p(S, \tau) + A_1 \left(\frac{S}{S^{**}}\right)^{q_1}, & \text{if } S > S^{**} \\ X - S, & \text{otherwise,} \end{cases} \quad (34)$$

where $A_1 = -\frac{(1-e^{(b-r_d)\tau})\mathcal{N}[-d_1(S^{**})]}{q_1}$.

3 Comparison of the Methods

Now we compare the efficiency of the different valuation procedures outlined before. We consider a Euro call USD put option with a strike of 0.9000, 3 months maturity. Market data are assumed to be 10% volatility, 3.5% Euro interest rate, 2% USD interest rate. In this scenario, the value of the European and American put are identical, so the European put can be taken as a benchmark for the American style value and Greeks. The value of the American call will be strictly larger than the value of the European call.

The parameters for Leisen-Reimer binomial trees are $N^{bin} = 2000$ time steps, the parameters for the finite differences are $N_S^{fd} = 1130$ spot steps, $N_T^{fd} = 1130$ time steps and $\theta = 0.5$ (Crank-Nicholson). We compare these methods with the approximation by Barone-Adesi and Whaley (BAW) and the Black-Scholes method.

3.1 Value Function

Figure 2 shows the value functions of a call as a function of the current spot. One of the weaknesses of BAW is that the precision can't be improved by changing a parameter.

We observe furthermore in Table 4 that in the BAW method the exercise boundary will be reached too early as compared to the binomial trees and finite differences. The computation of the Greeks will inherit this feature, whence we can't expect high accuracy for the Greeks using BAW near the optimal exercise boundary.

Spot	BS	BT	BAW	FD
0.97	0.06765478	0.07007488	0.0700066	0.07007446
0.971	0.06857433	0.0710532	0.07100095	0.07105275
0.972	0.06949664	0.0720353	0.072	0.07203482
0.973	0.07042164	0.07302112	0.073	0.07302061
0.974	0.07134923	0.07401058	0.074	0.07401004
0.975	0.07227935	0.07500358	0.075	0.07500305
0.976	0.0732119	0.07600002	0.076	0.076
0.977	0.07414681	0.077	0.077	0.077
0.978	0.075084	0.078	0.078	0.078

Table 4: Comparison of the values in USD using the methods near the optimal exercise boundary

The advantage of BAW is its speed, the method is superior if you need to price a vanilla far away from the optimal exercise boundary.

3.2 Delta and Gamma

We take these Greeks directly from the PDE grid or the tree as the information comes at no extra computational cost. Figures 3 and 4 show the expected behavior for BAW: delta approaches 1 to quickly and gamma approaches 0 to quickly. The Leisen-Reimer trees and finite differences yield equally good values for delta and gamma. However, we observed that the values of gamma near the optimal exercise boundary tend to be more stable using Leisen-Reimer trees.

3.3 Theta

In BAW and LR, theta has to be computed with difference quotients, in the finite differences we can take it from the grid. This implies about twice or trice the computational cost for trees depending on whether we use $\frac{1}{h}(f(x) - f(x - h))$ or $\frac{1}{2h}(f(x + h) - f(x - h))$. The accuracy of LR and finite differences appears identical (see Figure 5), so we would recommend finite differences to save on the computational cost.

The following Greeks can only be determined using the difference quotients in Tables 2 and 3. The choice of the parameter h is crucial. If we choose it too small, then the lack of precision in the value function will lead to a possibly larger error in the hedge parameter.

3.4 Rho (domestic) and Rho (foreign)

We compute these Greeks with the difference formulae of first order to keep the computation cost under control. In particular, we only need one more computation of $r_d - h$ or $r_f + h$ to compute the approximations

$$\rho_d \approx \frac{1}{h}(f(r_d) - f(r_d - h)) \quad \text{and} \quad (35)$$

$$\rho_f \approx \frac{1}{h}(f(r_f + h) - f(r_f)). \quad (36)$$

We take $h = 0.01$. Figures 7 and 8 show the approximations for ρ_d and ρ_f . Even for step size of $N^{bin} = 100$ in the binomial tree we obtain good approximations for these Greeks. Finite differences tend to oscillate for this grid size as illustrated in Figure 6, so that we would recommend LR trees for the rhos.

3.5 Vega and Volga

For the second derivative we need to choose an approximation from Table 3. Noticeably, the approximations of second order work better than the one of order four. We had specially good experience using

$$\text{Volga} \approx \frac{2f(x - 2h) - f(x - h) - 2f(x) - f(x + h) + 2f(x + 2h)}{14h^2}. \quad (37)$$

with $h = 0.05$. A smaller h would require more computational cost for the trees and finite differences, since the resulting value function would have to have higher precision. For small step size we find the precision of the LR trees superior to the finite differences, as illustrated in Figure 9 with $N^{bin} = N_T^{fd} = N_S^{fd} = 100$. Therefore we recommend LR trees for volga. To compute vega it is advisable to use the values already obtained for volga, i.e.

$$\text{Vega} \approx \frac{f(x - 2h) - f(x + 2h)}{4h}. \quad (38)$$

For small step sizes we find similar behavior for the vega as we found for volga. Therefore, we recommend LR trees to compute vega. The results are displayed in Figures 10 and 11.

3.6 Vanna

We approximate vanna using the second order derivative

$$\frac{f(S + h_S, \sigma + h_v) - f(S - h_S, \sigma + h_v) - f(S + h_S, \sigma - h_v) + f(S - h_S, \sigma - h_v)}{4h_S h_v}, \quad (39)$$

where h_S denotes the step size of the spot S and h_v the step size of the volatility σ . We take $h_S = 0.003$ and $h_v = 0.03$. Compared to the other Greeks, vanna requires very high accuracy in the finite difference based method, to avoid oscillatory behavior as illustrated in Figure 12. LR trees turn out to be the obviously better method here, although we need a fine grid near the optimal exercise boundary.

4 Summary

We analyzed three commonly used methods to determine the value of American style options with regard to their efficiency to compute the hedge parameters (Greeks), in particular: delta, gamma, theta, vega, vanna, volga, domestic and foreign rho. These were the analytic approximation by Barone-Adesi and Whaley, the finite difference method with Crank-Nicholson scheme and the binomial model in the variant of Leisen and Reimer.

The method by Barone-Adesi and Whaley is working with a fixed and non-improvable precision. Moreover, it lacks precision near the optimal exercise boundary. Its only strength lies in its speed.

We confirmed that using finite differences will deliver approximations for delta, gamma and theta directly from the grid without additional computational cost. Except for theta we obtain the same result for the binomial trees. Leisen Reimer trees yield more precise results for delta and gamma.

The remaining Greeks can not be taken from the grid, but have to be computed using finite difference quotients. We observed that Leisen Reimer trees are the superior method.

References

- [1] ANDERSEN, L. and BROTHERTON-RATCLIFFE, R. (1998). The Equity Option Volatility Smile - An Implicit Finite-Difference Approach. *The Journal of Computational Finance*, Vol. 1, No. 2
- [2] BARONE-ADESI, G. and WHALEY, R. E. (1987). Efficient Analytic Approximation of American Option Values. *Journal of Finance*, **42**pp. 301-320.
- [3] BENHAMOU, E. (2003). Optimal Malliavin Weighting Function for the Computation of the Greeks. *Mathematical Finance*, Volume 13, Issue 1.
- [4] BENHAMOU, E. (2004). Efficient Computation of Greeks for Discontinuous Payoffs by Transformation of the Payoff Function. *The Journal of Computational Finance*.
- [5] BLACK, F. and Scholes, M. (1973). The Pricing of Options and Other Corporate Liabilities. *Journal of Political Economy*, **81** pp. 637-654.
- [6] BROADIE, M., and GLASSERMAN, P. (1996). Estimating Security Price Derivatives using Simulation, *Management Science* **42**, 269-285.
- [7] CARR, P. (2001). Deriving Derivatives of Derivative Securities. *The Journal of Computational Finance*. Vol 4, Number 2, pp. 5-29.
- [8] COX, J. C., ROSS, S. A. and RUBINSTEIN, M. (1979). Option Pricing: A Simplified Approach. *Journal of Financial Economics*, **7** pp. 229-263.

- [9] GLASSERMAN, P., and ZHAO, X. (1999). Fast Greeks by Simulation in Forward LIBOR Models, *The Journal of Computational Finance*, Vol 3, Number 1, 5–39.
- [10] HAKALA, J. and WYSTUP, U. (2002). *Foreign Exchange Risk*. Risk Publications, London.
- [11] HULL, J. C. and WHITE, A. (1993). The Accelerated Binomial Option Pricing Modell. Working Paper. Faculty of Management University of Toronto.
- [12] JARROW, R. and RUDD, M. (1983). Option Pricing. *Homewood*, Illinois, pp. 183-188.
- [13] JOUBERT, A. and ROGERS, L.C.G. (1997). Fast, accurate and inelegant valuation of American options. In *Numerical Methods in Finance*, eds. L. C. G. Rogers & D. Talay, 88–92, Cambridge University Press.
- [14] KARATZAS, I. and SHREVE, S.E. (1998). *Methods of Mathematical Finance*. Springer.
- [15] LEISEN, D. and REIMER, M. (1996). Binomial Models For Option Valuation - Examining and Improving Convergence. *Applied Mathematical Finance* **3** pp. 319-346.
- [16] MACMILLAN, L. W. (1986). Analytic Approximation for the American Put Option. *Advances in Futures and Options Research*, **1** pp. 119-139.
- [17] PELSSER, A and VORST, T. (1994). The Binomial Model and the Greeks. *Journal of Derivatives*, pp. 45-48. Spring 1994.
- [18] The PREMIA2 project, see <http://www-rocq.inria.fr/mathfi/Premia/index.html>.
- [19] REISS, O. and WYSTUP, U. (2001). Efficient Computation of Option Price Sensitivities Using Homogeneity and other Tricks. *Journal of Derivatives* Vol. 9 No. 2.
- [20] ROGERS, L.C.G. and STAPLETON, J. (1998). Fast accurate binomial pricing of options. *Finance and Stochastics* **2**, 3–17.
- [21] SHREVE, S.E. (1996). *Stochastic Calculus and Finance*. Lecture notes, Carnegie Mellon University
- [22] TALEB, N. (1996). *Dynamic Hedging*. Wiley, New York.
- [23] TIAN, Y. (1993). A Modified Lattice Approach to Option Pricing. *Journal of Future Markets*, Vol. 13, No. 5 pp. 563-577.
- [24] WALLNER, C. (2002). *Effiziente Berechnung von Optionspreissensitivitäten für Optionen Amerikanischen Typs*. Diplomarbeit. Philipps-Universität Marburg, Department of Mathematics and Computer Science.
- [25] WYSTUP, U. (1999). *Vanilla Options*. Formula Catalogue of <http://www.MathFinance.de>.




Figure 1: Convergence Behavior of Tree Models with parameters $S = .81$, $K = .9$, $T = 1$, $r_d = .02$, $r_f = .035$, $\sigma = .3$




Figure 2: American call with $K = .9$, $T = 3m$, $r_d = .02$, $r_f = .035$, $\sigma = .1$




Figure 3: Delta of an American call with $K = .9$, $T = 3m$, $r_d = .02$, $r_f = .035$, $\sigma = .1$




Figure 4: Gamma of an American call with $K = .9$, $T = 3m$, $r_d = .02$, $r_f = .035$, $\sigma = .1$




Figure 5: Theta of an American call with $K = .9$, $T = 3m$, $r_d = .02$, $r_f = .035$, $\sigma = .1$




Figure 6: Behavior for the domestic rho of an American call with $N^{bin} = N_t^{fd} = N_S^{fd} = 100$




Figure 7: Rho (domestic) of an American call with $K = .9$, $T = 3m$, $r_d = .02$, $r_f = .035$, $\sigma = .1$




Figure 8: Rho (foreign) of an American call with $K = .9$, $T = 3m$, $r_d = .02$, $r_f = .035$, $\sigma = .1$




Figure 9: Behavior of an American call volga with $N^{bin} = N_T^{fd} = N_S^{fd} = 100$




Figure 10: Vega of an American call with $K = .9$, $T = 3m$, $r_d = .02$, $r_f = .035$, $\sigma = .1$




Figure 11: Volga of an American call with $K = .9$, $T = 3m$, $r_d = .02$, $r_f = .035$, $\sigma = .1$




Figure 12: Behavior of an American call vanna with $N^{bin} = N_T^{fd} = N_S^{fd} = 100$




Figure 13: Vanna of an American call with $K = .9$, $T = 3m$, $r_d = .02$, $r_f = .035$, $\sigma = .1$

Working Paper Series of HfB - Business School of Finance & Management: Centre for Practical Quantitative Finance

Previously released

No.	Author/Title	Year
01.	Wallner, Christian / Wystup, Uwe Efficient Computation of Option Price Sensitivities for Options of American Style	2004

Printed edition: € 25,-- +. € 2,50 shipping

Download: <http://www.hfb.de/Navigator/Fakultaet/Publikationen/Arbeitberichte/Show>

Working Paper Series of HfB - Business School of Finance & Management

Previously released

No.	Author/Title	Year
53.	Polleit, Thorsten The Slowdown in German Bank Lending - Revisited	2004
52.	Heidorn, Thomas / Siragusano, Tindaro Die Anwendbarkeit der Behavioral Finance im Devisenmarkt	2004
51.	Schütze, Daniel / Schallast, Christoph (Hrsg.) Wider die Verschleuderung von Unternehmen durch Pfandversteigerung	2004
50.	Gerhold, Mirko / Heidorn, Thomas Investitionen und Emissionen von Convertible Bonds (Wandelanleihen)	2004
49.	Chevalier, Pierre / Heidorn, Thomas / Krieger, Christian Temperaturderivate zur strategischen Absicherung von Beschaffungs- und Absatzrisiken	2003
48.	Becker, Gernot M. / Seeger, Norbert Internationale Cash Flow-Rechnungen aus Eigner- und Gläubigersicht	2003
47.	Boenkost, Wolfram / Schmidt, Wolfgang M. Notes on convexity and quanto adjustments for interest rates and related options	2003
46.	Hess, Dieter Determinants of the relative price impact of unanticipated Information in U.S. macroeconomic releases	2003
45.	Cremers, Heinz / Kluß, Norbert / König, Markus Incentive Fees. Erfolgsabhängige Vergütungsmodelle deutscher Publikumsfonds	2003
44.	Heidorn, Thomas / König, Lars Investitionen in Collateralized Debt Obligations	2003
43.	Kahlert, Holger / Seeger, Norbert Bilanzierung von Unternehmenszusammenschlüssen nach US-GAAP	2003
42.	Beiträge von Studierenden des Studiengangs BBA 012 unter Begleitung von Prof. Dr. Norbert Seeger Rechnungslegung im Umbruch - HGB-Bilanzierung im Wettbewerb mit den internationalen Standards nach IAS und US-GAAP	2003
41.	Overbeck, Ludger / Schmidt, Wolfgang Modeling Default Dependence with Threshold Models	2003
40.	Balthasar, Daniel / Cremers, Heinz / Schmidt, Michael Portfoliooptimierung mit Hedge Fonds unter besonderer Berücksichtigung der Risikokomponente	2002
39.	Heidorn, Thomas / Kantwill, Jens Eine empirische Analyse der Spreadunterschiede von Festsatzanleihen zu Floatern im Euroraum und deren Zusammenhang zum Preis eines Credit Default Swaps	2002
38.	Böttcher, Henner / Seeger, Norbert Bilanzierung von Finanzderivaten nach HGB, EstG, IAS und US-GAAP	2003
37.	Moormann, Jürgen Terminologie und Glossar der Bankinformatik	2002

36.	Heidorn, Thomas Bewertung von Kreditprodukten und Credit Default Swaps	2001
35.	Heidorn, Thomas / Weier, Sven Einführung in die fundamentale Aktienanalyse	2001
34.	Seeger, Norbert International Accounting Standards (IAS)	2001
33.	Stehling, Frank / Moormann, Jürgen Strategic Positioning of E-Commerce Business Models in the Portfolio of Corporate Banking	2001
32.	Strohhecker, Jürgen / Sokolovsky, Zbynek Fit für den Euro, Simulationsbasierte Euro-Maßnahmenplanung für Dresdner-Bank-Geschäftsstellen	2001
31.	Roßbach, Peter Behavioral Finance - Eine Alternative zur vorherrschenden Kapitalmarkttheorie?	2001
30.	Heidorn, Thomas / Jaster, Oliver / Willeitner, Ulrich Event Risk Covenants	2001
29.	Biswas, Rita / Löchel, Horst Recent Trends in U.S. and German Banking: Convergence or Divergence?	2001
28.	Löchel, Horst / Eberle, Günter Georg Die Auswirkungen des Übergangs zum Kapitaldeckungsverfahren in der Rentenversicherung auf die Kapitalmärkte	2001
27.	Heidorn, Thomas / Klein, Hans-Dieter / Siebrecht, Frank Economic Value Added zur Prognose der Performance europäischer Aktien	2000
26.	Cremers, Heinz Konvergenz der binomialen Optionspreismodelle gegen das Modell von Black/Scholes/Merton	2000
25.	Löchel, Horst Die ökonomischen Dimensionen der 'New Economy'	2000
24.	Moormann, Jürgen / Frank, Axel Grenzen des Outsourcing: Eine Exploration am Beispiel von Direktbanken	2000
23.	Heidorn, Thomas / Schmidt, Peter / Seiler, Stefan Neue Möglichkeiten durch die Namensaktie	2000
22.	Böger, Andreas / Heidorn, Thomas / Graf Waldstein, Philipp Hybrides Kernkapital für Kreditinstitute	2000
21.	Heidorn, Thomas Entscheidungsorientierte Mindestmargenkalkulation	2000
20.	Wolf, Birgit Die Eigenmittelkonzeption des § 10 KWG	2000
19.	Thiele, Dirk / Cremers, Heinz / Robé, Sophie Beta als Risikomaß - Eine Untersuchung am europäischen Aktienmarkt	2000
18.	Cremers, Heinz Optionspreisbestimmung	1999
17.	Cremers, Heinz Value at Risk-Konzepte für Marktrisiken	1999
16.	Chevalier, Pierre / Heidorn, Thomas / Rütze, Merle Gründung einer deutschen Strombörse für Elektrizitätsderivate	1999
15.	Deister, Daniel / Ehrlicher, Sven / Heidorn, Thomas CatBonds	1999
14.	Jochum, Eduard Hoshin Kanri / Management by Policy (MbP)	1999
13.	Heidorn, Thomas Kreditderivate	1999
12.	Heidorn, Thomas Kreditrisiko (CreditMetrics)	1999
11.	Moormann, Jürgen Terminologie und Glossar der Bankinformatik	1999
10.	Löchel, Horst The EMU and the Theory of Optimum Currency Areas	1998
09.	Löchel, Horst Die Geldpolitik im Währungsraum des Euro	1998

08.	Heidorn, Thomas / Hund, Jürgen Die Umstellung auf die Stückaktie für deutsche Aktiengesellschaften	1998
07	Moormann, Jürgen Stand und Perspektiven der Informationsverarbeitung in Banken	1998
06.	Heidorn, Thomas / Schmidt, Wolfgang LIBOR in Arrears	1998
05.	Jahresbericht 1997	1998
04.	Ecker, Thomas / Moormann, Jürgen Die Bank als Betreiberin einer elektronischen Shopping-Mall	1997
03.	Jahresbericht 1996	1997
02.	Cremers, Heinz / Schwarz, Willi Interpolation of Discount Factors	1996
01.	Moormann, Jürgen Lean Reporting und Führungsinformationssysteme bei deutschen Finanzdienstleistern	1995

Printed edition: € 25,-- +. € 2,50 shipping

Download: <http://www.hfb.de/Navigator/Fakultaet/Publikationen/Arbeitberichte/Show>

Special Working Paper Series of HfB - Business School of Finance & Management

Previously released

No.	Author/Title	Year
01.	Kahmer, Nicole / Moormann, Jürgen Studie zur Ausrichtung von Banken an Kundenprozessen am Beispiel des Internet (Fee: € 120,--)	2003

Order address / contact:

HfB ■ Business School of Finance & Management
Sonnemannstr. 9-11 ■ D-60314 Frankfurt/M. ■ Germany
Phone: +49 (0) 69 154 008-734 ■ Fax: +49 (0) 69 154 008-728
eMail: johannsen@hfb.de

Further informations about HfB ■ Business School of Finance & Management may be obtained at: <http://www.hfb.de>