

NOTA DI LAVORO

41.2010

Scaling up Ecosystem Services Values: Methodology, Applicability and a Case Study

By **Luke Brander**, Institute for
Environmental Studies, Amsterdam, the
Netherlands

Andrea Ghermandi, FEEM

Onno Kuik, Institute for Environmental
Studies, Amsterdam, the Netherlands

Anil Markandya, BC3 Basque Centre
for Climate Change, Bilbao, Spain

Paulo A.L.D. Nunes, FEEM, Università
Ca' Foscari di Venezia, Venice, Italy

Marije Schaafsma and **Alfred
Wagtendonk**, Institute for
Environmental Studies, Amsterdam, the
Netherlands

SUSTAINABLE DEVELOPMENT Series

Editor: Carlo Carraro

Scaling up Ecosystem Services Values: Methodology, Applicability and a Case Study

By Luke Brander, Institute for Environmental Studies, Amsterdam, the Netherlands

Andrea Ghermandi, FEEM

Onno Kuik, Institute for Environmental Studies, Amsterdam, the Netherlands

Anil Markandya, BC3 Basque Centre for Climate Change, Bilbao, Spain

Paulo A.L.D. Nunes, FEEM, Università Ca' Foscari di Venezia, Venice, Italy

Marije Schaafsma and Alfred Wagtendonk, Institute for Environmental Studies, Amsterdam, the Netherlands

Summary

The approach of using existing data on economic values of local ecosystem services for an assessment of these values at a larger geographical scale can be called “scaling up”. In a scaling-up exercise, economic values from a particular study site are transferred to another geographical setting, for instance to the regional, national or global scale. This paper proposes a methodology for scaling up ecosystem service values to a European level, assesses the availability of data for conducting this method, and illustrates the procedure with a case study on wetland values. The proposed methodology makes use of meta-analysis to produce a value function that is subsequently applied to individual European wetland sites. Site-specific, study-specific and context-specific variables are used to define a price vector that captures differences between sites and over time. The proposed method is shown to be practicable and to produce reasonably reliable aggregate value estimates.

Keywords: Ecosystem Services, Value Transfer, Meta-Analysis, Wetland Values

JEL Classification: C81, Q24, Q57

The authors wish to thank the European Environment Agency (EEA) for making this research possible. The researchers have also greatly benefited from discussions and cooperation with EEA staff, especially with the research coordinator Mr. Hans Vos. The usual disclaimers apply.

Address for correspondence:

Luke Brander
Institute for Environmental Studies
De Boelelaan 1087
1081 HV Amsterdam
The Netherlands
E-mail: luke.brand@ivm.vu.nl

Scaling up ecosystem services values: methodology, applicability and a case study

Luke Brander ^{1*}, Andrea Ghermandi ², Onno Kuik ¹, Anil Markandya ⁴, Paulo A.L.D. Nunes ^{2,3}, Marije Schaafsma ¹, Alfred Wagtendonk ¹

Abstract

The approach of using existing data on economic values of local ecosystem services for an assessment of these values at a larger geographical scale can be called “scaling up”. In a scaling-up exercise, economic values from a particular study site are transferred to another geographical setting, for instance to the regional, national or global scale. This paper proposes a methodology for scaling up ecosystem service values to a European level, assesses the availability of data for conducting this method, and illustrates the procedure with a case study on wetland values. The proposed methodology makes use of meta-analysis to produce a value function that is subsequently applied to individual European wetland sites. Site-specific, study-specific and context-specific variables are used to define a price vector that captures differences between sites and over time. The proposed method is shown to be practicable and to produce reasonably reliable aggregate value estimates.

Key words: ecosystem services, value transfer, meta-analysis, wetland values

JEL Classification: C81, Q24, Q57

¹ Institute for Environmental Studies, Amsterdam, the Netherlands

² Fondazione ENI Enrico Mattei, Milan, Italy

³ Università Ca' Foscari di Venezia, Venice, Italy

⁴ BC3 Basque Centre for Climate Change, Bilbao, Spain

* Corresponding author: luke.brander@ivm.vu.nl. The authors wish to thank the European Environment Agency (EEA) for making this research possible. The researchers have also greatly benefited from discussions and cooperation with EEA staff, especially with the research coordinator Mr. Hans Vos. The usual disclaimers apply.

1. Introduction

The approach of using existing data on economic values of local ecosystem services for an assessment of these values at a larger geographical scale can be called “scaling up”. In a scaling-up exercise, economic values from a particular study site are transferred to another geographical setting, for instance to the regional, national or global scale. Local values are thus not applied in another local context, but are used to estimate the values of all ecosystems (or ecosystem services) of similar characteristics in a larger region.

Scaling up builds on the methods and tools that have been developed for value transfer, and can be seen as an extension of value transfer. Value transfer is usually applied on a case-by-case basis. The transfer of economic values of individual ecosystem services from a particular study site to another – but similar – site (the policy site) has become a common tool in ecosystem assessment. In the scaling-up exercise, economic values from a particular study site (or sites) are extrapolated to a larger geographical setting.

Spatial scale is recognised as an important issue to the valuation of ecosystem services (Hein et al., 2006). The spatial scales at which ecosystem services are supplied and demanded contribute to the complexity of ecosystem valuation and management. On the supply-side, ecosystems themselves vary in spatial scale (e.g. small individual patches, large continuous areas, regional networks) and provide services at varying spatial scales. The services that ecosystems provide can be both on- and off-site. For example, a forest might provide recreational opportunities (on-site), downstream flood prevention (local off-site), and climate regulation (global off-site). On the demand-side, beneficiaries of ecosystem services also vary in terms of their locational distribution. The spatial scale over which ecosystem services are provided and received is determined by the spatial scale over which an ecosystem function has effect and the spatial scale of (potential) beneficiaries. For conceptualising the relationship between the supply and demand of ecosystem services one might imagine two overlaid maps – one representing the spatial extents of an ecosystem and the (potential) services it provides, and the other representing the spatial location of the (potential) beneficiaries of these services. It is important to recognise that ecosystem services result from the interaction of ecosystem functions and human activities. An ecosystem does not provide a service if no-one makes use of its potential to provide that service.

Ecosystem services often have different groups of beneficiaries (different in terms of spatial location and socio-economic characteristics). For example, the provision of recreational opportunities by an ecosystem will generally only benefit people in the immediate vicinity, whereas the existence of a high level of biodiversity may be valued by people at a much larger spatial scale. Differences in the size and characteristics of groups of beneficiaries per ecosystem service need to be taken into account in aggregating values for each service. The management of ecosystems may be further complicated in cases where the interests of different groups of beneficiaries (possibly at different spatial scales) are in conflict. This may occur when ecosystem services are mutually exclusive (e.g. timber extraction and carbon sequestration).

The values held by beneficiaries for ecosystem services may vary with a number of different factors that can be spatially defined (distance, availability of substitute and complementary sites, income, culture, and preferences). Use values are generally ex-

pected to decline with distance to an ecosystem – so called distance decay. Non-use values may also decline with distance between the ecosystem and beneficiary, although this relationship may be less related to distance than to cultural or political boundaries. The availability of substitute (complementary) sites within the vicinity of a selected ecosystem is expected to reduce (increase) the value of ecosystem services from that ecosystem. Socio-economic characteristics of beneficiaries (e.g. income, culture, and preferences) are not spatial variables per se, but differences in these variables between (groups of) beneficiaries can often be usefully defined in a spatial manner (e.g. by administrative area, region or country).

Consideration of the spatial scale of the provision and beneficiaries of ecosystem services is important for the calculation of the total economic value of these services (i.e. the aggregation of values across relevant areas and populations). In addition, accounting for spatial scale may be of further use in the formulation of policies to manage ecosystem services, for example in the identification of winners and losers, the need for compensation/incentives, and the design of policies such as payments for environmental services.

Regarding the estimation of ecosystem service values, there are a number of important issues to be considered related to spatial scale. In discussing these scale related issues we make a distinction between the estimation of values for an individual ecosystem site and for the entire stock of an ecosystem within a large geographic area. We have referred to the latter case as ‘scaling up’ ecosystem values when insufficiency of data requires applying value transfer methods.

At the level of an individual ecosystem site, unit values for ecosystem services are likely to vary with the characteristics of the ecosystem site (area, integrity, and type of ecosystem), beneficiaries (number, income, preferences), and context (availability of substitute and complementary sites and services). All of these variables have a spatial dimension that can be accounted for in estimating site-specific values. For example, in terms of ecosystem area, many ecosystem service values have been observed to exhibit diminishing returns to scale (i.e. adding an additional unit of area to a large ecosystem increases the total value of ecosystem services less than an additional unit of area to a smaller ecosystem). It is therefore important to account for the size of the ecosystem being valued.

For scaling up ecosystem values to estimate the total economic value of a change in the stock of ecosystems in a large geographic area, in addition to controlling for other spatial variables, it is necessary to account for the non-constancy of marginal values across the stock of an ecosystem. At the margin, a small change in ecosystem service provision (e.g. the loss of a small area) will not affect the value of services from other ecosystem sites. Non-marginal changes in ecosystem service provision, however, will affect the value of services from the remaining stock of ecosystems. As the ecosystem service becomes scarcer, its marginal and average values will tend to increase. This means that simply multiplying a constant per unit value by the total quantity of ecosystem service provision is likely to (substantially) underestimate the total value of a negative change. Appropriate adjustments to marginal values to account for large-scale changes in ecosystem service provision need to be made.

This paper discusses methods for scaling up existing estimates of ecosystem services' values to larger geographical scales (e.g., the European scale), and illustrates the meta-analytical value transfer method with a case study. Section 2 surveys the literature on value transfer methods as important building blocks for scaling-up applications. Section 3 discusses the practicability of methods for large-scale scaling up exercises. Section 4 illustrates the meta-analytical value transfer method by a case study on the valuation of European wetlands. Section 5 concludes.

2. Methods for value transfer

Value transfer is the procedure of estimating the value of an ecosystem (or goods and services from an ecosystem) by borrowing an existing valuation estimate for a similar ecosystem. The ecosystem of current policy interest is often called the "policy site" and the ecosystem from which the value estimate is borrowed is called the "study site". This procedure is often termed benefit transfer but since the values being transferred may also be estimates of costs or damages, the term value transfer is arguably more appropriate.

The use of value transfer to provide information for decision making has a number of advantages over conducting primary research to estimate ecosystem values. From a practical point of view it is generally less expensive and time consuming than conducting primary research. Value transfer can also be applied on a scale that would be unfeasible for primary research in terms of valuing large numbers of sites across multiple countries. Value transfer also has the methodological attraction of providing consistency in the estimation of values across policy sites.

Value transfer methods can be divided into four categories:

1. Unit value transfer;
2. Adjusted unit value transfer;
3. Value function transfer; and
4. Meta-analytic function transfer.

Unit value transfer involves estimating the value of an environmental good or service at a policy site by multiplying a mean unit value estimated at a study site by the quantity of that good or service at the policy site. Units values are generally either expressed as values per household or as values per unit of area. In the former case, aggregation of values is over the relevant population that hold values for the ecosystem in question. In the latter case, aggregation of values is over the area of the ecosystem.

Adjusted unit transfer involves making simple adjustments to the transferred unit values to reflect differences in site characteristics. The most common adjustments are for differences in income between study and policy sites and for differences in price levels over time or between sites.

Value or demand function transfer methods use functions estimated through valuation applications (travel cost, hedonic pricing, contingent valuation, or choice modelling) for a study site together with information on parameter values for the policy site to transfer values. Parameter values of the policy site are plugged into the value function to calculate a transferred value that better reflects the characteristics of the policy site.

Meta-analytic function transfer uses a value function estimated from multiple study results together with information on parameter values for the policy site to estimate values. The value function therefore does not come from a single study but from a collection of studies. This allows the value function to include greater variation in both site characteristics (e.g. socio-economic and physical attributes) and study characteristics (e.g. valuation method) that cannot be generated from a single primary valuation study. Rosenberger and Phipps (2007) identify the important assumptions underlying the use of meta-analytic value functions for value transfer:

1. There exists an underlying meta-valuation function that relates estimated values of a resource to site and study characteristics. Primary valuation studies provide point estimates on this underlying function that can subsequently be used in meta-analysis to estimate it;
2. Differences between sites can be captured through a price vector;
3. Values are stable over time, or vary in a systematic way; and
4. The sampled primary valuation studies provide “correct” estimates of resource value.

2.1 Markets for ecosystem services and distance decay effects

The distance between a person and an environmental good can be an important explanatory variable of this person’s willingness to pay (WTP) for that good. Transferring average WTP values from a study site where the relevant population is located close to the site to a policy site where the population lives much further away is likely to lead to overestimation of total WTP. Since the distribution of the population is likely to differ between the policy and study site, average distances between individuals and both sites are different, and value transfer studies should account for these differences.

Based on economic theory, the effect of distance on WTP is expected to be negative, indicating a distance-decay effect. Distance-decay (DD) implies that the WTP for a certain site decreases as the distance from the agent to the site increases. In other words, use values are expected to be decreasing with distance, because the cost of visiting a site increases with every kilometre one has to travel. The higher the distance, the higher are the costs, the lower the demand³. One of the main reasons to in-

³ Other tourism studies state that a longer journey does not necessarily create extra costs, as the trip itself can be enjoyed. Furthermore, a large distance is sometimes considered to be a positive character-

clude this distance-decay effect is to determine the size of the geographical boundaries (market size) of the environmental good in question. This relevant market is the population over which the willingness to pay (WTP) values can be aggregated to calculate the good's Total Economic Value.

Besides direct use values, non-use values are an important component of the Total Economic Value of any environmental good. The importance of distance for reliable value transfer or aggregation therefore depends on the type of value that a study site generates. There is no reason within standard economic theory why non-use values would also decrease with distance. The spatial discounting literature states that values that relate to what economists call non-use values (such as intrinsic and future values) should have much lower discount rates than use values (such as recreational, subsistence, therapeutic and aesthetic values) (Brown et al., 2002). The extent to which distance is important for reliable value transfer therefore also depends on the type of values generated by the study and policy sites.

Other cases in which a distance-decay effect is less likely to occur are for goods that have importance on a large scale. In this case the distance decay effects are likely to be very small or negligible, meaning that even very far from the site, people are willing to pay. The fact that something is either of national importance, of symbolic meaning or has the status of national park implies that (a) there are likely to be fewer substitutes leading to a protection status, or (b) that knowledge about the site is widely spread. Loomis (1996, 2000) find a low DD-effect for salmon, a symbolic species, and Pate and Loomis (1997) do not find any DD-effect at all for a National Park. On the other hand, whenever goods have a local importance due to some cultural association with the good, willingness to pay is likely to fall beyond that political or social border. Examples are distance-decay effects found for "local" goods, suggested to be due to a "sense of ownership" (Bateman et al., 2004) or "spatial identity" (Hanley et al., 2003).

For non-unique sites, such as a lake in a lake district, the availability of substitutes increases with distance, lowering the WTP for one particular site. As the distance to a site increases, the number of available substitutes is likely to increase as well – especially for local goods. However, substitution effects alone cannot always entirely explain DD-effects.

Distance can be specified in many different ways and for reliable transfer or aggregation, the specification should be consistent. Approaches differ in (a) objective versus perceptual or subjective distance; and (b) a straight line (as the crow flies) or based on the road net/travel distance, using more sophisticated GIS applications. Travel cost studies typically use GIS based distance calculations, assuming that people minimize their costs by choosing the shortest route. However, for non-use values, which form a large share of many environmental goods (Oglethorpe and Miliadou,

istic of a destination, as travellers associate a far away location with relaxation and 'being away far from busy day to day life' or a more adventurous trip.

2000; Kniivilä 2005), the least cost travel route does not matter and other specifications might be reliable. Another issue is to which part of the asset the distance should be measured. Ideally, the distance from individual A to the nearest access point of a site should be used for use-values. However, the larger the study area, the more difficult it becomes to determine the distance.

2.2 Substitute and complementary sites

One of the most important contextual factors in a value transfer exercise is the availability of substitutes. Ignoring substitutes means that if the transfer is performed between a landscape poor in ecosystem services to a landscape rich in ecosystem services WTP values are likely to be overestimated (Bateman et al., 1999). The question is what happens to the WTP for good A if the quality in a comparable good B increases. A substitution effect in economics is usually defined as the increase of demand for good A when the price of good B increases.

The consequence of disregarding substitutes is generally an overestimation of WTP, as the sum of the value of goods measured individually is higher than the value measured for all goods at once. For instance, respondents in an area with several lakes whose water quality is polluted will value cleaning up the first lake more than cleaning up the second lake, because (1) the first lake can be a substitute for the second lake, and (2) the respondent has a budget limitation which reduces the money available for cleaning up the second lake. Valuing goods separately and then adding up the values will overstate the true value, as every respondent will treat the ecosystem under study as if it were the first good.

Disregarding complementary sites causes underestimation of WTP. Complementarity occurs when goods are consumed jointly, for instance when two sites are visited during the same trip, or when there are synergy-effects in production, for instance when quality increases at one site automatically increase the quality of another site due to dependent ecosystems. The WTP of one site is therefore likely to be dependent on other available alternatives and their characteristics. As distance from the site or the geographical scale of the study increases, the number of substitutes is likely to increase.

In the economic geography literature, the spatial distribution of goods over the study area is addressed by including an indicator of accessibility. Fotheringham (1988) argues that if the WTP of both sites is dependent on distance, the substitution effect will be dependent on the relative distance between the sites. Just including distance from the agent to the substitutes therefore does not account for the proximity of substitutes, the spatial structure, and will lead to biased WTP estimates. However, no clear examples of environmental valuation studies account for such spatial structure.

Another important factor in a value transfer study is to determine the relevant substitutes for a certain environmental good. Different criteria have been used to determine the relevant alternatives:

- All available similar ecosystems in the study area or within a certain range; or

- All similar ecosystems known or visited by the respondent; or
- All nature sites in the study area; or even
- All possible recreation areas (not necessarily nature based).

2.3 Aggregation of values

Reliable value transfer should account for differences in socio-economic factors between the study site and the policy site. Large transfer errors can be introduced when value transfer from one region to another does not take into account the variations in relevant-socio-economic characteristics, such as income and demographics. Aggregation implies the estimation of the total WTP of a population by applying the individual WTP value-function from a representative sample to the entire population. As was first demonstrated by Smith (1975) and adopted later on by Loomis (2000), including distance in the WTP function that is used for aggregation can make an enormous difference in the total benefits estimate. The main question is what the size of the market is – i.e. to identify the population to which WTP should be aggregated and the spatial area in which this population lives. The DD effect can determine at what distance from a site people are no longer willing to pay anything for the ecosystem service in question.

A very illustrative example can be found in Bateman et al. (2006), who compare different aggregation methods and assess the effect of neglecting distance-effects. Since they found that response and WTP in principle were both negatively related to distance, they also account for this location effect, besides the socio-economic factors. Instead of aggregating sample means, they apply a spatially sensitive valuation function that takes into account the distance to the site and the socio-economic characteristics of the population in the calculation of values. Thereby, the variability of values across the entire economic market area is better represented in the total WTP. They found that not accounting for distance in the aggregation procedure can lead to overestimations of total benefits of up to 600%. This study shows that reliable aggregation should be based on information about socio-economic characteristics of the most spatially disaggregated level available and should account for distance-effects.

Aggregation sometimes refers to adding up the separately measured WTP values for different sites of a specific type of ecosystem to a Total Economic Value for all those sites together. However, as explained in the previous section, when these study sites function as substitutes or complements, summing up values without considering these interaction effects can lead to large aggregation biases.

Aggregation can also refer to summing up the WTP for different ecosystem services of the same good. This approach may lead to double counting. As long as the functions are entirely independent adding up the values is possible. However, ecosystem functions can be mutually exclusive, interacting or integral (Turner et al., 2004). The excludability or interaction of ecosystem functions and values can also be dependent on their relative geographical position, for instance with substitutes that are spatially dependent.

2.4 Geographic Information Systems

Geographic information systems can be used to help link valuation data with information on the physical (ecosystem size, availability of substitute sites etc.) and socio-economic (income, population, education) characteristics of the policy site.

Reliable value transfer has to account for differences in contextual factors that explain willingness to pay (WTP) for the study and policy site. There are two different sets of spatial attribute to be addressed in economic valuation of any environmental change: (a) the spatial pattern of the social, demographic and psychological characteristics of the affected population and (b) the physical characteristics of the goods and services under valuation. Ignoring the spatial aspects of the latter is assuming that they are randomly distributed over space in terms of quantity and quality. Assuming that population preferences are randomly distributed over space ignores demographic, socio-economic and cultural differences between regions, or the influence of location and distance on environmental values. The distribution will influence the substitution effects between ecosystem sites and determine interaction effects between ecosystem services, which affect aggregation possibilities.

2.5 Transfer Errors

For a number of reasons the application of any of the value transfer methods described above may result in significant transfer errors, i.e. that transferred values may differ significantly from the actual value of the ecosystem under consideration. There are three general sources of error in the values estimated using value transfer:

1. Errors associated with estimating the original measures of value at the study site(s). Measurement error in primary valuation estimates may result from weak methodologies, unreliable data, analyst errors, and the whole gamut of biases and inaccuracies associated with valuation methods.
2. Errors arising from the transfer of study site values to the policy site. So-called generalisation error occurs when values for study sites are transferred to policy sites that are different without fully accounting for those differences. Such differences may be in terms of population characteristics (income, culture, demographics, education etc.) or environmental/physical characteristics (quantity and/or quality of the good or service, availability of substitutes, accessibility etc.). This source of error is inversely related to the correspondence of characteristics of the

study and policy sites.⁴ There may also be a temporal source of generalisation error in that preferences and values for ecosystem services may not remain constant over time. Using value transfer to estimate values for ecosystem services under future policy scenarios may therefore entail a degree of uncertainty regarding whether future generations hold the same preferences as current or past generations.

3. Publication selection bias may result in an unrepresentative stock of knowledge on ecosystem values. Publication selection bias arises when the publication process through which valuation results are disseminated results in an available stock of knowledge that is skewed to certain types of results and that does not meet the information needs of value transfer practitioners. In the economics literature there is generally an editorial preference to publish statistically significant results and novel valuation applications rather than replications, which may result in publication bias.

Given the potential errors in applying value transfer, it is useful to examine the scale of these errors in order to inform decisions related to the use of value transfer. In making decisions based on transferred values or in choosing between commissioning a value transfer application or a primary valuation study, policy makers need to know the potential errors involved. In response to this need there is now a sizeable literature that tests the accuracy of value transfer. Rosenberger and Stanley (2006) and Eshet et al. (2007) provide useful overviews of this literature. Transfer errors are generally expressed as the Mean Absolute Percentage Error (MAPE), which is defined as the difference between observed value and predicted value divided by the observed value.

Table 2.1 summarises the results of a number of studies that measure transfer errors related to ecosystem values. The transfer errors presented in the table show an extremely large range from 0-7028 %. Although some studies find very high transfer errors (e.g. Downing and Ozuna, 1996; Kirchhoff, 1998) most studies find transfer errors in the range of 0-100%. Very high transfer errors may arise when the study and policy sites are very different or when the primary value to which the transferred value is compared is itself an outlier.

It should be noted that the measurement of transfer errors is itself inexact in that it involves a comparison between transferred values and primary valuation estimates,

⁴ In the context of meta-analytic function transfer, generalisation error can arise due to the common limitation of meta-analyses to capture differences in the quality and quantity of the services under consideration. It is often the case that the provision of goods and services is indicated in a meta-analysis merely with binary variables, and that quality is not captured at all. This limitation may translate into transfer errors, as the estimated transfer function cannot reflect important quality and quantity differences in characteristics across sites. A similar problem arises where non-identical services have been combined as one explanatory variable in the meta-analysis. Some level of aggregation across service types is often necessary in order to produce a manageable number of variables in the meta-regression, but at the cost of losing specific categories of services.

which are subject to inaccuracies and methodological flaws of their own. In general, primary values are treated as ‘true’ value observations and transferred values as approximations, whereas they are in fact both approximations. It should also be noted that a single prescribed acceptable level of transfer error is not meaningful because the level of error that is acceptable is likely to be context specific and related to other policy criteria.

There are a number of studies that specifically examine the transferability of value estimates between regions and countries. We summarize some of the main findings of each study below.

Loomis et al. (2005) examine the equivalency of contingent valuation (CV) results for forest fire prevention from studies in California, Florida, and Montana. They test the equality of variable coefficients across States using likelihood ratio tests. Over all tests they find mixed evidence for transferability.

Brouwer and Bateman (2005) transfer contingent valuation WTP estimates for reducing health risks associated with solar UV exposure between four countries (England, Scotland, Portugal, and New Zealand) to examine the sensitivity of transfer errors to differences in contexts. When contexts are similar, mean unit value transfers are actually found to perform better than value function transfer (e.g. when transfers are between England, Scotland, and New Zealand). When study and policy site contexts are different, however, and these differences can be controlled for, value function transfer is shown to produce lower transfer errors.

Kristofersson and Navrud (2007) use identical CV studies conducted in three countries (Norway, Sweden, and Iceland) to examine the validity of value transfers between those countries. The case study estimates use and non-use values for freshwater fish stocks. Values are transferred between study sites using both unit transfer and value function transfer. Equivalency analysis is applied to test the validity of value transfers. The study shows that the accuracy of value transfer relies heavily on the similarity of study sites.

Eshet et al. (2007) examine the accuracy of transferring values for the disamenity of housing locations close to waste transfer stations between four cities in Israel. Value functions derived from separate hedonic pricing studies are used to transfer values for each site. Transfer errors are observed to increase with dissimilarity between sites although errors remain relatively low (2-46%). In comparing the transfer functions estimated for separate study sites, the results of Chow and Wald tests did not indicate equality between value functions and estimated coefficients. The results of the value transfers using these functions did, nevertheless, result in very low transfer errors (particularly where site characteristics were highly similar). The authors therefore argue that a finding of statistical inequality between value functions does not necessarily robustly invalidate transfers of value between sites. In other words, even though valuation studies at different sites may produce different value functions, using these functions to transfer values across sites can still result in low transfer errors.

Table 2.1: Summary of studies measuring value transfer errors

Reference	Resource/activity	Method	Unit transfer error ¹	Function transfer error ¹
Loomis (1992)	Recreation		4–39	1–18
Parsons and Kealy (1994)	Water/recreation		4–34	1–75
Loomis et al. (1995)	Recreation	Nonlinear least-squares model	–	1–475
		Heckman model	–	1–113
Bergland et al. (1995)	Water quality		25–45	18–41
Downing and Ozuna (1996)	Fishing		0–577	–
Kirchhoff et al. (1997)	Whitewater rafting		36–56	87–210
	Birdwatching		35–69	2–35
Bowker et al. (1997)	Whitewater rafting	Pooled data (n–1)	–	14–160
		Pooled data (all)	–	16–57
Kirchhoff (1998)	Recreation/habitat	Benefit function transfer	–	2–475
		Meta-analysis transfer	–	3–7028
Brouwer and Spaninks (1999)	Biodiversity		27–36	22–40
Morrison and Bennett (2000)	Wetlands		4–191	–
Rosenberger and Loomis (2000)	Recreation		–	0–319
Piper and Martin (2001)	Rural water supply	Individual sites (similar)	–	6–20
		Individual sites (dissimilar)	–	89–149
		Pooled data	–	3–23
Van den Berg et al. (2001)	Water quality	Individual sites	1–239	0–298
		Pooled data (multi-state)	0–105	1–56
		Pooled data (state-level)	3–57	0–39
		Pooled data (contaminated sites)	3–100	2–50
Shrestha and Loomis (2001)	International recreation		–	1–81
Chattopadhyay (2003)	Air quality	N=304 (similar subgroups)	106–429	104–486
		N=609 (similar subgroups)	57–150	57–153
		N=913 (similar subgroups)	42–82	42–82
		N=1218 (similar subgroups)	36–67	36–67
		N=1522 (similar subgroups)	32–58	32–58
		N=913 (dissimilar subgroups)	89–128	65–110
Ready et al. (2004)	International air and water quality		20–81	20–83
Jeong and Haab (2004)	Marine recreational fishing	Access per trip	–	4–230
		Per one fish increase	–	2–457
Rozan (2004)	International air quality		–	19–44
Jiang et al. (2005)	Coastal land protection		–	53–85

Source: Rosenberger and Stanley (2006)

¹ The transfer errors are the mean absolute percentage error (MAPE)

Ready et al. (2004) analyse the transfer of contingent valuation WTP estimates for ill health avoidance for five European countries (Portugal, Spain, England, Norway, and the Netherlands). They explore transferring values using unit value transfer, adjusted unit transfer, and value function transfer and find similar transfer errors for all three methods (20-83%). The adjusted unit transfer involved adapting estimated WTP values using the ratio of average real income in the study and policy countries. The authors conclude that a single common value function for the countries included in the study does not exist (i.e. estimated coefficients in the value functions are not the same across countries).

Muthke and Holm-Mueller (2004) test the transferability of contingent valuation estimates for water quality for two German and two Norwegian lakes, thereby testing both national and international transferability. They examine unit value transfer, adjusted unit transfer, and value function transfer using the equivalence testing approach proposed by Kristofersson and Navrud (2005). They also perform Wald tests for equality of parameters. The study results show very high transfer errors for international value transfer suggesting that there was insufficient information available to fully adjust the study site values to the policy sites in another country. The authors argue that because economic factors, intrinsic values, tastes, and preferences of different cultures and societies show considerable variation, international unit value transfer is not feasible and that adjusted unit value transfer and value function transfer are also limited in the account they can take of differences in determining factors.

The existing evidence on regional and international value transfer suggests that there are significant differences between regions in the determinants of environmental values that are not being adequately controlled for in value transfer exercises. The results show that as study and policy sites become more different, transfer errors tend to increase.

Regarding the accuracy of meta-analytic value transfers there is more limited evidence. Rosenberger and Phipps (2001) compare transfer errors between demand function estimates and meta-analytic function estimates using travel cost data for hiking trips in Colorado. The meta-analytic function transfers are shown to result in lower transfer errors. Engel (2002) also specifically compares the performance of benefit function transfers and meta-analysis based function transfers. The results of this comparison are mixed but the conclusions produce an encouraging view of meta-analysis based transfers.

Eshet et al. (2007) describe their analysis as a “mini meta-analysis function transfer” because they use data and transfer functions from four separate samples and locations (including combinations of data from multiple sites). Their analysis is not, however, a meta-analysis in the sense that results from multiple samples and study sites are examined in a regression analysis.

Lindhjem and Navrud (2007) conduct a meta-analysis of contingent valuation results for non-timber forest benefits from Norway, Sweden, and Finland. They examine the accuracy of using alternative specifications of the meta-analytic value function to predict the value of selected observations in their dataset. The best model (a

restricted double-log model) produces mean and median transfer errors of 47% and 37% respectively. This transfer error is lower than that resulting from simple mean unit value transfer from studies from the same country (86%, 41%), and considerably lower than when the mean unit value transfer includes the results of studies from other countries (166%, 85%). These results provide some positive support for meta-analytic value transfer but also illustrate the differences in values between countries, even those with very similar economic, social, and institutional characteristics.

Brander and Florax (2006) use a meta-analytic value transfer function to estimate values for wetlands in the San Joaquin Valley (SJV) in California and for the Norfolk Broads in the UK. The lowest transfer error observed in this exercise is 29% for the valuation of water quality/nutrient retention, recreational hunting and fishing, other recreational activities and amenities in the SJV. Transfer errors of just over 50% are made for recreational hunting in the SJV, and for biodiversity and landscape maintenance and recreational activities in the Norfolk Broads. The transferred value for bird watching in the SJV, however, is over five times the primary value for this activity. Using a database of wetland values, Brander et al. (2006) employ an $n-1$ data splitting technique to estimate 200 meta-analytic value transfer functions and then test the accuracy of each function for predicting the value of the omitted observation in each case. The overall average transfer error is 74% with slightly less than 20% of the sample having transfer errors of 10% or less, and roughly 15% of observations showing transfer errors over 100%. Brander et al. (2007) perform a similar analysis using a database of coral reef recreation values. In this case the average transfer error for the sample of 73 value observations is 186%. The results of the above described studies that examine transfer errors resulting from meta-analytic function transfer are summarised in Table 2.2.

Table 2.2: Summary of studies measuring meta-analytic function transfer errors

Reference	Resource/activity	Method	Meta-analytic function transfer error ¹
Lindhjem and Navrud, 2007	Non-timber forest benefits	Restricted double-log model	47%
		Full double-log model	
Brander and Florax, 2006	Wetland, multiple services		29%
	Wetland, bird watching		433%
	Wetland, hunting		52%
	Wetland, biodiversity		53%
	Wetland, recreation		59%
	Wetland, non-use		99%
Brander et al, 2006	Wetland, multiple services		74%
Brander et al, 2007	Coral reef, recreation		186%

¹ The transfer errors are the mean absolute percentage error (MAPE)

3. Practicability of methods, data requirements, and data availability

3.1 Practicability of methods

The aim of this assessment of value transfer methodologies is to identify a practical procedure for scaling up estimates of ecosystem service values to the European level. Such a procedure should be feasible given the availability of existing data on ecosystem service values and ecosystem characteristics.

Navrud (2007) develops a practical approach for value transfer in the context of Danish environmental planning. The proposed approach is adjusted unit value transfer using estimated values per household or individual. Values are aggregated over the affected population at the policy site. This methodology is practical and straightforward for transferring values to specific policy sites but may be less suitable for large scale value transfers, for example in valuing all ecosystems at a regional, national, or European scale. First, this method relies on the identification of ecosystem valuation studies that correspond most closely with the policy site under consideration. This process could become burdensome as the number of policy sites increases. Second, transferring values per household or individual requires an assessment of the relevant affected population for each ecosystem, which again could become laborious with a large number of policy sites. Transferring values in per household/individual terms may also be problematic for ecosystem services that are generally not valued in these terms. Indirect use values, such as water filtration, are more likely to be valued as inputs in production (e.g. using production function or net factor income valuation methods) and are not expressed in per household terms.

Meta-analytic function transfer on the other hand is well suited to valuing large numbers of diverse policy sites in that the estimated value function can be applied to a database containing information on ecosystem and socio-economic characteristics of each site. It is a simple operation to “plug in” the characteristics of each policy site into a value function to estimate its value. If the value function is defined in terms of values per unit of area it is also a simple operation to aggregate values over spatial areas. Although this approach does not involve aggregation over the affected population, differences in ‘market size’ can still be taken into account through the inclusion of population in the vicinity of the ecosystem as an explanatory variable in the value function.

A clear limitation of meta-analytic function value transfer is related to the reliability of the estimated values. Evidence from the economic valuation literature shows that there are potentially very large transfer errors associated with this approach and that in some cases the relatively simple transfer of unit values may perform at least as well (see previous chapter). It is therefore advisable to test the transfer accuracy of a meta-analytic function in order to provide information about the reliability of the results.

A further potential drawback of using meta-analytic function value transfer is that it is likely to result in varying unit values across European regions. Due to differences in income levels and population densities across Europe, estimated ecosystem values per unit of area are likely to vary between regions. While this makes sense from an economic point of view, it may be politically sensitive, particularly if such information is used to make decisions regarding the allocation of conservation resources.

3.2 Data requirements

On balance, meta-analytic function value transfer offers the most practical approach to scaling up ecosystem service values to a European scale.⁵ The development of such functions requires sufficient data on the value of ecosystem services. Furthermore, the application of this value transfer approach requires sufficient data on the physical, spatial, and socio-economic characteristics of each ecosystem site under consideration.

A value transfer function is used to estimate a site specific ‘per hectare’ value based on the ecosystem site’s characteristics, such as size, type, abundance, and on the characteristics of the population that has a demand for the services of that site. In general form, a meta-analytic value function can be written as:

$$y_i = a + b_S X_{Si} + b_E X_{Ei} + b_C X_{Ci}$$

The variable y_i measures the value of ecosystem site i , based on three vectors of explanatory variables, namely characteristics of (i) the valuation study X_S , (ii) the valued ecosystem X_E , and (iii) the socio-economic and geographical context X_C . The coefficients b_S , b_E and b_C are the vectors containing the coefficients of the explanatory variables, and a is a constant. In order to estimate this function and to use it for value transfer, we need:

1. Data on actual values of $\{y, X\}$ for a sufficient number of ecosystem *study* sites (original valuation studies). With these data, a meta-analytical regression model is built to estimate the coefficients a and b . An example of such a meta-analytical regression model for wetland ecosystems is presented in the next chapter of this report.
2. Data on $\{X_E, X_C\}$ for all *policy* sites within the relevant area (e.g., Europe). The value transfer is done by plugging in these policy site data in the meta-analytical regression model.

The vector X_E includes ecosystem characteristics such as area and type. In our methodology we use hectares as the unit of area. For the ecosystem categorisation we propose to use the same ecosystem types as used in the EEA land cover data. These are given in Table 3.1. Differences between wetland sites in terms of the provision of

⁵ A brief discussion of alternative approaches and how these could be robustly compared with meta-analytic function transfer in future work is provided in section 5.

ecosystem services may be included in the X_E vector, either as dummies (yes/no) or as continuous variables.

The vector X_C includes socio-economic and geographical context variables such as income per capita of the population around the site, population density around the site, and some measure of ecosystem abundance at the local or regional level. Other variables that determine the population's demand for the ecosystem services could be included as well, depending on what the meta-analytical regression identified as significant explanatory variables.

Table 3.1: Ecosystem types used in EEA land cover data

Code	Ecosystem type	Code	Ecosystem type
3.1.1	Broad-leaved forest	4.1.1	Inland marshes
3.1.2	Coniferous forest	4.1.2	Peatbogs
3.1.3	Mixed forest	4.2.1	Salt marshes
3.2.1	Natural grassland	4.2.2	Salines
3.2.2	Moors and heathland	4.2.3	Intertidal mudflats
3.2.3	Sclerophyllous vegetation	5.1.1	Water courses
3.2.4	Transitional woodland shrub	5.1.2	Water bodies
3.3.1	Beaches, dunes and sand plains	5.2.1	Coastal lagoons
3.3.2	Bare rock	5.2.2	Estuaries
3.3.3	Sparsely vegetated areas		
3.3.4	Burnt areas		
3.3.5	Glaciers and perpetual snow		

The availability of the two types of data (study sites and policy sites) is discussed in the following sections.

3.3 Data availability

Availability of ecosystem service value data

Several good databases of environmental valuation results are available. The most comprehensive database is the Environmental Valuation Reference Inventory (available at the EVRI web-page <http://www.evri.ec.gc.ca/evri/>). Other useful online resources are Envalue (<http://www.environment.nsw.gov.au/envalue/>), the Ecosystem Services Database (<http://esd.uvm.edu/>). A number of country specific valuation databases have also been developed, such as the Environmental Valuation Source List for the UK (www.defra.gov.uk/environment/evslist/index.htm) and ValueBase for Sweden (<http://www.beijer.kva.se/valuebase.htm>). In addition, a number of ecosystem specific value databases have been constructed (e.g. for European forest valuation studies under the E45 Cost Action on European Forest Externalities project). These databases provide good starting points for the collection of economic valuation studies for the purposes of value transfer. The data usually comprises bibliographic

information and summaries of methods and results. Conducting meta-analyses using this information requires the construction of numerical databases specifically for this purpose.

Several meta-analyses have been conducted in the field of economic valuation of environmental resources, impacts, and services. Table 3.2 below lists a number of meta-analyses of ecosystem values.

Table 3.2: Meta-analyses of ecosystem values

Ecosystem/ecosystem service	Meta-analysis study
Wetlands	Brouwer et al., 1999 Woodward and Wui, 2001 Brander et al., 2006 Ghermandi et al., 2007
Groundwater	Boyle et al., 1994
Coral reef recreation	Brander et al., 2007
Woodland recreation	Bateman and Jones, 2003
Non-timber forest benefits	Lindhjem, 2007
Outdoor recreation	Smith and Karou, 1990 Rosenberger and Loomis, 2000 Shrestha and Loomis, 2001
Biodiversity	Nijkamp et al., 2008 Jacobsen and Hanley, 2007
Endangered species	Loomis and White, 1996
Urban air pollution	Kaoru and Smith, 1995
Marine and coastal water quality	Barton, 1999
Urban open space	Brander and Koetse, 2007

A comparison with Table 3.1 suggests that with respect to “type”, there is no complete mapping between study sites and policy sites. Additional research is needed to identify the most important “gaps”, and to examine whether additional meta-analyses could be carried out on the basis of existing original valuation studies. For some important ecosystems, such as wetlands and forests, meta-analyses of ecosystem values do exist and these could be used directly for the purpose of value transfer.

Spatial variables for meta-analytic value transfer

As discussed above, meta-analytic value functions are likely to include a number of variables that have a spatial dimension, including ecosystem size, ecosystem abundance, population within a given proximity of an ecosystem, and income per capita of that population. Table 3.3 presents data sources that could be used to construct these spatial variables on a European scale.

It should be noted that the proposed value transfer process estimates values for individual ecosystem ‘sites’ (distinct separate patches of a specific ecosystem type).⁶ In other words, ecosystem sites are the units of analysis in the value transfer exercise and therefore spatial variables need to be defined at this level.

Table 3.3: Spatial data sources

Data set	Download source / owner	GIS data- model / for- mat	Coordinate system and extent	Scale / resolution	Year
Corine land cover 2000 (CLC2000) seamless vector data- base Version 10 jan 2007	EEA Dataservice http://www.eea.europa.eu/ EEA	Vector poly- gon / ESRI Shape	Lambert Azimuthal Equal Area Europe	1:100 000	2000
Gridded Population of the World, version 3 (GPWv3)	Socioeconomic Data and Applications Center (SEDAC) operated by CIESIN http://sedac.ciesin.columbia.edu/gpw	GRID (ESRI)	WGS 1984 Europe	2.5 arc-minute	2000
Administrative land accounting units GISCO administrative boundaries (NUTS) v9	EEA Dataservice http://www.eea.europa.eu/	Vector poly- gon / ESRI Shape	Lambert Azimuthal Equal Area Europe	1:1 000 000	2004
Income per capita Gross domestic prod- uct (GDP) in 2003 at current market prices at NUTS level 2	Eurostat table reg_e2gdp	Table	Lambert Azimuthal Equal Area Europe	NUTS level 1- 3	2003

Ecosystems

The spatial data that need to be generated for ecosystem characteristics are:

- Locations of individual sites of each ecosystem type.
- Ecosystem size (in hectares) for each individual ecosystem site.

⁶ See section 4.5 for a full description of the proposed value transfer procedure. In brief, a value transfer function is used to estimate a site specific ‘per hectare’ value based on each ecosystem site’s characteristics (size, type, abundance, socio-economics). This ‘per hectare’ value is then multiplied by the area of the site to obtain a total value for that site. The values of individual sites can then be aggregated to obtain values for each ecosystem within a region, country, or Europe as a whole.

- Ecosystem abundance, which is defined as the total area of an ecosystem type within some radius (in km) of the centre of each ecosystem site. The degree to which a particular ecosystem type is scarce is in part determined by the scale of analysis. It is possible, for example, for a particular ecosystem type with a cluster of sites to be considered abundant at a local scale but very scarce at a European scale (if there is a small total area of this ecosystem type).

The necessary steps construct in ArcGIS a geo-referenced map of ecosystem areas in Europe from the Corine land cover database are illustrated by the construction of a wetland area map (ecosystem codes 4.1.1, 4.1.2, 4.2.1, 4.2.2, and 4.2.3 from Table 3.1). Figure 3.1 presents the resulting map of European wetlands.



Figure 3.1: Map of European wetlands

To determine the surface area of each ecosystem site we have used the Corine seamless vector land cover data (CLC2000). The sizes of the areas of individual ecosystem sites have been calculated with the calculate geometry function in ArcGIS.

We also calculated the centerpoints of the areas of the ecosystem sites. In the map example in Figure 3.2 the calculated centerpoint locations are displayed on top of the ecosystem site areas (polygons) for the type broad-leaved forest.





Figure 3.2: Centerpoints (red points) of ecosystem type broad-leaved forest (Code 3.1.1)

The *ecosystem abundance* is defined as the summed area of ecosystems within some radius of the centerpoint of each ecosystem site. In the examples used in this section, the radius is set at 50 km. This ArcGIS procedure has been modelled for a test area containing a large part of the Netherlands with ArcGIS model builder. It produces from the Corine (seamless vector) database for all sites of 21 different ecosystem types an abundance indicator value expressed in hectares. In the map example in Figure 3.3 the abundance of the ecosystem types broad-leaved forest and natural grassland are represented by the calculated background 100 meter raster map in which the highest scarcity is represented by the darkest areas.



Upper panel: Abundance of broad-leaved forest from less abundant (black) to abundant (white)

Lower panel: Abundance of natural grassland from less abundant (black) to abundant (white)

Figure 3.3: Ecosystem abundance indicators for broad-leaved forest and natural grasslands in the Netherlands

The map in Figure 3.4 shows an indicator of wetland abundance. It shows the centrepoint (red point) of an intertidal mudflat in Northern Ireland (ID 45970) with a

surface of 805 hectares, surrounded by 147,406 hectares of other wetland areas in a circular 50 km zone.




Figure 3.4: Wetland abundance in Northern Ireland

Population

The spatial data that need to be generated regarding population characteristics concern the population in the vicinity of the ecosystem site (usually defined within some radius of the centre of each ecosystem site). The process by which this data can be generated is described in Wagtenonk and Omtzigt (2003).

For the population data we can choose between two available population data sets for Europe:

1. The Gridded Population of the World dataset (GPWv3) of the Socioeconomic Data and Applications Center (SEDAC). File name: euds00ag (ESRI GRID format).
2. The population density dataset of the Joint Research Centre (EC-JRC). This is a disaggregated dataset (to 100 meter gridcells) based on the Corine land cover 2000 map.

A choice between these two datasets has to be made based on the required spatial resolution versus processing time needed for calculating average population densities in zones with a radius of 50 km around ecosystem sites.

We have performed a simple test using spatial statistics of Statistics Netherlands (Wijk- en buurtkaart CBS 2001), containing, among others, population density figures for all municipalities in the Netherlands. The test showed that on a municipal level the population density figures of the GPW dataset shows by far the best results for the tested municipalities (respectively 17.5 % difference and 233 % difference with the CBS figures, see Table 3.4). It has to be remarked however that the 100 meter gridcell values of the EC-JRC dataset show in general better values when zoomed in to the centres of separate towns. We are however more interested in average population density figures for larger areas and have therefore chosen to use the GPW dataset. It has to be noted that this test has only been performed on Dutch population density figures and not on figures for other countries in Europe. Still we expect this test to be representative.

A second reason to choose for the GPW dataset is the larger gridcell resolution of 2.5 arc-minute (circa 4.6 by 4.6 km) which will be much faster to process than the EC-JRC 100 meter gridcells.

Table 3.4: Comparison of EC-JRC and GPW population density datasets with official (CBS) density figures of some Dutch municipalities

	Amsterdam	Amstelveen	Texel	Utrecht	NO-polder	Bergen	% difference with CBS
CBS value	4449	1922	85	2664	95	326	
Popgrid EC-JRC	9412	5428	40	7313	40	15	233
Adjusted GPW	2966	1995	84	2591	91	149	17.5

As source data we used the population data provided by the International Earth Science Information Network (CIESIN), of Columbia University.

Income

The spatial data that need to be generated regarding economic characteristics concern income per capita of the population in the vicinity of each ecosystem site. Ideally the population to which the income data relates should be the population that hold values for the ecosystem in question. As it would be unfeasible to identify this population for each ecosystem site, we propose to use income per capita at the NUTS2 level. GIS can be used to link the location of each ecosystem site to the relevant NUTS2 region.

The income figures per capita for the NUTS areas in which the ecosystem sites are located can be calculated by combining Eurostat statistics for Gross Domestic Product (GDP) in 2003 (table `reg_e2gdp.xls`) at current market prices at NUTS level 2 with the administrative map units downloaded from the EEA dataservice: Administrative land accounting units GISCO (NUTS) v9.

4. Case study

4.1 Introduction

In this section the economic value of services and goods from wetlands in the European Union is estimated based on the meta-analytic value transfer methodology and scaling up procedure introduced in the previous sections. The preliminary work carried out by Ghermandi et al. (2007), who performed a meta-analysis of a very large number of wetland valuation studies provides the starting point for the scaling up valuation procedure. A wide range of relevant explanatory variables is included in the meta-analysis and value transfer exercise, including the abundance of wetlands, the type of ecosystems services provided, the population in the vicinity of each wetland, the GDP per capita (at NUTS2 level for European observations), the size of each wetland, and the economic valuation method used. In the scaling up exercise, the meta-analytic value transfer function is combined with the spatial information on 50,533 individual wetland sites generated with a GIS from the Corine land cover.

4.2 The wetland valuation literature

The monetary estimation of the market and non-market benefits of wetlands has been the subject of a large number of primary valuation studies. Since the publication of the first wetland valuation study in 1974 (Hammack and Brown, 1974), the number of studies aimed at estimating the value of wetlands has steadily grown. The most extensive review of the wetland valuation literature up to date is by Ghermandi et al. (2007) and counts 383 value observations from 166 independent valuation studies. This large number of closely related and comparable studies has stimulated the use of research synthesis techniques known as meta-analysis. Four meta-analyses of wetland valuation studies have been published:

1. Brouwer et al. (1999) analyze the results of contingent valuation method (CVM) studies of temperate climate zone wetlands. The definition of wetlands used in this study is very broad and the meta-analysis includes a number of valuation estimates for open water (rivers and lakes). The focus on estimates from CVM studies in developed countries, mainly the United States, narrows the sample size to 92 value observations from 30 studies.
2. Woodward and Wui (2001) similarly restrict the scope of their meta-analysis to include valuation studies for North American and European wetlands only. They use a narrower definition of wetland than Brouwer et al. (1999) while also including wetland values obtained with valuation techniques other than CVM. The resulting data set contains 65 value observations taken from 39 studies.
3. Brander et al. (2006) assembled a dataset of 215 value observations obtained from 80 studies. Their analysis includes studies from temperate and tropical regions, for different wetland types (including mangroves), and for a broader set of wetland functions and valuation methods. An important element of this meta-analysis is the addition of external socio-economic variables like GDP per capita and popula-

tion density. In spite of the broad geographical scope adopted, the distribution of primary valuation studies is still very much biased by the practice and availability of natural resource valuation studies rather than by the distribution of wetlands. In particular, studies from North America accounted for half of the total number of observations.

4. Ghermandi et al. (2007) greatly expanded the data set used in Brander et al. (2006) to include by far the largest number of primary valuation studies used in a meta-analysis of wetland values: namely, 383 independent observations derived from 166 studies. With respect to previous meta-analyses, there is an extension of the geographical coverage of the studies, which is less biased towards developed Western countries. Indeed, a clear increase in the number of studies from Africa, Asia and Europe in recent years is identified, while the number of new studies from North America – where wetland valuation was first widely used – shows a downward trend. In addition, man-made wetlands are included for the first time in a meta-analysis of wetland values. The innovative contributions of this model include the recognition of substitution effects between wetland sites and environmental pressure as important explanatory variables of wetland values. Furthermore, the presence of human pressures on the wetlands is taken into account in the analysis by means of an index of environmental stress and is recognized to lead to higher values.

4.3 Description of the data set and the meta-regression model

The data set of wetland values

The data set used for the determination of the meta-analytic value transfer function relies on the work conducted by Ghermandi et al. (2007). Figure 4.1 provides an overview of the location of the valued wetland sites and the year of publication of studies examined by Ghermandi et al. (2007).

Despite the focus of the case study on scaling European wetland values, the data set underlying the meta-regression analysis is not limited to European sites only. Reasons for this include the need to provide a sufficient number of observations for wetland services that are not frequently object of valuation studies (e.g. non-use values) and guarantee a sufficient degree of variability in the explanatory variables of the meta-regression. A large degree of variability in the explanatory variables is in fact not a limit to the meta-analysis as far as it can be assumed that there exists a single underlying function that links the size of a specific effect on the dependent variables to the explanatory variables. On the contrary, a sufficient degree of heterogeneity is necessary to robustly identify the size of a specific effect on the dependent variable.

All continents are represented in the data set. The largest number of observations is derived from North American studies (129), but a significant fraction comes from Asia (89), Europe (78) and Africa (53). South America (18) and Australasia (16) are less well represented. The studies included in the data set are all primary valuation

studies and, in order to limit the risk of introducing a publication bias, the analysis is not limited to publications from the “official scientific literature”, but also explores the complementary areas of “grey literature” (e.g. reports for both public and private institutions, consultancy studies) and unpublished research results. The average number of observations per study (2.3) and the maximum number of observations for a single study (10) is relatively low if compared to the total number of observations used in the analysis (383). As such, multiple sampling bias is expected to have a scarce influence on the results of the investigation.




Figure 4.1: Number of observations of wetland values in intervals of five years from 1972 to 2007 and geographical location

Since the goal of the meta-regression performed in this study is to provide a meta-analytic value transfer function to be applied to the spatial information derived from the CORINE map concerning of land uses in countries in the European Union, it was not possible to make use of the whole data set of valuations. Due to the large scope of the meta-analysis performed by Ghermandi et al. (2007) in fact, the definition of wetland upon which the selection of ecosystems types and valuation studies was based is a comprehensive one. It encompasses all ecosystem types embraced by the Ramsar definition⁷ with the exception of rice cultivations, coral reefs, sea-grass beds,

⁷ The definition of provided by the Ramsar Convention on Wetlands of International Importance identifies as wetland any area of “marsh, fen, peatland or water, whether natural or artificial, permanent or temporary, with water that is static of flowing, fresh, brackish or salt, including areas of marine water the depth of which at low tide does not exceed six meters” (art. 1.1)

rivers, and shallow lakes, which are implicitly included in the Ramsar definition but are seldom regarded as wetlands (Scott and Jones, 1995). The definition of wetland used in the EEA land cover data, however (see Table 3.1), is more restrictive and explicitly excludes wooded areas such as wet forests and forested floodplains, which are classified as forest ecosystems, and estuaries and coastal lagoons, which are regarded as water bodies. Furthermore, the data set by Ghermandi et al. (2007) includes a number of tropical ecosystems (e.g. mangroves), which do not naturally occur in European countries. For all the mentioned reasons, the original data set was restricted to include only ecosystems that are compatible with the definition of wetland used in the EEA land cover data. The total number of usable observations was reduced to 264.

The meta-regression model and the explanatory variables

The meta-analytical regression model used for the estimation of wetland values is illustrated in matrix notation, in equation 1.

$$\ln(y_i) = a + b_s X_{si} + b_w X_{wi} + b_c X_{ci} + u_i \quad (1)$$

The dependent variable (y) in the meta-regression equation is the vector of the wetland values standardized to 2003 US\$ per hectare per year. The subscript i assumes values from 1 to 264 (number of observations), a is the constant term, b_s , b_w and b_c are the vectors containing the coefficients of the explanatory variables and u is the vector of residuals. Table 4.1 provides an overview of the explanatory variables. They consist of three categories, namely characteristics of (i) the valuation study X_s , (ii) the valued wetland X_w and (iii) the socio-economic and geographical context X_c . The variable type (nominal, interval, or ratio) is also reported.

Study characteristics (X_s). Study characteristics accounted for in the model include the valuation method used and a dummy distinguishing between marginal and average values (Brander et al., 2006).

A wide array of valuation methods has been used in the primary studies for the assessment of the different values of wetlands. These include market-based methods – i.e., market prices (61), net factor income (34), opportunity cost (9), replacement cost (56) and production function (14) –, revealed preference methods – i.e., travel cost method (42) and hedonic pricing (5) –, and stated preference methods – i.e., contingent valuation method (62) and choice experiment (8). A dummy for each of the valuation methods is included in the meta-regression model to account for the heterogeneity of methods, as not all of them have a strong basis in welfare theory and as they produce estimates of different welfare measures.

In standardizing wetland values we face the problem of distinguishing between average and marginal values, both of which can be expressed as a monetary value per hectare. To distinguish between marginal and average per hectare values in the meta-regression, following Brander et al. (2006), a dummy variable is introduced, which takes a value equal to one for marginal values (36) and equal to zero for average values (228).

Table 4.1: Explanatory variables used in the meta-regression model

Group	Variable	Variable type	Levels / measurement unit	N		
Study (X_S)	Valuation method	Nominal	Contingent valuation method	62		
			Hedonic pricing	5		
			Travel cost method	42		
			Replacement cost	56		
			Net factor income	34		
			Production function	14		
			Market prices	61		
			Opportunity cost	9		
			Choice experiment	8		
			Marginal / average	Nominal	Average	228
Marginal	36					
Wetland (X_W)	Wetland type	Nominal	Inland marshes	182		
			Peatbogs	21		
			Salt marshes	64		
			Salines	0		
			Intertidal mudflats	41		
			Wetland size	Ratio	Hectares (ln)	264
			Service provided	Nominal	Flood control and storm buffering	34
					Surface and groundwater supply	33
					Water quality improvement	38
					Commercial fishing and hunting	53
					Recreational hunting	47
					Recreational fishing	49
					Harvesting of natural materials	39
					Fuel wood	13
					Non-consumptive recreation	70
					Amenity and aesthetics	34
Biodiversity	36					
Context (X_C)	GDP per capita ^a	Ratio	2003 US\$ person ⁻¹ year ⁻¹ (ln)	264		
			Population density	Ratio	Inhabitants in 50 km radius in year 2000 (ln)	264
			Wetland abundance	Ratio	Hectares in 50 km radius (ln)	264

N = number of observations for each variable or variable level

^a At NUTS 2 level for European observations, state level for observations from the U.S.A., and country level for all other observations

Wetland characteristics (X_W). Characteristics of the valued wetland site that are accounted for in the meta-regression model are the type and size of the wetland and the types of services provided.

The wetlands in the database are classified according to the EEA land cover nomenclature for wetland ecosystems. According to the EEA classification (see Table 3.1), inland wetlands include inland marshes and peatbogs, while coastal wetlands are classified into salt marshes, salines, and intertidal mudflats. As large wetlands may include areas with very different characteristics, the same observation may be classified under two or more wetland systems. The large majority of the wetlands in the database are inland marshes (182). A significantly lower number of observations

are available for the other wetland types: salt marshes (64), intertidal mudflats (41), and peatbogs (21). No valuation of salines is available.

Wetlands provide a number of services and goods that are of value to humans. The economic services of wetlands are derived from, but should not be confused with, their ecological and physical functions. The classification of wetland functions and services was the object of a large number of studies. Wetland values have generally been classified on the basis of the underlying wetland functions (Barbier, 2006), the characteristics of use and non-use values (Barbier et al., 1997), the provision of intermediate, final or future goods and services (Leschine et al., 1997), or private versus public or social values (Whitten and Bennett, 1998). In this study, we follow the approach proposed in the Millennium Ecosystem Assessment (2005), which is based on classification of ecosystem services into the categories of supporting, provisioning, regulating and cultural services. Table 4.2 illustrates the main wetland economic services and goods together with the valuation methods most commonly used for the estimation of their impact on human welfare. For some of the wetland services in Table 4.2 – i.e., appreciation of uniqueness to culture/heritage, educational, support of pollinators, sediment retention, micro-climate stabilization, regulation of greenhouse gases, and non-use values – no direct valuation study could be found in the literature. The largest number of observations is available for non-consumptive recreation (70). A relatively large number of observations are available for almost all other wetland services with the exception of fuel wood extraction for which only 13 observations are available. Care was taken in coding dummy variables for these wetland services in the database to avoid double counting of service categories. This is particularly the case for the supporting function services.

Context characteristics (X_C). Environmental valuation studies carried out at different geographical sites and involving populations with different socio-economic characteristics and consumer preferences typically produce different outcomes (Brouwer, 2000). Context characteristics are expected to significantly influence the value estimates (Brander et al., 2006). Three context variables are included in the meta-regression model: Gross Domestic Product (GDP) per capita, population living in the surroundings of the wetland, and wetland abundance.

The values of real GDP per capita used in the meta-regression are estimated in US\$ referring to the year 2003. Due to the different availability in the various contexts involved in the analysis, the data reflects the socio-economic characteristics of the population at different administrative levels: for European countries, real GDP per capita was calculated at NUTS2 level based on the data provided by Eurostat, for the US at state level (World Bank, 2006) and for the rest of the world at country level (World Bank, 2006). The socio-economic conditions of the population residing in proximity of the valued wetland sites vary largely across observations. This is reflected by the large variations in average real GDP per capita, which ranges from 616 to 47,547 UD\$ 2003 per person per year in Cambodia and Massachusetts, US, respectively.

Table 4.2: Principal services and goods provided by wetlands and valuation methods commonly used to estimate their value

Category	Wetland service	Valuation methods
Cultural	Amenity and aesthetics	CVM (22), HP (5), TCM (5)
	Non-consumptive recreational activities	CVM (44), TCM (18)
	Appreciation of uniqueness to culture/heritage	-
	Educational	-
	Recreational hunting	TCM (21), CVM (14)
	Recreational fishing	CVM (22), TCM (15)
	Non-use values	-
Supporting	Biodiversity	CVM (23), choice experiment (6), market prices (5)
	Support of pollinators	-
Provisioning	Commercial fishing and hunting	Market prices (18), NFI (18), CVM (10)
	Harvesting of natural materials	Market prices (18), NFI (11), CVM (8)
	Fuel wood	Market prices (7), NFI (4)
	Surface and groundwater supply	Replacement cost (15)
Regulating	Flood control and storm buffering	Replacement cost (20)
	Sediment retention	-
	Water quality improvement	Replacement cost (28), CVM (10)
	Micro-climate stabilization	-
	Regulation of greenhouse gases	-

HP = hedonic pricing; CVM = contingent valuation method; TCM = travel cost method; NFI = net factor income

Note: In brackets () the number of observations for each wetland service according to the most commonly used valuation methods

The total population living in a radius of 50 km around the wetland centre is estimated by means of a GIS based on the information reported in the Global Demography Project map (CIESIN, 2005), which contains geographically referenced information of world population in the year 2000. Population data for a range of different distance radii were generated and tested in the regression model. The 50 km radius variable was found to have the highest explanatory power. It is noted that the size of the market (i.e. the population within a certain distance of an ecosystem site that hold values for the ecosystem services from that site) may vary for different ecosystem services. In particular, non-use values are not expected to decrease with distance to the same extent as direct use values. In further analysis we may therefore consider interacting different measures of market size with different ecosystem service variables.

The total wetland area in a radius of 50 km around the wetland centre accounts for the uniqueness of a wetland environment and may help explaining the influence of people's perceptions and preferences due to the presence of other sites that can act as a substitute for some of the services provided. The area of nearby wetlands was esti-

mated by means of a GIS based on the information contained in the Global Lakes and Wetlands Database map (Lehner and Döll, 2004). As with the population variable, wetland abundance was calculated for a number of different distance radii and the 50 km variable was found to have the highest explanatory power.

Following Ghermandi et al. (2007), the geographical location of the wetland site is not included in the meta-regression model used in this study as it is significantly correlated to other variables such as the services and goods valued and the valuation method applied. For instance, valuation studies of the recreational hunting service tend to be concentrated in North America, while provision of materials and fuel wood are valued in South America, Asia and Africa more often than in North America and Europe. Other meta-analyses include geographical location of the primary studies in the meta-regression model as a set of dummy variables (Brander et al., 2006, Brouwer et al., 1999).

Standardization of values. To allow for a comparison between wetland values that have been calculated in different years and expressed in different currencies and metrics – e.g. willingness to pay (WTP) per household per year, capitalized values and marginal value per acre – standardization to common metric and currency is needed. WTP per household per year cannot be used as a common metric since several of the valuation methods used in the literature – e.g. net factor income, opportunity cost, replacement cost and market prices – do not produce WTP per person estimates. On the other hand WTP per person can be converted to a value per hectare per year if the relevant population is known. Following Ghermandi et al. (2007), values were thus standardized to 2003 US\$ per hectare per year. Values referring to different years were deflated using appropriate factors from the World Bank Development Indicators (World Bank, 2006), while differences in purchase power among the countries were accounted for by the Purchase Power Parity index provided by the Penn World Table (Heston et al., 2006). Some valuations are expressed in US\$ in the primary studies although they stem from other countries. Such values are first converted to local currency using exchange rates of the year of the study and only subsequently deflator factors and purchase power parity indexes are applied to obtain standardized values.

4.4 Results of the meta-regression model

The results obtained with the meta-regression model described in Table 4.1 using ordinary least squares (OLS) are presented in Table 4.3. In this (largely) semi-logarithmic model, the coefficients measure the constant proportional or relative change in the dependent variable for a given absolute change in the value of the explanatory variable. For the explanatory variables expressed as logarithms, the coefficients represent elasticities, that is, the percentage change in the dependent variable given a (small) percentage change in the explanatory variable. The values of R^2 (= 0.49) and adjusted R^2 (= 0.43) are reasonably high. We interpreted these results as signaling that the econometric model specification and the respective parameter es-

timates show a high goodness-of-fit. In other words, the estimation results approximate the real data.

Table 4.3: Results obtained with the meta-regression model of wetland values

	Variable	Coefficient	p-value
	(constant)	-3.078	0.187
Study variables	Contingent valuation methods	0.065	0.919
	Hedonic pricing	-3.286***	0.006
	Travel cost method	-0.974	0.112
	Replacement cost	-0.766	0.212
	Net factor income	-0.215	0.706
	Production function	-0.443	0.523
	Market prices	-0.521	0.317
	Opportunity cost	-1.889**	0.035
	Choice experiment	0.452	0.635
	Marginal	1.195***	0.008
	Wetland variables	Inland marshes	0.114
Peatbogs		-1.356**	0.014
Salt marshes		0.143	0.778
Intertidal mudflats		0.110	0.821
Wetland size		-0.297***	0.000
Flood control and storm buffering		1.102**	0.017
Surface and groundwater supply		0.009	0.984
Water quality improvement		0.893*	0.064
Commercial fishing and hunting		-0.040	0.915
Recreational hunting		-1.289***	0.004
Recreational fishing		-0.288	0.497
Harvesting of natural materials		-0.554	0.165
Fuel wood		-1.409**	0.029
Non-consumptive recreation		0.340	0.420
Amenity and aesthetics		0.752	0.136
Biodiversity		0.917*	0.053
Context variables	GDP per capita	0.468***	0.001
	Population in 50km radius	0.579***	0.000
	Wetland area in 50km radius	-0.023	0.583

OLS results. $R^2 = 0.49$; $Adj. R^2 = 0.43$. Significance is indicated with ***, **, and * for 1, 5, and 10% statistical significance levels respectively.

Of the study characteristics, the valuation methods, do not have statistically significant coefficient estimates with the exception of hedonic pricing and opportunity cost methods. The coefficient of hedonic pricing is significant, negative and large. The number of studies applying this method is however very small (5 observations). Table 4.3 shows relatively high coefficients for studies with stated preference methods (contingent valuation and choice experiment), although not statistically significant. This confirms the observation by Brander et al. (2006) who found high values for contingent valuation studies but contrasts with the results of Woodward and Wui (2001) who observed high values for studies using hedonic pricing and replacement cost as valuation method. Bearing in mind that contingent valuation and choice experiment are the only methods capable of shedding light on the non-use values and that

the respective parameters estimates are both not statistically significant, this signals that non-use values are important in explaining the wetland values, at the margin. One can interpret this result by the fact that non-use values have not been sufficiently addressed in the data set. In other words, a question of data availability. In fact, a closer look to the data sets shows that most of the surveys were performed answered by the visitors or the populations living in the neighbour region. Another element that supports this line of reasoning is that the constant term is not statistically significant either.

Several variables capturing wetland characteristics turn out to be statistically significant. Wetland types – as classified according to the EEA land cover nomenclature – are mostly not statistically significant, with the exception of peatbogs, whose negative coefficient is statistically significant at the 5% level. The coefficient of wetland size confirms the results of previous meta-analyses in indicating decreasing returns to scale. Of the wetland functions, the coefficients of hunting and fuel wood are negative and significant. This means that these factors reduce, at the margin, the monetary values of the wetlands. Alternatively, coefficients of flood control and storm buffering, biodiversity, water quality improvement, and amenity/aesthetics are positive. This means that these characteristics influence positively the value of the wetland.

Of the three context variables, only GDP per capita and population living in the surrounding of the wetland site are statistically significant and indicate a positive and inelastic effect. Contrarily to what observed by Ghermandi et al. (2007), in the subsample of the data set of wetland valuations selected for this study the coefficient of the variable accounting for wetland abundance is not statistically significant, which suggests that substitution effects due to the proximity of other wetlands to the valued site do not significantly contribute to determine of values in the considered wetland sites. Like before, this can also be interpreted in terms of the data availability: meaning that the econometric estimation results confirm the substitution effect in consumption – i.e. the higher is the presence of wetlands in my region, the lower is valuation of an additional hectare of wetland in the same region. The magnitude of this effect is, however, not statistically different from zero. A wide geographical coverage of wetlands in the European Union is not sufficiently addressed in the data set (which is concentrated in France, UK and North Europe).

As mentioned above, in order to allow a comparison between marginal and average wetland values, a dummy variable is introduced, taking a value equal to one for marginal values (36 observations) and equal to zero for average values (228 observations). According to the estimation results, marginal values are statistically significant and indicate a positive impact. This result reiterates the need to distinguish marginal, or incremental values, from the total values.

4.5 Proposed scaling up valuation procedure

The estimation results of the meta-regression function identify the variables that are statistically significant in explaining variation in wetland values. This value func-

tion can be combined with policy site data on the explanatory variables to estimate a per hectare value for the policy site(s) in question.

The proposed valuation procedure for scaling up wetland values to the European level consists of the following steps:

1. The meta-regression function will be used to estimate a per hectare annual value for each wetland in Europe given information on the characteristics of each individual wetland site in Europe. A database of wetland sites in Europe will be generated using a GIS to include information on: wetland type, size, location, wetland abundance, population in vicinity, income per capita of neighbouring population.
2. These site specific wetland values (in annual per hectare terms) will then be multiplied by the area of each wetland to obtain the annual value of services from each wetland.
3. The values of each wetland will then be aggregated to the regional, national, or European level. This will give us the annual value of services from the existing wetland network today.

These steps comprise the first stage of scaling up wetland values for use in policy analysis at the European level by providing an estimate of the aggregated annual value of European wetlands today. Using this procedure for policy analysis, however, requires a marginal analysis based on the comparison of alternative future policy scenarios. In other words, a second stage of analysis is required, which involves the definition of the policy scenarios and the estimation of wetland values under these scenarios. The definition of scenarios requires a description of the wetland network and the projection of the explanatory variables over time.

4.6 Value transfer results

A database of 53,743 European wetland sites was generated by the GIS analysis described in section 3. It was not, however, possible to generate the spatial variables required for the value transfer for all of these sites due to missing data in some of the data sets used. The number of wetlands for which all spatial variables are available is 50,533, which is 94% of the wetlands in the Corine database.

Using the approach described above, we applied a meta-analytic value transfer function to estimate annual per hectare values for each wetland site. Table 4.4 presents these values averaged by country. It can be seen that Sweden and Finland have the largest numbers of wetlands and the largest total areas of wetland. It is partly due to this relative abundance of wetlands, combined with low population densities, that the per-hectare values are low in these countries. By comparison, Belgium, Italy, and the Netherlands have much lower numbers and areas of wetlands but these wetlands have considerably higher values in per hectare terms.

Table 4.4: Number of wetland sites, wetland area, and mean value per hectare per year by country

Country	Number of wetlands	Wetland area (ha)	Mean value per ha per year (€)
Austria	211	31,748	5,052
Belgium	92	10,480	9,627
Bulgaria	81	11,584	3,110
Croatia	140	18,761	4,628
Cyprus	3	1,956	4,724
Czech Rep	105	8,987	4,435
Denmark	729	164,961	3,896
Estonia	1,146	197,786	837
Finland	14,140	1,971,961	224
France	1,419	358,163	5,693
Germany	1,391	418,945	4,353
Greece	302	64,766	3,992
Hungary	1,090	96,500	3,309
Ireland	2,173	1,210,044	676
Italy	344	68,891	9,125
Latvia	883	156,580	764
Lithuania	563	57,548	1,543
Malta	1	25	76,933
Netherlands	273	269,753	7,871
Poland	913	110,386	4,032
Portugal	162	28,293	7,686
Romania	1,532	384,611	2,615
Slovakia	74	4,293	5,792
Slovenia	13	3,249	7,340
Spain	392	112,684	6,647
Sweden	20,242	2,729,131	263
United Kingdom	2,119	753,691	2,480
Total	50,533	9,245,777	1,193

Table 4.5 presents the number of wetland sites, wetland area, and mean value per hectare per year by wetland type. By far the most prevalent type of wetland in Europe is peatbog, followed by inland marshes, salt marshes, and inter-tidal mudflats. There are very few saline wetlands included in the data. On average, salt marshes have the highest estimated values in per hectare terms. In comparison to other types of wetland, peatbogs have very low average per hectare values. This is likely to be due to their relative abundance and to the fact that a large proportion of this type of wetland is located in the sparsely populated north of Sweden and Finland.

In order to obtain an indication of the accuracy of this value transfer, we applied the same transfer approach to predict values for the wetlands included in the wetland value database. The average of the transferred values are 35% higher than the average of the observed values. This is comparable to the results of previous studies on

transfer error (see Table 2.1). According to the recent literature on formal validity testing of benefit transfer (see Kristofersson and Navrud, 2007 and 2005; Ready et al. 2004), this value is within the boundaries of technical acceptability and thus can be of support for policy design.

Table 4.5: Number of wetland sites, wetland area, and mean value per hectare per year by wetland type

Wetland type	Number of wetlands	Wetland area (ha)	Mean value per ha per year (€)
Inland marshes	8,842	1,159,153	4,129
Peatbogs	38,644	6,712,309	214
Salt marshes	1,621	306,754	5,734
Intertidal mudflats	1,180	995,094	4,112
Salines	246	72,467	5,475
Total	50,533	9,245,777	1,193

5. Discussion

This paper proposes a methodology for scaling up ecosystem service values to a European level, assesses the availability of data for conducting this method, and illustrates the procedure with a case study on wetland values. The proposed methodology makes use of meta-analytical value transfer to produce a value function that is subsequently applied to individual European wetland sites. Site-specific, study-specific and context-specific variables are used to define a price vector that captures differences between sites and over time. The proposed method is shown to be practicable and to produce reasonably reliable aggregate value estimates. There are, however, a number of issues that require further discussion and research.

Points of strength of the proposed meta-analytic value transfer technique are its ability to explicitly account for context variables which are relevant in understanding people's perceptions and preferences but are often neglected in primary valuation studies because constants of the analysis. Context variables included in the case study aim at assessing the influence of income effects, substitute sites, and population density on wetland values. Their spatial variability is captured on a scale based either on administrative boundaries (as for the variable real GDP per capita) or distance from the ecosystem centre (as for wetland abundance and population density). At the state of the art of meta-analytical value transfer, the choice of the spatial scale of analysis for context variables involves a certain degree of arbitrariness. The results of the econometric analysis can in fact provide *post hoc* information concerning the effect of different spatial scales on the value transfer function coefficients but do not help in understanding the mechanisms that determine the spatial scale that is relevant for each single study site. A possible improvement of the approach used in this study may rely on the distance from the site of interest as a weighting factor for calculating the value of a specific context variable. For instance in the case of the wetland abundance indicator used in the proposed analysis, the distance between the centre of the

valued site and the centre of each of its surrounding substitute sites can be used as a weight in the calculation of the abundance indicator for the econometric analysis.

An important distinction needs to be made between marginal and average values. In using a meta-analytic value function to estimate annual per hectare ecosystem values, and multiplying this estimate by ecosystem size, we are estimating average and total values. Average values may be useful for comparing the aggregate value of an ecosystem area relative to the size of the area and total values may be useful when considering non-marginal changes in ecosystem areas. Small changes in the extent of ecosystems should, however, be valued using marginal values. To assess marginal changes in ecosystem stocks at a European scale, we have proposed to estimate the associated marginal change in value by calculating the difference in total annual ecosystem service values between alternative scenarios. This step has not been taken in the present study but could be the subject of future work.

We warn that total annual ecosystem values should be presented with caution. In order to calculate total ecosystem values accurately we should account for non-constancy of marginal values, i.e., that as the area of an ecosystem increases, the added value of an additional unit of area is likely to decline. Indeed in the case of wetlands, per hectare values are shown to decline with the size of the wetland itself and with the abundance of wetlands in the vicinity. Conversely, as wetland area and abundance decline, per hectare values will increase. This means that multiplying a constant per hectare value by the total area of a wetland is likely to underestimate total value.

An important area for improvement in the proposed value transfer method is the treatment of ecosystem quality. As currently proposed, the method deals primarily with the quantity or area of ecosystems and does not deal well with changes in quality. To incorporate ecosystem quality into the value transfer process would require the definition and inclusion of quality variables in both the valuation data underlying the meta-analysis and in the data on European ecosystem networks. In the case of wetlands, several methods are available for assessing their ecological integrity (Fennessy et al., 2004). Most of them, however, are not practicable for the type of analysis that the present study aims at as they rely on local biological and physico-chemical measurements which are not available for most of the study and policy wetland sites. Some methods, however, use estimates of the anthropogenic pressures in the surrounding of the wetland as a proxy for its ecological status assuming that this strongly depends on them. Such methods are usually referred to as landscape assessments. Ghermandi et al. (2007) provide an example of how a GIS-based analysis can be used to construct such a landscape-based ecological integrity indicator. This can represent a point for development of the analysis conducted in the present study.

Another issue that requires further work is the incorporation of all components of economic value in estimating ecosystem values. Value transfer is inevitably restricted by the availability of primary valuation studies for the specific ecosystems and services of policy interest. While there may be a large and growing number of ecosystem service valuation studies available, there are still gaps in the knowledge.

As shown in the case study on wetlands, there are no valuation studies that explicitly capture non-use values for wetlands. Therefore the wetland value transfer function and resulting estimated values do not reflect non-use. Filling such gaps in the available value information would require targeted primary valuation studies.

In this report we have proposed the use of meta-analytic value transfer on the grounds that it is the most practical approach for scaling up values for a large number of ecosystem sites at a national or European scale. An alternative approach would be to automate an adjusted unit transfer method, in which the closest fit ecosystem value could be retrieved and adjusted for each ecosystem site. To make a detailed comparison of these alternative value transfer approaches in terms of practicality and associated transfer errors would require further work.

References

- Barbier, E.B. 2006. Valuing ecosystem services. *Economic Policy* 22 (49): 177–229.
- Barbier, E.B., M. Acreman, and D. Knowler. 1997. *Economic valuation of wetlands: A guide for policy makers and planners*. Gland, Switzerland: Ramsar Convention Bureau.
- Barton, D.N. 1999. The quick, the cheap and the dirty. Benefit transfer approaches to the non-market valuation of coastal water quality in Costa Rica. Doctor Scientiarum Dissertation 1999:03, Department of Economics and Social Sciences, Agricultural University of Norway.
- Bateman, I. J., and A. P. Jones. 2003. Contrasting conventional with multi-level modelling approaches to meta-analysis: expectation consistency in UK woodland recreation values. *Land Economics* 79:235–58.
- Bateman, I.J., P. Cooper, S. Georgiou, S. Navrud, G.L. Poe, R. Ready, P. Riera, M. Ryan and L.A. Wainger. 2004. Scope sensitivity tests for preference robustness: an empirical examination of economic expectations regarding the economic valuation of policies for reducing acidity in remote mountain lakes. EDM-2004-03. CSERGE Working Paper.
- Bateman, I.J., S. Georgiou and I.R. Lake. 2005. The aggregation of environmental benefit values: a spatially sensitive valuation function approach. EDM 05-04. CSERGE Working Paper.
- Bateman, I.J., I.H. Langford and N. Nishikawa. 1999. The aggregation of non-user benefits: a case study illustrating different approaches. GEC 99-18. CSERGE Working Paper.
- Bateman, I.A., B.H. Day, S. Georgiou and I. Lake. 2006. The aggregation of environmental benefit values: Welfare measures, distance decay and total WTP. *Ecological Economics* 60:450–60.
- Bergland, O., K. Magnussen and S. Navrud. 1995. Benefit transfer: Testing for accuracy and reliability. Discussion Paper D-03/95, Department of Economics, Agricultural University of Norway. Revised version in Florax, R.J.G.M., P. Nijkamp and K. Willis (eds.) 2000, Comparative environmental economic assessment: Meta analysis and benefit transfer. Kluwer Academic Publishers.
- Bowker, J.M., D.B.K. English and J.C. Bergstrom. 1997. Benefit transfer and count data travel cost models: An application and test of varying parameter approach with guided whitewater rafting. Working Paper FS 97-03, Department of Agricultural and Applied Economics, University of Georgia, Athen, Georgia.

- Boyle, K.J., G.L. Poe and J.C. Bergstrom. 1994. What do we know about groundwater values? Preliminary implications from a meta-analysis of contingent valuation studies. *American Journal of Agricultural Economics* 76 (5): 1055–61.
- Brander, L.M., and Florax, J.G.M. 2007. The valuation of wetlands: primary versus meta-analysis based value transfer. In: *Environmental Valuation: Interregional and Intraregional Perspectives*, J. I. Carruthers, and B. Mundy, eds., Ashgate, Aldershot: 231–52.
- Brander, L.M., R.J.G.M. Florax and J.E. Vermaat. 2006. The empirics of wetland valuation: a comprehensive summary and a meta-analysis of the literature. *Environmental and Resource Economics* 33:223–50.
- Brander, L.M., and M.J. Koetse. 2007. The value of urban open space: meta-analyses of contingent valuation and hedonic pricing results, IVM Working Paper 07/03.
- Brouwer, R. 2000. Environmental value transfer: state of the art and future prospects. *Ecological Economics* 32:137–52.
- Brouwer, R., and I.J. Bateman. 2005. Benefits transfer of willingness to pay estimates and functions for health-risk reductions: a cross-country study. *Journal of Health Economics* 24 (3): 591–611.
- Brouwer, R., and F.A. Spaninks. 1999. The validity of environmental benefits transfer: Further empirical testing. *Environmental and Resource Economics* 14 (1): 95.
- Brouwer, R., I.J. Langford, T.C. Bateman, T.C. Crowards and R.K. Turner. 1999. A meta-analysis of wetland contingent valuation studies. *Regional Environmental Change* 1:47–57.
- Brown, G., Reed, P. and Harris, C.C. (2002). Testing a Place-Based Theory for Environmental Evaluation: an Alaska Case Study. *Applied Geography*. 22.1:49-77.
- Chattopadhyay, S. 2003. A repeated sampling technique in assessing the validity of benefit transfer in valuing non-market goods. *Land Economics* 79 (4): 576–96.
- CIESIN, Center for International Earth Science Information Network, Columbia University; and Centro Internacional de Agricultura Tropical (CIAT). 2005. Gridded Population of the World Version 3 (GPWv3): Population Density Grids. Palisades, NY: Socioeconomic Data and Applications Center (SEDAC), Columbia University. Available at: <http://sedac.ciesin.columbia.edu/gpw> (download date: 26 February 2008).
- Downing, M., and J. Ozuna. 1996. Testing the reliability of the benefit function transfer approach. *Journal of Environmental Economics and Management* 30 (3): 316–22.
- Engel, S. 2002. Benefit function transfer versus meta-analysis as policy-making tools: a comparison. In R.J.G.M. Florax, P. Nijkamp and K.G. Willis (eds.), *Comparative Environmental Economic Assessment*. Cheltenham, Edward Elgar: 133–53.
- Eshet, T., M.G. Baron and M. Shechter. 2007. Exploring benefit transfer: disamenities of waste transfer stations. *Environmental and Resource Economics* 37:521–47.
- Fennessy, M.S., A.D. Jacobs, and M.E. Kentula. 2004. Review of rapid methods for assessing wetland conditions. EPA/620/R-04/009. Washington, D.C.: US Environmental Protection Agency.
- Fotheringham, A.S. 1988. Consumer store choice and choice set definition. *Marketing Science* 7 (3): 299–310.
- Ghermandi, A., J.C.J.M. van den Bergh, L.M. Brander, H.L.F. de Groot and P.A.L.D. Nunes. 2007. Exploring diversity: a meta-analysis of wetland conservation and creation.

- In Proc. of 9th International BIOECON Conference on Economics and Institutions for Biodiversity Conservation. Cambridge, UK, September 19–21, 2007.
- Hammack, J., and G.M. Brown. 1974. *Waterfowl and wetlands: toward a bio-economic analysis*. Washington, D.C.: Resources for the Future, John Hopkins University Press.
- Hanley, N., F. Schlapfer and J. Spurgeon. 2003. Aggregating the benefits of environmental improvements: distance-decay functions for use and non-use values. *Journal of Environmental Management* 68 (3): 297–304.
- Hein, L., van Koppen, K., de Groot, R.S., van Ierland, E.C. (2006). Spatial scales, stakeholders and the valuation of ecosystem services, *Ecological Economics* 57, 209–228.
- Heston, A., R. Summers and B. Aten. 2006. Penn World Table Version 6.2. Center for International Comparisons of Production, Income and Prices, University of Pennsylvania.
- Jacobsen, J., and N. Hanley. 2007. Are there income effects on global willingness to pay for biodiversity conservation? Paper presented at the European Association of Environmental and Resource Economists conference, Thessaloniki, Greece.
- Jeong, H., and T. Haab. 2004. The economic value of marine recreational fishing: Applying benefit transfer to marine recreational fisheries statistics survey. Department of Agricultural, Environmental, and Development Economics. The Ohio State University. Working Paper: AEDE-WP-0039-04.
- Jiang, Y., S.K. Swallow and M.P. McGonagle. 2005. Context-sensitive benefit transfer using stated choice models: specification and convergent validity for policy analysis. *Environmental and Resource Economics* 31 (4): 477–99.
- Kaoru, Y., and V.K. Smith. 1995. Can markets value air quality? A meta-analysis of hedonic property values models. *Journal of Political Economy* 103:209–27.
- Kirchhoff, S., B.G. Colby, and J.T. LaFrance. 1997. Evaluating the performance of benefit transfer: An empirical inquiry. *Journal of Environmental Economics and Management* 33 (1): 75–93.
- Kirchhoff, S. 1998. Benefit function transfer vs. meta-analysis as policy-making tools: a comparison. In Proc. of Workshop on Meta-Analysis and Benefit Transfer: State-of-the-art and Prospects. Tinbergen Institute, Amsterdam, April 6–7, 1998.
- Kniivilä, M. 2005. Users and non-users of conservation areas: Are there differences in WTP, motives and the validity of responses in CVM surveys? *Ecological Economics* 59 (4): 530–9.
- Kristofersson, D., and S. Navrud. 2005. Validity tests of benefit transfer: Are we performing the wrong tests? *Environmental and Resource Economics* 30 (3): 279–86.
- Kristofersson, D., and S. Navrud. 2007. Can use and non-use values be transferred across countries? In: *Environmental value transfer: issues and methods*, S. Navrud, and R. Ready, eds., Springer, Dordrecht: 207–25.
- Lehner, B., and P. Döll. 2004. Development and validation of a global database of lakes, reservoirs and wetlands. *Journal of Hydrology* 296 (1–4): 1–22.
- Leschine, T.M., K.F. Wellman and T.H. Green. 1997. The economic value of wetlands: wetlands' role in flood protection in Western Washington. Ecology Publication no. 97–100, Washington State Department of Ecology.
- Lindhjem, H. 2007. 20 years of stated preference valuation of non-timber benefits from Fennoscandian forests: A meta-analysis. *Journal of Forest Economics* 12:251–77.

- Lindhjem, H., and S. Navrud. 2007. How reliable are meta-analyses for international benefit transfer? *Ecological Economics*, forthcoming.
- Loomis, J.B. 1996. How large is the extent of the market for public goods: evidence from a nationwide contingent valuation survey. *Applied Economics* 28:779–82.
- Loomis, J. B., H.T. Le and A. Gonzales-Caban. 2005. Testing transferability of willingness to pay for forest fire prevention among three states of California, Florida and Montana. *Journal of Forest Economics* 11 (3): 125–40.
- Loomis, J.B., F. Ward and R. Ready. 1995. Testing transferability of recreation demand models across regions - a study of corps of engineers reservoirs. *Water Resources Research* 31 (3): 721–30.
- Loomis, J.B. 2000. Vertically summing public good demand curves: an empirical comparison of economic versus political jurisdictions. *Land Economics* 76: 312–21.
- Loomis, J.B., 1992. The evolution of a more rigorous approach to benefit transfer: benefit function transfer. *Water Resources Research* 28:701–5.
- Loomis, J.B., and D.S. White. 1996. Economic benefits of rare and endangered species: summary and meta-analysis. *Ecological Economics* 18:197–206.
- Millennium Ecosystem Assessment. 2005. Ecosystem and human well-being: synthesis. Washington, DC: Island Press.
- Morrison, M., and J. Bennett. 2000. Choice modelling, non-use values and benefit transfer. *Economic Analysis and Policy* 30 (1): 13–32.
- Muthke, T., and K. Holm-Mueller. 2004. National and international benefit transfer testing with a rigorous test procedure. *Environmental and Resource Economics* 29 (3): 323.
- Navrud, S. 2007. Practical tools for value transfer in Denmark – guidelines and an example. Working Report No. 28. Danish Environmental Protection Agency.
- Nijkamp, P., G. Vindigni and P.A.L.D. Nunes. 2008. Economics valuation of biodiversity: a comparative study. *Ecological Economics*, forthcoming.
- Oglethorpe, D.R., and D. Miliadou. 2000. Economic valuation of the non-use attributes of a wetland: A case-study for Lake Kerkini. *Journal of Environmental Planning and Management* 43:755–67.
- Parsons, G.R., and M.J. Kealy. 1994. Benefit transfer in a random utility model of recreation. *Water Resources Research*. 30: 2477–84.
- Pate, J. and J. Loomis. 1997. The effect of distance on willingness to pay values: a case study of wetlands and salmon in California. *Ecological Economics* 20 (3):199–207.
- Piper, S., and W.E. Martin. 2001. Evaluating the accuracy of the benefit transfer method: A rural water supply application in the USA. *Journal of Environmental Management* 63 (3): 223–35.
- Ready, R., S. Navrud, D. Brett, D. Richard, M. Fernando, M. Susana, S. Frank and X.R. Maria. 2004. Benefit transfer in Europe: How reliable are transfers between countries? *Environmental and Resource Economics* 29 (1): 67.
- Rosenberger, R.S., and R.J. Johnston. 2007. Selection effects in meta-valuation function transfers. Paper prepared for presentation at the Aarhus Colloquium of Meta-Analysis in Economics, September 27-30, 2007, Sonderborg, Denmark.
- Rosenberger, R.S., and T.D. Stanley. 2006. Measurement, generalization, and publication: Sources of error in benefit transfers and their management. *Ecological Economics* 60 (2): 372–8.

- Rosenberger, R.S., and J.B. Loomis. 2000. Using meta-analysis for benefit transfer: in-sample convergent validity tests of an outdoor recreation database. *Water Resources Research* 36: 1097–1107.
- Rosenberger, R.S., and J.B. Loomis. 2001. Benefit transfer of outdoor recreation use values. US Department of Agriculture, Forest Service.
- Rosenberger, R.S., and T.T. Phipps. 2007. Correspondence and convergence in benefit transfer accuracy: A meta-analytic review of the literature. In: *Environmental Values Transfer: Issues and Methods*, S. Navrud, and R. Ready, eds., Springer, Dordrecht.
- Rozan, A. 2004. Benefit transfer: A comparison of WTP for air quality between France and Germany. *Environmental and Resource Economics* 29 (3): 295.
- Scott, D.A., and T.A. Jones. 1995. Classification and inventory of wetlands: a global overview. *Vegetatio* 118:3–16.
- Shrestha, R.K., and J.B. Loomis. 2001. Testing a meta-analysis model for benefit transfer in international outdoor recreation. *Ecological Economics* 39:67–83.
- Smith, T.E. 1975. An axiomatic theory of spatial discounting behavior. *Papers in Regional Science* 35 (1): 31–44.
- Smith, V.K., and Y. Kaoru. 1990. Signals or noise – explaining the variation in recreation benefit estimates. *American Journal of Agricultural Economics* 72 (2): 419–33.
- Turner, K., S. Georgiou, R. Clark, R. Brouwer. 2004. Economic valuation of water resources in agriculture: From the sectoral to a functional perspective of natural resource management. FAO, Rome.
- Van den Berg, T.P., G.L. Poe and J.R. Powell. 2001. Assessing the accuracy of benefits transfers: Evidence from a multi-site contingent valuation study of groundwater quality. In J.C. Bergstrom, K.J. Boyle, and G.L. Poe, eds., *The economic value of water quality*. Northampton, MA: Edward Elgar.
- Wagtendonk, A., and N. Omtzigt. 2003. Analysis of population density around wetland areas in GPW (Gridded Population of the World) files. SPINlab document, Institute for Environmental Studies, Amsterdam.
- Whitten, S.M., and J.W. Bennett. 1998. Private and social values of wetlands. Research Report, School of Economics and Management, University College, The University of South Wales, Canberra.
- Woodward, R.T., and Y.S. Wui. 2001. The economic value of wetland services: a meta-analysis. *Ecological Economics* 37:257–70.
- World Bank. 2006. World development indicators. <http://devdata.worldbank.org/wdi2006/contents/index2.htm> (accessed December 2007).

NOTE DI LAVORO DELLA FONDAZIONE ENI ENRICO MATTEI

Fondazione Eni Enrico Mattei Working Paper Series

Our Note di Lavoro are available on the Internet at the following addresses:

<http://www.feem.it/getpage.aspx?id=73&sez=Publications&padre=20&tab=1>
http://papers.ssrn.com/sol3/JELJOUR_Results.cfm?form_name=journalbrowse&journal_id=266659
<http://ideas.repec.org/s/fem/femwpa.html>
<http://www.econis.eu/LNG=EN/FAM?PPN=505954494>
<http://ageconsearch.umn.edu/handle/35978>
<http://www.bepress.com/feem/>

NOTE DI LAVORO PUBLISHED IN 2010

GC	1.2010	Cristina Cattaneo: Migrants' International Transfers and Educational Expenditure: Empirical Evidence from Albania
SD	2.2010	Fabio Antoniou, Panos Hatzipanayotou and Phoebe Koundouri: Tradable Permits vs Ecological Dumping
SD	3.2010	Fabio Antoniou, Panos Hatzipanayotou and Phoebe Koundouri: Second Best Environmental Policies under Uncertainty
SD	4.2010	Carlo Carraro, Enrica De Cian and Lea Nicita: Modeling Biased Technical Change. Implications for Climate Policy
IM	5.2010	Luca Di Corato: Profit Sharing under the threat of Nationalization
SD	6.2010	Masako Ikefuji, Jun-ichi Itaya and Makoto Okamura: Optimal Emission Tax with Endogenous Location Choice of Duopolistic Firms
SD	7.2010	Michela Catenacci and Carlo Giupponi: Potentials and Limits of Bayesian Networks to Deal with Uncertainty in the Assessment of Climate Change Adaptation Policies
GC	8.2010	Paul Sarfo-Mensah and William Oduro: Changes in Beliefs and Perceptions about the Natural Environment in the Forest-Savanna Transitional Zone of Ghana: The Influence of Religion
IM	9.2010	Andrea Boitani, Marcella Nicolini and Carlo Scarpa: Do Competition and Ownership Matter? Evidence from Local Public Transport in Europe
SD	10.2010	Helen Ding and Paulo A.L.D. Nunes and Sonja Teelucksingh: European Forests and Carbon Sequestration Services : An Economic Assessment of Climate Change Impacts
GC	11.2010	Enrico Bertacchini, Walter Santagata and Giovanni Signorello: Loving Cultural Heritage Private Individual Giving and Prosocial Behavior
SD	12.2010	Antoine Dechezleprêtre, Matthieu Glachant and Yann Ménière: What Drives the International Transfer of Climate Change Mitigation Technologies? Empirical Evidence from Patent Data
SD	13.2010	Andrea Bastianin, Alice Favero and Emanuele Massetti: Investments and Financial Flows Induced by Climate Mitigation Policies
SD	14.2010	Reyer Gerlagh: Too Much Oil
IM	15.2010	Chiara Fumagalli and Massimo Motta: A Simple Theory of Predation
GC	16.2010	Rinaldo Brau, Adriana Di Liberto and Francesco Pigliaru: Tourism and Development: A Recent Phenomenon Built on Old (Institutional) Roots?
SD	17.2010	Lucia Vergano, Georg Umgieser and Paulo A.L.D. Nunes: An Economic Assessment of the Impacts of the MOSE Barriers on Venice Port Activities
SD	18.2010	ZhongXiang Zhang: Climate Change Meets Trade in Promoting Green Growth: Potential Conflicts and Synergies
SD	19.2010	Elisa Lanzi and Ian Sue Wing: Capital Malleability and the Macroeconomic Costs of Climate Policy
IM	20.2010	Alberto Petrucci: Second-Best Optimal Taxation of Oil and Capital in a Small Open Economy
SD	21.2010	Enrica De Cian and Alice Favero: Fairness, Credibility and Effectiveness in the Copenhagen Accord: An Economic Assessment
SD	22.2010	Francesco Bosello: Adaptation, Mitigation and "Green" R&D to Combat Global Climate Change. Insights From an Empirical Integrated Assessment Exercise
IM	23.2010	Jean Tirole and Roland Bénabou: Individual and Corporate Social Responsibility
IM	24.2010	Cesare Dosi and Michele Moretto: Licences, "Use or Lose" Provisions and the Time of Investment
GC	25.2010	Andrés Rodríguez-Pose and Vassilis Tselios (lxxvi): Returns to Migration, Education, and Externalities in the European Union
GC	26.2010	Klaus Desmet and Esteban Rossi-Hansberg (lxxvi): Spatial Development
SD	27.2010	Massimiliano Mazzanti, Anna Montini and Francesco Nicolli: Waste Generation and Landfill Diversion Dynamics: Decentralised Management and Spatial Effects
SD	28.2010	Lucia Ceccato, Valentina Giannini and Carlo Gipponi: A Participatory Approach to Assess the Effectiveness of Responses to Cope with Flood Risk
SD	29.2010	Valentina Bosetti and David G. Victor: Politics and Economics of Second-Best Regulation of Greenhouse Gases: The Importance of Regulatory Credibility
IM	30.2010	Francesca Cornelli, Zbigniew Kominek and Alexander Ljungqvist: Monitoring Managers: Does it Matter?
GC	31.2010	Francesco D'Amuri and Juri Marcucci: "Google it!" Forecasting the US Unemployment Rate with a Google Job Search index
SD	32.2010	Francesco Bosello, Carlo Carraro and Enrica De Cian: Climate Policy and the Optimal Balance between Mitigation, Adaptation and Unavoided Damage

- SD 33.2010 Enrica De Cian and Massimo Tavoni: [The Role of International Carbon Offsets in a Second-best Climate Policy: A Numerical Evaluation](#)
- SD 34.2010 ZhongXiang Zhang: [The U.S. Proposed Carbon Tariffs, WTO Scrutiny and China's Responses](#)
- IM 35.2010 Vincenzo Denicolò and Piercarlo Zanchettin: [Leadership Cycles](#)
- SD 36.2010 Stéphanie Monjon and Philippe Quirion: [How to Design a Border Adjustment for the European Union Emissions Trading System?](#)
- SD 37.2010 Meriem Hamdi-Cherif, Céline Guivarch and Philippe Quirion: [Sectoral Targets for Developing Countries: Combining "Common but Differentiated Responsibilities" with "Meaningful participation"](#)
- IM 38.2010 G. Andrew Karolyi and Rose C. Liao: [What is Different about Government-Controlled Acquirers in Cross-Border Acquisitions?](#)
- GC 39.2010 Kjetil Bjorvatn and Alireza Naghavi: [Rent Seekers in Rentier States: When Greed Brings Peace](#)
- GC 40.2010 Andrea Mantovani and Alireza Naghavi: [Parallel Imports and Innovation in an Emerging Economy](#)
- SD 41.2010 Luke Brander, Andrea Ghermandi, Onno Kuik, Anil Markandya, Paulo A.L.D. Nunes, Marije Schaafsma and Alfred Wagtendonk: [Scaling up Ecosystem Services Values: Methodology, Applicability and a Case Study](#)

(lxxxvi) *This paper was presented at the Conference on "Urban and Regional Economics" organised by the Centre for Economic Policy Research (CEPR) and FEEM, held in Milan on 12-13 October 2009.*