

A Service of

ZBW

Leibniz-Informationszentrum Wirtschaft Leibniz Information Centre for Economics

Groen, Jan J. J.; Kapetanios, George

## Working Paper Revisiting useful approaches to data-rich macroeconomic forecasting

Staff Report, No. 327

## Provided in Cooperation with:

Federal Reserve Bank of New York

*Suggested Citation:* Groen, Jan J. J.; Kapetanios, George (2008) : Revisiting useful approaches to datarich macroeconomic forecasting, Staff Report, No. 327, Federal Reserve Bank of New York, New York, NY

This Version is available at: https://hdl.handle.net/10419/60769

#### Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

#### Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.



# WWW.ECONSTOR.EU

## Federal Reserve Bank of New York Staff Reports

Revisiting Useful Approaches to Data-Rich Macroeconomic Forecasting

> Jan J. J. Groen George Kapetanios

Staff Report no. 327 May 2008 Revised September 2009

This paper presents preliminary findings and is being distributed to economists and other interested readers solely to stimulate discussion and elicit comments. The views expressed in the paper are those of the authors and are not necessarily reflective of views at the Federal Reserve Bank of New York or the Federal Reserve System. Any errors or omissions are the responsibility of the authors.

#### **Revisiting Useful Approaches to Data-Rich Macroeconomic Forecasting**

Jan J. J. Groen and George Kapetanios *Federal Reserve Bank of New York Staff Reports*, no. 327 May 2008; revised September 2009 JEL classification: C22, C53, E37, E47

#### Abstract

We compare a number of data-rich prediction methods that are widely used in macroeconomic forecasting with a lesser known alternative: partial least squares (PLS) regression. In this method, linear, orthogonal combinations of a large number of predictor variables are constructed such that the covariance between a target variable and these common components is maximized. We show theoretically that when the data have a factor structure, PLS regression can be seen as an alternative way to approximate this unobserved factor structure. In addition, we prove that when a large data set has a weak factor structure, which possibly vanishes in the limit, PLS regression still provides asymptotically the best fit for the target variable of interest. Monte Carlo experiments confirm our theoretical results that PLS regression performs at least as well as principal components regression and rivals Bayesian regression when the data have a factor structure. But when the factor structure in the data is weak, PLS regression outperforms both principal components and Bayesian regressions. Finally, we apply PLS, principal components, and Bayesian regressions to a large panel of monthly U.S. macroeconomic data to forecast key variables across different subperiods. The results indicate that PLS regression usually has the best out-of-sample performance.

Key words: macroeconomic forecasting, factor models, forecast combination, principal components, partial least squares, Bayesian ridge regression

Groen: Federal Reserve Bank of New York (e-mail: jan.groen@ny.frb.org). Kapetanios: Queen Mary University of London (email: g.kapetanios@qmul.ac.uk). The paper has benefited from helpful comments by Alexei Onatski, James Stock, and seminar participants at Brown University, Erasmus University Rotterdam, Queen Mary University of London, the "Time Series and Panel Modelling Workshop in honor of M. Hashem Pesaran" at the Goethe University of Frankfurt, the European Meeting of the Econometric Society in Milan, the NBER-NSF Time Series Conference in Aarhus, and the CEMMAP/UCL Conference on Unobserved Factor Models at University College London. The views expressed in this paper are those of the authors and do not necessarily reflect the position of the Federal Reserve Bank of New York or the Federal Reserve System.

### 1 Introduction

It has been a standard assumption in theoretical macroeconomic modeling that agents are processing all the available quantities of information when forming their expectations for the future. Also, policymakers traditionally have looked at a vast array of indicator series in the run-up to major policy decisions, or in the words of Lars Svensson (Svensson (2005)) about what central bankers do in practice: '(l)arge amounts of data about the state of the economy and the rest of the world ... are collected, processed, and analyzed before each major decision.' However, most traditional macroeconomic prediction approaches rarely consists of models that handle more than 10 variables, because it is either inefficient or downright impossible to incorporate a much larger number of variables in a single forecasting model and estimate it using standard econometric techniques. This failure of traditional macroeconomic forecasting methods prompted a new strand of research devoted to the theory and practice of alternative macroeconomic forecasting methods that utilize large data sets.

These alternative methods can be distinguished into two main categories. As, e.g., outlined in Hendry (1995), the methods of the first category involve inherently two steps: In the first step some form of variable selection is undertaken. The variables that are chosen are then used in a standard forecasting model. Recent developments in this line of research has focussed on automated model selection procedures in order to be better able to select the optimal predictors from large data sets; see Krolzig and Hendry (2001). An alternative group of forecasting methods consists of estimation strategies that allow estimation of a single equation model that utilizes *all* the information in a large data set and not just an 'optimal' subset of the available predictor series. This is a diverse group of forecasting methods ranging from factor-based methods to Bayesian regression and forecast combination. These two groups of methods inevitably overlap. However, we feel that the step of variable selection is, and involves methods that are, sufficiently distinct to merit separate mention and treatment. Instead, we focus in this paper on the latter group of data-rich forecasting methods.

Within the group of data-rich forecasting techniques, factor methods have gained a prominent place. These methods are related to the strict factor models used in finance, but, starting with Chamberlain and Rothschild (1983), they use weaker assumptions regarding the behavior of the idiosyncratic components, which allows the use of principal components in very large data sets to identify the common factors in such a data set. Stock and Watson (2002a) and Bai (2003) further formalized the underlying asymptotic theory. Stock and Watson (2002b) proved to be the starting point of a large empirical research output where, with mixed success, a limited number of principal components extracted from a large data set are used to forecast key macroeconomic variables. However, the use of principal components does not always guarantee that the information extracted from a large number of predictors is useful for forecasting. Boivin and Ng (2006) make it clear that if the forecasting power comes from a certain factor, this factor can be dominated by other factors in a large data set, as the principal components solely provide the best fit for the large data set and not for the target variable. This could explain why in some empirical applications principal components (PC) factor models are dominated by Bayesian regression and forecast combinations. Under Bayesian regression one essentially estimates a multivariate regression consisting of all predictor variables, but with the regression coefficients shrunken to a value close to zero. Starting with Bates and Granger (1969), forecast combination involves the use of subsets of predictor variables in distinct forecasting models and the production of multiple forecasts for the target variable, which are then averaged to produce a final forecast. The distinctive feature of these two approaches is that the information in a large data set is compressed such that this has explanatory power for the target variable. Note, however, that from an econometric perspective forecast combinations are *ad hoc* in nature, whereas it has been shown in De Mol et al. (2008) that Bayesian regression is theoretically related to PC-based factor models.

In this paper we revisit the use of principal components (PC) and Bayesian regression for data-rich macroeconomic forecasting and compare these with the lesser known method of partial least squares (PLS) regression. We propose PLS regression as an alternative data-rich approach that can be used for macroeconomic forecasting using very large data sets, irrespective of whether such a data set involves having a strong factor structure or not. PLS regression is implemented for large data sets through the construction of linear, orthogonal combinations of the predictor variables such that the linear combinations maximize the covariance between the target forecast variable and each of the common components constructed from the predictor variables. Although similar in spirit to PC regression, the explicit consideration of the target forecast variable addresses a major existing criticism towards PC regression as a forecasting technique.

One significant contribution of our paper is that we provide theoretical results that relates PLS to PC for large data sets. When a factor structure is a dominant feature of large data sets we can show that in the limit PLS and PC regressions will be equivalent, and thus PLS regression can be seen as a valid alternative technique to uncover the relevant, underlying unobserved factor structure. However, when this underlying factor structure is very weak, or even absent, we prove that PLS regression still provides asymptotically the best fit for the target variable of interest. Next, we argue that the range of forecast combination techniques can be seen as restricted versions of PLS. PLS, therefore, has explicit theoretical links with the currently used range of data-rich macroeconomic forecasting tools. More specifically, we provide a unification of forms of PLS, PC, forecast combination and, via the work of De Mol *et al.* (2008), Bayesian regression, thereby linking all major forecasting tools. Finally, we consider in detail the properties of PLS, PC and Bayesian regression for forecasting using both Monte Carlo analysis and an empirical application to gauge the potential of each of these data-rich approaches. Our work suggests a clear potential for PLS regression for a variety of contexts.

In the remainder of this paper we have the following structure: Section 2 discusses the most frequently used data-rich methods for macroeconomic forecasting. Then, in Section 3, we provide an overview of PLS regression, present some results on the asymptotic behavior of PLS regression, and report on an extensive Monte Carlo study that focuses on the out-of-sample properties of PLS, PC and Bayesian shrinkage regression. Section 4 presents an empirical application where PLS and the other data-rich forecasting methods are used on a large monthly US macroeconomic data set. Finally, Section 5 concludes.

## 2 Frequently Used Methods for Data-Rich Macroeconomic Forecasting

A useful framework for studying existing methods is provided by the following general forecasting equation

$$y_t = \alpha' x_t + \epsilon_t; \quad t = 1, \dots, T, \tag{1}$$

where  $y_t$  is the target of the forecasting exercise,  $x_t = (x_{1t} \cdots x_{Nt})'$  is a vector of dimension  $N \times 1$  and thus  $\alpha = (\alpha_1 \cdots \alpha_N)'$  is also  $N \times 1$ . It is assumed that the number of indicator variables N is too large for  $\alpha$  to be determined by standard methods such as ordinary least squares (OLS). The literature has proposed a number of ways how one can deal with this issue of large-dimensional data sets, of which we provide a selective review.

#### **Factor Methods**

The most widely used class of data-rich forecasting methods are factor methods. Factor methods have been at the forefront of developments in forecasting with large data sets and in fact started this literature with the influential work of Stock and Watson (2002a). The defining characteristic of most factor methods is that relatively few summaries of the large data sets are used in forecasting equations which thereby becomes a standard forecasting equation as they only involve a few variables. The assumption is that the co-movements across the indicator variables can be captured by a  $r \times 1$  vector of unobserved factors  $F_t = (F_{1t} \cdots F_{rt})'$ , i.e.

$$\tilde{x}_t = \Lambda' F_t + e_t \tag{2}$$

where  $\tilde{x}_t$  may be equal to  $x_t$  or may involve other variables such as, e.g., lags and leads of  $x_t$  and  $\Lambda$  is a  $r \times N$  matrix of parameters describing how the individual indicator variables relate to each of the r factors, which we denote with the terms 'loadings'. In (2)  $e_t$  represents a zero-mean I(0)vector of errors that represent for each indicator variable the fraction of dynamics unexplained by  $F_t$ , the 'idiosyncratic components'. The number of factors is assumed to be small, meaning  $r < \min(N, T)$ . So, implicitly, in (1)  $\alpha' = \tilde{\alpha}' \Lambda \tilde{x}_t$ , where  $F_t = \Lambda \tilde{x}_t$ , which means that a small, r, number of linear combinations of  $\tilde{x}_t$  represent the factors and act as the predictors for  $y_t$ . The main difference between different factor methods relate to how  $\Lambda$  is estimated. The use of principal components (PC) for the estimation of factor models is, by far, the most popular factor extraction method. It has been popularised by Stock and Watson (2002a,b), in the context of large data sets, although the idea had been well established in the traditional multivariate statistical literature. The method of principal components (PC) is simple. Estimates of  $\Lambda$  and the factors  $F_t$  are obtained by solving:

$$V(r) = \min_{\Lambda, F} \frac{1}{NT} \sum_{i=1}^{N} \sum_{t=1}^{T} (\tilde{x}_{it} - \lambda'_i F_t)^2,$$
(3)

where  $\lambda_i$  is a  $r \times 1$  vector of loadings that represent the N columns of  $\Lambda = (\lambda_1 \cdots \lambda_N)$ . One, nonunique, solution of (3) can be found by taking the eigenvectors corresponding to the r largest eigenvalues of the second moment matrix X'X, which then are assumed to represent the rows in  $\Lambda$ , and the resulting estimate of  $\Lambda$  provides the forecaster with an estimate of the r factors  $\hat{F}_t = \hat{\Lambda}\tilde{x}_t$ . To identify the factors up to a rotation, the data are usually normalized to have zero mean and unit variance prior to the application of principal components; see Stock and Watson (2002a) and Bai (2003)

PC estimation of the factor structure is essentially a static exercise as no lags or leads of  $x_t$  are considered. One alternative is dynamic principal components, which, as a method of factor extraction, has been suggested in a series of papers by Forni, Hallin, Lippi and Reichlin (see, e.g., Forni *et al.* (2000, 2004) among others) is designed to address this issue. Dynamic principal components are extracted in similar fashion to static principal components but, instead of the second moment matrix, the spectral density matrices of the data at various frequencies are used. These are then used to construct estimates of the common component of the data set which is a function of the unobserved factors. This method uses leads of the data and as a result its application to forecasting has been slow for obvious reasons. Recent work by the developers of the method has addressed this issue (see, e.g., Forni *et al.* (2005)).

#### **Bayesian** (Shrinkage) Regression

Bayesian regression is a standard tool for providing inference for  $\alpha$  in (1) and there exist a large variety of approaches for implementing Bayesian regression. We will provide a brief exposition of this method. A starting point is the specification of a prior distribution for  $\alpha$ . Once this is in place standard Bayesian analysis proceeds by incorporating the likelihood from the observed data to obtain a posterior distribution for  $\alpha$  which can then be used for a variety of inferential purposes, including, of course, forecasting.

A popular and simple implementation of Bayesian regression results in a shrinkage estimator for  $\alpha$  in (1) given by

$$\hat{\alpha}_{BRR} = (X'X + vI)^{-1}X'y \tag{4}$$

where  $X = (x_1, ..., x_T)'$ ,  $y = (y_1, ..., y_T)'$  and v is a shrinkage scalar parameter. The shrinkage estimator (4) shrinks the OLS estimator, given by  $(X'^{-1}X'y)$  towards zero, thus enabling a reduction in the variance of the resulting estimator. This is a major feature of Bayesian regression that makes it useful in forecasting when large data sets are available. This particular implementation of Bayesian regression implies that elements of  $\alpha$  are small but different from zero ensuring that all variables in  $x_t$  are used for forecasting. In this sense, Bayesian regression can be linked to other data-rich approaches. When a certain factor structure is assumed in the data, Bayesian regression through (4) will forecast  $y_t$  by projecting it on a weighted sum of all Nprincipal components of X, with decaying weights, instead of projecting it on a limited number of r principal components with equal weights as in PC regression; see De Mol *et al.* (2008).

### 3 An Alternative: Partial Least Squares Regression

Partial least squares (PLS) is a relatively new method for estimating regression equations, introduced in order to facilitate the estimation of multiple regressions when there is a large, but finite, amount of regressors.<sup>1</sup> The basic idea is similar to principal component analysis in that factors or components, which are linear combinations of the original regression variables, are used, instead of the original variables, as regressors. A major difference between PC and PLS is that, whereas in PC regressions the factors are constructed taking into account only the values of the  $x_t$  variables, in PLS, the relationship between  $y_t$  and  $x_t$  is considered as well in constructing the factors. PLS regression does not seem to have been explicitly considered for data sets with a very large number of series, i.e., when N is assumed in the limit to converge to infinity. The

<sup>&</sup>lt;sup>1</sup>Herman Wold and co-workers introduced PLS regression between 1975 and 1982, see, e.g., Wold (1982). Since then it has received much attention in a variety of disciplines, especially in chemometrics, outside of economics.

latter assumption has motivated the use of PC regression for macroeconomic forecasting, as the principal components under that assumption can under certain conditions identify the common factors in the data set (see, e.g., Bai (2003)). One of our contributions is, therefore, to develop the asymptotic properties of PLS regression under similar assumptions that  $N, T \rightarrow \infty$  for data sets that have a common factor structure, and we do that in Section 3.1. In addition, we also investigate in that subsection the asymptotic properties of PLS regression when large data sets have a weak factor structure that possibly vanishes asymptotically. Section 3.2 then describes how PLS regression ties up with other data-rich techniques. Finally, Section 3.3 reports on a number of Monte Carlo experiments in which we compare PLS regression with PC-based factor modeling.

#### 3.1 PLS Regression: Methodology and Theory

There are a variety of definitions for PLS and accompanying specific PLS algorithms that inevitably have much in common. A conceptually powerful way of defining PLS is to note that the PLS factors are those linear combinations of  $x_t$ , denoted by  $\Upsilon x_t$ , that give maximum covariance between  $y_t$  and  $\Upsilon x_t$  while being orthogonal to each other. Of course, in analogy to PC factors, an identification assumption is needed, to construct PLS factors, in the usual form of a normalization.

A simple algorithm to construct k PLS factors is discussed among others, in detail, in Helland (1990). Assuming for simplicity that  $y_t$  has been demeaned and  $x_t$  have been normalized to have zero mean and unit variance, a simplified version of the algorithm is given below

#### Algorithm 1

- 1. Set  $u_t = y_t$  and  $v_{i,t} = x_{i,t}$ , i = 1, ...N. Set j = 1.
- 2. Determine the  $N \times 1$  vector of indicator variable weights or loadings  $w_j = (w_{1j} \cdots w_{Nj})'$ by computing individual covariances:  $w_{ij} = Cov(u_t, v_{it}), i = 1, ..., N$ . Construct the j-th PLS factor by taking the linear combination given by  $w'_j v_t$  and denote this factor by  $f_{j,t}$ .

- 3. Regress  $u_t$  and  $v_{i,t}$ , i = 1, ..., N on  $f_{j,t}$ . Denote the residuals of these regressions by  $\tilde{u}_t$  and  $\tilde{v}_{i,t}$  respectively.
- 4. If j = k stop, else set  $u_t = \tilde{u}_t$ ,  $v_{i,t} = \tilde{v}_{i,t}$  i = 1, ..., N and j = j + 1 and go to step 2.

This algorithm makes clear that PLS is computationally tractable for very large data sets. Once PLS factors are constructed  $y_t$  can be modeled or forecast by regressing  $y_t$  on  $f_{j,t}$  j = 1, ..., k. Helland (1988, 1990) provide a general description of the partial least squares (PLS) regression problem. Helland (1988) shows that the estimates of the coefficients  $\alpha$  in the regression of  $y_t$  on  $x_t$ , as in (1), obtained implicitly via PLS Algorithm 1 and a regression of  $y_t$  on  $f_{j,t}$  j = 1, ..., k, are mathematically equivalent to

$$\hat{\alpha}_{PLS} = V_k (V_k' X' X V_k)^{-1} V_k' X' y \tag{5}$$

with  $V_{k_1} = (X'y \quad X'XX'y \quad \cdots \quad (X'X)^{k-1}X'y), X = (x_1 \cdots x_T)'$  and  $y = (y_1 \cdots y_T)'$ . Thus, (5) suggests that the PLS factors that result from Algorithm 1 span the Krylov subspace generated by X'X and X'y, resulting in valid approximations of the covariance between  $y_t$  and  $x_t$ .

Next, we undertake a theoretical analysis of PLS when  $T, N \to \infty$ . Note that previous work focused exclusively on the case of a finite N. We consider two mutually exclusive frameworks. In the first, a factor structure exists for X. In the second there is only a very weak factor structure that can actually disappear as  $N \to \infty$  but some cross-sectional dependence is allowed. We start by analysing the factor case. Here, our goal is to show how PLS regression behaves relative to PC regression, as  $T, N \to \infty$ . Within the general regression (1), which relates  $y_t$  to  $x_t$ , denote the PLS and PC implied regression estimates by  $\hat{\alpha}_{PLS}$  and  $\hat{\alpha}_{PC}$  respectively, based on (5) and

$$\hat{\alpha}_{PC} = \Lambda' (\Lambda X' X \Lambda')^{-1} \Lambda X' y \tag{6}$$

where  $\Lambda$  is the  $r \times N$  matrix of linear combinations of the  $x_{it}$ 's that result from minimizing (3). Let the matrix norm we use be  $||A|| = tr(A'A)^{1/2}$ . We follow Stoica and Söderström (1998) who assume the following :

#### Assumption 1 Let

$$X'X = S\Psi S' + C_{\delta}$$

where  $\Psi = diag(\psi_1, ..., \psi_r)$  for some r < N, S'S = I,  $C_{\delta}$  denotes a term whose matrix norm is  $O_p(\delta)$  and  $\delta \to 0$ . This assumption implies for (5) and (6) within (1):

$$||\hat{\alpha}_{PLS} - \hat{\alpha}_{PC}|| = O_p(\delta). \tag{7}$$

An interesting aside that comes out of the above setup of Stoica and Söderström (1998) is that if the definition of k, suggested in the previous paragraph, is adopted then, for finite N at least, and assuming a reduced rank structure as in Assumption 1 we have that  $k \leq r$ . We will comment on that later in more detail.

For our large data set framework, with  $T, N \to \infty$ , we now assume the following factor structure for the explanatory variables  $x_t$ :

Assumption 2 Let  $\Sigma = \Sigma_N = [\sigma_{ij}]$  denote the  $N \times N$  second moment matrix of X.  $\Sigma$  can be factorized as follows:

$$\Sigma = \tilde{S}\tilde{\Psi}\tilde{S}' + R$$

where  $\tilde{S}\tilde{S}' = I$ ,  $\tilde{\Psi} = diag(\tilde{\psi}_{N1}, ..., \tilde{\psi}_{Nr})$  is a  $r \times r$  matrix, r < N and  $||R|| = o_p(N)$ .

Finally, we need an additional assumption regarding the second moments of  $x_t$ , i.e.,

Assumption 3 For all i, j = 1, ..., N

$$\sum_{t=1}^{T} (x_{i,t} x_{j,t} - \sigma_{i,j}) = O_p(T^{-1/2})$$

**Remark 1** Assumptions 2 and 3 deserve some comment. Assumption 2 states that the variables in X are asymptotically with respect to N collinear. Under a standard factor assumption there can be only a finite number of unbounded eigenvalues, and hence unbounded singular values, for the covariance matrix of the data. In case of Assumption 2 we can have an infinity of unbounded eigenvalues as long as the sum of all but the first r eigenvalues is  $o_p(N)$ . In particular the remainder term, R, can, in fact, be parameterized as a neglected 'weak' factor model whose eigenvalue characterization allows for unbounded eigenvalues which, however, have to grow at a rate slower than N. Assumption 3 is a mild, high level, assumption. It is sufficient to have a central limit theorem for  $(x_{i,t}x_{j,t} - \sigma_{i,j})$  for this assumption to hold.

Given the above set-up we now end-up with the following Theorem

**Theorem 1** Under Assumptions 1-3, and as  $N, T \to \infty$  sequentially, we have for the PLS and PC regression-implied estimates of the individual coefficients  $\alpha$  in the general regression framework (1):

$$||\hat{\alpha}_{PLS} - \hat{\alpha}_{PC}|| = o_p\left((NT)^{-1/2}\right).$$
 (8)

**Proof:** In order the prove this result in our case we need to show that as  $T, N \to \infty$ , Assumption 1 holds in probability asymptotically; for details see Appendix A.

Theorem 1 therefore indicates that under an assumed factor structure the PLS and PC regression approaches provide an asymptotically equivalent modeling of the relationship between  $y_t$  and  $x_t$  in a general, large-dimensional, regression set-up as in (1). The intuition behind this Theorem can be described as follows. The principal components extracted from  $x_t$  in, say, (1) provide the basis vectors that span the r-dimensional space of the underlying dominant factor model for  $x_t$ . Then, the PLS factors extracted from  $x_t$  reflect an immediate rotation of this r-dimensional space that provides the best fit for  $y_t$  and hence  $k \leq r$  in (5). Of course, in the limit that part of the r-dimensional space spanned by the principal components that is not relevant for the fit of  $y_t$  becomes redundant in (6). But exactly because the PLS factors yield in one step the most relevant rotation of this space one can expect a more efficient modeling of  $y_t$  in finite samples using PLS regression than using PC regression.

Next, we consider the theoretical properties of PLS when there exists a weak factor structure in X that can disappear as  $N \to \infty$ . We make explicit the dependence of the coefficient and variable vector on N and write the model in (1) as

$$y_t = \alpha'_N x_{N,t} + \epsilon_t \tag{9}$$

where  $x_{N,t} = (x_{1,N,t}, ..., x_{N,N,t})'$ . We make the following assumptions

#### Assumption 4

**Part I:** Let  $\Sigma = \Sigma_N = [\sigma_{ij}]$  denote the  $N \times N$  second moment matrix of X. Then,

$$\left\|\Sigma_{XX} - I\right\|_1 = o(N),$$

where  $\|.\|_1$  denotes the Minkowski 1-norm.

Part II: Further, we have

$$0 < \lim_{N \to \infty} \sup_{i,j} \frac{\alpha_{N,i}}{\alpha_{N,j}} < \infty.$$

Assumption 5 Uniformly over j = 1, ..., N

$$\frac{1}{T} \sum_{t=1}^{T} x_{j,t} y_t - \sigma_{Xy,j} = O_p \left( T^{-1/2} \right)$$

where  $\sigma_{Xy,j} = E(x_{j,t}y_t).$ 

**Remark 2** Again, Assumptions 4 and 5 deserve comment. Part I of Assumption 4 states that the covariance matrix of X becomes relatively close in a particular sense to the identity matrix as  $N \to \infty$ . It allows, e.g., for structures such as

$$\Sigma_{XX} = \begin{pmatrix} B & 0\\ 0 & I \end{pmatrix} \tag{10}$$

where B is a  $N_1 \times N_1$  symmetric positive definite matrix  $(N_1 < N)$  that can have a factor structure as outlined in Assumption 2,  $N_1 \to \infty$  and  $N_1^2/N \to 0$ . In such a set-up relatively more variables are added to X, as N grows, that are unrelated to the factor structure implied by B, which therefore gets more and more diluted. Another possibility consistent with Part I of Assumption 4 is a covariance matrix where non-diagonal elements are non-zero but tending to zero as the dimension of the matrix increases. For example, denoting the representative element of  $\Sigma_{XX}$  by  $\sigma_{ij}$ , we can set  $\sigma_{ij} = \frac{1}{i^{\kappa}j^{\kappa}}$ . Then, the Assumption holds as long as  $\kappa > 0.5$ . The same result holds if we set  $\sigma_{ij} = \frac{1}{N^{2\kappa}}$ . Note that such a covariance matrix can be obtained by specifying a factor model where the loadings depend on and tend to zero with N. Part II of Assumption 4 states that no variable in  $x_{N,t}$  has a dominant effect in the forecasting regression. This assumption can certainly be relaxed to allow, e.g., for a small subset of variables to be redundant. Further extensions are also possible but we choose to have this assumption so as to provide a relatively simple theoretical analysis while illustrating the desirable properties of PLS in a lower collinearity setting than assumed under a standard factor structure. Assumption 5 is a mild assumption of a similar nature to Assumption 3

Based on such a weak factor structure we now can formulate the following theorem:

**Theorem 2** Let  $\sigma^2$  denote the true variance of  $\epsilon_t$  in the general regression framework (1) (and thus (9)) and  $\hat{\sigma}^2$  its estimate from a PLS regression, given by steps 1-3 of Algorithm 1, using the N predictor variables that correspond with (1). Under Assumptions 4-5, and as  $N, T \to \infty$  sequentially,  $\hat{\sigma}^2 - \sigma^2 = o_p(1)$ .

**Proof:** The proof is given in Appendix B.

Theorem 2 suggests that under a weak factor (or 'near factor') structure PLS regression is still able to estimate a model that has asymptotically the best fit, as the theorem implies that the PLS regression  $R^2$  will converge to the population  $R^2$  of the general regression (1). Hence, PLS regression should continue to do well in modeling a target variable even if the collinearity amongst the predictor variables is not as strong as it is assumed to be for conventional factor models.

#### **3.2** Linking Alternative Data Rich Methods

In this subsection, we briefly consider links between alternative data rich methods such as PLS, PC and Bayesian regression given the results presented in the previous subsection. We first note that Garthwaite (1994) provides a rationale to cast (*ad hoc*) forecast combinations in terms of the above described PLS framework. Essentially what Garthwaite (1994) shows is that a general PLS algorithm like Algorithm 1 can be expressed in terms of sequences of univariate regressions, i.e.,

#### Algorithm 2

- 1. Set  $u_t = y_t$  and  $v_{i,t} = x_{i,t}$ , i = 1, ...N. Set j = 1.
- Regress u<sub>t</sub> on v<sub>i,t</sub>, i = 1,...,N and denote the OLS estimate of the coefficient of each regression by β<sub>i</sub>. Construct the j-th PLS factor by taking the weighted average of β<sub>i</sub>v<sub>it</sub>:
   f<sub>j,t</sub> = w̃'<sub>j</sub>v<sub>t</sub> with w̃<sub>j</sub> = ((β<sub>1</sub>w<sub>1j</sub>) ··· (β<sub>N</sub>w<sub>Nj</sub>))' where (w<sub>1j</sub> ··· w<sub>Nj</sub>) are given.
- 3. Regress  $u_t$  and  $v_{i,t}$ , i = 1, ..., N on  $f_{j,t}$ . Denote the residuals of these regressions by  $\tilde{u}_t$  and  $\tilde{v}_{i,t}$  respectively.
- 4. If  $j = k_1$  stop, else set  $u_t = \tilde{u}_t$ ,  $v_{i,t} = \tilde{v}_{i,t}$  i = 1, ..., N and j = j + 1 and go to step 2.

Therefore, when in this algorithm one sets  $(w_{1j} \cdots w_{Nj}) = (Var(v_{1t}) \cdots Var(v_{Nt}))$  Algorithm 1 follows, but if one assumes  $(w_{1j} \cdots w_{Nj}) = (\frac{1}{N} \cdots \frac{1}{N})$  than in the one-factor case the Capistrán and Timmermann (2008) projection on equal-weighted mean (PEW) forecast combination approach follows. In general, forecast combinations can be interpreted through Algorithm 2 as restricted approximations to one-factor PLS regression, with alternative specifications for  $(w_{1j} \cdots w_{Nj})$  and often with zero intercept and slope coefficients in the final forecast regression.<sup>2</sup> By interpreting forecast combinations as a form of PLS regression, Theorems 1 and 2 provide the underpinning for the relatively good performance of forecast combinations vis-à-vis PC regressions within different data environments.<sup>3</sup> Note, though, that PLS is much more general and it allows for several factors to be included in the forecast regression.

In addition, De Mol *et al.* (2008) prove the existence of a form of asymptotic equivalence between PC regression and Bayesian regression when the underlying data comply with a factor structure.<sup>4</sup> Thus, given that structure, Bayesian regression should, via Theorem 1, be asymptotically equivalent to PLS regression and, under the one-factor assumption, forecast combinations.

 $<sup>^{2}</sup>$ Granger and Ramanathan (1984) suggest regressing individual forecasts on the target variable with the resulting parameter estimates serving as combination weights. More generally, by specifying a loss function one can derive combination weights that are optimal under that specific loss function (Elliott and Timmermann (2004)). Timmermann (2006) provides a comprehensive survey of the forecast combination literature.

<sup>&</sup>lt;sup>3</sup>For example, Faust and Wright (2007) show that forecast combination methods provide better out-of-sample performance than factor methods when applied to high-dimensional panels of U.S. macroeconomic data.

<sup>&</sup>lt;sup>4</sup>It is worth noting that the assumption made by De Mol *et al.* (2008) on the idiosyncratic part of their entertained factor model bears similarities to our Assumption 2. In particular, just like the implication of our Assumption 2, discussed in Remark 1, the idiosyncratic component of their factor model can accommodate a residual 'weak' factor model in the sense that the eigenvalues implied by that factor model can be unbounded but have to grow at a rate slower than N.

Therefore, the introduction of the PLS regression framework provide a means to asymptotically tie together different existing data-rich forecasting methods and provides a theoretical rationale for the common empirical finding that different data-rich approaches have similar performance.

#### 3.3 Monte Carlo Analysis

In this subsection, we explore through Monte Carlo experiments the finite sample performance of PLS regression relative to PC regression and Bayesian regression. We consider both the case when the data have a factor structure and the case where there is no factor structure.

#### Monte Carlo Set-up

For our Monte Carlo experiments we consider the following data generating processes (DGPs):

$$y_{t} = \alpha'(x_{1,t} \cdots x_{N,t})' + \epsilon_{t} = \alpha' x_{t} + \epsilon_{t}, \quad t = 1, ..., T,$$

$$x_{t} = \Lambda' f_{t} + u_{t},$$

$$f_{t} = (f_{1,t} \cdots f_{r,t})' \sim \text{iid}N(\mathbf{0}, I_{r}), \quad r << N,$$

$$\epsilon_{t} = \sqrt{cN}\varepsilon_{t},$$
(11)

with the  $N \times 1$  vector of regression parameters  $\alpha = (\alpha_1 \cdots \alpha_N)'$ ,  $\epsilon_t$  is a zero-mean disturbance term that we discuss in more detail later, and  $\Lambda = (\lambda_1 \cdots \lambda_N)$  is a  $r \times N$  matrix of factor loadings that corresponds with the  $r \times 1$  vector of factors  $f_t$  with  $\lambda_i = (\lambda_{i,1} \cdots \lambda_{i,r})'$ . The DGP for  $x_t$  in (11) uses a  $N \times 1$  vector of zero-mean disturbances  $u_t = (u_{1,t} \cdots u_{N,t})'$ . The individual regression coefficients in (11) are determined as  $\alpha_i \sim \text{iid } N(0,1)$  and the disturbances for the Nexplanatory variables are determined in a similar manner:  $u_{i,t} \sim \text{iid } N(0,1)$ .

We consider a number of cases for the factor loadings, which we summarize as:

Case I: 
$$\lambda_{i,j} \sim \operatorname{iid} N(0,1)$$
 for  $i = 1, \dots, N$  &  $j = 1, \dots, r$ .  
Case II:  $\lambda_{i,j} \begin{cases} = \tilde{\lambda}_{i,j} \sim \operatorname{iid} N(0,1) & \text{for } i = 1, \dots, N_1, N_1 = N^{\kappa_1} \ (\kappa_1 = 0.25, \ 0.75) \ \& \ j = 1, \dots, r. \\ = 0 & \text{for } i = N_1 + 1, \dots, N \ \& \ j = 1, \dots, r. \end{cases}$   
Case III:  $\lambda_{i,j} = \frac{\tilde{\lambda}_{i,j}}{N^{\kappa_2}}$  and  $\tilde{\lambda}_{i,j} \sim \operatorname{iid} N(0,1)$  for  $i = 1, \dots, N, \ j = 1, \dots, r \ \& \ \kappa_2 = 0.25, \ 0.75.$   
Clearly, Case I represents a standard factor model where the predictor variables are driven by

Clearly, Case I represents a standard factor model where the predictor variables are driven by r common factors. Cases II and III, on the other hand, imply much weaker factor structures

in  $x_t$  than assumed under Case I and these become progressively weaker as the cross-section dimension N increases. Under Case II we assume a structure of the form (10) where we set  $N_1 = N^{\kappa_1}$  and  $\kappa_1 = 0.25$ , 0.75. This is operationalized by using a factor model where the factors are not pervasive but affect a subset  $N_1$  of the variables in  $x_t$ . The subset of predictor variables  $(N_1 - N)$  are non-informative for the factors and this subset will dominate the panel of N predictor variables as  $N \to \infty$ , where  $\kappa_1$  determines the speed with which this occurs. Case III assumes that the representative non-diagonal element of the covariance matrix of  $x_t$  is given by  $\frac{1}{N^{2\kappa_2}}$ . We obtain this by using factor loadings that tend to zero as  $N \to \infty$ . As such Case III represents a case where we have a 'near-factor model' in  $x_t$  in which  $\kappa_2$  determines how close this structure is to a standard factor model, and the larger  $\kappa_2$  the further away it is from such a standard factor model.

An important parameter for the Monte Carlo study is the population  $R^2$  of the  $y_t$  regression equation in (11). We control this by controlling the variance of the  $y_t$  disturbance term  $\epsilon_t$  in (11) through  $c = (1 + r)\tilde{c}$ , where  $\varepsilon_t \sim iidN(0, 1)$ . Setting  $\tilde{c} = 1, 4, 9$  gives a population  $R^2$ equal to 0.5, 0.2 and 0.1 in case of the standard factor model (Case I). For the case of weak factors (Cases II and III) we assume that r = 0 for the purposes of setting c. Therefore, in this case, the calibrated population  $R^2$  is slightly lower than for the standard factor case but this is a minor deviation since the factor loadings under Cases II and III are small. These values for  $R^2$  provide, in our view, reasonable representations of empirically relevant situations.

When we assume a standard factor structure, we generate data through (11) for r = 1, 6 and we set that the assumed number of PC factors,  $k_2$ , is equal to the true number of factors, r, when carrying out PC regression. For the case of PLS regression we argued in Section 3.1 that the number of PLS factors,  $k_1$ , is at the most equal and very likely smaller than the number of PC factors, as not all factors have to be relevant for the target variable. This is more so the case here, as both the factor loadings as well as the regression parameters  $\alpha$  are randomly generated in our simulations and thus the factors need not be of equal strength for  $y_t$  (which indeed is also true in reality). Therefore, for PLS regression we set the number of factors  $k_1$  to 1 and 3 to correspond to the respective  $k_2$  PC factors. For the weak factor cases, Cases II and III, we generate data based on r = 1 and assume in case of PLS and PC regressions  $k_1 = k_2 = 1$  in order to focus on how a decreasing amount of collinearity within  $x_t$  affects the relative performance of these methods rather than differences in relevance for  $y_t$  across a multiple of factors. For Bayesian regression we follow De Mol *et al.* (2008) and set the shrinkage parameter proportional to the cross-section dimension N of the explanatory series: qN, where q = 1, 5.

We evaluate the competing methods using the relative out-of-sample mean squared prediction error (MSE) compared to PC regression. To construct these relative MSEs, we generate T + 100data points according to (11) and estimate all models over the first T observations. We then use the implied regression weights to do one-step ahead forecasting and get forecast errors for T + $1, \ldots, T+100$ . The results across all variants are computed for N, T = 20, 30, 50, 100, 200, 400and are each based on 1000 Monte Carlo replications.

#### Monte Carlo Results

Starting with the standard factor case in Tables 1-2, one notes that the performance of PLS and PC regressions are fairly similar. Note, though, that the relative performance of PLS regression improves when the number of entertained PC and PLS factors increases, which reflects the theoretical result of Section 3.1. The only other noteworthy feature for this case, given already available Monte Carlo work on the relative performance of PC *versus* Bayesian regressions, is the confirmation of the good predictive performance of Bayesian regression in a large number of cases.

For the intermediate cases of weak factor models a dichotomy in performance emerges between PC regression on one side and both PLS and Bayesian regression on the other. Already with moderate degrees of factor weakness under both cases II and III, see Tables 3 and 4, it becomes clear that PC regression performs poorly when the dimensions of the underlying panel of predictor variables become large, as it is almost always outperformed by the two other methods. PLS regression, on the other hand, performs in a large number of cases as well as Bayesian regression, which in Tables 3 and 4 has a slight edge vis-à-vis PLS regression when an appropriate shrinkage parameter is chosen. Tables 5 and 6 also report simulation results under Cases II and III for the factor loadings, but now with much more severe factor weakness than in Tables 3 and 4. What becomes clear from Tables 5 and 6 is that the relative performance of PLS regression improves substantially to a point that it now performs at least as well, and in a lot of cases better, as Bayesian regression.<sup>5</sup>

To conclude we see that our Monte Carlo study suggests a great advantage for PLS regression compared to all other methods we considered in terms of forecasting performance. Especially in the case where the data do not have a standard parsimonious representation such as a (standard) common factor structure.

## 4 Empirical Applications

In this section we further analyze the properties of PLS, PC and Bayesian regressions within an empirical context. We describe in Section 4.1 how we implement the different methods on the data. In Section 4.2 we provide details on the utilized large data set and on how we construct our predictor and explanatory variables. Finally, the results of the different forecast exercises are reported in Section 4.3.

#### 4.1 Implementation of the Data-Rich Methods and the Forecast Comparison

We follow standard practice in the macroeconomic forecasting literature and use as benchmark for our data-rich based forecasts an autoregressive (AR) model

$$\Delta y_{t+h,t} = \alpha^{h} + \sum_{i=1}^{p} \rho_{i} \Delta y_{t-i+1,t-i} + \epsilon_{t+h,t}, \quad t = 1, \dots, T$$
(12)

with  $\Delta y_{t+h,t} = y_{t+h} - y_t$  for h > 0 and  $\Delta y_{t-i+1,t-i} = y_{t-i+1} - y_{t-i}$  for  $i = 1, \ldots, p$ . The number of lagged first differences p in (12) is determined by sequentially applying the standard Schwarz (1978)'s BIC starting with a maximum lag order of  $p = p_{max}$  down to p = 1. Next, we use as a benchmark the unconditional mean,

$$\Delta y_{t+h,t} = \alpha^h + \epsilon_{t+h,t},\tag{13}$$

which implies a random walk (RW) forecast for the level of the forecast variable  $y_t$ . Our assessment of the forecasting performance of the data-rich methods relative to pure AR-based and random walk-based forecasts is based on the square root of the mean of the squared forecast

 $<sup>{}^{5}</sup>$ Indeed, if one completely turns off the factor structure in (11) one would get a similar result as in Tables 5 and 6. These simulation results are available upon request from the authors.

errors (RMSE). In Section 4.3 we will report ratios of the RMSE of the respective data-rich forecasting approaches relative to the RMSE based on either (12) or (13). Superior out-of-sample performance of a data-rich method relative to these benchmarks is, obviously, indicated by a RMSE ratio smaller than one.

Our data-rich forecasts of h period-ahead changes in  $y_t$  are generated using a model that adds the information extracted from the N explanatory variables in the  $N \times 1$  vector  $X_t = (x_{1,t} \cdots x_{N,t})$  to the benchmark models (12) and (13), i.e., respectively

$$\Delta y_{t+h,t} = \alpha^h + \beta^{h'} F(X_t) + \sum_{i=1}^p \rho_i \Delta y_{t-i+1,t-i} + \epsilon_{t+h,t}$$
(14)

and

$$\Delta y_{t+h,t} = \alpha^h + \beta^{h'} F(X_t) + \epsilon_{t+h,t}.$$
(15)

where  $\beta^h$  is  $r \times 1$ . In (14) and (15)  $F(X_t)$  represents a  $r \times 1$  function of  $X_t$  that compresses the information in the N indicator variables, i.e. through principal components (PC), partial least squares (PLS) or by estimating the  $\beta^h$ 's through Bayesian regression (BR, where r = N). We operationalize the construction of  $F(X_t)$  on our data sets as follows:

#### **Principal Components Regression**

Following Stock and Watson (2002b) we take our  $T \times N$  matrix of N indicator variables  $X = (X'_1 \cdots X'_T)'$  and normalize this such that the variables are in zero-mean and unity variance space, which results in the  $T \times N$  matrix  $\tilde{X}$ . We then compute the r eigenvectors of the  $N \times N$  matrix  $\tilde{X}'\tilde{X}$  that correspond to the first r largest eigenvalues of that matrix, which we assemble in the  $N \times r$  matrix  $\Lambda^r$ . These eigenvectors are then used to approximate the common factors F that determine the series in X, i.e.,  $F = \tilde{X}\Lambda^r$ , which gives us  $F(X_t)$  in (14) and (15).

Our forecasting models will be updated based on an expanding window of historical data, which in case of the principal components-based models evolves as follows:

- 1. First forecast for all h is generated on  $t_0$ .
- 2. Extract r principal components  $F_t$  from the N indicator variables over the sample  $t = 1, \ldots, t_0 h$ .

- 3. Estimate either (14) or (15) with  $F(X_t) = F_t$  over the sample  $t = 1, \ldots, t_0 h$  for each h.
- 4. Extract r principal components  $F_t$  from the N indicator variables N over the sample  $t = 1, \ldots, t_0$ .
- 5. Generate for h the forecast  $\Delta \hat{y}_{t+h,t}$  using the parameter estimates from step 3 and  $F_t$  from step 4.
- 6. Repeat for  $t_0 + 1, \ldots, T h$  for each h.

#### **Bayesian Regression**

When Bayesian regression is used to compress the forecast information in the N indicator variables,  $F(X_t)$  in (14) and (15) simply equals  $X_t$ , whereas  $\beta^h$  is estimated with the shrinkage estimator (4). As in the case of principal components-based regressions we use a normalized version of the  $T \times N$  matrix of explanatory variables  $X = (X'_1 \cdots X'_T)'$ , indicated with  $\tilde{X}$ , and we also demean  $\Delta y_{t+h,t}$  first before we estimate  $\beta^h_{BRR}$ . By doing this we follow De Mol *et al.* (2008), as the regression can then be interpreted as a Bayesian regression with a Gaussian prior. In case of (14) we first regress both the demeaned  $\Delta y_{t+h,t}$  and the  $\tilde{X}$  on  $\Delta y_{t-i+1,t-i}$  for  $i = 1, \ldots, p$ , and use the resulting residuals in (4) to estimate  $\beta^h$ . Estimates for the intercept term and the  $\rho_i$ 's in (14) can then be trivially recovered.

The Bayesian regression forecasts are updated using an expanding window of data:

- 1. First forecast for all h is generated on  $t_0$ .
- 2. Estimate either (15) or (14) with (4) for  $\beta^h$  using  $\tilde{X}$  over the sample  $t = 1, \ldots, t_0 h$  for each h.
- 3. Generate for h the forecast  $\Delta \hat{y}_{t+h,t}$  using the parameter estimates from step 2.
- 4. Repeat for  $t_0 + 1, \ldots, T h$  for each h.

#### Partial Least Squares Regression

With partial least squares (PLS) regression,  $F(X_t)$  in (14) and (15) is constructed by computing r orthogonal combinations from the N indicator variables, where the weights of the individual

indicator variables in the respective combinations are chosen such that the covariance with  $\Delta y_{t+h,t}$  is maximized. The general PLS algorithm from Section 3.1 can be implemented for macroeconomic forecasting as follows:

#### Algorithm 3

 Denote, as before, the T × N matrix of indicator variables, each normalized to have a zero mean and unit variance, as X and demean the predictor variable, i.e.

$$\Delta \dot{Y}_{h} = \left( I_{T} - \iota(\iota'^{-1}\iota') \begin{pmatrix} \Delta y_{h+1,1} \\ \vdots \\ \Delta y_{T,T-h} \end{pmatrix} \right).$$

2. The r PLS factors  $F_{1,t}^{PLS}, \ldots, F_{r,t}^{PLS}$  and their loadings  $w_1, \ldots, w_r$  are iteratively build up through projections on lower order PLS factors followed by computing the covariances between the resulting residuals of the columns from  $\tilde{X}$  and those of  $\Delta \dot{Y}_h$ :

$$F_l^{PLS} = \tilde{X}_{l|l-1} w_l; \quad w_l = \frac{1}{T-1} \tilde{X}'_{l|l-1} \Delta \dot{Y}_{h,l|l-1} \quad for \quad l = 1, \dots, r$$
(16)

where for l = 1 (the first PLS factor)

$$\tilde{X}_{1|0} = \tilde{X}, \quad \Delta \dot{Y}_{h,1|0} = \Delta \dot{Y}_h,$$

and for l > 1

$$\tilde{X}_{l|l-1} = \left(I_T - F_{l-1}^{PLS} (F_{l-1}^{PLS'} F_{l-1}^{PLS})^{-1} F_{l-1}^{PLS'} \right) \tilde{X}_{l-1|l-2} \quad and$$
$$\Delta \dot{Y}_{h,l|l-1} = \left(I_T - F_{l-1}^{PLS} (F_{l-1}^{PLS'} F_{l-1}^{PLS})^{-1} F_{l-1}^{PLS'} \right) \Delta \dot{Y}_{h,l-1|l-2}.$$

3. Finally, we simply plug in the r PLS factors  $F_t^{PLS} = (F_{1,t}^{PLS} \cdots F_{r,t}^{PLS})'$  from (16) in the predictive regression (15) which we estimate in the standard way:

$$\Delta y_{t+h,t} = \alpha^h + \beta^{h'} F_t^{PLS} + \epsilon_{t+h,t}.$$
(17)

When lagged predictor variables are included in the predictive regression, as in (14), one needs to control for the effect of  $\Delta y_{t,t-1}, \ldots, \Delta y_{t-p+1,t-p}$  on the covariances between  $\Delta y_{t+h,t}$ and  $x_{1,t}, \ldots, x_{N,t}$ . Like in the BR case we do that by projecting the demeaned  $\Delta y_{t+h,t}$  as well as the columns of  $\tilde{X}$  on  $\Delta y_{t,t-1}, \ldots, \Delta y_{t-p+1,t-p}$ , and then using the resulting residuals in Algorithm 3 in order to be able to construct a model like

$$\Delta y_{t+h,t} = \alpha^h + \beta^{h'} F_t^{PLS} + \sum_{i=1}^p \rho_i \Delta y_{t-i+1,t-i} + \epsilon_{t+h,t}.$$
(18)

Finally, forecasts from (17) and (18) are generated as follows, again using an expanding window of historical data:

- 1. First forecast for all h is generated on  $t_0$ .
- 2. Extract r PLS factors  $F_t^{PLS}$  from the N indicator variables over the sample  $t = 1, \ldots, t_0 h$  for each h based on Algorithm 3.
- 3. Estimate either (17) or (18) over the sample  $t = 1, ..., t_0 h$  for each h.
- 4. Extract r PLS factors  $F_t^{PLS}$  from the N indicator variables over the sample  $t = 1, ..., t_0$  for each h using the corresponding loadings  $w_r$  from step 2 based on Algorithm 3.
- 5. Generate for h the forecast  $\Delta \hat{y}_{t+h,t}$  using the parameter estimates from step 3 and  $F_t^{PLS}$  from step 4.
- 6. Repeat for  $t_0 + 1, \ldots, T h$  for each h.

This leaves us with one more issue: either the appropriate number of factors r for PC or PLS regression, or in case of Bayesian regression the appropriate value for the shrinkage parameter v in (4). In the latter case, we are not aware of the availability of a theoretically justified approach to select the optimal shrinkage parameter v for a given data set. Instead, De Mol *et al.* (2008) suggest that an appropriate Bayesian regression for a large data set under a factor structure should be based on a shrinkage parameter that is proportional to the cross-section dimension of the data set. Therefore, for BR-based estimation of (14) and (15) we use in (4) v = qN with q = 1 and 5, which is in the range for v that De Mol *et al.* (2008) found useful in their

applications. To facilitate a fair comparison between the BR-based forecasts and both the PCbased and PLS-based forecasts, we therefore estimate the PC-based and PLS-based versions of (14) and (15) using a range of fixed numbers of factors. In case of PC-based forecasts, we will consider the predictive performance of (14) and (15) using r = 2, 4 and 6 PC factors, whereas in case of PLS-based forecasts we consider the performance using r = 1, 2 and 3 PLS factors.

#### 4.2 The Data Set and Variable Construction

Stock and Watson (2007) reorganize the large panel of macroeconomic, financial and surveybased predictor variables for the United States from Stock and Watson (2002b) and update the span of the data to the end of 2006. Both our forecast variables and our panel of indicator variables are extracted from Stock and Watson (2007) <sup>6</sup> and we focus on the 109 monthly series from this U.S. data set, which before transformation span a sample starting in January 1959 and ending in December 2006.

The panel of predictor variables consist of 105 series spanning real variables, labor market data, data on price indices (and subcomponents) and wages, money and credit series, asset prices and surveys. The predictor variables are transformed such that they are I(0), which in general means that the real variables are expressed in log first differences and we use simply first differences of series expressed in rates, such as interest rates; see Appendix C for more details. With respect to nominal series we transform these into first differences of annual growth rates in order to guarantee that the dynamic properties of these transformed series are comparable to those of the rest of the panel, as for example motivated in D'Agostino and Giannone (2006, Appendix B).<sup>7</sup> Hence, after transforming the indicator variables we end up with an effective span of the data that starts in February 1960 (i.e. 1960.2) and ends in December 2006 (i.e. 2006.12).

This predictor variables panel will be used to forecast appropriate transformations of CPI inflation, industrial production, the unemployment rate and the federal funds rate. These fore-

 $<sup>^{6}</sup>$ We are very grateful to Mark Watson who provided us with the underlying raw data from Stock and Watson (2007).

<sup>&</sup>lt;sup>7</sup>This particular transformation acknowledges that series like log price levels and log money aggregate levels behave as if they are I(2), possibly because of mean growth shifts due to policy regime shifts, financial liberalizations and other phenomena.

cast variables are not part of the panel of predictors and are transformed such to guarantee stationarity:

$Y_t$	$\Delta y_{t,t-1}$	$\Delta y_{t+h,t}$
CPI index Industrial Production index Unemployment rate Federal Funds rate	$ \begin{array}{c} \Delta \ln Y_{t,t-12} - \Delta \ln Y_{t-1,t-13} \\ \Delta \ln Y_{t,t-1} \\ \Delta Y_{t,t-1} \\ \Delta Y_{t,t-1} \end{array} $	$ \Delta \ln Y_{t+h,t+h-12} - \Delta \ln Y_{t,t-12}  \Delta \ln Y_{t+h,t}  \Delta Y_{t+h,t}  \Delta Y_{t+h,t}  \Delta Y_{t+h,t} $

As described in the previous subsection, the forecasting models are updated based on an expanding window of data and all forecasts are direct forecasts for 4 horizons (in months): h = 1, h = 3, h = 12 and h = 24, which are horizons commonly analyzed in the literature. The forecast evaluation spans three samples: January 1972 - December 2006, January 1972 - December 1984 and January 1985 - December 2006. The latter two sub-samples split the first sample in two around the start of the 'Great Moderation', e.g., McConnell and Perez-Quiros (2000) and Sensier and van Dijk (2004) find evidence for a downward, exogenous, shift in the volatility of a large number of U.S. macroeconomic time series around 1985. This sample split is of particular importance for forecasting U.S. economic time series, as it has been shown that it is difficult for a lot of data-rich, approaches (including *Greenbook* projections from the Federal Reserve Board) to beat simple, non-structural benchmarks like RW and AR models after the occurrence of the 'Great Moderation'.<sup>8</sup>

#### 4.3 Forecasting Results

As discussed in Section 4.1, we will assess the forecasting performance of our three data-rich forecast methods with two simple benchmark forecasts: those based on an autoregressive (AR) specification and those based on the unconditional mean or random walk (RW) model (respectively (12) and (13)). The first set of evaluation results can be found in Table 7 and relate to forecasting changes in annual CPI inflation. Across the three evaluation samples, the full 1972-2006 evaluation sample and the two sub-samples, PLS regression does dominates in 11 out of

<sup>&</sup>lt;sup>8</sup>See, for example, D'Agostino *et al.* (2006) who compare PC-based, VAR-based and *Greenbook* forecasts for U.S. inflation and economic growth with simple benchmarks for both pre- and post-Great Moderation samples.

the 24 cases (i.e., 3 evaluation samples times 2 benchmark models times 4 forecasting horizons). In fact, only Bayesian regression can, up to a certain extent compete with PLS regression. Note, though, that when it loses out the PLS approach still provides a close second best. Of course, the overall forecasting performance of the data-rich approaches is less over the post-Great Moderation period, especially  $vis-\hat{a}-vis$  the AR benchmark over longer horizons, which is consistent with findings elsewhere in the literature.

The Great Moderation has less of a negative effect on the predictive performance of our data-rich methods for industrial production growth and unemployment rate changes; see the results for 1985-2006 in Tables 8-9. PLS regression is clearly the overall winner in this case, as it dominates the other methods in, respectively, 13 and 20 out of the total of 24 forecasting exercises in each of these tables. In particular in case of unemployment PC regression performs poorly, and essentially only for the one-month horizon Bayesian regression is able to perform better than PLS regression.

Finally, we turn to the results for the federal funds rate in Table 10. As the federal funds rate is determined by the Federal Reserve Board, which sets the target for the federal funds rate by taking into account both nominal and real developments, data-rich methods, which feed off of both nominal and real series, are expected to perform well in predicting fed funds rate changes. This seems certainly to be the case for PLS and Bayesian regressions, although Bayesian regression performs relatively poorly for the post-Great Moderation 1985-2006 period. PLS regression, however, performs well throughout the different evaluation samples: it outperforms the other two approaches in 18 out of the 24 evaluations.

The empirical forecast evaluations in this subsection lead to a number of general observations. First, it is clear that the PLS-based forecast models are, generally speaking, amongst the best performing models. And even in the minority of cases that they are outperformed by either PC or Bayesian regression approaches, the results in Tables 7-10 indicate that they are close competitors. Note also that in Tables 7-10 the performance of methods that use PLS factors and Bayesian shrinkage estimators are pretty close, with PLS usually having the edge.

### 5 Conclusions

In this paper we have revisited a number of approaches for prediction with many predictors that are widely used in macroeconomic forecasting and we compare these with a less widely known alternative approach: partial least squares (PLS) regression. Under PLS regression, one constructs a number of linear combinations of the predictor variables such that the covariances between the target variable and each of these linear combinations are maximized.

We provide theoretical arguments for the asymptotic similarity between principal components (PC) regression and PLS regression when the underlying data has a common factor structure. When the factor structure in a large data set is weak, and possibly vanishes as  $N \to \infty$ , we prove that PLS regression will continue to provide a model with the best asymptotic fit for a target variable. We also argue that forecast combinations can be considered as a specific form of PLS regression. Hence, whether or not a large panel of predictors has a clear factor structure, we would expect PLS regression, like Bayesian ridge regression, to do well in macroeconomic forecasting.

An extensive Monte Carlo analysis, which compares PC regression and Bayesian regression with PLS regression, yields a number of interesting insights. Firstly, when we assume that the predictors relate to the target variable through a standard common factor structure, PLS regression is shown to have an out-of-sample performance that is at least comparable to, and often better than PC regression. PLS regression also compares well to Bayesian regression under this data specification, especially when the number of relevant factors for the target variable increases. When the relation between the predictors and the target variable only has a weak factor structure, PLS regression clearly has the edge in terms of out-of-sample forecasting performance.

Finally, we apply PC, PLS and Bayesian regression on a panel of 105 U.S. monthly macroeconomic and financial variables to forecast CPI inflation, industrial production, unemployment and the federal funds rate, where these forecasts are evaluated across several sub-samples. PLS regression turns out to be generally the best performing method, and even in the few cases when it is outperformed by PC or Bayesian regression, PLS regression remains a close competitor.

		TOPT		ידורמידי			actor	(nem)	TACTON	TAT OA		n modu			2m (In)	-	, 1)		
				$R^2$ =	=0.5					$R^2 = ($	0.33					$R^2 =$	=0.2		
	T/N	20	30	50	100	200	400	20	30	50	100	200	400	20	30	50	100	200	400
	20	0.92	0.96	1.00	1.02	1.04	1.05	1.01	1.02	1.05	1.06	1.07	1.07	1.06	1.07	1.08	1.09	1.09	1.10
	30	0.90	0.94	0.98	1.01	1.03	1.03	0.99	1.01	1.02	1.03	1.05	1.05	1.03	1.04	1.05	1.06	1.06	1.07
PLS	50	0.88	0.93	0.96	0.99	1.01	1.02	0.96	0.98	1.00	1.02	1.03	1.03	1.01	1.02	1.03	1.03	1.04	1.04
	100	0.86	0.87	0.91	0.95	0.96	0.96	0.91	0.91	0.98	0.96	0.96	0.97	0.97	0.96	0.99	0.98	0.99	0.98
	200	0.83	0.87	0.91	0.94	0.96	0.97	0.91	0.92	0.94	0.95	0.96	0.96	0.94	0.95	0.96	0.97	0.96	0.97
	400	0.84	0.88	0.92	0.96	0.97	0.98	0.92	0.95	0.96	0.97	0.98	0.98	0.95	0.96	0.97	0.98	0.98	0.98
	20	0.71	0.78	0.86	0.93	0.96	0.98	0.94	0.95	0.97	0.98	0.99	0.99	1.04	1.03	1.02	1.01	1.00	1.00
	30	0.62	0.70	0.80	0.89	0.94	0.97	0.90	0.94	0.96	0.98	0.99	0.99	1.04	1.04	1.03	1.02	1.00	1.00
BR(1)	50	0.53	0.59	0.70	0.83	0.90	0.95	0.86	0.90	0.93	0.96	0.98	0.99	1.01	1.04	1.04	1.03	1.01	1.01
	100	0.38	0.38	0.41	0.48	0.52	0.57	0.67	0.67	0.69	0.63	0.63	0.61	0.86	0.80	0.81	0.71	0.72	0.68
	200	0.33	0.31	0.29	0.28	0.27	0.26	0.62	0.59	0.54	0.45	0.36	0.31	0.75	0.72	0.64	0.53	0.40	0.33
	400	0.34	0.33	0.31	0.29	0.27	0.27	0.65	0.63	0.60	0.53	0.44	0.36	0.79	0.77	0.72	0.64	0.52	0.40
	20	0.86	0.90	0.94	0.97	0.99	1.00	0.93	0.95	0.96	0.98	0.99	0.99	0.97	0.97	0.98	0.99	0.99	0.99
	30	0.79	0.85	0.90	0.95	0.98	0.99	0.91	0.93	0.96	0.98	0.99	0.99	0.96	0.97	0.98	0.99	0.99	0.99
BR(5)	50	0.70	0.77	0.85	0.92	0.96	0.98	0.87	0.90	0.93	0.96	0.98	0.99	0.94	0.96	0.97	0.98	0.99	0.99
	100	0.51	0.56	0.63	0.73	0.77	0.81	0.73	0.75	0.78	0.80	0.83	0.83	0.86	0.84	0.85	0.84	0.86	0.85
	200	0.40	0.42	0.48	0.55	0.62	0.65	0.66	0.66	0.66	0.67	0.68	0.69	0.78	0.77	0.74	0.73	0.71	0.71
	400	0.37	0.38	0.40	0.47	0.55	0.61	0.67	0.66	0.65	0.66	0.67	0.68	0.80	0.78	0.76	0.74	0.72	0.71
$\mathbf{Note}$	s: The	entrie	s are a	verage	one-pe	riod ał	iead, o	ut-of-sa	mple N	ASE rat	tios rel	ative t	o princ	ipal co	mponeı	nts-bas	ed one-	period	ahead
forec	ists acr	oss 1,00	JO Mor	nte Car	lo repli	cations	. The	target	and in	dicator	variab]	les are	genera	thr thr	ough tl	he DG	Ps in (	11 who	ere we
consid	ler a fa	ctor loa	iding st	ructure	e as out	lined u	nder Ca	as I as	suming	$r = k_2$	factors	; we ge	nerate	T + 100	) observ	rations	on the	predict	or and
expla	natory	variable	ss, use t	the first	T  obset	ervatior	is to est	imate 1	the mod	dels and	use th	e result	ing par	ameter	estima	tes to f	orecast	the pre	edictor
varial	ole over	T + 1,	$\dots, T$	-100.	We imp	ose diff	erent le	vels of	for the	asympt	otic fit	of the	predict	ion reg	ression,	symbc	lized b	y the $R$	<sup>2</sup> 's. In
case (	of PC r	egressio	$n k_2$ fa	ctors a	re extra	acted, in	n case o	of PLS	regressi	on $k_1$ f	actors a	are exti	acted,	and in	case of	Bayesi	an regr	ession (	BR) a
shrinl	cage pa	rameter	$q \times N$	is used	ł with g	q = 1, 5													

Table 1: Standard Common Factor Case, Relative MSE compared to PC for  $(k_1, k_2) = (1, 1)$ 

		400	1.18	1.17	1.13	0.98	0.82	0.88	0.97	0.99	1.00	0.66	0.33	0.40	0.93	0.96	0.99	0.85	0.71	0.71
		200	1.17	1.19	1.15	0.91	0.86	0.92	0.97	1.00	1.00	0.69	0.41	0.52	0.92	0.96	0.98	0.85	0.72	0.73
(2,5)	=0.2	100	1.17	1.16	1.11	0.90	0.90	0.96	0.98	1.00	1.02	0.77	0.54	0.65	0.92	0.96	0.97	0.87	0.74	0.75
	$R^2=$	50	1.13	1.12	1.07	0.94	0.90	0.95	0.99	1.01	1.04	0.83	0.67	0.75	0.93	0.95	0.97	0.87	0.77	0.78
		30	1.11	1.08	1.03	0.95	0.91	0.94	0.99	1.02	1.04	0.82	0.75	0.80	0.91	0.94	0.96	0.86	0.79	0.81
		20	1.13	1.08	1.01	0.97	0.93	0.94	0.99	1.02	1.00	0.85	0.81	0.83	0.90	0.94	0.94	0.87	0.83	0.84
		400	1.15	1.12	1.10	0.84	0.84	0.87	0.97	0.98	0.99	0.67	0.31	0.36	0.98	0.98	0.99	0.86	0.69	0.68
o moder		200	1.14	1.11	1.08	0.84	0.88	0.92	0.97	0.98	0.98	0.65	0.37	0.45	0.97	0.98	0.98	0.84	0.69	0.68
	0.33	100	1.13	1.06	1.09	0.90	0.89	0.89	0.96	0.97	0.97	0.66	0.46	0.55	0.97	0.97	0.96	0.82	0.69	0.67
	$R^2 =$	50	1.09	1.06	1.02	0.94	0.87	0.89	0.95	0.96	0.94	0.71	0.56	0.62	0.95	0.96	0.94	0.82	0.68	0.68
		30	1.08	1.03	0.97	0.95	0.88	0.89	0.95	0.94	0.90	0.71	0.64	0.67	0.94	0.94	0.91	0.79	0.70	0.70
(Samo)		20	1.06	0.96	0.95	0.90	0.89	0.89	0.93	0.93	0.89	0.80	0.70	0.72	0.94	0.93	0.90	0.83	0.73	0.73
		400	1.06	1.03	1.07	0.91	0.86	0.89	0.97	0.97	0.96	0.59	0.27	0.27	1.07	1.03	1.00	0.83	0.66	0.62
		200	1.08	1.05	1.02	1.03	0.88	0.88	0.96	0.95	0.92	0.57	0.28	0.28	1.06	1.02	0.98	0.80	0.62	0.56
	0.5	100	1.00	0.98	1.04	0.86	0.86	0.92	0.93	0.91	0.83	0.49	0.29	0.30	1.03	1.00	0.94	0.75	0.57	0.48
	$R^2 =$	50	1.00	0.96	0.91	0.85	0.82	0.83	0.88	0.82	0.72	0.45	0.32	0.33	1.02	0.97	0.88	0.68	0.50	0.42
		30	0.93	0.89	0.89	0.89	0.81	0.78	0.81	0.74	0.62	0.46	0.35	0.36	1.00	0.93	0.81	0.63	0.47	0.41
		20	0.93	0.85	0.84	0.77	0.73	0.76	0.75	0.67	0.56	0.44	0.40	0.41	0.98	0.89	0.76	0.59	0.47	0.43
		T/N	20	30	50	100	200	400	20	30	50	100	200	400	20	30	50	100	200	400
					PLS						BR(1)						BR(5)			

Table 2: Standard Common Factors Case. Relative MSE compared to PC for  $(k_1, k_2) = (3, 6)$ 

Notes: See the notes for Table 1.

12001			100000												-		(7~, (T^	(+)	-
				$R^2$	=.5					$R^2 =$	.33					$R^{2}=$	=.2		
	T/N	20	30	50	100	200	400	20	30	50	100	200	400	20	30	50	100	200	400
	20	0.84	0.89	0.94	0.99	1.01	1.03	1.00	1.03	1.04	1.06	1.06	1.05	1.07	1.09	1.11	1.10	1.09	1.07
	30	0.80	0.86	0.91	0.98	1.01	1.03	0.97	1.00	1.03	1.06	1.07	1.06	1.04	1.07	1.09	1.11	1.11	1.09
PLS	50	0.75	0.83	0.89	0.95	1.00	1.03	0.92	0.96	1.00	1.04	1.07	1.07	1.01	1.03	1.07	1.08	1.11	1.11
	100	0.66	0.72	0.75	0.83	0.85	0.84	0.83	0.84	0.88	0.88	0.89	0.85	0.91	0.93	0.90	0.87	0.86	0.90
	200	0.64	0.69	0.75	0.78	0.80	0.80	0.80	0.82	0.83	0.83	0.83	0.80	0.87	0.87	0.87	0.85	0.82	0.77
	400	0.65	0.72	0.76	0.82	0.84	0.86	0.82	0.84	0.87	0.88	0.88	0.87	0.89	0.90	0.90	0.91	0.89	0.86
	20	0.71	0.78	0.86	0.92	0.96	0.98	0.94	0.95	0.97	0.98	0.99	0.99	1.02	1.02	1.02	1.01	1.00	1.00
	30	0.63	0.70	0.80	0.89	0.94	0.97	0.91	0.94	0.96	0.98	0.99	0.99	1.03	1.04	1.03	1.01	1.01	1.00
BR(1)	50	0.52	0.60	0.70	0.83	0.91	0.95	0.86	0.89	0.94	0.96	0.98	0.99	1.01	1.03	1.04	1.03	1.01	1.01
	100	0.36	0.37	0.42	0.52	0.56	0.58	0.70	0.66	0.70	0.64	0.64	0.62	0.83	0.85	0.76	0.73	0.65	0.68
	200	0.33	0.31	0.30	0.27	0.27	0.26	0.62	0.59	0.54	0.44	0.36	0.31	0.75	0.71	0.64	0.52	0.40	0.33
	400	0.34	0.33	0.31	0.29	0.27	0.27	0.65	0.63	0.60	0.53	0.44	0.36	0.79	0.77	0.73	0.64	0.51	0.40
	20	0.86	0.90	0.95	0.97	0.99	1.00	0.94	0.95	0.97	0.98	0.99	0.99	0.96	0.97	0.98	0.98	0.98	0.98
	30	0.80	0.85	0.91	0.95	0.98	0.99	0.91	0.94	0.96	0.98	0.99	0.99	0.95	0.97	0.98	0.98	0.99	0.99
BR(5)	50	0.70	0.77	0.85	0.92	0.96	0.98	0.87	0.90	0.93	0.96	0.98	0.99	0.94	0.96	0.97	0.98	0.99	0.99
	100	0.51	0.56	0.65	0.75	0.79	0.81	0.74	0.74	0.80	0.80	0.82	0.83	0.85	0.86	0.83	0.85	0.84	0.85
	200	0.40	0.43	0.48	0.55	0.62	0.65	0.66	0.66	0.66	0.67	0.68	0.69	0.78	0.76	0.75	0.73	0.71	0.71
	400	0.36	0.38	0.40	0.47	0.55	0.61	0.66	0.66	0.65	0.66	0.67	0.68	0.80	0.78	0.76	0.74	0.72	0.71
Note	s: The	entrie	s are s	werage	one-pe	riod ał	nead, o	ut-of-se	ample 1	MSE ra	tios rel	ative t	o princ	ipal co	mponei	nts-base	ed one-	period	ahead
foreca	sts acr	oss 1,00	00 Moi	nte Car	lo repli	ications	. The	target	and in	dicator	variab	les are	genera	ted thr	ough tl	he DGI	Ps in (	(11) wh	ere we
consid	ler a fé	actor los	ading s	tructur	e as ou	utlined	under (	Case II	assum	ing $N_1$	$= N^{0.7}$	<sup>5</sup> and	r = 1 f	actors;	we ger	ierate 7	r + 100	) observ	rations
on the	e predi	ctor and	d expla	natory	variabl	les, use	the fir:	st $T$ ob	servati	ons to e	stimat	e the n	10dels 5	nd use	the re	sulting	parame	eter est	imates
to for	ecast t.	he pred	ictor va	ariable	over $T$	$+1, \ldots$	., T + 1	00. We	e impos	ie differe	ent leve	els of fo	or the a	sympto	tic fit e	of the p	predicti	on regr	ession,
symbc	bized k	by the <i>I</i>	$\mathbb{R}^{2}$ 's. Ir	n case c	of both	PC reg	ression	and Pl	LS regr	ession 1	factor	is extr	acted, a	and in e	case of	Bayesia	an regr	ession (	BR) a
$_{ m shrink}$	tage pa	rametei	$r q \times N$	is used	d with g	q = 1, 5													

Table 3: Weak Factor Case II (Weak Pervasiveness  $\kappa_1 = 0.75$ ), Relative MSE compared to PC for  $(k_1, k_2) = (1, 1)$ 

5		XTMD AA				XTMD AA		7.1 . 91	1 5 	~~· · ( ^	0.1001		odinoo	5 5 7 7 7		Tarl To	1211	- ( + ) -	
				$R^2$	=.5					$R^2 =$	.33					$R^{2}=$	=.2		
	T/N	20	30	50	100	200	400	20	30	50	100	200	400	20	30	50	100	200	400
	20	0.81	0.87	0.91	0.96	0.98	0.99	1.00	1.02	1.05	1.05	1.04	1.03	1.09	1.12	1.12	1.09	1.06	1.04
	30	0.76	0.82	0.88	0.94	0.97	0.99	0.96	0.99	1.03	1.05	1.05	1.03	1.06	1.09	1.10	1.11	1.08	1.06
PLS	50	0.72	0.76	0.83	0.90	0.96	0.98	0.92	0.95	0.99	1.04	1.06	1.06	1.01	1.05	1.09	1.11	1.11	1.08
	100	0.61	0.63	0.64	0.67	0.68	0.65	0.79	0.83	0.82	0.77	0.71	0.67	0.90	0.90	0.88	0.85	0.82	0.73
	200	0.57	0.60	0.60	0.60	0.55	0.48	0.76	0.76	0.74	0.69	0.61	0.49	0.84	0.83	0.80	0.73	0.62	0.49
	400	0.58	0.60	0.63	0.65	0.66	0.61	0.79	0.78	0.78	0.77	0.73	0.66	0.87	0.86	0.85	0.82	0.76	0.66
	20	0.71	0.79	0.86	0.93	0.96	0.98	0.94	0.95	0.97	0.98	0.99	0.99	1.03	1.03	1.02	1.00	1.00	1.00
	30	0.63	0.71	0.80	0.89	0.94	0.97	0.91	0.93	0.96	0.98	0.99	0.99	1.03	1.04	1.02	1.01	1.00	1.00
BR(1)	50	0.52	0.59	0.70	0.82	0.91	0.95	0.87	0.90	0.93	0.97	0.98	0.99	1.01	1.04	1.04	1.02	1.01	1.01
	100	0.37	0.40	0.40	0.48	0.55	0.58	0.68	0.69	0.66	0.66	0.62	0.61	0.84	0.84	0.79	0.70	0.70	0.66
	200	0.33	0.31	0.30	0.28	0.27	0.26	0.62	0.59	0.54	0.45	0.36	0.31	0.75	0.71	0.64	0.52	0.41	0.33
	400	0.34	0.32	0.31	0.29	0.27	0.26	0.66	0.63	0.60	0.53	0.44	0.36	0.79	0.76	0.73	0.64	0.52	0.40
	20	0.86	0.91	0.94	0.97	0.99	1.00	0.94	0.95	0.97	0.98	0.99	0.99	0.96	0.97	0.98	0.98	0.99	0.99
	30	0.80	0.86	0.91	0.96	0.98	0.99	0.91	0.93	0.96	0.97	0.99	0.99	0.95	0.97	0.98	0.98	0.99	0.99
BR(5)	50	0.70	0.77	0.85	0.92	0.96	0.98	0.87	0.90	0.93	0.96	0.98	0.99	0.94	0.96	0.97	0.98	0.99	0.99
	100	0.51	0.59	0.64	0.73	0.79	0.82	0.75	0.76	0.78	0.82	0.82	0.83	0.85	0.86	0.84	0.83	0.85	0.85
	200	0.40	0.43	0.48	0.56	0.62	0.66	0.66	0.66	0.66	0.67	0.68	0.69	0.78	0.76	0.75	0.73	0.71	0.71
	400	0.37	0.37	0.40	0.47	0.55	0.61	0.67	0.65	0.66	0.66	0.67	0.68	0.80	0.78	0.77	0.74	0.72	0.71
$\mathbf{Note}$	s: The	entries	are ave:	rage on	e-period	d ahead	l, out-oi	f-sample	e MSE	ratios re	lative t	o princ	ipal con	aponen	ts-based	d one-p	eriod al	head fo	recasts
acros	; 1,000	Monte	Carlo	replicat	tions.	The tar	rget and	d indica	ator va	riables	are gen	erated	throug	h the I	DGPs in	n (11) -	where	we cons	sider a
factor	loadin	g struct	ture as	outline	ad unde	r Case	III assu	uming )	$\lambda_{i,j} =  ilde{\lambda}$	$_{i,j}/(N^{0})$	$^{25}$ ) and	l r = 1	factors	; we ge	nerate (	T + 100	) observ	vations	on the
predie	ctor and	l explaı	natory .	variable	es, use t	the first	T  obset	ervation	is to est	imate t	he mod	els and	use the	e result	ing para	ameter	estima	tes to fo	orecast
the p	redicto	r variab	le over	T + 1,	$\dots, T$	+ 100.	We im <sub>l</sub>	oose dif	ferent 1	evels of	for the	asym]	ptotic fi	t of th	e predic	ction re	gressio	n, symł	olized
by th	$R^{2}$ 's.	In cas	e of bo	th PC	regress	ion and	l PLS r	egressic	on 1 fac	ctor is e	extracte	d, and	in case	e of Ba	yesian 1	egressi	on (BR	t) a shr	inkage
paran	neter q	$\timesN$ is	used w	ith $q =$	: 1, 5.														

Table 4: Weak Factor Case III (Weak Loadings  $\kappa_2 = 0.25$ ), Relative MSE compared to PC for  $(k_1, k_2) = (1, 1)$ 

				$R^2$ :					4	$R^2_{=}$	: 33					$R^2$ =	=.2	-	
	T/N	20	30	50	100	200	400	20	30	50	100	200	400	20	30	50	100	200	400
	20	0.77	0.82	0.89	0.94	0.97	0.99	1.00	1.02	1.04	1.04	1.02	1.01	1.09	1.12	1.11	1.08	1.05	1.03
	30	0.71	0.76	0.83	0.91	0.96	0.97	0.95	0.98	1.01	1.03	1.03	1.02	1.08	1.11	1.11	1.09	1.07	1.04
PLS	50	0.62	0.68	0.76	0.85	0.92	0.96	0.90	0.93	0.98	1.02	1.03	1.02	1.02	1.06	1.09	1.11	1.09	1.06
	100	0.47	0.48	0.54	0.51	0.51	0.49	0.68	0.73	0.72	0.71	0.66	0.66	0.89	0.82	0.83	0.80	0.71	0.64
	200	0.41	0.40	0.41	0.37	0.33	0.25	0.67	0.64	0.61	0.53	0.42	0.30	0.78	0.75	0.71	0.60	0.46	0.32
	400	0.41	0.40	0.40	0.39	0.37	0.32	0.69	0.67	0.66	0.60	0.53	0.41	0.81	0.79	0.77	0.70	0.60	0.46
	20	0.71	0.78	0.86	0.92	0.96	0.98	0.94	0.96	0.97	0.98	0.99	1.00	1.03	1.03	1.02	1.01	1.00	1.00
	30	0.63	0.71	0.80	0.89	0.94	0.97	0.91	0.94	0.96	0.98	0.99	0.99	1.04	1.04	1.03	1.02	1.01	1.01
BR(1)	50	0.53	0.61	0.70	0.82	0.91	0.95	0.87	0.90	0.94	0.96	0.98	0.99	1.01	1.04	1.03	1.03	1.01	1.01
	100	0.37	0.39	0.42	0.45	0.52	0.56	0.66	0.68	0.64	0.65	0.64	0.65	0.86	0.80	0.79	0.73	0.69	0.65
	200	0.33	0.31	0.30	0.28	0.27	0.26	0.62	0.59	0.54	0.45	0.36	0.31	0.75	0.71	0.64	0.52	0.41	0.33
	400	0.34	0.33	0.31	0.29	0.27	0.27	0.65	0.63	0.60	0.53	0.44	0.36	0.79	0.77	0.73	0.64	0.52	0.40
	20	0.86	0.90	0.94	0.97	0.98	0.99	0.93	0.95	0.97	0.98	0.99	1.00	0.96	0.98	0.98	0.99	0.99	1.00
	30	0.80	0.86	0.91	0.95	0.97	0.99	0.91	0.93	0.96	0.97	0.99	0.99	0.96	0.97	0.98	0.99	0.99	1.00
BR(5)	50	0.70	0.78	0.85	0.92	0.96	0.98	0.87	0.90	0.93	0.96	0.98	0.99	0.94	0.96	0.97	0.98	0.99	0.99
	100	0.53	0.58	0.65	0.72	0.77	0.81	0.72	0.76	0.77	0.81	0.83	0.85	0.86	0.84	0.84	0.85	0.85	0.84
	200	0.40	0.43	0.49	0.56	0.62	0.66	0.66	0.66	0.67	0.67	0.69	0.69	0.78	0.76	0.75	0.73	0.71	0.71
	400	0.37	0.38	0.41	0.47	0.55	0.62	0.67	0.66	0.65	0.66	0.67	0.68	0.80	0.79	0.77	0.74	0.72	0.71
Note	s: See	the not	es for <b>J</b>	able 3,	but nc	w with	$N_1 = .$	$N^{0.25}$ .											

$\sim$	
-	I
<u> </u>	
(0)	1
14	-
4	
ŗ	
Ę	
РС	
Q	
ц Т	
re	
D2	)
no	
ں اتا	
ISI	
2	
ΪVθ	
ati	
6	
Щ.	-
25)	) 
Ū.	
 2	
$\mathbb{S} \mathcal{K}_1 \equiv$	
$\text{ness} \kappa_1 =$	222
veness $\kappa_1 =$	
siveness $\kappa_1 =$	· · · · · · · · · · · · · · · · · · ·
vasiveness $\kappa_1 =$	
Pervasiveness $\kappa_1 =$	
$\epsilon$ Pervasiveness $\kappa_1 =$	· · · · · · · · · · · · · · · · · · ·
eak Pervasiveness $\kappa_1 =$	
Weak Pervasiveness $\kappa_1 =$	
I (Weak Pervasiveness $\kappa_1 =$	
$=$ II (Weak Pervasiveness $\kappa_1 =$	
ase II (Weak Pervasiveness $\kappa_1 =$	
Case II (Weak Pervasiveness $\kappa_1 =$	
or Case II (Weak Pervasiveness $\kappa_1 =$	
actor Case II (Weak Pervasiveness $\kappa_1 =$	
Factor Case II (Weak Pervasiveness $\kappa_1 =$	
ak Factor Case II (Weak Pervasiveness $\kappa_1 =$	
Veak Factor Case II (Weak Pervasiveness $\kappa_1 =$	
Weak Factor Case II (Weak Pervasiveness $\kappa_1 =$	
5: Weak Factor Case II (Weak Pervasiveness $\kappa_1 =$	
ole 5: Weak Factor Case II (Weak Pervasiveness $\kappa_1 =$	
Pable 5: Weak Factor Case II (Weak Pervasiveness $\kappa_1 =$	
Table 5: Weak Factor Case II (Weak Pervasiveness $\kappa_1 =$	

		20 20 20 20 20 20 20 20 20 20 20 20 20 2	30 30 1. acto	$\frac{1}{50}$		200 200	400	20 20 100 1 00	30 30	$\frac{0}{R^2}$ , 100 $\frac{R^2}{50}$	100 100	200 200	400 400	20 113	30 30	$\frac{01}{R^2} = \frac{R^2}{80}$	, <sup>w2</sup> ) - =.2 100	200 1 05	400
	30	0.70	0.76 0.76	0.84 0.84	0.91	0.95	0.98 0.98	0.96 0	0.99	1.03	1.04	1.02	1.01 1.02	$1.12 \\ 1.08$	1.13	1.12 1.12	1.10	1.07	1.03
PLS	50 100	0.61	0.67	0.75	0.84	0.92	0.96	0.89	0.94	0.98	1.02	1.04	1.03	1.05	1.07	1.10	1.10	1.09 0.69	1.06 0.64
	200	0.38	0.37	0.37	0.35	0.31	0.24	0.66	0.63	0.59	0.52	0.42	0.30	0.77	0.74	0.69	0.59	0.46	0.32
	400	0.37	0.37	0.37	0.37	0.35	0.31	0.67	0.66	0.63	0.59	0.51	0.41	0.80	0.78	0.75	0.69	0.58	0.45
	20	0.72	0.79	0.87	0.93	0.96	0.98	0.94	0.96	0.98	0.99	0.99	1.00	1.04	1.03	1.02	1.01	1.00	1.00
	30	0.63	0.71	0.81	0.89	0.94	0.97	0.92	0.94	0.96	0.98	0.99	1.00	1.04	1.04	1.03	1.02	1.01	1.00
BR(1)	50	0.53	0.60	0.70	0.83	0.91	0.95	0.87	0.91	0.93	0.96	0.98	0.99	1.03	1.04	1.04	1.02	1.02	1.01
	100	0.38	0.39	0.44	0.48	0.52	0.57	0.67	0.68	0.68	0.65	0.65	0.65	0.79	0.79	0.75	0.72	0.68	0.65
	200	0.34	0.32	0.30	0.28	0.27	0.27	0.63	0.59	0.54	0.45	0.37	0.31	0.76	0.71	0.65	0.52	0.41	0.33
	400	0.34	0.33	0.31	0.29	0.27	0.27	0.66	0.63	0.60	0.53	0.44	0.36	0.79	0.76	0.73	0.64	0.52	0.40
	20	0.87	0.91	0.94	0.97	0.98	0.99	0.94	0.95	0.97	0.98	0.99	1.00	0.97	0.98	0.99	0.99	1.00	1.00
	30	0.80	0.86	0.91	0.95	0.97	0.99	0.91	0.94	0.96	0.98	0.99	0.99	0.96	0.97	0.98	0.99	0.99	1.00
BR(5)	50	0.71	0.78	0.85	0.92	0.96	0.98	0.87	0.90	0.93	0.96	0.98	0.99	0.95	0.96	0.97	0.98	0.99	1.00
	100	0.53	0.58	0.67	0.73	0.79	0.82	0.73	0.76	0.78	0.81	0.84	0.85	0.83	0.84	0.84	0.84	0.85	0.84
	200	0.41	0.44	0.49	0.56	0.62	0.66	0.67	0.67	0.67	0.68	0.69	0.69	0.78	0.76	0.75	0.73	0.72	0.71
	400	0.37	0.38	0.41	0.48	0.56	0.62	0.67	0.66	0.65	0.66	0.67	0.68	0.80	0.78	0.77	0.74	0.72	0.71
$\operatorname{Note}$	s: See	the not	es for <b>T</b>	able 4,	but nc	ow with	$\lambda_{i,j} =$	$ ilde{\lambda}_{i,j}/(N$	$r^{0.75}$ ).										

1
Ļ
$k_{i}$
دی
C
Gr
5
2
0
Ę.
ared
Õ
E
8
Ē
$\mathbf{SI}$
2
Ve
÷Ę
la
ž
်ဂ
0.75
= 0.75
2 = 0.75
$\kappa_2 = 0.75$
gs $\kappa_2 = 0.75$
ings $\kappa_2 = 0.75$
adings $\kappa_2 = 0.75$
oadings $\kappa_2 = 0.75$
: Loadings $\kappa_2 = 0.75$
ak Loadings $\kappa_2 = 0.75$
Veak Loadings $\kappa_2 = 0.75$
(Weak Loadings $\kappa_2 = 0.75$ )
II (Weak Loadings $\kappa_2 = 0.75$ )
III (Weak Loadings $\kappa_2 = 0.75$ )
se III (Weak Loadings $\kappa_2 = 0.75$ )
Case III (Weak Loadings $\kappa_2 = 0.75$ )
r Case III (Weak Loadings $\kappa_2 = 0.75$ )
tor Case III (Weak Loadings $\kappa_2 = 0.75$ )
actor Case III (Weak Loadings $\kappa_2 = 0.75$ )
Factor Case III (Weak Loadings $\kappa_2 = 0.75$ )
k Factor Case III (Weak Loadings $\kappa_2 = 0.75$ )
eak Factor Case III (Weak Loadings $\kappa_2 = 0.75$ )
Weak Factor Case III (Weak Loadings $\kappa_2 = 0.75$ )
: Weak Factor Case III (Weak Loadings $\kappa_2 = 0.75$ )
5 6: Weak Factor Case III (Weak Loadings $\kappa_2 = 0.75$ )
ble 6: Weak Factor Case III (Weak Loadings $\kappa_2 = 0.75$ )
Table 6: Weak Factor Case III (Weak Loadings $\kappa_2 = 0.75$ )

	Р	C Regressi	ion	$_{\rm PL}$	S Regressi	on	Bayesian	Regression
h	2 PC	4 PC	6 PC	$1 PL\overline{S}$	2 PLS	3 PLS	N	5N
			Janua	ary 1972 -	December	2006		
_	0.0001	0.0100		Benchma	irk: RW		0.0010	
1	0.9291	0.9138	0.9293	0.9114	0.9208	0.9333	0.9218	0.9017
3	0.9064	0.8951	0.8975	0.8678	0.9003	0.9213	0.9213	0.8885
12	0.8411	0.8308	0.8559	0.8101	0.8233	0.8446	0.8398	0.8185
24	0.9102	0.8659	0.8933	0.8441	0.8476	0.8636	0.8321	0.8291
				Benchme	ark: AR			
1	0.9804	0.9795	0.9847	0.9643	1.0044	1.0565	0.9862	0.9640
3	0.9582	0.9692	0.9754	0.9229	0.9769	1.0059	0.9636	0.9330
12	0.8541	0.8201	0.8360	0.8032	0.8031	0.8251	0.8461	0.8244
24	0.8223	0.7694	0.7942	0.7709	0.7701	0.7884	0.8642	0.8457
			-	1008	- ·			
			Janua	ary 1985 -	December	2006		
				Domohour	mh. DW			
1	0.0504	0.0205	0.0491	0.0510	0 0279	0.0509	0.0496	0 0201
1	0.9394	0.9393	0.9451	0.9510	0.9572	0.9008	0.9480 1.0272	1 0009
ა 10	1.0149	1.0000	0.9972	1.0159	1.0230 1.0720	1.0203	1.0373	1.0008
12	1.0180	1.1100	1.1323	1.0108	1.0729	1.1001	1.1005	1.0144
24	1.0205	1.0884	1.1555	1.0127 David	1.0340	1.1322	1.0742	0.9287
1	0.0777	0.0507	0.0491	Denchmo	ark: AR	1 0940	0.0000	0.0794
1	0.9777	0.9587	0.9481	0.9707	0.9487	1.0340	0.9988	0.9724
3 10	0.9733	0.9602	0.9580	0.9696	0.9853	1.0681	1.0231	0.9852
12	0.9165	0.9798	0.9918	0.9024	0.9433	0.9859	1.0053	0.9289
24	0.9737	1.0086	1.1228	0.9446	1.0047	1.0787	1.0158	0.8738
			Ionus	mr 1079	Docombor	1094		
			Janua	ary 1972 -	December	1964		
				Benchma	ırk: RW			
1	0.8985	0.8881	0.9156	0.8710	0.9045	0.9160	0.8948	0.8742
3	0.8244	0.8082	0.8229	0.7703	0.8067	0.8432	0.8337	0.8033
12	0.7679	0.7098	0.7417	0.7242	0.7221	0.7413	0.7406	0.7418
24	0.8657	0.7984	0.8039	0.7791	0.7615	0.7570	0.7299	0.7726
				Benchmed	ark: AR			
1	0.9830	0.9995	1.0193	0.9579	1.0559	1.0780	0.9738	0.9558
3	0.9461	0.9757	0.9883	0.8848	0.9705	0.9552	0.9157	0.8910
12	0.8083	0.7374	0.7589	0.7504	0.7336	0.7472	0.7976	0.8104
24	0.7826	0.7134	0.7148	0.7196	0.6922	0.6959	0.7976	0.8104

Table 7: Forecast evaluation for CPI inflation

**Notes:** The table reports the ratio of the RMSE of either a version of (14) vis-àvis autoregressive model (12) or a version of (15) vis-à-vis the random walk model (13) for CPI inflation (see Section 4.2)) at each horizon h (in months). Versions of (14) and (15) depend on the usage of principal components (PC), partial least squares (PLS) or Bayesian regression (BR) to compress the information in the panel of predictor variables; see Section 4.1. In each case we use several sub-variants, depending either on the number of principal components, PLS factors or shrinkage parameters (BRR), where the shrinkage parameters is assumed to be proportional to the number of predictors in the panel (N = 104). The best performing method relative to the benchmarks are highlighted in **bold**.

	PC	C Regressio	on	PI	S Regress	ion	Bayesian	Regression
h	2 PC	4 PC	6 PC	1 PLS	2 PLS	3 PLS	N	5N
			-					
			Janu	ary 1972 -	December	2006		
				Benchm	ark: RW			
1	0.8600	0.8557	0.8547	0.8437	0.8589	0.8615	0.8553	0.8292
3	0.8216	0.8287	0.8412	0.7784	0.7668	0.7836	0.7832	0.7665
12	1.0414	1.0621	1.0657	0.7912	0.8558	0.8630	0.8541	0.8186
24	1.0734	1.0875	1.0929	0.9037	0.8575	0.8963	0.8749	0.8534
				Benchm	nark: AR			
1	0.9388	0.9291	0.9303	0.9239	0.9246	0.9664	0.9525	0.9217
3	0.9289	0.9310	0.9411	0.8870	0.8752	0.9036	0.9279	0.9076
12	1.1922	1.2204	1.1878	0.8768	0.9598	0.9514	0.9786	0.9265
24	1.1505	1.1803	1.1840	1.0030	0.9657	0.9994	0.9956	0.9607
			т	1005		2000		
			Janu	ary 1985 -	December	2006		
				Benchm	ark: RW			
1	0.9758	0.9822	0.9751	0.9610	0.9876	0.9810	0.9854	0.9423
3	0.8486	0.8562	0.8628	0.8196	0.8737	0.9021	0.9260	0.8384
12	0.7791	0.7878	0.7980	0.7422	0.8619	0.9127	0.9125	0.8008
24	0.7044	0.7216	0.7312	0.7699	0.7432	0.8405	0.8524	0.7432
				Benchm	nark: AR			
1	0.9685	0.9874	0.9831	0.9528	0.9650	1.0141	1.0120	0.9576
3	0.9784	0.9874	0.9880	0.9443	1.0287	1.0490	1.1338	1.0013
12	1.0794	1.0815	1.0794	0.9759	1.2166	1.2446	1.3238	1.1329
24	1.0725	1.0761	1.0867	1.1274	1.1642	1.2677	1.3147	1.1475
			_					
			Janu	ary 1972 -	December	: 1984		
				Benchm	ark: RW			
1	0.7950	0.7839	0.7867	0.7776	0.7858	0.7941	0.7812	0.7655
3	0.8086	0.8161	0.8307	0.7605	0.7181	0.7309	0.7181	0.7354
12	1.1660	1.1946	1.1940	0.8043	0.8338	0.8139	0.7989	0.8166
24	1.4260	1.4413	1.4439	1.0704	0.9961	0.9845	0.9297	0.9994
				Benchm	nark: AR			
1	0.9186	0.8884	0.8937	0.9043	0.8969	0.9336	0.9110	0.8972
3	0.9066	0.9070	0.9206	0.8634	0.8081	0.8420	0.8373	0.8693
12	1.2216	1.2589	1.2169	0.8299	0.8448	0.8187	0.8189	0.8378
24	1.1835	1.2218	1.2184	0.9508	0.8605	0.8604	0.8269	0.8734

Table 8: Forecast evaluation for industrial production

**Notes:** See the notes for Table 7.

	PC	C Regressi	ion	$_{\rm PL}$	S Regressi	on	Bayesian	Regression
h	2 PC	4 PC	6 PC	1 PLS	2 PLS	3 PLS	N	5N
			Janu	ary 1972 -	December	2006		
				Benchm	$ark \cdot BW$			
1	0.8927	0.8977	0.9073	0.8856	0.8879	0.8923	0.8866	0.8693
3	0.8095	0.8119	0.8227	0.7964	0.7705	0.8018	0.7870	0.7758
12	1.0483	1.0516	1.0424	0.8338	0.8434	0.8759	0.8872	0.8509
24	1 1398	1 1686	1 1453	0.9195	0.8471	0.8950	0.8871	0.8896
- 1	1.1000	1.1000	1.1100	Benchm	ark: AR	0.0000	0.0011	0.0000
1	0.9057	0.9169	0.9228	0.9061	0.9009	0.9182	0.9507	0.9326
3	0.9031	0.9065	0.9352	0.8889	0.8616	0.8935	0.9403	0.9100
12	1.0542	1.0494	1.0438	0.8606	0.8645	0.8873	0.9182	0.8791
24	1 0761	1 1071	1 0861	0.8926	0.8221	0.8618	0.8543	0.8624
	110101	111011	1.0001	0.0010	0.0221	0.0010	0.0010	0.0021
			Janu	arv 1985 -	December	2006		
				Benchm	ark: RW			
1	0.9850	0.9848	0.9867	0.9761	0.9781	0.9875	1.0125	0.9597
3	0.8959	0.8918	0.8897	0.8741	0.8926	0.9357	0.9464	0.8666
12	0.9602	0.9393	0.9332	0.8257	0.9385	0.9403	0.9749	0.8685
24	1.0090	1.0537	1.0204	1.0248	0.9305	1.0132	1.0247	0.9484
				Benchm	ark: AR			
1	0.9584	0.9596	0.9592	0.9545	0.9534	0.9721	1.0410	0.9929
3	0.9484	0.9477	0.9415	0.9095	0.9483	0.9839	1.0875	0.9945
12	1.0078	0.9825	0.9543	0.8646	1.0096	0.9726	1.0629	0.9537
24	0.9999	1.0305	0.9874	1.0153	0.9224	0.9844	1.0044	0.9360
			Janu	ary 1972 -	December	1984		
				Benchm	ark: RW			
1	0.8294	0.8384	0.8536	0.8237	0.8261	0.8269	0.7972	0.8073
3	0.7757	0.7806	0.7966	0.7665	0.7211	0.7480	0.7212	0.7404
12	1.0740	1.0850	1.0765	0.8353	0.8030	0.8486	0.8530	0.8426
24	1.2186	1.2438	1.2390	0.8714	0.7931	0.8218	0.8008	0.8541
				Benchm	ark: AR			
1	0.8667	0.8856	0.8963	0.8705	0.8621	0.8782	0.8816	0.8877
3	0.8833	0.8884	0.9321	0.8801	0.8222	0.8527	0.8709	0.8718
12	1.0667	1.0686	1.0701	0.8586	0.8086	0.8550	0.8662	0.8532
24	1.1536	1.1878	1.1822	0.8395	0.7597	0.7914	0.7716	0.8269

Table 9: Forecast evaluation for unemployment

**Notes:** See the notes for Table 7.

	PC Regression			PLS Regression			Bayesian Regression					
h	2 PC	4 PC	6 PC	1 PLS	2 PLS	3 PLS	N	5N				
January 1972 - December 2006												
Donahar and DW												
1	0.0175	0.0107	0.0007	Benchmo	ark: KW	OOFFF	0.0501	0 991 4				
1	0.9175	0.9107	0.8897	0.8541	0.8374	0.8555	0.8581	0.8314				
び 10	0.9279	0.9560	0.9747	0.8920	0.9048	0.9196	0.9194	0.8873				
12	0.9490	0.9281	0.9345	0.8939	0.9424	0.9757	1.0182	0.9398				
24	1.0426	0.9947	1.0146	0.9262 Benchm	1.0004 ark: A R	1.0154	1.0251	0.9656				
1	1 0387	0.0664	0.0325	0.0713	0 8/00	0.8503	0.0088	0.8028				
1	1.0307	1.0525	0.9525 0.0705	0.9713 0.0717	0.0400	0.0093	0.9088	0.8928				
0 19	1.0200	1.0525 0.8474	0.9795	0.9111	0.9211	0.9214	0.9100	0.8985				
24	0.0002	0.0474	0.0344	0.8130	0.0054	0.0344 0.0665	0.9444	0.0794				
24	0.9159	0.9200	0.9420	0.0124	0.3010	0.3005	0.9709	0.9288				
			Janu	ary 1985 -	December	2006						
				Benchmed	ark: RW							
1	0.8528	1.2544	1.2335	0.8909	1.1798	1.3448	1.2528	1.0070				
3	0.7880	1.0271	1.0339	0.7546	1.1476	1.2513	1.2157	0.9454				
12	0.8613	0.9935	0.9964	0.8102	1.0153	1.0762	1.0696	0.9481				
24	0.9447	0.9363	0.9516	0.8836	0.9970	1.0492	1.0174	0.9477				
				Benchm	ark: AR							
1	1.0731	1.3156	1.2996	1.0316	1.3079	1.4715	1.3617	1.0669				
3	0.9620	1.0767	1.0707	0.9482	1.1327	1.3825	1.1911	0.9495				
12	0.8188	0.9445	0.9398	0.7939	0.9752	1.0251	1.0518	0.9308				
24	0.8839	0.8847	0.8820	0.8404	0.9461	0.9928	0.9700	0.9000				
			Ianu	ome 1079	December	1094						
			Janu	ary 1972 -	December	1984						
				Benchme	ark: RW							
1	0.9223	0.8781	0.8569	0.8511	0.8044	0.8046	0.8191	0.8160				
3	0.9394	0.9492	0.9692	0.9040	0.8773	0.8819	0.8861	0.8819				
12	0.9590	0.8920	0.8985	0.9000	0.9031	0.9293	0.9860	0.9226				
24	1.0409	0.9758	0.9800	0.8812	0.9310	0.9179	0.9395	0.9111				
				Benchm	ark: AR							
1	1.0364	0.9397	0.9039	0.9673	0.8009	0.8045	0.8719	0.8805				
3	1.0296	1.0504	0.9710	0.9726	0.9002	0.8704	0.8897	0.8932				
12	0.8548	0.8044	0.7886	0.8003	0.8234	0.8411	0.8969	0.8511				
24	0.9740	0.9104	0.9153	0.8326	0.8910	0.8789	0.9089	0.8912				

Table 10: Forecast evaluation for the federal funds rate

**Notes:** See the notes for Table 7.

### References

- Bai, J., 2003, Inferential Theory for Factor Models of Large Dimensions, *Econometrica* 71, 135– 173.
- Bates, J. M. and C. W. J. Granger, 1969, The Combination of Forecasts, Operations Research Quarterly 20, 451–468.
- Boivin, J. and S. Ng, 2006, Are More Data Always Better for Factor Analysis?, Journal of Econometrics 132, 169–194.
- Capistrán, C. and A. Timmermann, 2008, Forecast Combination With Entry and Exit of Experts, *Journal of Business & Economic Statistics*. forthcoming.
- Chamberlain, G. and M. Rothschild, 1983, Arbitrage, Factor Structure, and Mean-Variance Analysis in Large Asset Markets, *Econometrica* **51**, 1305–1324.
- D'Agostino, A. and D. Giannone, 2006, Comparing Alternative Predictors Based on Large-Panel Factor Models, *Working Paper 680*, European Central Bank.
- D'Agostino, A., D. Giannone and P. Surico, 2006, (Un)Predictability and Macroeconomic Stability, *Working Paper 605*, European Central Bank.
- De Mol, C., D. Giannone and L. Reichlin, 2008, Forecasting Using a Large Number of Predictors: Is Bayesian Regression a Valid Alternative to Principal Components?, *Journal of Econometrics* 146, 318–328.
- Elliott, G. and A. Timmermann, 2004, Optimal Forecast Combinations Under General Loss Functions and Forecast Error Distributions, *Journal of Econometrics* **122**, 47–79.
- Faust, J. and J. H. Wright, 2007, Comparing Greenbook and Reduced Form Forecasts using a Large Realtime Dataset, mimeo, Federal Reserve Board and Johns Hopkins University.
- Forni, M., M. Hallin, M. Lippi and L. Reichlin, 2000, The Generalized Dynamic Factor Model: Identification and Estimation, *Review of Economics and Statistics* 82, 540–554.

- Forni, M., M. Hallin, M. Lippi and L. Reichlin, 2004, The Generalised Factor Model: Consistency and Rates, *Journal of Econometrics* 119, 231–255.
- Forni, Mario, Marc Hallin, Marco Lippi and Lucrezia Reichlin, 2005, The Generalized Dynamic Factor Model: One-Sided Estimation and Forecasting, *Journal of the American Statistical* Association 100(471), 830–840.
- Garthwaite, P. H., 1994, An Interpretation of Partial Least Squares, Journal of the American Statistical Association 89, 122–127.
- Granger, C. W. J. and R. Ramanathan, 1984, Improved Methods of Combining Forecasts, Journal of Forecasting 3, 197–204.
- Helland, I. S., 1988, On the Structure of Partial Least Squares Regression, Communications in Statistics - Simulation and Computation 17, 581–607.
- Helland, I. S., 1990, Partial Least Squares Regression and Statistical Models, Scandinavian Journal of Statistics 17, 97–114.
- Hendry, D. F., 1995, Dynamic Econometrics, Oxford University Press.
- Krolzig, H.-M. and D. F. Hendry, 2001, Computer Automation of General-to-Specific Model Selection Procedures, Journal of Economic Dynamics and Control 25, 831–866.
- McConnell, M. and G. Perez-Quiros, 2000, Output Fluctuations in the United States: What has Changed Since the Early 1980's?, *American Economic Review* **90**, 1464–1476.
- Schwarz, G., 1978, Estimating the Dimension of a Model, Annals of Statistics 6, 461–464.
- Sensier, M. and D. J. C. van Dijk, 2004, Testing for Volatility Changes in U.S. Macroeconomic Time Series, *Review of Economics and Statistics* 86, 833–839.
- Stock, J. H. and M. W. Watson, 2002a, Forecasting Using Principal Components from a Large Number of Predictors, Journal of the American Statistical Association 97, 1167–1179.
- Stock, J. H. and M. W. Watson, 2002b, Macroeconomic Forecasting Using Diffusion Indexes, Journal of Business & Economic Statistics 20, 147–162.

- Stock, J. H. and M. W. Watson, 2007, Forecasting in Dynamic Factor Models Subject to Structural Instability, *mimeo*, Harvard University and Princeton University.
- Stoica, P. and T. Söderström, 1998, Partial Least Squares: A First-Order Analysis, Scandinavian Journal of Statistics 25, 17–24.
- Svensson, Lars E O, 2005, Monetary Policy with Judgment: Forecast Targeting, International Journal of Central Banking 1(1), 1–54.
- Timmermann, A., 2006, Forecast Combinations, in G. Elliott, C. W. J. Granger and A. Timmermann (editors), Handbook of Economic Forecasting, North Holland, Amsterdam, pp. 135–196.
- Wold, H., 1982, Soft Modeling. The Basic Design and Some Extensions, in K.-G. Jöreskog andH. Wold (editors), Systems Under Indirect Observation, Vol. 2, North-Holland, Amsterdam.

# Appendices

## A Proof of Theorem 1

We first note the following obvious fact: If

$$||\hat{\alpha}_{PLS}^n - \hat{\alpha}_{PC}^n|| = o_p(a_{N,T}), \tag{A.1}$$

for some sequence  $a_{N,T}$ , then

$$||\hat{\alpha}_{PLS} - \hat{\alpha}_{PC}|| = o_p(a_{N,T}/b_{N,T}), \tag{A.2}$$

where  $\hat{\alpha}_{PLS}^n$  and  $\hat{\alpha}_{PC}^n$  are the PLS and PC regression estimates (5) and (6) obtained when

$$X^n = \frac{1}{b_{N,T}}X$$

are used as regressors. Given this fact, we focus on  $\hat{\alpha}_{PLS}^n$  and  $\hat{\alpha}_{PC}^n$  where we set  $b_{N,T} = (NT)^{1/2}$ . First, we have that

$$X^{n'}X^n = \frac{1}{N}\Sigma + \frac{1}{N}\left(\frac{X^{n'}X^n}{T} - \Sigma\right)$$

But, by the factor assumption,

$$\frac{1}{N}\Sigma = \frac{1}{N}\tilde{S}\tilde{\Psi}\tilde{S}' + o(1)$$

where

$$\tilde{\psi}_{Ni} = O(N), \quad i = 1, \dots, r.$$

The proof is complete if we show that

$$\left\|\frac{1}{N}\left(\frac{X^{n'}X^n}{T} - \Sigma\right)\right\| = o_p(1),$$

since, then, by Assumption 1 and (7), it follows that

$$||\hat{\alpha}_{PLS}^n - \hat{\alpha}_{PC}^n|| = o_p(1),$$

and then, by (A.1) and (A.2), (8) follows. But,

$$\left\| \frac{1}{N} \left( \frac{X^{n'} X^n}{T} - \Sigma \right) \right\| = \left( \frac{1}{N^2} \sum_{i=1}^N \sum_{j=1}^N \left( \frac{1}{T} \sum_{t=1}^T (x_{i,t} x_{j,t} - \sigma_{i,j}) \right)^2 \right)^{1/2} = O_p(T^{-1/2}),$$

where the second equality follows by Assumption 2, proving the result.

## **B** Proof of Theorem 2

For the proof, we first focus on population quantities and deal with the estimation problem as a last step. Although not necessary, we assume without loss of generality that the variance of  $y_t$ is finite. This implies that  $\alpha'_N \Sigma_{XX} \alpha_N < \infty$ . Then, Assumption 4 implies that

$$\alpha_{N,i} = \frac{\tilde{\alpha}_{N,i}}{\sqrt{N}}$$

for some  $\tilde{\alpha}_{N,i}$ , such that

$$0 < \sup_{i} |\tilde{\alpha}_{N,i}| < \infty$$

Of course,  $\alpha_{N,i}$  can be equivalently written as

$$\alpha_{N,i} = \Sigma_{XX}^{-1} \sigma_{Xy}.$$

Using the above we can write

$$\alpha'_N \Sigma_{XX} \alpha_N = \sigma'_{Xy} \Sigma_{XX}^{-1} \sigma_{Xy}$$

and therefore there exist  $\tilde{\sigma}_{Xy,i}$  such that

$$\sigma_{Xy,i} = \frac{\tilde{\sigma}_{Xy,i}}{\sqrt{N}}$$

where

$$0 < \sup_{i} |\tilde{\sigma}_{Xy,i}| < \infty. \tag{B.3}$$

For PLS where for one factor we have that

$$p \lim_{N \to \infty} \hat{\alpha}_N = \sigma_{Xy}$$

it follows that

$$p \lim_{N \to \infty} \frac{1}{T} \hat{\alpha}'_N X' X \hat{\alpha} \hat{\beta}_N = \sigma'_{Xy} \Sigma_{XX} \sigma_{Xy}$$

The result we wish to prove follows if we show that

$$\sigma'_{Xy} \Sigma_{XX} \sigma_{Xy} - \sigma'_{Xy} \Sigma_{XX}^{-1} \sigma_{Xy} = o(1)$$

For a square matrix  $A = [a_{ij}]$ , define the following matrix norm:

$$||A||_{\tilde{\sigma}} = \sum_{i=1}^{N} \sum_{j=1}^{N} |\tilde{\sigma}_{Xy,1,i}| |a_{ij}| |\tilde{\sigma}_{Xy,1,j}|$$

This is easily seen to satisfy the standard conditions for it to be a valid matrix norm. Then, we wish to prove that  $\|\Sigma_{XX} - \Sigma_{XX}^{-1}\|_{\tilde{\sigma}} = o(N)$ . But we immediately have that

$$\left\|\Sigma_{XX} - \Sigma_{XX}^{-1}\right\|_{\sigma} \le \left\|\Sigma_{XX} - I\right\|_{\tilde{\sigma}} + \left\|I - \Sigma_{XX}^{-1}\right\|_{\tilde{\sigma}} \tag{B.4}$$

Using the Sherman Morrison formula, and for any  $N \times N$  nonnsingular matrix A, any  $N \times N$ matrix M such that  $(A + M)^{-1}$  exists, any matrix norm and some constant c > 0, it can be easily proven that

$$\left\| A^{-1} - (A+M)^{-1} \right\| \le c \left\| M \right\|$$

Thus, it follows that as long as  $\|\Sigma_{XX} - I\|_{\tilde{\sigma}} = o(N)$ , (B.4) holds. But,  $\|\Sigma_{XX} - I\|_{\tilde{\sigma}} = o(N)$  is equivalent to  $\|\Sigma_{XX} - I\|_1 = o(N)$ , for the more familiar Minkowski 1-norm  $\|.\|_1$  which holds by Assumption 4. Finally, we need to prove that

$$\left\|\frac{1}{T}\sum_{t=1}^{T}x_{j,t}y_t - \sigma_{Xy,i}\right\| = o_p(1)$$
(B.5)

But, by assumption 5

$$\frac{1}{T}\sum_{t=1}^{T} x_{j,t} y_t - \sigma_{Xy,i} = O_p\left(T^{-1/2}\right), \text{ uniformly over } i. \tag{B.6}$$

Hence, (B.5) follows if  $N/T \rightarrow 0$ .

### C Data Set

The data set used for forecasting are the monthly series from the panel of U.S. predictor series as employed in Stock and Watson (2007), but excluding our four forecast variables: CPI inflation, (aggregate) industrial production, (aggregate) unemployment rate and the (effective) federal funds rate. In order to have I(0) predictor variables, the underlying raw series need to be appropriately transformed; generally we employ the same transformation as Stock and Watson (2007), except for the nominal series where we follow, e.g., D'Agostino and Giannone (2006) and

Transformation code	Transformation $X_t$ of raw series $Y_t$
$     \begin{array}{c}       1 \\       2 \\       3 \\       4 \\       5 \\       6     \end{array} $	$\begin{aligned} X_t &= Y_t \\ X_t &= \Delta Y_{t,t-1} \\ X_t &= \Delta Y_{t,t-12} - \Delta Y_{t-1,t-13} \\ X_t &= \ln Y_t \\ X_t &= \Delta \ln Y_{t,t-1} \\ X_t &= \Delta \ln Y_{t,t-12} - \Delta \ln Y_{t-1,t-13} \end{aligned}$

Table C.1: Transformation of the predictor variables

use first differences of twelve-month transformations of the raw series. Table C.1 summarizes our potential transformations for the raw series.

Hence, we are using as predictor variables the following 105 series, which span before transformation the sample January 1959 - December 2006 and we refer to Stock and Watson (2007) for more details regarding data construction and sources:

Series $Y_t$	$\frac{\text{Transformation:}}{(See Table C.1)}$
INDUSTRIAL PRODUCTION INDEX - FINAL PRODUCTS	5
INDUSTRIAL PRODUCTION INDEX - CONSUMER GOODS	5
INDUSTRIAL PRODUCTION INDEX - DURABLE CONSUMER GOODS	5
INDUSTRIAL PRODUCTION INDEX - NONDURABLE CONSUMER GOODS	5
INDUSTRIAL PRODUCTION INDEX - BUSINESS EQUIPMENT	5
INDUSTRIAL PRODUCTION INDEX - MATERIALS	5
INDUSTRIAL PRODUCTION INDEX - DURABLE GOODS MATERIALS	5
INDUSTRIAL PRODUCTION INDEX - NONDURABLE GOODS MATERIALS	5
INDUSTRIAL PRODUCTION INDEX - MANUFACTURING (SIC)	5
INDUSTRIAL PRODUCTION INDEX - RESIDENTIAL UTILITIÉS	5
INDUSTRIAL PRODUCTION INDEX - FUELS	5
NAPM PRODUCTION INDEX (PERCENT)	1
CAPACITY UTILIZATION - MÀNUFACTÚRING (SIC)	1
AVG HRLY EARNINGS, PROD WRKRS, NONFARM - GOODS-PRODUCING	6
AVG HRLY EARNINGS, PROD WRKRS, NONFARM - CONSTRUCTION	6
AVG HRLY EARNINGS, PROD WRKRS, NONFARM - MFG	6
REAL AVG HRLY EARNINGS, PROD WRKRS, NONFARM - GOODS-PRODUCING	5
REAL AVG HRLY EARNINGS, PROD WRKRS, NONFARM - CONSTRUCTION	5
REAL AVG HRLY EARNINGS, PROD WRKRS, NONFARM - MFG	5
EMPLOYEES, NONFARM - TOTAL PRIVATE	5
EMPLOYEES, NONFARM - GOODS-PRODUCING	5
EMPLOYEES, NONFARM - MINING EMPLOYEES, NONFARM - CONSTRUCTION	5
EMPLOYEES, NONFARM - MFG	5
EMPLOYEES, NONFARM - DURABLE GOODS	5
EMPLOYEES, NONFARM - NONDURABLE GOODS	5
EMPLOYEES, NONFARM - SERVICE-PROVIDING	5
EMPLOYEES, NONFARM - TRADE, TRANSPORT, UTILITIES	5
EMPLOYEES, NONFARM - WHOLESALE TRADE	5
EMPLOYEES, NONFARM - RETAIL TRADE	5
EMPLOYEES, NONFARM - FINANCIAL ACTIVITIES	5
EMPLOYEES, NONFARM - GOVERNMENT	5

INDEX OF HELP-WANTED ADVERTISING IN NEWSPAPERS (1967=100;SA) EMPLOYMENT: RATIO; HELP-WANTED ADS:NO. UNEMPLOYED CLF CIVILIAN LABOR FORCE: EMPLOYED, TOTAL (THOUS.,SA) CIVILIAN LABOR FORCE: EMPLOYED, NONAGRIC.INDUSTRIES (THOUS.,SA) UNEMPLOY.BY DURATION: PERSONS UNEMPL.LESS THAN 5 WKS (THOUS.,SA) UNEMPLOY.BY DURATION: PERSONS UNEMPL.5 TO 14 WKS (THOUS., SA) UNEMPLOY.BY DURATION: PERSONS UNEMPL.15 WKS + (THOUS.,SA) UNEMPLOY.BY DURATION: PERSONS UNEMPL.15 TO 26 WKS (THOUS.,SA) UNEMPLOY.BY DURATION: PERSONS UNEMPL.27 WKS + (THOUS,SA) AVG WKLY HOURS, PROD WRKRS, NONFARM - GOODS-PRODUCING AVG WKLY OVERTIME HOURS, PROD WRKRS, NONFARM - MFG HOUSING AUTHORIZED: TOTAL NEW PRIV HOUSING UNITS (THOUS., SAAR) HOUSING STARTS:NONFARM(1947-58);TOTAL FARM&NONFARM(1959-)(THOUS.,U)SA HOUSING STARTS:NORTHEAST (THOUS.U.)S.A. HOUSING STARTS:MIDWEST(THOUS.U.)S.A. HOUSING STARTS:SOUTH (THOUS.U.)S.A. HOUSING STARTS:WEST (THOUS.U.)S.A. INTEREST RATE: U.S.TREASURY BILLS, SEC MKT, 3-MO. (% PER ANN, NSA) INTEREST RATE: U.S. TREASURY BILLS, SEC MKT, 6-MO. (% PER ANN, NSA) INTEREST RATE: U.S.TREASURY CONST MATURITIES,1-YR.(% PER ANN,NSA) INTEREST RATE: U.S.TREASURY CONST MATURITIES,5-YR.(% PER ANN,NSA) INTEREST RATE: U.S.TREASURY CONST MATURITIES,10-YR.(% PER ANN,NSA) BOND YIELD: MOODY'S AAA CORPORATE (% PER ANNUM) BOND YIELD: MOODY'S BAA CORPORATE (% PER ANNUM) INTEREST RATE SPREAD: 6-MO. TREASURY BILLS MINUS 3-MO. TREASURY BILLS INTEREST RATE SPREAD: 1-YR. TREASURY BONDS MINUS 3-MO. TREASURY BILLS INTEREST RATE SPREAD: 10-YR. TREASURY BONDS MINUS 3-MO. TREASURY BILLS INTEREST RATE SPREAD: AAA CORPORATE MINUS 10-YR. TREASURY BONDS INTEREST RATE SPREAD: BAA CORPORATE MINUS 10-YR. TREASURY BONDS MONEY STOCK: M1(CURR,TRAV.CKS,DEM DEP,OTHER CK'ABLE DEP)(BIL\$,SA) MZM (SA) FRB St. Louis MONEY STOCK:M2(M1+O'NITE RPS,EURO\$,G/P&B/D MMMFS&SAV&SM TIME DEP)(BIL\$,SA) MONETARY BASE, ADJ FOR RESERVE REQUIREMENT CHANGES(MIL\$,SA) DEPOSITORY INST RESERVES: TOTAL, ADJ FOR RESERVE REQ CHGS(MIL\$, SA) DEPOSITORY INST RESERVES:NONBORROWED, ADJ RES REQ CHGS(MIL\$, SA) Commercial and Industrial Loans at All Commercial Banks (FRED) Billions \$ (SA) CONSUMER CREDIT OUTSTANDING - NONREVOLVING(G19) Personal Consumption Expenditures, Price Index (2000=100), SAAR Personal Consumption Expenditures - Durable Goods, Price Index (2000=100), SAAR Personal Consumption Expenditures - Nondurable Goods, Price Index (2000=100), SAAR Personal Consumption Expenditures - Services, Price Index (2000=100), SAAR PCE Price Index Less Food and Energy (SA) Fred PRODUCER PRICE INDEX: FINISHED GOODS (82=100,SA) PRODUCER PRICE INDEX: FINISHED CONSUMER GOODS (82=100,SA) PRODUCER PRICE INDEX:INTERMED MAT.SUPPLIES & COMPONENTS(82=100,SA) PRODUCER PRICE INDEX:CRUDE MATERIALS (82=100,SA) Real PRODUCER PRICE INDEX:CRUDE MATERIALS (82=100,SA) SPOT MARKET PRICE INDEX:BLS & CRB: ALL COMMODITIES (1967=100) Real SPOT MARKET PRICE INDEX:BLS & CRB: ALL COMMODITIES(1967=100) PRODUCER PRICE INDEX: CRUDE PETROLEUM (82=100,NSA) PPI Crude (Relative to Core PCE) NAPM COMMODITY PRICES INDEX (PERCENT) UNITED STATES; EFFECTIVE EXCHANGE RATE(MERM)(INDEX NO.) FOREIGN EXCHANGE RATE: SWITZERLAND (SWISS FRANC PER U.S.\$) FOREIGN EXCHANGE RATE: JAPAN (YEN PER U.S.\$) FOREIGN EXCHANGE RATE: UNITED KINGDOM (CENTS PER POUND) FOREIGN EXCHANGE RATE: CANADA (CANADIAN \$ PER U.S.\$) S&P'S COMMON STOCK PRICE INDEX: COMPOSITE (1941-43=10) S&P'S COMMON STOCK PRICE INDEX: INDUSTRIALS (1941-43=10) S&P'S COMPOSITE COMMON STOCK: DIVIDEND YIELD (% PER ANNUM) S&P'S COMPOSITE COMMON STOCK: PRICE-EARNINGS RATIO (%,NSA) COMMON STOCK PRICES: DOW JONES INDUSTRIAL AVERAGE S&P'S COMPOSITE COMMON STOCK: DIVIDEND YIELD (% PER ANNUM) U. OF MICH. INDEX OF CONSUMER EXPECTATIONS(BCD-83) PURCHASING MANAGERS' INDEX (SA)

 $\mathbf{2}$ 

2

 $\mathbf{5}$ 

5

5

5

5

5

5

NAPM NEW ORDERS INDEX (PERCENT)	
NAPM VENDOR DELIVERIES INDEX (PERCENT)	
NAPM INVENTORIES INDEX (PERCENT)	
NEW ORDERS (NET) - CONSUMER GOODS & MATERIALS, 1996 DOLLARS (BCI)	
NEW ORDERS, NONDEFENSE CAPITAL GOODS, IN 1996 DOLLARS (BCI)	

 $\begin{array}{c}
 1 \\
 1 \\
 5 \\
 5
 \end{array}$