

Butucea, Cristina; Nussbaum, Michael

**Working Paper**

## Asymptotic equivalence of discretely observed geometric Brownian motion to a Gaussian shift

SFB 373 Discussion Paper, No. 1999,59

**Provided in Cooperation with:**

Collaborative Research Center 373: Quantification and Simulation of Economic Processes,  
Humboldt University Berlin

*Suggested Citation:* Butucea, Cristina; Nussbaum, Michael (1999) : Asymptotic equivalence of discretely observed geometric Brownian motion to a Gaussian shift, SFB 373 Discussion Paper, No. 1999,59, Humboldt University of Berlin, Interdisciplinary Research Project 373: Quantification and Simulation of Economic Processes, Berlin,  
<https://nbn-resolving.de/urn:nbn:de:kobv:11-10046497>

This Version is available at:

<https://hdl.handle.net/10419/61778>

**Standard-Nutzungsbedingungen:**

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

**Terms of use:**

*Documents in EconStor may be saved and copied for your personal and scholarly purposes.*

*You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.*

*If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.*

# Asymptotic equivalence of discretely observed geometric Brownian motion to a Gaussian shift

Cristina Butucea, Michael Nussbaum  
SFB 373, Humboldt-Universität zu Berlin,  
Weierstrass Institute Berlin, Germany

## Abstract

Financial models consider often stochastic processes satisfying certain differential equations. We show that the solution of a particular geometric Brownian motion observed in discrete time is asymptotically equivalent with a Gaussian white noise model.

## 1 Introduction

In finance, stock prices are modelled by geometric Brownian motion since the well known Black-Scholes model, classically written as follows

$$dS_t = \mu S_t dt + \sigma S_t dW_t,$$

where  $\{W_t, t \in [0, 1]\}$  is a Brownian motion, with appreciation rate  $\mu \in \mathbb{R}$  and volatility  $\sigma > 0$ . The process  $S_t$  describes the evolution of the stock price at continuous time  $t$ .

This model is further used in computing the price of derivative securities (e.g. option valuation). Thus, a good knowledge of the volatility structure, in particular, is needed in the analysis of financial analysis and forecasting. For further details of financial calculations we refer to Musiela and Rutkowski [4].

We also put an accent here on the study of the time-dependent volatility. The following geometric Brownian motion is considered

$$dS_t = \sigma(t) S_t dW_t, \tag{1}$$

where the volatility function  $\sigma$  is time-dependent,  $t \in [0, 1]$ , belongs to some set  $H$  and is bounded away from 0 ( $\sigma(t) \geq \varepsilon > 0$  for all  $t$  in  $[0, 1]$ ). Without loss of generality we assume that the initial value is  $S_0 = 1$ .

Consider  $Y_t = \log S_t$  and substitute in equation (1). The volatility is a continuous function and then a solution to this last equation is known to satisfy

$$dY_t = -\frac{\sigma^2(t)}{2} dt + \sigma(t) dW_t. \tag{2}$$

Remark that, from a practical point of view, the continuous time model (2) is obviously unrealistic. We have at our disposal discrete time observations. Therefore, we consider discrete observations of the process  $Y_t$  in (2) at times  $t_i = i/n$ , for  $i = 1, \dots, n$  and the following normalized independent increments

$$Z_i = \sqrt{n} (Y_{t_i} - Y_{t_{i-1}}) = -\frac{\sqrt{n}}{2} \int_{t_{i-1}}^{t_i} \sigma^2(u) du + \sqrt{n} \int_{t_{i-1}}^{t_i} \sigma(u) dW_u. \quad (3)$$

We show that the statistical experiment (as defined later on) induced by observations in (3) has the same asymptotic behavior as the following white noise shift model:

$$dX_t = \frac{1}{\sqrt{2}} \log \sigma(t) dt + \frac{1}{\sqrt{n}} dW_t, \quad (4)$$

with  $t \in [0, 1]$ .

The approach is in the sense of Le Cam theory of asymptotic equivalence of statistical experiments, in a nonparametric setup. We assume that the volatility belongs to a Hölder class of functions,  $H(\beta, L)$ , with  $\beta \in (1/2, 1]$  of functions that are bounded away from 0:

$$H(\beta, L) = \left\{ \sigma : [0, 1] \rightarrow [\varepsilon, \infty) \mid \varepsilon > 0, |\sigma(x) - \sigma(y)| \leq L |x - y|^\beta \right\}.$$

Note that we prove the asymptotic equivalence without indicating constructive procedures (Markov kernels) which allow to pass from one model to the other (see Nussbaum and Klemelä [6] for such results).

In this paper, we obtain the result that the geometric Brownian motion (1), observed over a regular grid is asymptotically equivalent with the white noise model drifted observations in (4), in the sense of Le Cam's deficiency measure tending to 0 with  $n \rightarrow \infty$ . In Section 2 we introduce briefly the asymptotic equivalence notion and the main tools in our proofs. Results are stated and proven in several consecutive steps in Section 3.

## 2 Notation, definitions

Let  $E_{i,n} = \{\Omega, \mathcal{A}, (P_{i,n} = P_{i,n}(\sigma) : \sigma \in H)\}$ ,  $i = 1, 2$  be two statistical experiments defined on the same probability space  $(\Omega, \mathcal{A})$  and indexed by the same parameter space.

In order to evaluate how close the two experiments are, we consider Le Cam deficiency measure or the  $\Delta$ -pseudodistance. The experiments  $E_{i,n}$  are said to be asymptotically equivalent if this distance,  $\Delta(E_{1,n}, E_{2,n})$ , tends to 0 with  $n$ . For more details about the asymptotic equivalence theory we refer to Le Cam and Yang [3], Nussbaum [5]. In the following, we describe upper bounds of this distance by means of Hellinger distance.

Assume that there exists  $\sigma_0$  in  $H$  such that the associated probability measure dominates all other probability measures of the same experiment. If the likelihood processes  $\Lambda_{i,n}(\sigma) = dP_{i,n}(\sigma) / dP_{i,n}(\sigma_0)$  are defined on the same probability space (or have such versions), then the  $\Delta$ -distance satisfies

$$\Delta(E_{1,n}, E_{2,n}) \leq \frac{1}{2} \sup_{\sigma \in H} E |\Lambda_{1,n}(\sigma) - \Lambda_{2,n}(\sigma)|.$$

This upper bound can be expressed by means of Hellinger distance as follows. The Hellinger distance between two likelihood processes is by definition

$$H^2(\Lambda_{1,n}, \Lambda_{2,n}) = \frac{1}{2} E \left( \sqrt{\Lambda_{1,n}} - \sqrt{\Lambda_{2,n}} \right)^2.$$

Then we deduce easily that

$$\Delta(E_{1,n}, E_{2,n}) \leq \sup_{\sigma \in H} H(\Lambda_{1,n}, \Lambda_{2,n}). \quad (5)$$

In particular, we use the following formulas, where  $N(\mu, \sigma^2)$  denotes the Gaussian law with mean  $\mu$  and variance  $\sigma^2$ ,

$$H^2(N(\mu_1, \sigma^2), N(\mu_2, \sigma^2)) \leq \frac{(\mu_1 - \mu_2)^2}{4\sigma^2}, \quad (6)$$

$$H^2(N(\mu, \sigma_1^2), N(\mu, \sigma_2^2)) \leq 2 \frac{(\sigma_1 - \sigma_2)^2}{\sigma_1^2 + \sigma_2^2}. \quad (7)$$

### 3 Equivalence results

Let  $P_n^Y$  and  $P_n^X$  denote the distribution of  $\{Y_i = Y_{t_i}\}_{i=1, \dots, n}$  in (2) and  $\{X_t(n)\}_{t \in [0,1]}$  in (4), respectively, and  $E_n^Y = \left\{ [0, 1]^n, \mathcal{B}_{[0,1]}^n, (P_n^Y(\sigma), \sigma \in H(\beta, L)) \right\}$ ,  $E_n^X = \left\{ C_{[0,1]}, \mathcal{B}_{C_{[0,1]}}, (P_n^X(\sigma), \sigma \in H(\beta, L)) \right\}$  the associated experiments. We prove that these experiments are equivalent in a few steps.

Consider the following random variables

$$Z_i^* = -\frac{\sigma^2(t_i)}{2\sqrt{n}} + \sigma(t_i) \xi_i, \quad (8)$$

where  $\xi_i$  are independent, having standard Gaussian law.

**Lemma 1** *If  $E_{1,n}$  is the experiment generated by  $\{Z_i\}_{i=1, \dots, n}$  in (3) and  $E_{1,n}^*$  by  $\{Z_i^*\}_{i=1, \dots, n}$  in (8), then these experiments are asymptotically equivalent, i.e.*

$$\lim_{n \rightarrow \infty} \Delta(E_{1,n}, E_{1,n}^*) = 0.$$

**Proof.** Let us remark that observations  $Z_i$  and  $Z_i^*$  in (3) and (8) are independent and have Gaussian law,

$$P_i : N \left( -\frac{h_i(\sigma^2)}{2\sqrt{n}}, h_i(\sigma^2) \right), \text{ where } h_i(\sigma^2) = n \int_{t_{i-1}}^{t_i} \sigma^2(u) du,$$

respectively,

$$P_i^* : N \left( -\frac{\sigma^2(t_i)}{2\sqrt{n}}, \sigma^2(t_i) \right).$$

In order to prove that  $\Delta(E_{1,n}, E_{1,n}^*) \rightarrow 0$ , when  $n \rightarrow \infty$ , it suffices by (5) to prove that  $H^2(P_{1,n}, P_{1,n}^*) \rightarrow 0$ , as  $n \rightarrow \infty$ , where  $P_{1,n} = \bigotimes_{i=1}^n P_i$  and  $P_{1,n}^* = \bigotimes_{i=1}^n P_i^*$ . We have

$$\begin{aligned} H^2(P_{1,n}, P_{1,n}^*) &\leq 2 \sum_{i=1}^n H^2(P_i, P_i^*) \\ &\leq 2 \sum_{i=1}^n \left[ H^2(P_i, \tilde{P}_i) + H^2(\tilde{P}_i, P_i^*) \right], \end{aligned}$$

where  $\tilde{P}_i$  denotes the law of the following independent random variables

$$\tilde{Z}_i = -\frac{h_i(\sigma^2)}{2\sqrt{n}} + \sigma(t_i) \xi_i.$$

Using formulas (6) and (7), we get

$$\begin{aligned} H^2(P_{1,n}, P_{1,n}^*) &\leq 2 \sum_{i=1}^n \left[ 2 \frac{\left( \sqrt{h_i(\sigma^2)} - \sigma(t_i) \right)^2}{h_i(\sigma^2) + \sigma^2(t_i)} + \frac{1}{4\sigma^2(t_i)} \frac{(h_i(\sigma^2) - \sigma^2(t_i))^2}{4n} \right] \\ &\leq \sum_{i=1}^n (h_i(\sigma^2) - \sigma^2(t_i))^2 \left[ \frac{1}{\varepsilon^2} + \frac{1}{8n\varepsilon^2} \right]. \end{aligned}$$

Remark that for the continuous function  $\sigma$  there exists a value  $u_i \in [t_{i-1}, t_i]$  such that  $h_i(\sigma^2) = \sigma^2(u_i)$  and then

$$\begin{aligned} |h_i(\sigma^2) - \sigma^2(t_i)| &= |\sigma^2(u_i) - \sigma^2(t_i)| \\ &\leq 2B(\beta, L) |u_i - t_i|^\beta = O(1) n^{-\beta}, \end{aligned}$$

where  $B(\beta, L)$  is an upper bound of functions  $\sigma$  in  $H(\beta, L)$ . Finally,

$$H^2(P_{1,n}, P_{1,n}^*) \leq n^{1-2\beta} (1 + o(1)),$$

which tends to 0 for all  $\beta > 1/2$ . ■

**Remark 1** In addition to the main result we notice the following facts. Similarly, let us consider the process  $M_t$  satisfying

$$dM_t = \sigma(t) dW_t$$

and discrete observations of this process at times  $t_i = i/n$ , for  $i = 1, \dots, n$ , together with

$$U_i = \sqrt{n} (M_{t_i} - M_{t_{i-1}}) = \sqrt{n} \int_{t_{i-1}}^{t_i} \sigma(u) dW_u. \quad (9)$$

Analogously to Lemma 1, it is stated in the following Lemma that the Gaussian random variables

$$U_i^* = \sigma(t_i) \xi_i \quad (10)$$

introduce an equivalent experiment to (9). Via the same steps as for the observations  $Z_i^*$  we obtain in our Theorem the equivalence of (10) to observations  $X_i$  in (4). This proves the fact that the drift part of the discrete observations  $Y_i$  in (2) is asymptotically non significant.

**Lemma 2** If  $E_{2,n}$  is the experiment associated to  $\{U_i\}_{i=1,\dots,n}$  in (9) and  $E_{2,n}^*$  to  $\{U_i^*\}_{i=1,\dots,n}$  in (10) then

$$\lim_{n \rightarrow \infty} \Delta(E_{2,n}, E_{2,n}^*) = 0.$$

**Proof.** The proof goes similarly to that of Lemma 1, by writing the laws of  $\{U_i\}_{i=1,\dots,n}$  and  $\{U_i^*\}_{i=1,\dots,n}$ ,  $P_{2,n} : \bigotimes_{i=1}^n N(0, h_i(\sigma^2))$  and  $P_{2,n}^* : \bigotimes_{i=1}^n N(0, \sigma^2(t_i))$ , respectively. By inequalities (6) and (7) on the Hellinger distance and using the Hölder property of the volatility function  $\sigma$ , we conclude immediately. ■

The following result states that the solution  $Y_t$  of the geometric Brownian motion, discretely observed and the Gaussian drifted model  $X_t$  are equivalent. Moreover, the non random expectation term in (8) is asymptotically non informative and all the information can be 'extracted' from the variance term, in model (10).

**Theorem 1** The experiments  $E_n^Y$  and  $E_n^X$  associated to  $\{Y_i\}_{i=1,\dots,n}$  in (3) and  $\{X_t(n)\}_{t \in [0,1]}$  in (4), respectively, are asymptotically equivalent, i.e.

$$\lim_{n \rightarrow \infty} \Delta(E_n^Y, E_n^X) = 0.$$

Moreover, the drift part of observations  $\{Y_i\}_{i=1,\dots,n}$  in (3) is asymptotically non significant in this model.

**Proof.** We use several times the transitivity property of the asymptotic equivalence property. It is sufficient to prove that  $E_{1,n}^*$  (see Lemma 1), respectively  $E_{2,n}^*$  (see Lemma 2) are equivalent with the same global experiment  $E_n^X$ .

Let us consider first  $E_{2,n}^* = \left\{ [0, 1]^n, \mathcal{B}_{[0,1]}^n, (P_{2,n}^*(\sigma), \sigma \in H(\beta, L)) \right\}$  and remark that  $P_{2,n}^* = \bigotimes_{i=1}^n P_{\sigma(t_i)}^2$  can be written as a product measure of distributions from an exponential family having parameters  $\sigma(t_i)$ . By following the same steps as in Grama and Nussbaum [2] (see Example 4.2, p 189), we deduce that this model is equivalent to

$$Z_i^{**} = \frac{\log \sigma(t_i)}{\sqrt{2}} + \varepsilon_i, \quad (11)$$

where  $\sigma \in H(\beta, L)$  and  $\varepsilon_i$  are i.i.d. standard normal distributed.

The proof that  $E_{1,n}^* = \left\{ [0, 1]^n, \mathcal{B}_{[0,1]}^n, (P_{1,n}^*(\sigma), \sigma \in H(\beta, L)) \right\}$  with  $P_{1,n}^*(\sigma) = \bigotimes_{i=1}^n P_{\sigma(t_i)}^1$  is asymptotically equivalent to (11) follows the same arguments slightly differently, as given below.

The final step is immediate and consists of applying Brown and Low [1] in order to deduce that the experiment induced by observations  $Z_i^{**}$  is asymptotically equivalent to  $E_n^X$  in (4). These facts finish the proof of our theorem.

The global experiment  $E_{1,n}^*$  has to be reduced to the local experiment of the same observations with  $\sigma \in H_{\sigma_0, \gamma_n}(\beta, L)$ , a shrinking neighborhood of  $\sigma_0$ . For a fixed function  $\sigma_0$  in  $H(\beta, L)$  define

$$H_{\sigma_0, \gamma_n}(\beta, L) = \{ \sigma \in H(\beta, L) : \|\sigma - \sigma_0\|_\infty \leq \gamma_n \},$$

where  $\gamma_n = (n / \log n)^{-\beta/(2\beta+1)}$ .

Observations  $Z_i^*$ , (8), have a normal density written as an exponential model as follows

$$\begin{aligned} p_i(x) &= \frac{1}{\sigma(t_i) \sqrt{2\pi}} \exp \left\{ -\frac{1}{2\sigma^2(t_i)} \left( x + \frac{\sigma^2(t_i)}{2\sqrt{n}} \right)^2 \right\} \\ &= \exp \left\{ \lambda_i \frac{x^2}{2} - \frac{x}{2\sqrt{n}} + \frac{1}{8\lambda_i n} + \frac{1}{2} \log \left( -\frac{\lambda_i}{2\pi} \right) \right\}, \end{aligned}$$

where  $\lambda_i = -1/\sigma^2(t_i)$ . We write this exponential model as

$$p_i(x) = \exp \{ \lambda_i U(x) - V_n(\lambda_i) \} \exp \left\{ -\frac{x}{2\sqrt{n}} \right\},$$

where  $U(x) = x^2/2$  and

$$\begin{aligned} V_n(\lambda_i) &= -\frac{1}{2} \log \left( -\frac{\lambda_i}{2\pi} \right) - \frac{1}{8\lambda_i n}, \\ V_n'(\lambda_i) &= -\frac{1}{2\lambda_i} + \frac{1}{8\lambda_i^2 n}, \\ V_n''(\lambda_i) &= \frac{1}{2\lambda_i^2} \left( 1 - \frac{1}{2\lambda_i n} \right) = I_n(\lambda_i). \end{aligned}$$

Via Theorem 3.2 in Grama and Nussbaum [2], the local  $E_{1,n}^*$  is equivalent to the discrete Gaussian experiment:

$$Z_i^{(1)} = \sigma(t_i) + I_n(\sigma_0(t_i))^{-1/2} \varepsilon_i, \sigma \in H_{\sigma_0, \gamma_n}(\beta, L). \quad (12)$$

Denote  $I(\lambda_i) = 1/(2\lambda_i^2)$  and consider the variables

$$Z_i^{(2)} = \sigma(t_i) + I(\sigma_0(t_i))^{-1/2} \varepsilon_i, \sigma \in H_{\sigma_0, \gamma_n}(\beta, L). \quad (13)$$

We prove that the experiments generated by  $Z^{(1)} = \{Z_i^{(1)}\}_{i=1, \dots, n}$  and  $Z^{(2)} = \{Z_i^{(2)}\}_{i=1, \dots, n}$  are asymptotically equivalent, in order to bring our problem back to the case of Grama and Nussbaum [2] and apply the variance-stabilizing transformation.

We show that the Hellinger distance tends to 0 with  $n$ , as follows

$$\begin{aligned} H^2(Z^{(1)}, Z^{(2)}) &\leq 2 \sum_{i=1}^n H^2(N(\sigma(t_i), I_n(\sigma_0(t_i))^{-1}), N(\sigma(t_i), I(\sigma_0(t_i))^{-1})) \\ &\leq 4 \sum_{i=1}^n \frac{\left(I_n(\sigma_0(t_i))^{-1/2} - I(\sigma_0(t_i))^{-1/2}\right)^2}{I_n(\sigma_0(t_i))^{-1} + I(\sigma_0(t_i))^{-1}} \\ &\leq 4 \sum_{i=1}^n \frac{\left(\left(1 - \frac{1}{2n\sigma_0(t_i)}\right)^{-1/2} - 1\right)^2}{\left(1 - \frac{1}{2n\sigma_0(t_i)}\right)^{-1} + 1}. \end{aligned}$$

If we denote  $a_i = 1/(2n\sigma_0(t_i))$  then

$$\frac{\left((1 - a_i)^{-1/2} - 1\right)^2}{(1 - a_i)^{-1} + 1} = \frac{\left(1 - (1 - a_i)^{-1/2}\right)^2}{2 - a_i} \leq a_i^2$$

and

$$H^2(Z^{(1)}, Z^{(2)}) \leq \frac{1}{n^2} \sum_{i=1}^n \frac{1}{\sigma_0^2(t_i)} \leq \frac{1}{n\varepsilon^2} \xrightarrow{n \rightarrow \infty} 0.$$

The last step is to apply a variance-stabilizing transformation  $\Gamma$ , solution of the differential equation  $\Gamma'(\lambda) = \sqrt{I(\lambda)}$ , which gives  $\Gamma(\lambda) = \log \lambda / \sqrt{2}$ . We obtain the equivalent local Gaussian experiment

$$Z_i^{(3)} = \frac{\log \sigma(t_i)}{\sqrt{2}} + \varepsilon_i, \sigma \in H_{\sigma_0, \gamma_n}(\beta, L).$$

By globalization arguments of Grama and Nussbaum [2] (Theorem 3.8) we obtain the global experiment in (11) (where  $\sigma$  in the class  $H(\beta, L)$ ). ■



## References

- [1] Brown, L. D. and Low, M. G. (1996) Asymptotic equivalence of nonparametric regression and white noise. *Ann. Statist.* **24**, 2384-2398
- [2] Grama, I. and Nussbaum, M. (1998) Asymptotic equivalence for nonparametric generalized linear models. *Probab. Th. Rel. Fields* **111**, 167-214
- [3] Le Cam, L. and Yang, G. (1990) *Asymptotics in Statistics*. Springer-Verlag, New-York
- [4] Musiela, M. and Rutkowski, M. (1997) *Martingale Methods in Financial Modelling*. Springer-Verlag Berlin Heidelberg
- [5] Nussbaum, M. (1996) Asymptotic equivalence of density estimation and Gaussian white noise. *Ann. Statist.* **24**, 2399-2430
- [6] Nussbaum, M. and Klemelä, J. (1998) Constructive asymptotic equivalence of density estimation and Gaussian white noise. Disc. paper 53, SFB 373, Humboldt-Universität zu Berlin