

Golubev, Georgi; Härdle, Wolfgang

**Working Paper**

## On adaptive smoothing in partial linear models

SFB 373 Discussion Paper, No. 2001,48

**Provided in Cooperation with:**

Collaborative Research Center 373: Quantification and Simulation of Economic Processes,  
Humboldt University Berlin

*Suggested Citation:* Golubev, Georgi; Härdle, Wolfgang (2001) : On adaptive smoothing in partial linear models, SFB 373 Discussion Paper, No. 2001,48, Humboldt University of Berlin, Interdisciplinary Research Project 373: Quantification and Simulation of Economic Processes, Berlin, <https://nbn-resolving.de/urn:nbn:de:kobv:11-10049963>

This Version is available at:

<https://hdl.handle.net/10419/62684>

**Standard-Nutzungsbedingungen:**

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

**Terms of use:**

*Documents in EconStor may be saved and copied for your personal and scholarly purposes.*

*You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.*

*If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.*

# On adaptive smoothing in partial linear models

G. Golubev<sup>1\*</sup> and W. Härdle<sup>2†</sup>

<sup>1</sup>CMI, Université de Provence, 39 rue F. Joliot-Curie, 13453 Marseille Cedex 13, France  
e-mail: golubev@gyptis.univ-mrs.fr

<sup>2</sup>Humboldt-Universität zu Berlin, Spandauer Straße 1, 10178 Berlin, Germany  
e-mail: haerdle@wiwi.hu-berlin.de

*We consider a problem of estimation of parametric components in a partial linear model. Suppose that a finite set  $\mathcal{E}$  of linear estimators is given. Our goal is to mimic the estimator in  $\mathcal{E}$  that has the smallest risk. Using a second order expansion of the risk of linear estimators we propose a practically feasible adaptive procedure for choice of smoothing parameters based on the principle of unbiased risk estimation.*

*Key words:* Partial linear model, second order minimax risk, adaptive estimation.

*AMS 1991 Subject Classification:* Primary 62G05, 62G20; secondary 62C20.

---

\*The author was partially supported by RFFI grant 001–00266.

†The author was supported by the Deutsche Forschungsgemeinschaft, Sonderforschungsbereich 373 "Quantifikation und Simulation Ökonomischer Prozesse", Humboldt-Universität zu Berlin.

# 1 Introduction

Partial linear models represent now a flexible and growing class of models for statistical applications. In the present paper we will deal with the simplest partial linear model

$$Y_i = \theta^T Z_i + m(X_i) + \xi_i, \quad i = 1, \dots, n, \quad (1)$$

where  $\theta \in \mathbf{R}^d$  is a column-vector of unknown parameters,  $\xi_i$  are i.i.d. random variables with zero mean and the finite variance  $\sigma^2 = \mathbf{E} \xi_i^2$ . The regressors  $X_i \in [0, 1]$  are assumed to be i.i.d. random variables with a strictly positive density  $q(x)$  on  $[0, 1]$ . We will also assume that they do not depend on  $\xi_i$ . The nuisance function  $m(x)$ ,  $x \in [0, 1]$  is unknown but such that the random variables  $m(X_i)$  have zero mean.

It is well known that if  $m(x)$ ,  $x \in [0, 1]$  is sufficiently smooth then  $\theta$  can be estimated with the ordinary parametric rate Heckman (1986), Rice (1986), Robinson (1987), Speckman (1988), Bhattacharia and Zhao (1997), Mammen and Van de Geer (1997), Chen (1998). Further references and applications of partially linear models can be found in the recent book by Härdle, Liang and Gao (2000). Roughly speaking almost all first order effects in these models can be explained at a heuristic level if one assumes that the nuisance function  $m(x)$  is known. In Golubev and Härdle (2000) the second order term of quadratic minimax risk was found up to a constant (see Theorems 1, 2 below).

This paper continues Golubev and Härdle (2000) concentrating on data driven choice of smoothing parameters for penalized least-square estimators. Usually this problem is considered as a problem of minor importance in the modern theory of semi-parametric estimation. But from a practical point of view this problem plays the same role as a data driven choice of the bandwidth in density estimation problem. There is a vast mathematical literature on data driven choice of smoothing parameters (see Akaike (1973), Mallows (1973), Efroimovich and Pinsker (1984), Lepski (1991), Lepski (1992), Golubev and Nussbaum (1992), Kneip (1994), Lepski and Spokoiny (1997), Nemirovskii (1998), Tsybakov (1998), Barron, Birgé and Massart (1999), among others). The problem of adaptive choice of smoothing parameters in the framework of the second order minimax theory for distribution function estimation was considered in Golubev and Levit (1996).

To simplify some technical details we will assume that the nuisance function  $m(x)$  belongs to the Sobolev ball

$$\mathbf{W}_0 = \left\{ m : \int_0^1 [m^{(\beta)}(x)]^2 dx \leq L, \quad \int_0^1 q(x)m(x) dx = 0 \right\},$$

where the smoothness  $\beta$  is positive integer.

Our consideration is based on the so-called orthogonal series approach. The cornerstone idea of this approach is to parameterize the functional class  $\mathbf{W}_0$ . We do this by constructing an orthonormal system in the Hilbert space  $\mathbf{L}_q^2[0, 1]$  which is equipped with

the norm  $\|\cdot\|_q$  and with the inner product  $\langle \cdot, \cdot \rangle_q$  defined by

$$\|f\|_q^2 = \int_0^1 q(x) f^2(x) dx, \quad \langle f, g \rangle_q = \int_0^1 q(x) f(x) g(x) dx.$$

Let  $\psi_k$ ,  $k = 0, \dots, \beta - 1$  be the first orthonormal polynomials in  $\mathbf{L}_q^2[0, 1]$ . The remainder functions  $\psi_k$ ,  $k \geq \beta$  are defined as solutions of the following boundary value problem (see e.g. Speckman (1985))

$$\begin{aligned} (-1)^\beta \frac{d^{2\beta}}{dx^{2\beta}} \psi_s(x) &= \lambda_s q(x) \psi_s(x), \\ \frac{d^k}{dx^k} \psi_s(x) \Big|_{x=0} &= \frac{d^k}{dx^k} \psi_s(x) \Big|_{x=1} = 0, \quad k = \beta, \dots, 2\beta - 1. \end{aligned} \quad (2)$$

The eigen functions  $\psi_s(x)$  are uniformly bounded in  $x$  and  $s$ . The asymptotic behavior of the eigen values  $\lambda_k$  plays a very important role in splines theory and it is well-known that

$$\lambda_s = [1 + o(1)] (\pi s)^{2\beta} \left[ \int_0^1 q^{1/(2\beta)}(x) dx \right]^{2\beta}, \quad s \rightarrow \infty. \quad (3)$$

We refer the interested reader to the book by Tikhomirov (1986) for geometric interpretation of this basis. Details about asymptotic behavior of the eigen values can be found in Utreras (1980), Speckman (1985). Notice that any function  $m \in \mathbf{W}_0$  can be represented as the Fourier series

$$m(t) = \sum_{k=1}^{\infty} \nu_k \psi_k(t), \quad \text{where } \nu_k = \langle m, \psi_k \rangle_q \quad \text{and} \quad \sum_{k=1}^{\infty} \nu_k^2 \lambda_k \leq L. \quad (4)$$

Now we are ready to construct a penalized least-square estimator. Let  $\Sigma$  be an arbitrary diagonal matrix with the entries  $\Sigma_{kk} = \sigma_k \neq 0$  and let the matrix  $\Psi$  be defined as  $\Psi_{ki} = \psi_k(X_i)$ . We estimate the parameter  $\theta$  by

$$\hat{\theta} = \arg \min_{\theta \in R^d} \min_{\nu} \left\{ \|Y - Z^T \theta - \Psi^T \nu\|^2 + \|\Sigma^{-1} \nu\|^2 \right\}. \quad (5)$$

Our further considerations will be essentially based on the second order theory of semi-parametric estimation. The next two theorems describe the performance of  $\hat{\theta}$  up to the second order terms (for more details see Golubev and Härdle (2000)). Denote

$$H = n \Sigma^{-2} (n \Sigma^{-2} + E)^{-1}, \quad (6)$$

where  $E$  is identity matrix, and let

$$IMSE[H, \nu] = \|(E - H)\nu\|^2 + \frac{\sigma^2}{n} \text{tr } H^2$$

be the integrated mean-square error of recovering the function  $m(x)$  in the model (1) provided that  $\theta$  is known. In other words

$$IMSE[H, \nu] = [1 + o(1)] \mathbf{E} \sum_{k=1}^{\infty} (\nu_k - H_{kk} \bar{\nu}_k)^2,$$

where

$$\bar{\nu}_k = \frac{1}{n} \sum_{i=1}^n (Y_i - \theta^T Z_i) \psi_k(X_i).$$

**Theorem 1** Suppose that  $\mathbf{E} |\xi_k|^{2+\delta} < \infty$  for some  $\delta > 0$  and

$$\lim_{n \rightarrow \infty} \log^{1/2} n \max_{i,k} Z_{ki}^2 / \sum_{i=1}^n Z_{ki}^2 = 0, \quad \max_n \|(ZZ^T)^{-1}\| \max_k \sum_{i=1}^n Z_{ki}^2 < \infty, \quad (7)$$

$$\lim_{n \rightarrow \infty} \text{tr}^2 H \log^{1/2} n / n = 0.$$

Then as  $n \rightarrow \infty$  uniformly in  $m \in \mathbf{W}_0$

$$\mathbf{E}(\hat{\theta} - \theta)(\hat{\theta} - \theta)^T = (ZZ^T)^{-1} \{ \sigma^2 + [1 + o(1)] IMSE[H, \nu] \}.$$

This theorem demonstrates how the risk of  $\hat{\theta}$  depends on the nuisance function  $m(\cdot)$  and the penalization matrix  $\Sigma^{-1}$ . We see that the first order term of the risk does not depend on the nonparametric nuisance function whereas the second order term in the risk expansion coincides with the integrated mean square error of recovering  $m(x)$  in the model (1) provided  $\theta$  is known. This is why we are interested in the analysis of the second order theory. One can maximize the second order term over all nuisance functions from  $\mathbf{W}_0$  and then minimize it over all penalizations  $\Sigma$  or equivalently over all  $H$ . Thus one obtains the following result about the minimax penalization (see also Pinsker (1980)).

**Theorem 2** Let  $\hat{\theta}$  be the estimator defined by (5) with  $\Sigma = H^{1/2}(E - H)^{-1/2} \sigma / \sqrt{n}$ , where  $H$  is the diagonal matrix

$$H_{ss} = \left[ 1 - \omega \sqrt{\lambda_s} \right]_+, \quad (8)$$

where  $[x]_+ = \max(x, 0)$ , and  $\omega$  be a root of the equation

$$\frac{\sigma^2}{n} \sum_{s=1}^{\infty} \lambda_s \left[ \frac{1}{\omega \sqrt{\lambda_s}} - 1 \right]_+ = L. \quad (9)$$

Under the conditions of Theorem 1 as  $n \rightarrow \infty$

$$\sup_{m \in \mathbf{W}_0} \mathbf{E}(\hat{\theta} - \theta)(\hat{\theta} - \theta)^T = (ZZ^T)^{-1} \sigma^2 \{1 + [1 + o(1)]n^{-1} \text{tr} H\}.$$

If  $\xi_i$  are Gaussian then for any  $B > 0$

$$\inf_{\tilde{\theta}} \sup_{m \in \mathbf{W}_0} \sup_{\theta \in \mathbf{R}^d} \mathbf{E}(\tilde{\theta} - \theta)(\tilde{\theta} - \theta)^T = (ZZ^T)^{-1} \sigma^2 \{1 + [1 + o(1)]n^{-1} \text{tr} H\},$$

where  $\inf$  is taken over all estimators of the parameter  $\theta$ .

This result shows that the optimal regularization matrix  $\Sigma$  strongly depends on the parameter  $L$ , which defines the functional class  $\mathbf{W}_0$ . In practice this parameter is hardly known. Therefore our next step is to construct a practically feasible data-driven method for adaptive penalization.

## 2 An adaptive estimator

The goal of adaptation is to choose the regularization matrix  $\Sigma$  in (5) based on the observations in order to minimize the covariance matrix  $\mathbf{E}(\hat{\theta} - \theta)(\hat{\theta} - \theta)^T$ . Theorem 1 plays an essential role in a such choice since it says that the second order risk of  $\hat{\theta}$  is controlled by  $IMSE[H, \nu]$ . Thus we see that in order to minimize the second order risk we have to minimize  $IMSE[H, \nu]$  with respect to  $H$ . This functional depends on the nuisance parameters  $\nu$ , which are of course not known. The ordinary way to perform minimization in this situation is based on the principle of unbiased risk estimation. The main idea is well known and commonly used in nonparametric estimation (see e.g. Akaike (1973), Mallows (1973)). Heuristic arguments for adaptive choice of  $H$  are the following. Let  $\mu = (\theta_1, \dots, \theta_d, \nu_1, \dots)^T$ . Define an estimator of this vector by

$$\hat{\mu}(H) = \arg \min_{\mu} \left\{ \|Y - Q^T \mu\|^2 + n \|H^{-1/2}(E - H)^{1/2} \mu\|^2 \right\}, \quad Q = \begin{pmatrix} Z \\ \Psi \end{pmatrix}. \quad (10)$$

Simple algebra reveals that  $\hat{\mu}(H)$  can be computed as

$$\hat{\mu}(H) = [QQ^T + nH^{-1}(E - H)]^{-1} QY. \quad (11)$$

With  $U(H) = Q^T[QQ^T + nH^{-1}(E - H)]^{-1}Q$  we easily see that

$$\mathbf{E} \|Y - Q^T \hat{\mu}(H)\|^2 = \mathbf{E} \|Q^T [\hat{\mu}(H) - \mu]\|^2 - 2\sigma^2 \text{tr} U(H). \quad (12)$$

Noticing that by the law of large numbers

$$QQ^T \approx \begin{pmatrix} ZZ^T & 0 \\ 0 & nE \end{pmatrix}$$

we arrive at

$$[QQ^T + nH^{-1}(E - H)]^{-1}QQ^T \approx \begin{pmatrix} E & 0 \\ 0 & H \end{pmatrix}$$

and

$$U(H) \approx Z^T(ZZ^T)^{-1}Z + \Psi^T H \Psi / n.$$

So with this in mind we obtain omitting the terms of the order  $O(1)$

$$\begin{aligned} \mathbf{E} \|Q^T[\hat{\mu}(H) - \mu]\|^2 &\approx \mathbf{E} \|\Psi^T(E - H)\nu\|^2 + \sigma^2 \mathbf{E} \text{tr} U^2(H) \\ &\approx n \sum_{k=1}^{\infty} (1 - H_{kk})^2 \nu_k^2 + \sigma^2 \text{tr} H^2 + \sigma^2 \text{tr} Z^T(ZZ^T)^{-1}Z(1 + 2\text{tr} H/n) \\ &\approx \text{IMSE}(H)n. \end{aligned} \quad (13)$$

Thus to minimize the risk of the estimator one could choose (cf. (12))

$$\hat{H} = \arg \min_{h \in \mathcal{H}} \left\{ \|Y - Q^T \hat{\mu}(H)\|^2 + 2\sigma^2 \text{tr} U(H) \right\}, \quad (14)$$

where  $\mathcal{H}$  is a set in  $\mathbf{l}_2(0, \infty)$ , which must not be very rich. For instance we will assume that  $\mathcal{H}$  is a finite set. We discuss the required properties  $\mathcal{H}$  in more detail in the sequel. Examples of commonly used classes  $\mathcal{H}$  are projection smoothers with  $H_{kk} = \mathbf{1}\{k \leq w\}$ , for integer parameter  $w \in [1, n]$ , the Pinsker (1980) or minimax filters (see Theorem 2)

$$H_{kk} = [1 - (k/w)^\beta]_+, \quad w \in [1, n] \quad (15)$$

and smoothing splines Wahba (1990)

$$H_{kk} = (1 + \mu \lambda_k)^{-1}, \quad \mu > 0, \quad (16)$$

where  $\lambda_k$  are the eigen values of the boundary value problem (2). We would like to remind that in the last case the estimator has the form

$$\hat{\theta} = \arg \min_{\theta} \min_m \left\{ \sum_{i=1}^n [Y_i - \theta^T Z_i - m(X_i)]^2 + \mu \int_0^1 [m^{(\beta)}(x)]^2 dx \right\} \quad (17)$$

and there exist very fast computational algorithms for finding  $\hat{\theta}$  (see for more details Green and Silverman (1994) and Schimek (2000)).

Unfortunately in partial linear models the empirical risk  $n(\hat{\theta} - \theta)(\hat{\theta} - \theta)^T$  is non-degenerate. The variance of this random variable is of the order of the mean. This leads to some difficulties in evaluation of the performance of  $\hat{\theta}$  with  $\hat{H}$  from (14). To overcome these difficulties we use the idea of splitting by Nemirovskii (1998). Only a part of the whole sample say  $Y_1, \dots, Y_N$ , where  $N \ll n$  will be used to construct the matrix  $H$ .

Once  $H$  is chosen the whole sample  $Y_1, \dots, Y_n$  is used to compute the estimator. Thus our main idea is to estimate the right-hand side in (14) based on the data  $Y_1, \dots, Y_N$ . We do this in the following way. Let

$$\hat{\mu}_N(H) = \arg \min_{\mu} \left\{ \|Y - Q_N^T \mu\|_N^2 + N \|H^{-1/2}(E - H)^{1/2} \mu\|^2 \right\}$$

be the estimator of the parameter  $\mu$  based on the data  $Y_1, \dots, Y_N$ . With (12) and (13) we obtain

$$\begin{aligned} \mathbf{E} \|Y - Q^T \hat{\mu}_N(H)\|^2 + 2\sigma^2 \text{tr} U(H) &\approx n \sum_{k=1}^{\infty} (1 - H_{kk})^2 \nu_k^2 + \sigma^2 \text{tr} U^2(H) \\ &= \frac{n}{N} \left[ N \sum_{k=1}^{\infty} (1 - H_{kk})^2 \nu_k^2 + \sigma^2 \text{tr} U_N^2(H) \right] + \left(1 - \frac{n}{N}\right) \sigma^2 \text{tr} U_N^2(H) \\ &\approx \frac{n}{N} \left[ \mathbf{E} \|Y_N - Q_N^T \hat{\mu}_N(H)\|_N^2 + 2\sigma^2 \text{tr} U_N(H) \right] + \left(1 - \frac{n}{N}\right) \sigma^2 \text{tr} U_N^2(H). \end{aligned}$$

Thus we can replace (14) by

$$H^* = \arg \min_{H \in \mathcal{H}} \left\{ \|Y - Q_N^T \hat{\mu}_N(H)\|_N^2 + 2\sigma^2 \text{tr} U_N(H) - \sigma^2 \text{tr} U_N^2(H) (1 - N/n) \right\} \quad (18)$$

and the adaptive estimator of  $\theta$  is defined now by

$$\theta^* = \arg \min_{\theta} \min_{\nu} \left\{ \|Y - Z^T \theta - \Psi^T \nu\|^2 + n \|(H^*)^{-1/2}(E - H^*)^{1/2} \nu\|^2 \right\}. \quad (19)$$

Our further analysis is essentially based on some properties of the smoothing matrix  $H$ . We will assume that the set  $\mathcal{H}$  is finite, its cardinality is less then  $O(n)$  and uniformly in  $H \in \mathcal{H}$ :

$$\begin{aligned} 0 &\leq H_{kk} \leq 1, \\ H_{kk} &= 1, \text{ for } k = 1, \dots, \beta - 1, \\ \text{tr} H &\leq C_{\mathcal{H}} \text{tr} H^2, \\ \text{tr}^2 H &\leq C_{\mathcal{H}} n / \log n, \\ \sum_{k > n}^{\infty} H_{kk}^2 k^2 &\leq C_{\mathcal{H}} n, \end{aligned} \quad (20)$$

where  $C_{\mathcal{H}}$  is some constant.

In order to simplify some technical details we assume also that the regressors are such that for some sufficiently large constant  $C_Z$  the following conditions are hold

$$\max_{i \in [1, N]} Z_{ki}^2 \leq \frac{C_Z}{N} \sum_{i=1}^N Z_{ki}^2 \leq \frac{2C_Z}{n} \sum_{i=1}^n Z_{ki}^2, \quad \|(ZZ^T)^{-1}\| \sum_{k,i=1}^n Z_{ki}^2 \leq C_Z. \quad (21)$$

The following theorem is the main result of the paper.



**Theorem 3** *Let  $\xi_i$  be Gaussian and  $N = n/\log^{1+\kappa} n$ ,  $\kappa > 0$ . Then under conditions (20-21) uniformly in  $m \in \mathbf{W}_0$  and such that  $\|m\|_q \leq M < \infty$  as  $n \rightarrow \infty$*

$$\mathbf{E}(\theta^* - \theta)(\theta^* - \theta)^T = (ZZ^T)^{-1} \{ \sigma^2 + [1 + o(1)] \inf_{H \in \mathcal{H}} IMSE[H, \nu] \}.$$

This theorem could be interpreted as an oracle inequality (see Nemirovski (1998)) in the following way. Suppose we are allowed to make use of only the estimators  $\hat{\theta}$  defined by (5), (6) with  $H \in \mathcal{H}$ . Let us assume that there is an oracle which says the nuisance function  $m(\cdot)$ . If we know this function or equivalently  $\nu$  then according to Theorem 1 we can minimize the risk of  $\hat{\theta}$  up to a second order term choosing  $H^* = \arg \min_{H \in \mathcal{H}} IMSE[H, \nu]$ . It is clear with this family of estimators we cannot do better. On the other hand Theorem 3 says that we can achieve almost the same performance without oracles.

**Remark 1.** There is no other way to compute  $H^*$  from (18) except the complete search. Thus from a computational point of view it is better to have cardinality of  $\mathcal{H}$  as small as possible. But if  $\mathcal{H}$  is not sufficiently rich the risk of the adaptive estimator  $\theta^*$  may increase substantially. So there is a compromise between computational and statistical efficiency. The simplest way to resolve this problem is to use so-called exponential grids. As an example consider the family of minimax smoothers defined by (15). Let us chose a sufficiently small number  $\varepsilon$ , say  $\varepsilon = 0.1$  and consider the grid of bandwidths

$$w_s = (1 + \varepsilon)^s, \quad s = 0, \dots, \log n/\varepsilon.$$

Let  $\mathcal{H}^\varepsilon$  be a corresponding class of smoothers having the elements

$$H_{kk} = [1 - (k/w_s)^\beta]_+, \quad s = 0, \dots, \log n/\varepsilon.$$

Cardinality of  $\mathcal{H}^\varepsilon$  is  $\log n/\varepsilon$ , which is much smaller than the cardinality  $\mathcal{H}$ . On the other hand for any  $H \in \mathcal{H}$  we can find  $H^\varepsilon \in \mathcal{H}^\varepsilon$  such that uniformly in  $\nu \in \mathbf{l}_2(0, \infty)$

$$IMSE[H^\varepsilon, \nu] \leq (1 + \varepsilon) IMSE[H, \nu].$$

Indeed let  $H_{kk} = [1 - (k/w)^\beta]_+$  for some  $w \in [1, n]$ . Take  $H_{kk} = [1 - (k/w^\varepsilon)^\beta]_+$ , where  $w^\varepsilon = \min\{w_s : w_s \geq w\}$ . Since  $H_{kk}^\varepsilon \geq H_{kk}$  we evidently have  $\|(1 - H^\varepsilon)\nu\|^2 \leq \|(1 - H)\nu\|^2$  and it is easy to see that  $\|H^\varepsilon\|^2 \leq (1 + \varepsilon)\|H\|^2$ . Thus using  $\mathcal{H}^\varepsilon$  instead of  $\mathcal{H}$  we may have only a little increment of the risk but we improve significantly the computational efficiency. The same remark concerns of course the spline estimator.

**Remark 2.** In order to construct  $\theta^*$  we divide the sample in to two parts. From a practical point of view this idea is of course not very attractive. As a rule we use the estimator

$$\hat{\theta} = \arg \min_{\theta} \min_{\nu} \{ \|Y - Z^T \theta - \Psi^T \nu\|^2 + \sigma^2 n \|\hat{H}^{-1/2} (E - \hat{H})^{1/2} \nu\|^2 \}$$

with  $\widehat{H}$  from (14). So it would be very interesting to find out whether this estimator is adaptive in the sense of Theorem 3. Unfortunately, our arguments applied in the proof of this theorem cannot be used to answer this question because of a strong dependence between the estimators of parametric and nonparametric parts in the partial linear model.

### 3 Proof of Theorem 3

We start the proof of Theorem 3 with some auxiliary results. First, using the Taylor formula we will find an asymptotic expansion for the risk of the estimator  $\widehat{\theta}$  defined by (5). Rewrite (11) as

$$\widehat{\mu}(H) - \mu = nS^{-1}H^{-1}(E - H) + S^{-1}Q\xi, \quad (22)$$

where  $S = A + B$ , with

$$A = \begin{pmatrix} ZZ^T & 0 \\ 0 & nE + \sigma^2\Sigma^{-2} \end{pmatrix}, \quad B = \begin{pmatrix} 0 & Z\Psi^T \\ \Psi Z^T & \Psi\Psi^T - nE \end{pmatrix}. \quad (23)$$

In order to compute

$$S^{-1} = A^{-1/2}(E + A^{-1/2}BA^{-1/2})^{-1}A^{-1/2}. \quad (24)$$

we use the Taylor expansion with respect to  $A^{-1/2}BA^{-1/2}$  in the right-hand side of the above equation. Denote for brevity  $H = (E + \sigma^2\Sigma^{-2}/n)^{-1}$ . Thus we have to check that the operator norm of the matrix

$$A^{-1/2}BA^{-1/2} = \begin{pmatrix} 0 & (ZZ^T)^{-1/2}Z\Psi^TH^{1/2}/\sqrt{n} \\ H^{1/2}\Psi Z^T(ZZ^T)^{-1/2}/\sqrt{n} & H^{1/2}(\Psi\Psi^T/n - E)H^{1/2} \end{pmatrix} \quad (25)$$

is sufficiently small. Denote also by  $\#\mathcal{H}$  cardinality of  $\mathcal{H}$  and  $\|Z_s\|_N^2 = \sum_{i=1}^N Z_{si}^2$ .

**Lemma 1** *For any  $x \leq n$*

$$\mathbf{P}\left\{\|A^{-1/2}BA^{-1/2}\|^2 > C(1+x)\text{tr}^2 H/n\right\} \leq \#\mathcal{H}\exp(-x^2/C),$$

where  $C$  is a sufficiently large constant.

**Proof** directly follows from Lemma 2 in Golubev and Härdle (2000). ■

The next lemma gives an asymptotic expansion for the risk of  $\widehat{\theta}$  defined by (5) with the penalization  $\Sigma = H^{1/2}(E - H)^{-1/2}/\sqrt{n}$  depending on the data  $(Y_i, X_i)$ .

**Lemma 2** *Uniformly in  $\|m\|_q \leq M < \infty$  as  $n \rightarrow \infty$  we have*

$$\mathbf{E}(\hat{\theta} - \theta)(\hat{\theta} - \theta)^T = \mathbf{E} \Delta_H \Delta_H^T + n^{-1} \|(ZZ^T)^{-1}\|^{-1} O(E), \quad (26)$$

where

$$\begin{aligned} \Delta_H &= (ZZ^T)^{-1} Z\xi + [E + o(E)](ZZ^T)^{-1} Z\Psi^T(E - H)\nu \\ &\quad + n^{-1}[E + o(E)](ZZ^T)^{-1} Z\Psi^T H\Psi Z^T (ZZ^T)^{-1} Z\xi \\ &\quad - n^{-1}[E + o(E)](ZZ^T)^{-1} Z\Psi^T H\Psi\xi \\ &\quad + n^{-1}[E + o(E)](ZZ^T)^{-1} Z\Psi^T H(\Psi\Psi^T/n - E)H\Psi\xi, \end{aligned} \quad (27)$$

$O(E)$  is a bounded  $d \times d$ -matrix and  $o(E)$  is a  $d \times d$ -matrix with the operator norm tending to 0 as  $n \rightarrow \infty$ .

**Proof.** Notice that in view of (5)  $\|Z^T(\hat{\theta} - \theta)\| \leq \|\xi + \Psi^T\nu\|$ . Therefore for any  $\delta > 0$  we have

$$\mathbf{E}\|\hat{\theta} - \theta\|^{2+2\delta} \leq C(\delta)n^{1+\delta}\|(ZZ^T)^{-1/2}\|^{-1-\delta}$$

and we get by the Hölder inequality and Lemma 1 with  $x = (2C \log(n\#\mathcal{H}))^{1/2}$

$$\begin{aligned} \mathbf{E}(\hat{\theta} - \theta)(\hat{\theta} - \theta)^T &= \mathbf{E}(\hat{\theta} - \theta)(\hat{\theta} - \theta)^T \mathbf{1} \{ \|A^{-1/2}BA^{-1/2}\| \leq \varepsilon \} \\ &\quad + \mathbf{E}(\hat{\theta} - \theta)(\hat{\theta} - \theta)^T \mathbf{1} \{ \|A^{-1/2}BA^{-1/2}\| > \varepsilon \} \\ &\leq \mathbf{E}(\hat{\theta} - \theta)(\hat{\theta} - \theta)^T \mathbf{1} \{ \|A^{-1/2}BA^{-1/2}\| \leq \varepsilon \} \\ &\quad + n\|(ZZ^T)^{-1/2}\|^{-2} \mathbf{P}^{\delta/(1+\delta)} \{ \|A^{-1/2}BA^{-1/2}\| > \varepsilon \} O(E) \\ &\leq \mathbf{E}(\hat{\theta} - \theta)(\hat{\theta} - \theta)^T \mathbf{1} \{ \|A^{-1/2}BA^{-1/2}\| \leq \varepsilon \} + n^{-1}\|(ZZ^T)^{-1}\|^{-1} O(E), \end{aligned} \quad (28)$$

Next one obtains by (24), (25) and by the Taylor formula when  $\|A^{-1/2}BA^{-1/2}\| \leq \varepsilon$

$$\begin{aligned} S^{-1} &= \begin{pmatrix} (ZZ^T)^{-1} & 0 \\ 0 & n^{-1}H \end{pmatrix} - \frac{1}{n} \begin{pmatrix} 0 & (ZZ^T)^{-1} Z\Psi^T H \\ H\Psi Z^T (ZZ^T)^{-1} & H(\Psi\Psi^T/n - E) \end{pmatrix} \\ &\quad + (E + \varepsilon O(E))n^{-1}(ZZ^T)^{-1} Z\Psi^T H \begin{pmatrix} \Psi Z^T (ZZ^T)^{-1} & (\Psi\Psi^T/n - E) \\ * & * \end{pmatrix}, \end{aligned} \quad (29)$$

where  $*$  denotes a matrix that is not needed in further calculations. Thus once again using Lemma 1, which says that  $\mathbf{P} \{ \|A^{-1/2}BA^{-1/2}\| \geq \varepsilon \}$  is exponentially small, we arrive from (22) and (27-29) at the assertion of the lemma.  $\square$

In the sequel we will use the following very simple auxiliary fact. Let  $V$  be a finite set in  $\mathbf{l}_2(1, \infty)$  with cardinality  $\#V$  and let  $\eta_k$  be zero mean random variables, such that for any given  $v \in V$

$$D(v) = \mathbf{E} \left( \sum_{k=1}^{\infty} v_k \eta_k \right)^2 = \sum_{k,l=1}^{\infty} v_k v_l \mathbf{E} \eta_k \eta_l < \infty.$$

Suppose now that  $v \in V$  are random variables depending on  $\eta_k$ ,  $k = 0, \dots$  and we want to evaluate from above  $\mathbf{E} \left( \sum_{k=0}^{\infty} v_k \eta_k \right)^2$ . The next lemma provides a solution of this problem.

**Lemma 3** Assume that for  $\lambda = \sqrt{\log(S\#V)/2}$  ( $S \geq 1$ ) and for any given  $v \in V$

$$\mathbf{E} \exp \left[ \frac{\lambda}{\sqrt{D(v)}} \sum_{k=1}^{\infty} v_k \eta_k \right] \leq \exp(2\lambda^2). \quad (30)$$

Then

$$\mathbf{E} \left( \sum_{k=1}^{\infty} v_k \eta_k \right)^2 \leq 8 \log(S\#V) \mathbf{E} D(v) + 2 \max_{v \in V} D(v) / S. \quad (31)$$

**Proof.** By the Markov inequality for any  $Q > 0$  we have

$$\begin{aligned} \mathbf{E} \left[ \sum_{k=1}^{\infty} v_k \eta_k \right]^2 &= \mathbf{E} D(v) \left[ \frac{1}{\sqrt{D(v)}} \sum_{k=1}^{\infty} v_k \eta_k \right]^2 \leq \mathbf{E} D(v) \max_{v \in V} \left[ \frac{1}{\sqrt{D(v)}} \sum_{k=1}^{\infty} v_k \eta_k \right]^2 \\ &\leq Q^2 \mathbf{E} D(v) + \max_{v \in V} D(v) \sum_{v \in V} \mathbf{P} \left\{ \left| \frac{1}{\sqrt{D(v)}} \sum_{k=1}^{\infty} v_k \eta_k \right| > Q \right\} \\ &\leq Q^2 \mathbf{E} D(v) + 2\#V \exp(-\lambda Q + 2\lambda^2) \max_{v \in V} D(v). \end{aligned}$$

Finally choosing  $Q = 2\sqrt{2 \log(S\#V)}$  we arrive at (31). ■

**Lemma 4** Let  $\eta_{sk} = \sum_{i=1}^N Z_{si} \psi_k(X_i)$ . Then uniformly in  $\nu \in \mathbf{W}_0$  and in  $H \in \mathcal{H}$

$$\mathbf{E} \left[ \sum_{k=1}^{\infty} (1 - H_{kk}) \nu_k \eta_{sk} \right]^2 \leq C \log(n\#\mathcal{H}) \|Z_s\|_N^2 \left[ \mathbf{E} \sum_{k=1}^{\infty} (1 - H_{kk})^2 \nu_k^2 + \frac{L}{n} \right]. \quad (32)$$

**Proof.** Since  $|1 - H_{kk}| \leq 1$  one obtains by the Cauchy-Schwarz inequality that for any  $M > 1$

$$\mathbf{E} \left[ \sum_{k=1}^{\infty} (1 - H_{kk}) \nu_k \eta_{sk} \right]^2 \leq 2 \mathbf{E} \left[ \sum_{k=1}^M (1 - H_{kk}) \nu_k \eta_{sk} \right]^2 + 2CM^{-2\beta+1} L \|Z_s\|_N^2. \quad (33)$$

Next we apply Lemma 3 to the first term in right-hand side of (33). So we put

$$v_k = (1 - H_{kk}) \nu_k, \quad \eta_k = \sum_{i=1}^N Z_{si} \psi_k(X_i), \quad D(v) = \|Z_s\|_N^2 \sum_{k=1}^M (1 - H_{kk})^2 \nu_k^2.$$

Check now (30). With  $\lambda = \sqrt{\log(n\#\mathcal{H})/2}$  and  $M = N/[C_Z \log(n\#\mathcal{H})]$  we get by the Cauchy-Schwarz inequality and (21)

$$\frac{\lambda \max_i |Z_{si}| \sum_{k=1}^M (1 - H_{kk}) |\nu_k|}{\|Z_s\|_N \left( \sum_{k=1}^M (1 - H_{kk})^2 \nu_k^2 \right)^{1/2}} \leq \frac{\lambda \max_i |Z_{si}| M^{1/2}}{\|Z_s\|_N} \leq \lambda C_Z^{1/2} M^{1/2} N^{-1/2} < \frac{1}{2}.$$

With this in mind since  $X_i$  are independent we have by the Taylor formula

$$\begin{aligned} \mathbf{E} \exp \left[ \frac{\lambda}{\sqrt{D(v)}} \sum_{k=1}^{\infty} v_k \eta_k \right] &= \prod_{i=1}^N \mathbf{E} \exp \left\{ \frac{\lambda Z_{si}}{\sqrt{D(v)}} \sum_{k=1}^M (1 - H_{kk}) \nu_k \psi_k(X_i) \right\} \\ &\leq (1 + \sqrt{e} \lambda^2 / 2) \leq \exp(\lambda^2). \end{aligned}$$

Hence using (33) and Lemma 3 we finish the proof.  $\square$

**Lemma 5** *Let  $\xi_i$  be i.i.d.  $\mathcal{N}(0, \sigma^2)$ . Then*

$$\mathbf{E} \left\{ \sum_{k=1}^{\infty} H_{kk} \left[ \left( \sum_{i=1}^N Z_{si} \psi_k(X_i) \right)^2 - \|Z_s\|_N^2 \right] \right\}^2 \leq C \log(n\#\mathcal{H}) \|Z_s\|_N^4 \mathbf{E} \operatorname{tr} H^2, \quad (34)$$

$$\mathbf{E} \left\{ \sum_{k=1}^{\infty} H_{kk} \left[ \left( \sum_{i=1}^N \xi_i \psi_k(X_i) \right)^2 - N \sigma^2 \right] \right\}^2 \leq C \log(n\#\mathcal{H}) N^2 \mathbf{E} \operatorname{tr} H^2, \quad (35)$$

$$\mathbf{E} \left\{ \sum_{k=1}^{\infty} H_{kk}^2 \left[ \left( \sum_{i=1}^N \xi_i \psi_k(X_i) \right)^2 - N \sigma^2 \right] \right\}^2 \leq C \log(n\#\mathcal{H}) N^2 \mathbf{E} \operatorname{tr} H^2, \quad (36)$$

$$\mathbf{E} \left\{ \sum_{k=1}^{\infty} H_{kk} \sum_{i=1}^N [\psi_k^2(X_i) - 1] \right\}^2 \leq C \log(n\#\mathcal{H}) N \mathbf{E} \operatorname{tr} H, \quad (37)$$

$$\mathbf{E} \left\{ \sum_{k=1}^{\infty} H_{kk}^2 \sum_{i=1}^N [\psi_k^2(X_i) - 1] \right\}^2 \leq C \log(n\#\mathcal{H}) N \mathbf{E} \operatorname{tr} H^2, \quad (38)$$

$$\mathbf{E} \left\{ \sum_{k=1}^{\infty} H_{kk} \sum_{i \neq j}^N \psi_k(X_j) \psi_k(X_i) \xi_i \xi_j \right\}^2 \leq C \log(n\#\mathcal{H}) N \mathbf{E} \operatorname{tr} H^2, \quad (39)$$

$$\mathbf{E} \left\{ \sum_{k=1}^{\infty} H_{kk} \sum_{j,i=1}^N [\psi_k(X_i) \psi_k(X_j) - \delta_{ij}] [Z_N^T (Z_N Z_N^T)^{-1} Z_N]_{ij} \right\}^2 \quad (40)$$

$$\leq C \log(n\#\mathcal{H}) \mathbf{E} \operatorname{tr} H^2.$$

**Proof.** We prove only the first inequality since the others can be checked in the similar way. Once again use Lemma 3. Putting

$$v_k = H_{kk}, \quad \eta_k = \left( \sum_{i=1}^N Z_{si} \psi_k(X_i) \right)^2 - \|Z_s\|_N^2$$

we have by (20 - 21) that for any given  $v$

$$D(v) = \mathbf{E} \left( \sum_{k=1}^{\infty} v_k \eta_k \right)^2 = 2[1 + o(1)] \operatorname{tr} H^2 \|Z_s\|_N^4.$$

Since  $\operatorname{tr} H^2 \leq n$  we choose  $S = n^2$ . Thus to complete the proof of the lemma it remains to check (30). Using the Cauchy-Schwarz inequality one obtains

$$\begin{aligned} & \mathbf{E} \exp \left\{ \frac{\lambda}{\sqrt{D(v)}} \sum_{k=1}^{\infty} v_k \eta_k \right\} \\ & \leq \mathbf{E}^{1/2} \exp \left\{ \frac{4\lambda}{\sqrt{D(v)}} \sum_{j < i}^N Z_{si} Z_{sj} \sum_{k=1}^{\infty} v_k \psi_k(X_i) \psi_k(X_j) \right\} \\ & \quad \times \mathbf{E}^{1/2} \exp \left\{ \frac{2\lambda}{\sqrt{D(v)}} \sum_{i=1}^N Z_{si}^2 \sum_{k=1}^{\infty} v_k [\psi_k^2(X_i) - 1] \right\}. \end{aligned} \quad (41)$$

Notice that in view of (20-21) one obtains for  $\lambda = \sqrt{\log(n^2 \# \mathcal{H})}$

$$\frac{\lambda |Z_{si} Z_{sj}|}{\sqrt{D(v)}} \left| \sum_{k=1}^{\infty} v_k \psi_k(X_i) \psi_k(X_j) \right| = o(1), \quad \frac{\lambda Z_{si}^2}{\sqrt{D(v)}} \left| \sum_{k=1}^{\infty} v_k [\psi_k^2(X_i) - 1] \right| = o(1).$$

Therefore we can use the Taylor formula to compute the right-hand side of (41). Hence acting in this way we arrive at

$$\mathbf{E} \exp \left\{ \frac{\lambda}{\sqrt{D(v)}} \sum_{k=1}^{\infty} v_k \eta_k \right\} \leq \exp(2\lambda^2).$$

Now (31) implies the inequality (34).  $\square$

**Lemma 6** *Let  $\eta_{ks} = N^{-1/2} \sum_{i=1}^N [\psi_k(X_i) \psi_l(X_i) - \delta_{kl}]$ . For any  $M > 1$  uniformly in  $\nu \in \mathbf{W}_0$*

$$\mathbf{E} \left| \sum_{k,l=0}^{\infty} \nu_k \nu_l (1 - H_{kk})(1 - H_{ll}) \eta_{ks} \right| \leq C M \log^{1/2}(M) \left[ \mathbf{E} \sum_{k=1}^{\infty} (1 - H_{kk})^2 \nu_k^2 + L M^{-2\beta} \right] \quad (42)$$

and

$$\mathbf{E} \sum_{k,s=1}^{\infty} H_{kk} H_{ss} \eta_{ks}^2 \leq C \log(n) \mathbf{E} \operatorname{tr}^2 H. \quad (43)$$

Let  $\zeta_{ks} = \sum_{i,j=1}^N \psi_k(X_i) \psi_s(X_j) [Z_N^T (Z_N Z_N^T)^{-1} Z_N]_{ij}$ . Then uniformly in  $\nu \in \mathbf{W}_0$

$$\mathbf{E} \left( \sum_{k,s=1}^{\infty} |\nu_s| H_{kk} \zeta_{ks} \right)^2 \leq C \log n \left( \sum_{s=1}^{\infty} |\nu_s| \right)^2 \mathbf{E} \operatorname{tr}^2 H + C L n^{-2\beta+1}. \quad (44)$$

**Proof.** Note that  $\psi_k(X_i)\psi_l(X_i) - \delta_{kl}$  are i.i.d. bounded random variables having zero mean. Therefore

$$\mathbf{P}\left\{|\eta_{ks}| \geq x\right\} \leq \exp(-Cx^2). \quad (45)$$

On the other hand one obtains by the Cauchy-Schwarz inequality

$$\begin{aligned} & \mathbf{E} \left| \sum_{k,l=0}^{\infty} \nu_k \nu_l (1 - H_{kk})(1 - H_{ll}) \eta_{kl} \right| \\ & \leq \left( \sum_{k=1}^M |\nu_k| |1 - H_{kk}| \right)^2 \max_{k,l \leq M} |\eta_{kl}| + L \left( \sum_{k,l > M} k^{-2\beta} l^{-2\beta} \mathbf{E} \eta_{kl}^2 \right)^{1/2} \\ & \leq MQ \mathbf{E} \sum_{k=1}^M \nu_k^2 (1 - H_{kk})^2 + CL \sum_{k,l \leq M} \mathbf{P}\left\{|\eta_{kl}| \geq Q\right\} + CLM^{-2\beta+1}. \end{aligned}$$

Putting  $Q = \sqrt{C' \log(M)}$ , where  $C'$  is a sufficiently large constant, we arrive with (45) at (42).

The second inequality (43) can be proved in the similar way. Using the Cauchy-Schwarz inequality, (20) and (45) we have

$$\begin{aligned} \mathbf{E} \sum_{k,s=1}^{\infty} H_{kk} H_{ss} \eta_{ks}^2 & \leq \mathbf{E} \max_{k,s \leq n} \eta_{ks}^2 \sum_{k,s=1}^n H_{kk} H_{ss} + \mathbf{E} \sum_{k,s > N} H_{kk} H_{ss} \eta_{ks}^2 \\ & \leq Q^2 \mathbf{E} \operatorname{tr}^2 H + n^2 \mathbf{P}\left\{\max_{k,s \leq n} |\eta_{ks}| > Q\right\} + \mathbf{E} \left( \sum_{k > n} H_{kk}^2 k^2 \right) \left( \sum_{k,s > n} k^{-2} s^{-2} \eta_{ks}^4 \right)^{1/2} \\ & \leq Q^2 \mathbf{E} \operatorname{tr}^2 H + n^4 \exp(-CQ^2) + C. \end{aligned}$$

We complete the proof of the lemma choosing  $Q = 2\sqrt{\log n/C}$ . The proof of (44) is similar and we omit it. ■

Now we are ready to evaluate the performance of the estimator  $\theta^*$ .

**Lemma 7** *Let*

$$\lim_{n \rightarrow \infty} \frac{N \log n}{n} = 0. \quad (46)$$

*Then uniformly in  $m \in \mathbf{W}_0$  as  $n \rightarrow \infty$*

$$\mathbf{E}(\theta^* - \theta)(\theta^* - \theta)^T = (ZZ^T)^{-1} \left[ \sigma^2 + (1 + o(1)) \mathbf{E} \left( \|(E - H^*)\nu\|^2 + \frac{\sigma^2}{n} \operatorname{tr} H^{*2} \right) \right]. \quad (47)$$

**Proof.** In order to compute the right-hand side of (26) we represent  $\Psi Z^T$ ,  $\Psi \xi$ ,  $Z \xi$  in the following form

$$\begin{aligned}\Psi Z^T &= \sum_{i=1}^N \psi_k(X_i) Z_{si} + \sum_{i=N+1}^n \psi_k(X_i) Z_{si} = \Psi Z_0^T + \Psi Z_1^T, \\ \Psi \xi &= \sum_{i=1}^N \psi_k(X_i) \xi_i + \sum_{i=N+1}^n \psi_k(X_i) \xi_i = \Psi \xi_0 + \Psi \xi_1, \\ Z \xi &= \sum_{i=1}^N Z_{ki} \xi_i + \sum_{i=N+1}^n Z_{ki} \xi_i = Z \xi_0 + Z \xi_1.\end{aligned}$$

and note that  $H^*$  does not depend on  $\Psi Z_1^T$ ,  $\Psi \xi_1$  and  $Z \xi_1$ . The dependence between  $H^*$  and  $\Psi Z_0^T$ ,  $\Psi \xi_0$ ,  $Z \xi_0$  can be evaluated by Lemmas 4 – 6. So we have for example by Lemma 4 and (46), (21)

$$\begin{aligned}\mathbf{E} [Z \Psi^T (E - H^*) \nu] [Z \Psi^T (E - H^*) \nu]^T &= \mathbf{E} [(Z \Psi_0^T + Z \Psi_1^T) (E - H^*) \nu] [(Z \Psi_0^T + Z \Psi_1^T) (E - H^*) \nu]^T \\ &= \mathbf{E} [Z \Psi_0^T (E - H^*) \nu] [Z \Psi_0^T (E - H^*) \nu]^T + \mathbf{E} [Z \Psi_1^T (E - H^*) \nu] [Z \Psi_1^T (E - H^*) \nu]^T \\ &\leq \left[ \sum_{i=N+1}^n Z_{ki} Z_{si} + O(E) \log n \sum_{s=1}^d \sum_{i=1}^N Z_{si}^2 \right] \mathbf{E} \sum_{k=1}^\infty (1 - H_{kk}^*)^2 \nu_k^2 + \frac{C \log n}{n} \sum_{s=1}^d \|Z_s\|_N^2 \\ &\leq \left[ Z Z^T + O(E) \frac{N \log n}{n} \sum_{s=1}^d \sum_{i=1}^n Z_{si}^2 \right] \mathbf{E} \sum_{k=1}^\infty (1 - H_{kk}^*)^2 \nu_k^2 + \frac{C \log n}{n} \sum_{s=1}^d \|Z_s\|_N^2 \\ &\leq Z Z^T [E + o(E)] \left[ \mathbf{E} \sum_{k=1}^\infty (1 - H_{kk}^*)^2 \nu_k^2 + \frac{o(1)}{n} \right].\end{aligned}$$

Similar arguments can be used in order to compute the remainder terms in the right-hand side of (26).  $\square$

**Proof of Theorem 3.** Denote

$$L_N[H] = \|Y - Q_N^T \hat{\mu}_N(H)\|_N^2 + 2\sigma^2 \text{tr} U_N(H) - (1 - N/n) \sigma^2 \text{tr} U_N^2(H).$$

In view of Lemma 7 it suffices to show that uniformly in  $H \in \mathcal{H}$

$$\mathbf{E} L_N[H] = [1 + o(1)] N \mathbf{E} \left[ \|(E - H) \nu\|^2 + \sigma^2 \text{tr} H^2 / n \right] + R, \quad (48)$$

where  $R$  is a constant which does not depend on  $H$ . Let  $S_N = Q_N Q_N^T + N H^{-1} (E - H) = A_N + B_N$ , with

$$A_N = \begin{pmatrix} Z_N Z_N^T & 0 \\ 0 & N H^{-1} \end{pmatrix}, \quad B_N = \begin{pmatrix} 0 & Z_N \Psi_N^T \\ \Psi_N Z_N^T & \Psi_N \Psi_N^T - N E \end{pmatrix}.$$



Then by Lemma 1 (see also (29)) we have

$$S_N^{-1} = \begin{pmatrix} (Z_N Z_N^T)^{-1} & 0 \\ 0 & N^{-1} H \end{pmatrix} - [E + o(E)] N^{-1} \begin{pmatrix} 0 & (Z_N Z_N^T)^{-1} Z_N \Psi_N^T H \\ H \Psi_N Z_N^T (Z_N Z_N^T)^{-1} & H(\Psi_N \Psi_N^T / N - E) H \end{pmatrix}. \quad (49)$$

Therefore

$$S_N^{-1} Q_N Q_N^T = \begin{pmatrix} E & 0 \\ 0 & H \end{pmatrix} - [E + o(E)] \begin{pmatrix} (Z Z^T)^{-1} Z \Psi_N^T H \Psi_N Z^T / N & -(Z Z^T)^{-1} Z \Psi_N^T (E - H) \\ 0 & H \Psi_N Z^T (Z Z^T)^{-1} Z \Psi_N^T / N \end{pmatrix}. \quad (50)$$

and

$$Q_N S_N^{-1} Q_N^T = U_N(H) = Z_N^T (Z_N Z_N^T)^{-1} Z_N + N^{-1} \Psi_N H \Psi_N^T + N^{-1} [E + o(E)] [Z_N^T (Z_N Z_N^T)^{-1} Z_N \Psi_N H \Psi^T + \Psi_N^T H \Psi_N Z_N^T (Z_N Z_N^T)^{-1} Z_N + \Psi_N^T H (\Psi_N \Psi_N^T / N - E) H \Psi_N]. \quad (51)$$

First we estimate the traces of  $U_N(H)$  and  $U_N^2(H)$ . By a simple algebra we obtain

$$N^{-1} \mathbf{E} \operatorname{tr} \Psi_N^T H \Psi_N = \mathbf{E} \left\{ \operatorname{tr} H + \frac{1}{N} \sum_{i=1}^N \sum_{k=1}^{\infty} H_{kk} [\psi_k^2(X_i) - 1] \right\}, \quad (52)$$

$$\begin{aligned} N^{-2} \mathbf{E} \operatorname{tr} \Psi_N^T H \Psi_N \Psi_N^T H \Psi_N &= N^{-1} \mathbf{E} \operatorname{tr} \Psi_N^T H^2 \Psi_N + N^{-1} \mathbf{E} \operatorname{tr} \Psi_N^T H (\Psi_N \Psi_N^T / N - E) H \Psi_N \\ &= (1 + N^{-1}) \mathbf{E} \operatorname{tr} H^2 + \mathbf{E} \frac{2}{N} \sum_{i=1}^N \sum_{k=1}^{\infty} H_{kk}^2 [\psi_k^2(X_i) - 1] \\ &\quad + \frac{1}{N} \sum_{k,s=1}^{\infty} H_{kk} H_{ss} \left[ \frac{1}{\sqrt{N}} \sum_{i=1}^N \psi_k(X_i) \psi_s(X_i) - \delta_{ks} \right]^2 \end{aligned} \quad (53)$$

and

$$\begin{aligned} N^{-1} \mathbf{E} \operatorname{tr} Z_N^T (Z_N Z_N^T)^{-1} Z_N \Psi_N^T H \Psi_N &= N^{-1} \operatorname{tr} Z_N^T (Z_N Z_N^T)^{-1} Z_N \mathbf{E} \operatorname{tr} H \\ &\quad + \frac{1}{N} \sum_{k=1}^{\infty} H_{kk} \sum_{j,i=1}^N [\psi_k(X_i) \psi_k(X_j) - \delta_{ij}] [Z_N^T (Z_N Z_N^T)^{-1} Z_N]_{ij}. \end{aligned} \quad (54)$$

Since  $\text{tr } Z_N^T(Z_N Z_N^T)^{-1} Z_N \leq C$  we obtain by (37), (40), (43) and (51-54)

$$\begin{aligned}\mathbf{E} \text{tr } U_N(H) &= [1 + o(1)] \mathbf{E} \text{tr } H + \text{tr } Z_N^T(Z_N Z_N^T)^{-1} Z_N, \\ \mathbf{E} \text{tr } U_N^2(H) &= [1 + o(1)] \mathbf{E} \text{tr } H^2 + \text{tr } Z_N^T(Z_N Z_N^T)^{-1} Z_N.\end{aligned}\quad (55)$$

Next notice

$$\begin{aligned}\|Y - Q_N^T \hat{\mu}_N(H)\|_N^2 &= \|(E - Q_N^T S_N^{-1} Q_N) \xi_N + (E - Q_N^T S_N^{-1} Q_N) Q_N^T \mu\|_N^2 \\ &= \|(E - Q_N^T S_N^{-1} Q_N) \xi_N\|_N^2 + \|Q_N^T (E - S_N^{-1} Q_N Q_N^T) \mu\|_N^2 \\ &\quad + 2 \xi_N^T (E - Q_N S_N^{-1} Q_N^T) Q_N^T (E - S_N^{-1} Q_N Q_N^T) \mu.\end{aligned}\quad (56)$$

The first term in the right-hand side of (56) is evaluated with (49) as

$$\begin{aligned}\mathbf{E} \|(E - Q_N^T S_N^{-1} Q_N) \xi_N\|_N^2 - N \sigma^2 &= -2 \mathbf{E} \xi_N^T Q_N^T S_N^{-1} Q_N \xi_N \\ &\quad + \mathbf{E} \xi_N^T Q_N S_N^{-1} Q_N^T Q_N^T S_N^{-1} Q_N \xi_N \\ &= -2 \mathbf{E} \xi_N^T Z_N^T (Z_N Z_N^T)^{-1} Z_N \xi_N - 2[1 + o(1)] N^{-1} \mathbf{E} \xi_N^T \Psi_N^T H \Psi_N \xi_N \\ &\quad + \mathbf{E} \xi_N^T [Z_N^T (Z_N Z_N^T)^{-1} Z_N + [1 + o(1)] N^{-1} \Psi_N^T H \Psi_N]^2 \xi_N.\end{aligned}\quad (57)$$

Using (36) we get

$$\begin{aligned}N^{-1} \mathbf{E} \xi_N^T \Psi_N^T H \Psi_N \xi_N &= \mathbf{E} \sum_{k=1}^{\infty} H_{kk} \frac{1}{N} \sum_{i,j=1}^N \psi_k(X_j) \psi_k(X_i) \xi_i \xi_j \\ &= \sigma^2 \mathbf{E} \sum_{k=1}^{\infty} H_{kk} + \mathbf{E} \sum_{k=1}^{\infty} H_{kk} \left[ \left( \frac{1}{\sqrt{N}} \sum_{i=1}^N \psi_k(X_i) \xi_i \right)^2 - \sigma^2 \right] = [1 + o(1)] \sigma^2 \mathbf{E} \sum_{k=1}^{\infty} H_{kk}.\end{aligned}\quad (58)$$

Next with (36), (43) we obtain

$$\begin{aligned}N^{-2} \mathbf{E} \xi_N^T (\Psi_N^T H \Psi_N)^2 \xi_N &= \mathbf{E} \text{tr } H^2 + \mathbf{E} \sum_{k=1}^{\infty} H_{kk}^2 \left[ \left( \frac{1}{\sqrt{N}} \sum_{i=1}^N \psi_k(X_i) \xi_i \right)^2 - \sigma^2 \right] \\ &\quad + \frac{1}{\sqrt{N}} \mathbf{E} \sum_{s,k=1}^{\infty} H_{kk} H_{ss} \left( \frac{1}{\sqrt{N}} \sum_{i=1}^N \psi_k(X_i) \xi_i \right) \left( \frac{1}{\sqrt{N}} \sum_{i=1}^N \psi_s(X_i) \xi_i \right) \\ &\quad \times \left( \frac{1}{\sqrt{N}} \sum_{i=1}^N [\psi_k(X_i) \psi_s(X_i) - \delta_{ks}] \right) = (1 + o(1)) \mathbf{E} \text{tr } H^2 \\ &\quad + \frac{O(1)}{\sqrt{N}} \mathbf{E} \left[ \sum_{s,k=1}^{\infty} H_{kk} \left( \frac{1}{\sqrt{N}} \sum_{i=1}^N \psi_k(X_i) \xi_i \right)^2 \right]^2 \\ &\quad + \frac{O(1)}{\sqrt{N}} \mathbf{E} \sum_{s,k=1}^{\infty} H_{kk} H_{ss} \left( \frac{1}{\sqrt{N}} \sum_{i=1}^N [\psi_k(X_i) \psi_s(X_i) - \delta_{ks}] \right)^2 = (1 + o(1)) \mathbf{E} \text{tr } H^2.\end{aligned}\quad (59)$$

Therefore noticing that  $\text{tr } Z_N^T(Z_N Z_N^T)^{-1} Z_N \leq C$  we arrive with (57)–(59) at

$$\mathbf{E} \|(E - Q_N^T S_N^{-1} Q_N) \xi_N\|_N^2 = N\sigma^2 + (1 + o(1)) \mathbf{E} (\text{tr } H^2 - 2 \text{tr } H) \quad (60)$$

$$- \text{tr } Z_N^T (Z_N Z_N^T)^{-1} Z_N. \quad (61)$$

Our next step is to estimate  $\mathbf{E} \|Q_N^T (E - S_N^{-1} Q_N Q_N^T) \mu\|_N^2$  in (56). Starting with the first order term (see (50)) we have by (42) with  $M = N/\log^{1/2+\delta}(n)$

$$\begin{aligned} \mathbf{E} \|\Psi^T(E - H)\nu\|_N^2 &= \mathbf{E} \sum_{i=1}^N \left[ \sum_{k=1}^{\infty} (1 - H_{kk}) \nu_k \psi_k(X_i) \right]^2 \\ &= N \mathbf{E} \sum_{k=1}^{\infty} (1 - H_{kk})^2 \nu_k^2 + \mathbf{E} \sum_{k,l=\beta}^{\infty} (1 - H_{kk}) \nu_k \nu_l (1 - H_{ll}) \sum_{i=1}^N [\psi_k(X_i) \psi_l(X_i) - \delta_{kl}] \\ &= [1 + o(1)] N \mathbf{E} \sum_{k=1}^{\infty} (1 - H_{kk})^2 \nu_k^2 + \log^{1+\delta}(n)/N. \end{aligned} \quad (62)$$

The remainder terms in  $\mathbf{E} \|Q_N^T (E - S_N^{-1} Q_N Q_N^T) \mu\|_N^2$  associated with the second order terms in (50) are evaluated by the similar arguments. We have by (34) (replacing there  $Z_{si}$  by  $[Z_N^T (Z_N Z_N^T)^{-1} Z_N]_{si}$  and by  $[Z_N \theta]_i$ )

$$\begin{aligned} &N^{-2} \mathbf{E} \|Z_N^T (Z_N Z_N^T)^{-1} Z_N \Psi_N^T H \Psi_N^T Z_N^T \theta\|_N^2 \\ &= N^{-2} \mathbf{E} \sum_{s=1}^N \left\{ \sum_{k=1}^{\infty} H_{kk} \left( \sum_{i=1}^N \psi_k(X_i) [Z_N^T (Z_N Z_N^T)^{-1} Z_N]_{si} \right) \left( \sum_{j=1}^N \psi_k(X_j) [Z_N \theta]_j \right) \right\}^2 \\ &\leq 4N^{-2} \mathbf{E} \sum_{s=1}^N \left\{ \sum_{k=1}^{\infty} H_{kk} \left( \sum_{i=1}^N \psi_k(X_i) [Z_N^T (Z_N Z_N^T)^{-1} Z_N]_{si} \right)^2 \right\}^2 \\ &\quad + 4N^{-2} \mathbf{E} \sum_{s=1}^N \left\{ \sum_{k=1}^{\infty} H_{kk} \left( \sum_{j=1}^N \psi_k(X_j) [Z_N \theta]_j \right)^2 \right\}^2 \\ &\leq C N^{-2} \log(n \# \mathcal{H}) \left\{ \|Z_N \theta\|^2 + \sum_{i,s=1}^N [Z_N^T (Z_N Z_N^T)^{-1} Z_N]_{si}^2 \right\} \mathbf{E} \text{tr}^2 H = o(1) \mathbf{E} \text{tr } H, \end{aligned}$$

$$\begin{aligned} &\mathbf{E} \|Z_N^T (Z_N Z_N^T)^{-1} Z_N \Psi_N^T (E - H) \nu\|_N^2 \\ &= \mathbf{E} \sum_{i=1}^N \left\{ \sum_{k=1}^{\infty} (1 - H_{kk}) \nu_k \left( \sum_{j=1}^N \psi_k(X_j) [Z_N^T (Z_N Z_N^T)^{-1} Z_N]_{ij} \right) \right\}^2 \\ &\leq \log(n \# \mathcal{H}) \sum_{i,j=1}^N [Z_N^T (Z_N Z_N^T)^{-1} Z_N]_{ij}^2 \left[ \mathbf{E} \sum_{k=1}^{\infty} (1 - H_{kk})^2 \nu_k^2 + \frac{C}{n} \right] \end{aligned}$$

$$= o(1)N \left[ \mathbf{E} \sum_{k=1}^{\infty} (1 - H_{kk})^2 \nu_k^2 + \frac{C}{n} \right]$$

and by (44)

$$\begin{aligned} & N^{-2} \mathbf{E} \left\| \Psi_N^T H \Psi_N Z_N^T (Z_N Z_N^T)^{-1} Z_N \Psi_N^T \nu \right\|_N^2 \\ &= N^{-2} \mathbf{E} \sum_{i=1}^N \left\{ \sum_{k=1}^{\infty} \sum_{s=1}^{\infty} H_{kk} \psi_k(X_i) \nu_s \sum_{l,m=1}^N [Z_N^T (Z_N Z_N^T)^{-1} Z_N]_{lm} \psi_k(X_l) \psi_s(X_m) \right\}^2 \\ &\leq C N^{-1} \mathbf{E} \left\{ \sum_{k=1}^{\infty} \sum_{s=1}^{\infty} H_{kk} |\nu_s| \sum_{l,m=1}^N [Z_N^T (Z_N Z_N^T)^{-1} Z_N]_{lm} \psi_k(X_l) \psi_s(X_m) \right\}^2 \\ &\leq C N^{-1} \log n \left( \sum_{s=1}^{\infty} |\nu_s| \right)^2 \mathbf{E} \operatorname{tr}^2 H + C n^{-1} N^{-1} = o(1) \mathbf{E} \operatorname{tr} H. \end{aligned}$$

Thus using the above inequalities and (62) we arrive at

$$\mathbf{E} \left\| Q_N^T (E - S_N^{-1} Q_N Q_N^T) \mu \right\|_N^2 = [1 + o(1)] N \mathbf{E} \sum_{k=1}^{\infty} (1 - H_{kk})^2 \nu_k^2 + o(1) \mathbf{E} \sum_{k=1}^{\infty} H_{kk}^2. \quad (63)$$

The interference term in (56) is evaluated by the similar way. So one gets

$$\mathbf{E} \left| \xi_N^T (E - Q_N S_N^{-1} Q_N^T) Q_N^T (E - S_N^{-1} Q_N Q_N^T) \mu \right| \leq o(1) \mathbf{E} \left[ N \sum_{k=1}^{\infty} (1 - H_{kk})^2 \nu_k^2 + \sum_{k=1}^{\infty} H_{kk}^2 \right].$$

The proof of (48) follows now from the above equation (56) and (55), (63). ■

## 4 Concluding remarks

In practical applications of statistical methods one always compromises between theoretical efficiency and numerical complexity. Partial linear models provide us with a good example of that. In the analysis of this model we have assumed that the smoothness  $\beta$  is known. This assumption is not crucial from the theoretical point of view since if  $\beta$  is unknown we can use the adaptation method based on unbiased risk estimation (see Golubev and Nussbaum (1992) or Kneip (1994)) with respect to the family of bases parameterized by the parameter  $\beta$ . We have intentionally considered the simplest case in order to avoid even more complicated proofs. On the other hand the standard practical recommendation  $\beta = 2$  seems to be a very reasonable tradeoff between optimal rates of convergence and complexity of computation.

Assuming that the smoothness is known we still encounter problems of computation the second order minimax estimator since in order to do that we need to:

- estimate the density  $q(x)$  of  $X_i$  if it is unknown
- compute the eigen functions and eigen values of (2)
- compute the empirical Fourier coefficients
- solve the minimization problem.

The first three steps require so many numerical resources that the exact implementation of the second order minimax estimator becomes almost impractical. We have already mentioned that one can avoid all these problems organizing the computation in the time domain. That means that we compute the spline estimator defined by (17). Its computation can be reduced to manipulations with band matrices and therefore can be done very fast (see Green and Silverman (1994)). This method of estimation is not optimal but we shall see that it is nearly optimal. In order to clarify this point let us look at the second order minimax efficiency  $EF(\beta)$  defined by

$$EF(\beta) = \lim_{n \rightarrow \infty} \frac{\sup_{\nu \in V} \inf_{H \in \mathcal{H}_{min}} IMSE[H, \nu]}{\sup_{\nu \in V} \inf_{H \in \mathcal{H}_{spl}} IMSE[H, \nu]}, \quad (64)$$

where

$$\begin{aligned} V &= \left\{ \nu_k : \sum_{k=1}^{\infty} \nu_k^2 \lambda_k \leq 1 \right\}, \\ \mathcal{H}_{min} &= \left\{ H : H_{kk} = [1 - \mu \lambda_k]_+, \mu > 0 \right\}, \\ \mathcal{H}_{spl} &= \left\{ H : H_{kk} = (1 + \mu \lambda_k)^{-1}, \mu > 0 \right\}. \end{aligned}$$

The motivation of (64) comes from Theorem 3, which shows how the performance of the estimator depends on the set  $\mathcal{H}$ . In particular, if one uses splines then  $\mathcal{H} = \mathcal{H}_{spl}$ ; for the second order minimax estimator one has to take  $\mathcal{H} = \mathcal{H}_{min}$ . The value of  $EF(\beta)$  can be computed analytically but it is more instructive to look at the plot presented on Figure 1. For comparison we placed here the plot of the minimax efficiency of the projection method. We see that even for  $\beta = 2$  minimax efficiency of the spline method is close to 0.92. This fact together with fast algorithms make splines very a powerful numerical tool for practical applications in partial linear models.




Figure 1: Minimax efficiency as function of the smoothness  $\beta$  (solid line: minimax efficiency of the spline method, dash line: minimax efficiency of the projection estimator).

## References

- [1] Akaike, H. (1973) Information theory and an extension of the maximum likelihood principle. In *Proceedings 2nd International Symposium on Information Theory*, P. N. Petrov and F. Csaki (Eds.) Akademia Kiado, Budapest, 267-281.
- [2] Barron, A., Birgé, L., Massart, P. (1999). Risk bounds for model selection via penalization. *Probab. Theory and Related Fields* **113** 301–413.
- [3] Bhattacharia P.K. and Zhao, P.-L. (1997). Semiparametric inference in a partial linear models. *Ann. Statist.* **25** 244–262.
- [4] Chen, H. (1998). Convergence rates for parametric components in a partly linear model. *Ann. Statist.* **16** 136–146.
- [5] Golubev, G., Härdle, W. (2000). On the second order minimax estimation in partial linear models. *Math. Methods Statist.* **9**, No 2, 160–175.
- [6] Golubev G., Levit B. (1996). Distribution function estimation: adaptive smoothing. *Math. Methods Statist.* **5** 383–403.

- [7] Golubev, G., Nussbaum, M. (1992). An adaptive spline estimates in nonparametric regression model. *Theory of Probab. Appl.* **3** 553–560. (In Russian)
- [8] Green, P. J., Silverman, B. W. (1994). *Nonparametric regression and generalized linear models. A roughness penalty approach*. Chapman & Hall, London.
- [9] Härdle, W., Liang, H., Gao, J. (2000). *Partially linear models*. Physica Verlag & Springer Verlag Germany.
- [10] Efroimovich, S.Y. and Pinsker, M.S. (1984). A learning algorithm for nonparametric filtering. *Automat. i Telemekh.* **11** 58–65. (In Russian).
- [11] Heckman, N.E. (1986). Spline smoothing in a partly linear models. *Journal of the Royal Statistical Society, Series B*, **48** 244–248.
- [12] Kneip, A. (1994). Ordered linear smoothers. *Ann. Statist.* **22**, 835–866.
- [13] Lepski, O. V. (1991). Asymptotically minimax adaptive estimation I: upper bounds. Optimal adaptive estimates. *Theory Probab. Appl.* **37** 682–697.
- [14] Lepski, O. V. (1992). On problems of adaptive estimation in white Gaussian noise. In *Topics in Nonparametric Estimation. Advances in Soviet Math.*, Khasminskii R. Z. ed. Amer. Math. Soc., Providence, R. I. **12** 87–106.
- [15] Lepski, O. V., Spokoiny, V. G. (1997). Optimal pointwise adaptive methods in nonparametric estimation. *Ann. Statist.* **25** 2512–2546.
- [16] Mallows, C. L. (1973). Some comments on  $C_p$ . *Technometrics*, **15**, 661–675.
- [17] Mammen, E., Van de Geer, S. (1997). Penalized quasi-likelihood estimation in partial linear models. *Annals Statist.* **25**, 1014–1035.
- [18] Nemirovskii, A. (1998). *Topics in Nonparametric Statistics*. Ecole d’été de probabilités de St. Flour. Lecture Notes.
- [19] Pinsker, M. S. (1980). Optimal filtering of square integrable signals in Gaussian in Gaussian noise. *Problems Inform. Transmission*, **16** 120–133.
- [20] Robinson, P.M. (1987). Asymptotically efficient estimation in the presence of heteroskedasticity of unknown form. *Econometrica*, **55** 875–891.
- [21] Rice, J.A. (1986). Convergence rates for partially splined models. *Statistics and Probability Letters*, **4** 203–208.

- [22] Schimek, M. G. (2000). Estimation and inference in partially linear models with splines. *Journal of Statistical Planning & Inferences*, **91**, 525–540.
- [23] Speckman, P. (1985). Spline smoothing and optimal rates of convergence in non-parametric regression models. *Ann. Statist.* **13** 970–983.
- [24] Speckman, P. (1988). Kernel smoothing in partial linear models. *Journal of Royal Statistical Society, Series B*, **50** 413–416.
- [25] Tsybakov, A. B. (1998). Pointwise and sup-norm sharp adaptive estimation of functions on the Sobolev classes. *Ann. Statist.* **26** 2420–2469.
- [26] Tikhomirov, V. (1986). *Fundamental principles of the theory of extremal problems* John Wiley & Sons.
- [27] Utreras, F. (1980). Sur le choix des paramètres d’ajustement dans le lissage par fonction spline. *Numer. Math.* **34**, 15–28.
- [28] Wahba, G. (1990). *Spline Models for Observational Data*. *S.I.A.M.*, Philadelphia.