

Seierstad, Atle

Working Paper

Conditions implying the vanishing of the Hamiltonian at the infinite horizon in optimal control problems

Memorandum, No. 2002,22

Provided in Cooperation with:

Department of Economics, University of Oslo

Suggested Citation: Seierstad, Atle (2002) : Conditions implying the vanishing of the Hamiltonian at the infinite horizon in optimal control problems, Memorandum, No. 2002,22, University of Oslo, Department of Economics, Oslo

This Version is available at:

<https://hdl.handle.net/10419/63014>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.

MEMORANDUM

No 22/2002

Conditions implying the vanishing of the Hamiltonian at the infinite horizon in optimal control problems

*By
Atle Seierstad*

ISSN: 0801-1117

Department of Economics
University of Oslo

This series is published by the
University of Oslo
Department of Economics

P. O.Box 1095 Blindern
N-0317 OSLO Norway
Telephone: + 47 22855127
Fax: + 47 22855035
Internet: <http://www.oekonomi.uio.no/>
e-mail: econdep@econ.uio.no

In co-operation with
**The Frisch Centre for Economic
Research**

Gaustadalleén 21
N-0371 OSLO Norway
Telephone: +47 22 95 88 20
Fax: +47 22 95 88 25
Internet: <http://www.frisch.uio.no/>
e-mail: frisch@frisch.uio.no

List of the last 10 Memoranda:

No 21	Morten Søberg The Duhem-Quine thesis and experimental economics: A reinterpretation. 22 pp.
No 20	Erling Barth, Bernt Bratsberg and Oddbjørn Raaum Local Unemployment and the Relative Wages of Immigrants: Evidence from the Current Population Surveys. 53 pp.
No 19	Erling Barth, Bernt Bratsberg and Oddbjørn Raaum Local Unemployment and the Earnings Assimilation of Immigrants in Norway. 46 pp.
No 18	Gunnar Bårdsen, Eilev S. Jansen and Ragnar Nymoen Testing the New Keynesian Phillips curve. 38 pp.
No 17	Morten Søberg Voting rules and endogenous trading institutions: An experimental study. 36 pp.
No 16	Gabriela Mundaca A Drift of the "Drift Adjustment Method". 35 pp.
No 15	Oddbjørn Raaum, Hege Torp and Tao Zhang Do individual programme effects exceed the costs? Norwegian evidence on long run effects of labour market training. 60 pp.
No 14	Oddbjørn Raaum, Hege Torp and Tao Zhang Business cycles and the impact of labour market programmes. 52 pp.
No 13	Geir B. Asheim, Anne Wenche Emblem and Tore Nilssen Deductibles in Health Insurances: Pay or Pain? 15 pp.
No 12	Oddbjørn Raaum and Knut Røed Do Business Cycle Conditions at the Time of Labour Market Entry Affect Future Unemployment?. 22 pp.

A complete list of this memo-series is available in a PDF® format at:
<http://www.oekonomi.uio.no/memo/>

Conditions implying the vanishing of the Hamiltonian at the infinite horizon in optimal control problems.

by

Atle Seierstad, University of Oslo

Abstract. In an infinite horizon optimal control problem, the Hamiltonian vanishes at the infinite horizon, when the differential equation is autonomous. The integrand in the integral criterion may contain the time explicitly, but it has to satisfy certain integrability conditions. A generalization of Michels (1982) result is obtained.

Introduction. Michel (1982) proved that the Hamiltonian vanishes when the time goes to infinity in problems where the differential equation is autonomous and the criterion is of the form $\int_0^\infty e^{rt} g(x, u) dt$. Below, a generalization of this result to the case where the criterion is of the form $\int_0^\infty f_0(t, x, u) dt$ is obtained, using a slightly modified proof.

Results. Consider the following problem.

$$\max_{u(\cdot)} \int_0^\infty f_0(t, x, u) dt, \tag{1}$$

subject to

$$\dot{x} = f(x, u) \text{ v.e.}, \quad u \in U, \quad x(0) = x_0, \tag{2}$$

$$\begin{aligned} \lim_{t \rightarrow \infty} x_i(t) &= x^i, & i &= 1, \dots, m' \\ \liminf_{t \rightarrow \infty} x_i(t) &\geq x^i, & i &= m' + 1, \dots, m. \end{aligned} \tag{3}$$

Here, U is a given set in \mathbb{R}^m , $f_i : \mathbb{R}^n \rightarrow \mathbb{R}$, $i = 0, \dots, n$, and f_i and $\partial f_i / \partial x$, $i = 0, \dots, n$, are continuous, as well as $\partial f_0 / \partial t$, and *v.e.* (virtually everywhere) means everywhere, except at a countable number of points.

Assume that $(x^*(t), u^*(t))$ is optimal among all pairs $(x(\cdot), u(\cdot))$ satisfying (2), (3), and for which the integral in (1) is convergent, with $u(\cdot)$ piecewise continuous on each finite interval. Assume also that $\int_0^\infty |f_0(t + \delta, x^*(t), u^*(t))| dt$ exists for all δ in an interval $(-\varepsilon, \varepsilon)$, and that, for some piecewise continuous function $\alpha(t)$, $|\partial f_0(t + \delta, x^*(t), u^*(t)) / \partial t| \leq \alpha(t)$ for all $\delta \in (-\varepsilon, \varepsilon)$, $t \in [0, \infty)$ and $\int_0^\infty \alpha(t) dt < \infty$. Then the following theorem holds:

Theorem. There exists a pair $(p_0, p(t))$, $p_0 \geq 0$, $(p_0, p(t)) \neq 0$ for all t , such that $p(t)$ is a continuous solution of the adjoint equation

$$\dot{p} = -H_x(t, x^*(t), u^*(t), p(t)) \text{ v.e.}, \quad (4)$$

where $H = p_0 f_0(t, x, u) + p(t) f(x, u)$. Moreover,

$$\max_{u \in U} H(t, x^*(t), u, p(t)) = H(t, x^*(t), u^*(t), p(t)) \text{ v.e.} \quad (5)$$

Finally,

$$\lim_{t \rightarrow \infty} H(t, x^*(t), u^*(t), p(t)) = 0. \quad (6)$$

Proof. Let S be a given positive number at which $u^*(\cdot)$ is continuous. Consider the auxiliary problem

$$\max_{u(\cdot), T > 0} \left\{ \int_0^T f_0(t, x, u) dt + g(T) \right\}, \quad (7)$$

subject to (2) and $x(T) = x^*(S)$, where $g(T) = \int_T^\infty f_0(t, x^*(t - (T - S)), u^*(t - (T - S))) dt$. Let $(x(\cdot), u(\cdot))$, defined on $[0, T]$, be admissible in the auxiliary problem (i.e. satisfying (2) and $x(T) = x^*(S)$). Extend $(x(\cdot), u(\cdot))$ to all $[0, \infty)$, by letting $x(t) = x^*(t - (T - S))$, $u(\cdot) = u^*(t - (T - S))$ for $t > T$. An admissible pair in the original problem is then obtained (i.e. satisfying (2),(3)), hence $\int_0^S f_0(t, x^*(t), u^*(t)) dt + g(S) = \int_0^S f_0(t, x^*(t), u^*(t)) dt + \int_S^\infty f_0(t, x^*(t), u^*(t)) dt = \int_0^\infty f_0(t, x^*(t), u^*(t)) dt \geq \int_0^\infty f_0(t, x(t), u(t)) dt = \int_0^T f_0(t, x(t), u(t)) dt + \int_T^\infty f_0(t, x(t), u(t)) dt = \int_0^T f_0(t, x(t), u(t)) dt + g(T)$. Thus, $(x^*(\cdot), u^*(\cdot))$, restricted to $[0, S]$, is optimal in the auxiliary problem. Now, $g(T) = \int_S^\infty f_0(s + T - S, x^*(s), u^*(s)) ds$. Hence, by the standard finite horizon maximum principle, a pair $(p_0^S, p^S(\cdot))$ exists, such that $|(p_0^S, p^S(0))| = 1$, $p_0^S \geq 0$, (4) and (5) are satisfied on $[0, S]$ and $[H(T, x^*(T), u^*(T), p(T)) + p_0^S dG/dT]_{T=S} = 0$. (The continuous differentiability of $G(T)$ near S follows from the existence of $\alpha(\cdot)$.) Let S_k be an increasing sequence converging to ∞ , such that $(p_0^{S_k}, p^{S_k}(0))$ is convergent. Then for each t , $p^{S_k}(t)$ is convergent, and $p_0 := \lim_k p_0^{S_k}$ and $p(t) := \lim_k p^{S_k}(t)$ satisfies $(p_0, p(0)) = 1$ and (4) and (5) on $[0, \infty)$. Finally, as $(d/dt) H(t, x^*(t), u^*(t), p^S(t)) = [(\partial/\partial t) H(t, x, u, p)]_{(x,u,p)=(x^*(t),u^*(t),p^S(t))} = p_0^S f_{0t}(t, x^*(t), u^*(t))$ and $H(S, x^*(S), u^*(S), p^S(S)) = -p_0^S [dG(T)/dT]_{T=S}$, then $H(S, x^*(S), u^*(S), p^S(S)) - H(t, x^*(t), u^*(t), p^S(t)) = \int_t^S p_0^S f_{0t}(t, x^*(t), u^*(t)) dt$ and $H(t, x^*(t), u^*(t), p^S(t)) = -\int_t^\infty p_0^S f_{0t}(t, x^*(t), u^*(t)) dt$. Letting $S = S_k$ and $k \rightarrow \infty$, gives, for each t , $H(t, x^*(t), u^*(t), p(t)) = -\int_t^\infty p_0 f_{0t}(t, x^*(t), u^*(t)) dt$. Then the existence of $\alpha(\cdot)$ yields (6).

Note: Statement and/or proofs of the finite horizon control results applied above can be

found in the references number 2, 3 and 4 below.

References:

1. Michel, Ph. (1982), On the transversality condition in infinite horizon optimal problems., *Econometrica*, Vol. 50. No 4, 975-985.

2. Neustadt, L.W. (1976), *Optimization, A Theory of Necessary Conditions*, Princeton University Press, New Jersey.

3. Pontryagin, L.S. et. al. (1962) *The mathematical Theory of Optimal Processes*, Interscience, New York.

4. Seierstad, A, and K. Sydsaeter. (1987), *Optimal Control Theory with Economic Applications*, North-Holland, Amsterdam.

5. Ye, J.J, (1993), Nonsmooth maximum principles for infinite-horizon problems, *Journal of Optimization Theory and Applications*, Vol 76, No. 3, 485-500.