

Hecq, Alain; Palm, Franz; Urbain, Jean-Pierre

Working Paper

Testing for Common Cyclical Features in Var Models with Cointegration

CESifo Working Paper, No. 451

Provided in Cooperation with:

Ifo Institute – Leibniz Institute for Economic Research at the University of Munich

Suggested Citation: Hecq, Alain; Palm, Franz; Urbain, Jean-Pierre (2001) : Testing for Common Cyclical Features in Var Models with Cointegration, CESifo Working Paper, No. 451, Center for Economic Studies and ifo Institute (CESifo), Munich

This Version is available at:

<https://hdl.handle.net/10419/75715>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.



Working Papers

TESTING FOR COMMON CYCLICAL FEATURES IN VAR MODELS WITH COINTEGRATION

Alain Hecq
Franz C. Palm
Jean-Pierre Urbain*

CESifo Working Paper No. 451

April 2001

CESifo

Center for Economic Studies & Ifo Institute for Economic Research
Poschingerstr. 5, 81679 Munich, Germany

Tel.: +49 (89) 9224-1410

Fax: +49 (89) 9224-1409

e-mail: office@CESifo.de



An electronic version of the paper may be downloaded

- from the SSRN website: www.SSRN.com
- from the CESifo website: www.CESifo.de

* The authors want to thank Peter Boswijk, Jörg Breitung, Bertrand Candelon, Niels Haldrup, Cheng Hsiao, Horst Zank, an associate editor and two anonymous referees for very useful comments on a previous draft of this paper. The usual disclaimer applies.

TESTING FOR COMMON CYCLICAL FEATURES IN VAR MODELS WITH COINTEGRATION

Abstract

We consider VAR models for variables exhibiting cointegration and common cyclical features. While the presence of cointegration reduces the rank of the long-run multiplier matrix, other types of common features lead to rank reduction of the short-run dynamics. We distinguish between strong and weak form reduced rank structures. Strong form reduced rank structures analyzed by Engle and Kozicki (1993) arise when a linear combination of the first differenced variables in a cointegrated VAR is white noise whereas in the presence of a weak form reduced rank structure, linear combinations of the first differenced variables corrected for the long-run effects are white noise. The weak form has an interest in its own. For instance, it is a necessary condition for the existence of first order codependent cycles in a VAR(2). Also, it is a necessary condition for the strong form. We also consider the mixed form which combines strong and weak forms. We discuss the model selection issues which arise from this distinction and propose a simple approach to testing for these structures using a sequence of likelihood ratio test statistics. The finite sample behavior of the sequential approach is analyzed in a Monte Carlo experiment. Finally, we illustrate the relevance of the different forms of reduced ranks with an empirical analysis of US business fluctuations over the period 1954-1996.

JEL Classification: C32

Keywords: Serial correlation common features, reduced rank structure, cointegration

Alain Hecq
Maastricht University
Fac. of Economics & Business Admin.
6200 MD Maastricht
The Netherlands

<i>Franz C. Palm</i>	<i>Jean-Pierre Urbain</i>
<i>Maastricht University</i>	<i>Maastricht University</i>
<i>Fac. of Economics & Business Admin.</i>	<i>Fac. of Economics & Business Admin.</i>
<i>6200 MD Maastricht</i>	<i>6200 MD Maastricht</i>
<i>The Netherlands</i>	<i>The Netherlands</i>
<i>F.Palm@ke.unimaas.nl</i>	

1 Introduction and Motivation

The interest in comovements between economic variables leading to common cyclical features has arisen for instance because economic theory predicts such comovements and many economic variables exhibit strong correlation at various frequencies. The vast literature on cointegration has focussed on long-run comovements. More recently, some authors have analyzed the existence of short-run comovements between stationary time series or between first differenced cointegrated $I(1)$ series (see Engle and Kozicki 1993; Gouriéroux and Peaucelle, 1989; Tiao and Tsay, 1989). Among these approaches, the concept of serial correlation common features (SCCF hereafter) introduced by Engle and Kozicki (1993) appears to be useful. It means that stationary time series move together in a way such that there exist linear combinations of these variables which yield white noise processes. In general, imposing these common features restrictions when they are appropriate will induce an increase in estimation efficiency (Lütkepohl, 1991) and accuracy of forecasts (Vahid and Issler, 1999). The associated cofeature vectors measure the intensity of short-run relationships between economic variables and they often have a straightforward economic interpretation. Under these restrictions a set of time series can be decomposed into their permanent and transitory components (see *inter alia* Vahid and Engle, 1993). Notice however that in these decompositions the number of common features and cointegrating vectors is assumed to be equal to (seldomly smaller than) the number of variables.¹

The aim of this paper is to analyze common cyclical features in relation with cointegration. The strong assumption that some linear combination of the first differences of the variables in the model is white noise will be called a strong form reduced rank structure (SF). It corresponds to the case of serial correlation common features of the variables in first differences and assumes that the left null spaces of the short-run dynamic matrices and cointegrating matrix overlap. Of course, in line with other authors, when SCCF appears to be too strong, one could test for the existence of cofeatures in the form of linear combinations of the variables differenced once, that are not white noise but have lower order dynamics than the individual variables. Tiao and Tsay (1989) for example, study this type of structure in a multivariate ARMA model and they call it scalar component model (SCM). We consider a natural weaker alternative assumption under which the common cyclical part is reduced to a white noise

¹To avoid confusion, it should be noticed from the outset that the term common cyclical features refers to a particular type of commonality leading to specific reduced rank structures. This concept should not be confused with the concept of cycle used in business cycle analyses (see the discussion in Cubadda, 1999). On the other hand, the concept of common cycles (in contrast to common cyclical features) refers to the common transitory component in particular permanent-transitory decompositions (see Vahid and Engle, 1993; Hecq, Palm and Urbain, 2000).

by taking a linear combination of the variables in the first differences corrected for long-run effects. This case will be termed weak form reduced rank structures (WF). The WF is attractive as it allows for different common factors generating respectively the long-run and short-run dynamics of economic variables. It is a necessary condition for the existence of first order codependent cycles in a VAR(2) as studied by Vahid and Engle (1997). As it is also a necessary condition for the SCCF, it is a natural hypothesis to be tested in sequential model specification.

Our framework is similar to that of Vahid and Engle (1993), but less restrictive as we explicitly consider the WF, implying linear combinations of the first differenced $I(1)$ variables to be predictable at low frequencies. In the presence of WF only, the lower bound to the number of common cycles is one whereas under SF, there have to be at least r common cycles in the system, with r being equal to the cointegration rank. Notice that Reinsel and Ahn (1992) briefly discuss a form similar to the WF. In general, they impose a nesting structure on the null spaces of the model dynamics. They do not discuss all the implications for the admissible number of common features. We study both the WF and the SF, taking into account the implications of the WF for the SF in modeling. Thereby, we do not impose a nesting structure on the null spaces of the model dynamics.

The paper is organized as follows. In Section 2 we present different forms of reduced rank structures that arise in empirical work. We focus on the partially non-stationary vector autoregression that will be reparametrized as an Vector Error Correction Model (VECM). The relationships between the strong and weak form reduced rank structures will be analyzed. The mixed form (MF) combining SF and WF will also be considered. Our model representation follows the lines of Ahn (1997) and Reinsel and Ahn (1992) but focusses on the constraints between the number of cointegrating and common cyclical feature vectors. Section 3 presents simple statistical procedures based on a two-step canonical correlation and maximum likelihood analyzes that allow to test various kinds of reduced rank structures, in particular to check whether short and long-run matrices have a common left null space. In Section 4, we study the small sample behavior of common feature tests using Monte Carlo simulations. We show why the number of common feature vectors can be artificially bounded by a wrong assumption about the nature of the reduced rank structure. We present a testing strategy that allows us to study cointegration and other common features of unknown order in an integrated framework. Finally, Section 5 illustrates the relevance of different forms of reduced structures, in particular of the WF for macroeconomic applications. It also demonstrates the use of the tests in the search for long-run and short-run relationships among real consumption, investment and gross domestic product in the US, in the period 1954-1996. A

final section concludes.

2 Reduced rank structures

Let us consider a Gaussian Vector Autoregression of finite order p (VAR(p)) model for an n -vector time series $\{y_t, t = 1, \dots, T\}$:

$$y_t = \sum_{i=1}^p \Phi_i y_{t-i} + \varepsilon_t, \quad t = 1, \dots, T, \quad (1)$$

for fixed values of y_{-p+1}, \dots, y_0 and where ε_t is a n -dimensional homoskedastic Gaussian mean innovation process relative to $\mathfrak{F}_t = \{y_{t-1}, y_{t-2}, \dots, y_1\}$ with nonsingular covariance matrix Ω . Let L denote the lag operator and define $\Phi(L) = I_n - \sum_{i=1}^p \Phi_i L^i$. We make the following assumption

Assumption 1 (Cointegration): *In the VAR model (1), we assume that*

1. $\text{rank}(\Phi(1)) = r, 0 < r < n$, so that $\Phi(1)$ can be expressed as $\Phi(1) = -\alpha\beta'$, with α and β both $(n \times r)$ matrices of full column rank r ;
2. the characteristic equation $|\Phi(\xi)| = 0$ has $n - r$ roots equal to 1 and all other roots outside the unit circle.

Assumption 1 implies (see Johansen, 1995) that the process y_t is cointegrated of order (1,1). The columns of β span the space of cointegrating vectors, and the elements of α are the corresponding adjustment coefficients or factor loadings. Decomposing the matrix lag polynomial $\Phi(L) = \Phi(1)L + \Phi^*(L)(1 - L)$, and defining $\Delta = (1 - L)$, we obtain the vector error correction model:

$$\Delta y_t = \alpha\beta' y_{t-1} + \sum_{j=1}^{p-1} \Phi_j^* \Delta y_{t-j} + \varepsilon_t, \quad t = 1, \dots, T, \quad (2)$$

where $\Phi_0^* = I_n$, $\Phi_j^* = -\sum_{k=j+1}^p \Phi_k$ ($j = 1, \dots, p-1$). Note that for notational convenience, deterministic terms (constants, trends, ...) are omitted at this level of presentation. With the exception of some simulation results in Section 4, throughout this paper we will also assume that p is known. Serial correlation common feature (see Engle and Kozicki, 1993) holds for the VECM (2), if there exists a $(n \times s)$ matrix $\tilde{\beta}$, whose columns span the cofeature space, such that $\tilde{\beta}' \Delta y_t = \tilde{\beta}' \varepsilon_t$ is a s -dimensional vector mean innovation process with respect to the information available at time t , \mathfrak{F}_t .

Consequently, serial correlation common features arise if there exists a cofeature matrix $\tilde{\beta}'$ such that the following two conditions are satisfied:

$$\textbf{Assumption 2:} \quad \tilde{\beta}' \Phi_j^* = 0_{(s \times n)}, \quad j = 1 \dots p-1 \quad (3)$$

$$\textbf{Assumption 3:} \quad \tilde{\beta}' \Phi(1) = -\tilde{\beta}' \alpha \beta' = 0_{(s \times n)} \quad (4)$$

Assumption 2 implies that $\tilde{\beta}'$ must lie in the intersection of the left null spaces of the matrices describing the short-run dynamics. Given that $\Phi_j^* = -\sum_{k=j+1}^p \Phi_k$, $j = 1, \dots, p-1$ and $\Phi_p^* = -\Phi(1) = -(I_n - \sum_{j=1}^p \Phi_j)$, Assumption 3 implies that $\tilde{\beta}'(I_n - \Phi_1) = 0_{(s \times n)}$, e.g. Φ_1 must have eigenvalues equal to one with multiplicity s and the corresponding eigenvectors must lie in the intersection of the left null spaces of the Φ_j^* matrices. Note that if the ranges of the Φ_j^* 's matrices are nested, i.e. if $\text{range}(\Phi_{j+1}^*) \subseteq \text{range}(\Phi_j^*)$, a nested reduced rank structure arises (see e.g. Ahn and Reinsel, 1988). We consider the restrictions implied by (3) or by (3) and (4) without imposing further nesting of the ranges of the Φ_j^* 's. This leads us to distinguish the following two concepts:

Definition 1 (Strong Form Reduced Rank Structure): *If in addition to Assumption 1 (cointegration) both Assumptions 2 and 3 hold, the implied reduced rank structure of the VECM (2) will be labelled a strong form reduced rank structure (SF). Under SF, there exists a $(n \times s)$ matrix $\tilde{\beta}$, whose columns span the cofeature space, such that $\tilde{\beta}' \Delta y_t = \tilde{\beta}' \varepsilon_t$ is a s -dimensional vector mean innovation process with respect to \mathfrak{S}_t .*

Definition 2 (Weak Form Reduced Rank Structure): *If in addition to Assumption 1 (cointegration) only Assumption 2 holds, the implied reduced rank structure of the VECM (2) will be labelled a weak form reduced rank structure (WF). Under WF, there exists a $(n \times s)$ matrix $\tilde{\beta}$, whose columns span the cofeature space, such that $\tilde{\beta}'(\Delta y_t - \alpha \beta' y_{t-1}) = \tilde{\beta}' \varepsilon_t$ is a s -dimensional vector mean innovation process with respect to \mathfrak{S}_t .*

Remark (a) The SF is usually considered in the literature (see inter alia Engle and Kozicki, 1993, Vahid and Engle, 1993 among others). It leads to serial correlation common features (SCCF). We however prefer to use the concept of SF in order to enable a formal comparison with the WF and to highlight the fact that the concept of SCCF generally applies to stationary vector processes irrespective of the presence or absence of cointegration. Under the SF, we may define a $(n(p-1) + r) \times 1$ vector $X_{t-1}^* = [\Delta y'_{t-1}, \dots, \Delta y'_{t-p+1}, y'_{t-1} \beta']'$ and a $n \times (n(p-1) + r)$ matrix $\Phi^* = [\Phi_1^*, \dots, \Phi_{p-1}^*, \alpha]$, so that (2) is written as

$$\Delta y_t = \Phi^* X_{t-1}^* + \varepsilon_t, \quad t = 1, \dots, T. \quad (5)$$

Under the assumption of a SF, Φ^* is of reduced rank $n - s$ and can be written as $\Phi^* = A^*[C_1^*, \dots, C_{p-1}^*, C_p^*] = A^*C^*$, where A^* is $n \times (n - s)$ full column rank matrix and C^* is $(n - s) \times (n(p - 1) + r)$ and $\tilde{\beta}' A^* C^* X_{t-1}^* = 0$, e.g. $\tilde{\beta} \in sp(A_\perp^*)$ where A_\perp^* is the orthogonal complement² of A^* . Consequently, as pointed out by Vahid and Engle (1993), in a n -dimensional $I(1)$ vector process y_t with $r < n$ cointegrating vectors, if the elements of y_t have common cyclical features (given by $f_t = C^* X_{t-1}^*$) there can be at most $n - r$ linearly independent cofeature vectors that eliminate the common cyclical features since the cofeature matrix must³ lie in $sp(\alpha_\perp)$. The SF implies that $s \leq n - r$ and that the common dynamic factors f_t consist of linear combinations of the elements of X_{t-1}^* . The implications of the SF can be stated more formally as:

Lemma 1: For the SF, $sp(\alpha) \subseteq sp(\tilde{\beta}_\perp)$.

The proof follows directly from the linear independence between the vectors β and $\tilde{\beta}$ (see Vahid and Engle, 1993) so that $\text{rank}[\beta : \tilde{\beta}] = r + s \leq n$. Hence we have that $\dim[sp(\alpha)] \leq \dim[sp(\tilde{\beta}_\perp)]$ or that $\text{rank}(\alpha) \leq \text{rank}(\tilde{\beta}_\perp)$ implying that $r \leq n - s$.

Remark (b) In the case of WF, we analogously define a $n(p - 1) \times 1$ vector $X_{t-1} = [\Delta y'_{t-1}, \dots, \Delta y'_{t-p+1}]'$ and the $n \times n(p - 1)$ matrix $\Phi = [\Phi_1^*, \dots, \Phi_{p-1}^*]$, so that (2) becomes

$$\Delta y_t = \alpha \beta' y_{t-1} + \Phi X_{t-1} + \varepsilon_t, \quad t = 1, \dots, T. \quad (6)$$

Under the assumption of a WF, Φ is of reduced rank $n - s$ and can be written as $\Phi = A[C_1, \dots, C_{p-1}] = AC$, where A is $n \times (n - s)$ full column rank matrix and C is $(n - s) \times n(p - 1)$ such that $\tilde{\beta}' AC X_{t-1} = 0$. The cofeature matrix $\tilde{\beta}$ must lie in $space(A_\perp)$ but not necessarily in $space(\alpha_\perp)$.

It is important to stress the difference between SF and WF. Firstly, the assumption of a SF reduced rank rules out predictability at any frequency and hence implies common cycles at all frequencies. On the contrary, by allowing for linear combinations that are predictable in the long run, the WF reduced rank structure restricts the short-run dynamics. Secondly, in the WF case, both the possible number and the nature of the common cyclical features change: s may be greater than $n - r$ but has to remain $\leq n - 1$ and the corresponding $n - s$

²In the sequel, space will be denoted by sp . We shall always denote the orthogonal complement of any $n \times s$ -dimensional matrix B , with $n > s$ and $\text{rank}(B) = s$, by the $n \times (n - s)$ matrix B_\perp such that $B' B_\perp = 0$ with $\text{rank}(B_\perp) = n - s$ and $\text{rank}(B : B_\perp) = n$. We then say that B_\perp spans the null space of B and B' spans the left null space of B_\perp .

³Consider the VAR(2) model $\Delta y_t = -(I - \Phi_1 - \Phi_2)y_{t-1} - \Phi_2 \Delta y_{t-1} + \varepsilon_t$ with $n = 4$, and $\text{rank}(I - \Phi_1 - \Phi_2) = r = 2$. In this case, $\text{rank}(\Phi_2)$ should be necessarily equal to 2, 3 or 4. Otherwise in the strongly nested structure, $\text{rank}(\Phi_2) = 1$ would mean that $s + r = 5 > n$ which is not possible.

common dynamic factors consist of linear combinations of the elements of X_{t-1} , $f_t = CX_{t-1}$, which only contain lagged first differences of the process. It is important to notice that the existence of s weak form common feature vectors with $s > r$, implies the existence of $s - r$ strong form common features as is shown in Lemma 2.

Lemma 2: In the VAR model (1) under Assumption 1 with $s > r$, Assumption 2 implies the existence of $s - r$ SF common feature vectors.

Proof: Denote by $\tilde{\beta}$ the $n \times s$ matrix of linearly independent WF common feature vectors. Any nonsingular transformation of $\tilde{\beta}$, $\tilde{\beta}A$, with A being an $s \times s$ nonsingular matrix, also forms a basis of the space spanned by the columns of $\tilde{\beta}$ and therefore is also a basis of the WF common feature space. The matrix $\tilde{\beta}'\Phi(1) = -\tilde{\beta}'\alpha\beta'$ has rank $\min(r, s)$. Therefore, if $s > r$, there are $s - r$ linearly independent column vectors such that there is an $n \times (s - r)$ matrix B with full column rank such that $B'\alpha = 0$. B can be constructed as $B = \tilde{\beta}A^*$ by choosing the $s \times (s - r)$ matrix A^* with rank $s - r$ such that B forms a basis for the left null space of α . Note that we can always normalize B such that the upper part equals I_{s-r} . \square

Remark (c) As pointed out, the WF has an interest in its own as it is a necessary condition for the existence of first order codependent cycles in a VAR(2) (see e.g. Vahid and Engle, 1997; Hecq, 2000) and of the SF. The WF restrictions are generally not invariant to alternative vector error correction representations such as that where y_{t-p} appears in levels instead of y_{t-1} . The implications of the lack of invariance are that the results from a reduced rank analysis of short-run dynamics are parametrization-specific.⁴ Invariance may be obtained at the price of assuming a SCCF or that the ranges of Φ_j^* 's are nested (see e.g. Ahn and Reinsel, 1988). The methods put forward in this paper can be applied to any of these alternative parametrizations. We present the analysis for the VECM (2) with y_{t-1} appearing in levels, first, because this parametrization is frequently used in empirical work; second because if a reduced rank structure is found it will imply a lower order SCM than for other parametrizations; third, the WF is more likely to be appropriate as it applies to the coefficients of the higher order lags of Δy_t in the VECM, which are usually less significant than those of small order lags of Δy_t (for non-seasonal processes). Alternatively, when modeling series for which there are no strong reasons to a priori prefer any of the VECM parametrizations, one can test the WF restrictions for each parametrization. Next, one can test the SF restrictions for those parametrizations for which the WF restrictions are not rejected. This sequential testing is likely to lead to detecting useful structures in the data.

⁴For instance, for the VAR(2) model in footnote 3, the WF is implied by $\tilde{\beta}'\Phi_2 = 0$, whereas when y_{t-2} is included in the error-correction term, WF restrictions require $\tilde{\beta}'(I - \Phi_1) = 0$.

When $n > 2$, and $s - r > 0$, besides the $s - r$ SF common features implied by s WF common features, the mixed form (MF) reduced rank restrictions may arise. They combine the SF and the WF in the following way.

Definition 3 (Mixed Form Reduced Rank Structure): *If in addition to Assumption 1 (cointegration) Assumption 2 holds for s common feature vectors $\tilde{\beta} = (\tilde{\beta}_1, \tilde{\beta}_2)$, with $\tilde{\beta}_1$ and $\tilde{\beta}_2$ being $n \times s_1$ and $n \times s_2$ full rank matrices respectively, with $s_1 + s_2 = s$, and in addition Assumption 3 holds for s_1 common feature vectors $\tilde{\beta}_1$ with $s > s_1$ and $n - r > s_1 > \max(0, s - r)$, then the implied reduced rank structure of the VECM (2) will be labelled a mixed form reduced rank structure (MF). Under MF, the $(n \times s)$ matrix $\tilde{\beta}$ spans the co-feature space, such that $\tilde{\beta}_1' \Delta y_t = \tilde{\beta}_1' \varepsilon_t$ is a s_1 -dimensional vector mean innovation process with respect to \mathfrak{F}_t and $\tilde{\beta}_2' (\Delta y_t - \alpha \beta' y_{t-1}) = \tilde{\beta}_2' \varepsilon_t$ is a s_2 -dimensional vector mean innovation process to \mathfrak{F}_t .*

Remark (d) Under the MF, there are $s_1 - \max(0, s - r) > 0$ SF common feature vectors which are not implied by the WF and yield testable restrictions on the parameters of the VECM (2). The matrix $\tilde{\beta}_1$ consists of $s - r$ columns which are linear combinations of $\tilde{\beta}$ and $s_1 - \max(0, s - r)$ columns of $\tilde{\beta}$ which satisfy Assumption 3.

Remark (e) Note that in the mixed case s_1 and s_2 have to satisfy the inequalities $s_1 + s_2 \leq n - 1$ and $s_1 \leq n - r$. Also, along the lines of lemma 1, we get $sp(\alpha) \supseteq sp(\tilde{\beta}_\perp)$.

Notice that we could easily extend these representations in order to analyze models in which only a part of short-run components disappears. This type of reduced rank structures has been studied by Ahn and Reinsel (1988) for stationary processes, Tiao and Tsay (1989) for VARMA models and by Reinsel and Ahn (1992) and Ahn (1997) for partially non-stationary processes.

3 Testing Different Forms of Reduced Rank Structures

3.1 Reduced rank hypotheses

The difference between the SF and the WF can be illustrated in terms of two competing models where we assume both cointegration and the existence of a $(n \times s)$ common feature matrix $\tilde{\beta}$. Under the assumption of SF the following model holds

$$\tilde{\beta}' \Delta y_t = \tilde{\beta}' \varepsilon_t, \quad t = 1, \dots, T, \quad (7)$$

while under WF we have

$$\tilde{\beta}' (\Delta y_t - \alpha \beta' y_{t-1}) = \tilde{\beta}' \varepsilon_t, \quad t = 1, \dots, T. \quad (8)$$

Let us first assume that the cointegrating rank r is known and fixed. For a given maintained reduced rank structure (WF or SF), we may consider the sequence of hypotheses (or models) in each column separately in order to test $H_0 : rank(\tilde{\beta}) \geq s$ against $H_a : rank(\tilde{\beta}) < s$ for the different values of s starting with $s = 0$ for the model without common features. In the SF case, the maximum number of common feature vectors is $n - r$. For the WF s has an upper bound⁵ of $n - 1$.

INSERT TABLE 1 ABOUT HERE

For each value of $s (\leq n - r)$ we can also compare the SF against the nesting alternative of a WF. The resulting structure of the various hypotheses of interest is summarized in Table 1, where \subset indicates the direction of the nesting between the different implied models. Table 1 shows that the hypotheses are nested "horizontally" and "vertically". In empirical work, one will usually start by considering "vertical" sequences of nested hypotheses for the WF and the SF respectively and for each sequence determine the value of s for which the null hypothesis is not rejected. Denote these values by s and s_1 respectively and by $H_{s_1, s}$ the hypothesis that the number of SF and WF common features is s_1 and s respectively. Next, for the values of s larger than $\max(1, s - r + 1)$ and for which the SF is not rejected, one will usually test horizontally the SF against the WF. All other "diagonal" comparisons, such as $H_{0, s}$ versus $H_{1, 1}$ for instance, involve non-nested hypotheses. Table 1 presents the structure of reduced rank hypotheses for $s_1 = 0, \dots, n - r$; $s = 0, \dots, n$ and $r = 1, \dots, n - 1$ and $n = 4$. The table is easily extended if we also consider the cointegrating rank as unknown.

3.2 Testing

Given that the hypotheses to be tested are nested, we rely on ML estimation of the underlying models following the approaches by Reinsel and Ahn (1992), Ahn (1997), Ahn and Reinsel (1988), Reinsel (1993) among others. Usually, when r and s are unknown, it appears impossible to find an explicit solution for the likelihood equations (see Johansen, 1995; Ahn, 1997). There are essentially two approaches to the determination of r, s and to the estimation of the parameters of interest. The first approach proposed and investigated by Ahn (1997), Ahn and Reinsel (1988) is to exploit the nested reduced rank structures and to compute numerically a Gaussian reduced-rank estimator based on iterative solution of approximate Newton-Raphson equations. Alternatively, one may follow a two-step approach in which r is first determined, while ignoring restrictions on the short-run dynamics of the model. Once r is determined and

⁵ $s = n$ implies that $\Delta y_t - \alpha \beta' y_{t-1}$ is already a n dimensional vector white noise process.

β (the cointegrating matrix) is estimated, s may be determined using the approach proposed by Vahid and Engle (1993) for example. The rationale behind this simple two-step analysis is that the determination of r and the efficiency of estimation of β are not affected asymptotically by the presence of the reduced rank structure on the short-run dynamics (see also Ahn, 1997; Phillips, 1991).

We use the two-step approach, although one may reasonably suspect small sample efficiency losses compared to using a one-step full information estimation method. As pointed out by various authors, a convenient way to test for reduced rank structures within the VECM is based on canonical correlation analysis. Let us first assume that r and β are known or that superconsistent estimates are available so that we may essentially consider them to be fixed and given.

Define the $T \times n$ matrices $W_1 = \Delta Y = (\Delta y_1, \dots, \Delta y_T)'$, $Y_{-1} = (y_0, \dots, y_{T-1})'$, $Z_1 = \Delta Y^*$ with ΔY^* being the LS residuals from the multivariate regression of ΔY on $Y_{-1}\beta$ and the $T \times (n(p-1) + r)$ matrix $W_2 = [Z_2, Y_{-1}\beta]$ with Z_2 being the $T \times n(p-1)$ matrix $(\Delta Y_{-1}^*, \dots, \Delta Y_{-p+1}^*)$. Under the maintained hypothesis of a SF reduced rank structure, the sequence of common feature Gaussian likelihood ratio test statistics for $H_0 : \text{rank}(\Phi^*) \leq n-s$ against $H_a : \text{rank}(\Phi^*) > n-s$, where Φ^* is defined in (5), or equivalently for $H_0 : \text{rank}(\tilde{\beta}) \geq s$ against $H_a : \text{rank}(\tilde{\beta}) < s$ can be shown (see Lütkepohl, 1991; Velu et al, 1986)) to be

$$\xi_S = -T \sum_{i=1}^s \log(1 - \lambda_i), \quad s = 1, \dots, n-r, \quad (9)$$

where $0 \leq \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_{n-r} < 1$ are the ordered eigenvalues of the symmetric matrix $(W_1' W_1)^{-1/2} W_1' W_2 (W_2' W_2)^{-1} W_2' W_1 (W_1' W_1)^{-1/2}$. The test statistic (9) can also be interpreted as the minimum of the objective function of the GMM estimator of $\tilde{\beta}$ subject to the normalization $(1/T) \tilde{\beta}' W_1' W_1 \tilde{\beta} = I_s$ (see Anderson and Vahid, 1998). For known r and β , under the null the test statistic ξ_S is asymptotically χ^2 -distributed with $s(n(p-1) + r) - s(n-s)$ degrees of freedom (Vahid and Engle, 1993).

In the case of WF reduced rank structure, this likelihood ratio test for $H_0 : \text{rank}(\tilde{\beta}) \geq s$ against $H_a : \text{rank}(\tilde{\beta}) < s$ reads as

$$\xi_W = -T \sum_{i=1}^s \log(1 - \tilde{\lambda}_i), \quad s = 1, \dots, n-1, \quad (10)$$

where $0 \leq \tilde{\lambda}_1 \leq \tilde{\lambda}_2 \leq \dots \leq \tilde{\lambda}_{n-1} < 1$ are the ordered eigenvalues of the symmetric matrix $(Z_1' Z_1)^{-1/2} Z_1' Z_2 (Z_2' Z_2)^{-1} Z_2' Z_1 (Z_1' Z_1)^{-1/2}$. This statistic has an asymptotic χ^2 -distribution with $s(n(p-1)) - s(n-s)$ degrees of freedom under the null.

A MF reduced rank structure hypothesis $H_0 : \text{rank}(\tilde{\beta}_1) \geq s_1$, for $\min(n-r, s) \geq s_1 > \max(0, s-r)$ and $\text{rank}(\tilde{\beta}_2) \geq s_2$ against $H_a : \text{rank}(\tilde{\beta}_1) < s_1$ or $\text{rank}(\tilde{\beta}_2) < s_2$, with $s_1 + s_2 = s$, can be tested in several ways. One way is to test SF restrictions for $s_1 = 1, \dots, n-r$, using the statistic ξ_S in (9). As this test ignores the restrictions implied by the existence of s_2 weak form common features, some power might be lost as will be illustrated in the next section.

Alternatively, the parameters $\tilde{\beta}$ and α from the WF can be estimated jointly by FIML for given s and β , e.g. by maximizing the likelihood function based on the $(s \times 1)$ subsystem (8), normalized on the first s variables of Δy_t by setting $\tilde{\beta}' = (I_s \ \tilde{\beta}_{s \times (n-s)}^*)'$, and completed by adding $(n-s)$ "reduced form" equations for the remaining $(n-s)$ variables in Δy_t

$$B' \Delta y_t = \begin{pmatrix} 0_{s \times n} & 0_{s \times n} & \dots & 0_{s \times n} & \alpha_1 \\ \Phi_{21}^* & \Phi_{22}^* & \dots & \Phi_{2p-1}^* & \alpha_2 \end{pmatrix} \hat{X}_{t-1}^* + B' \varepsilon_t, \quad (11)$$

with

$$B' = \begin{pmatrix} I_s & \tilde{\beta}_{s \times (n-s)}^* \\ 0_{(n-s) \times s} & I_{n-s} \end{pmatrix},$$

$\hat{X}_{t-1}^* = X_{t-1}^*$ with β replaced by the first stage superconsistent estimate, the Φ_{2i}^* matrices, $i = 1, \dots, p-1$, indicate the $n-s$ bottom rows of the Φ_i^* matrices in (2) and $(\alpha_1' \ \alpha_2')$ is the partition of $\alpha' B$. Under a MF structure, for given β , s and s_1 , we can specify a similar pseudo-structural system:

$$B' \Delta y_t = \begin{pmatrix} 0_{s_1 \times n} & 0_{s_1 \times n} & \dots & 0_{s_1 \times n} & 0_{s_1 \times r} \\ 0_{s_2 \times n} & 0_{s_2 \times n} & \dots & 0_{s_2 \times n} & \alpha_2 \\ \Phi_{31}^* & \Phi_{32}^* & \dots & \Phi_{3p-1}^* & \alpha_3 \end{pmatrix} \hat{X}_{t-1}^* + B' \varepsilon_t, \quad (12)$$

where

$$B' = \begin{pmatrix} I_{s_1} & \tilde{\beta}_{1, s_1 \times (n-s_1)}^* \\ 0_{s_2 \times s_1} & \tilde{\beta}_{2, s_2 \times (n-s_1)}^* \\ 0_{(n-s) \times s_1} & A_{(n-s) \times (n-s_1)} \end{pmatrix},$$

$\tilde{\beta}_{2, s_2 \times (n-s_1)}^* = (I_{s_2} \ \tilde{\beta}_{2, s_2 \times (n-s)}^*)'$, $A_{(n-s) \times (n-s_1)} = (0_{(n-s) \times s_2} \ I_{n-s})$, the Φ_{3i}^* matrices, $i = 1, \dots, p-1$, indicate the $n-s$ bottom rows of the Φ_i^* matrices in (2) and $\alpha' B = (0_{s_1 \times r}' \ \alpha_2' \ \alpha_3')$ with α_2 and α_3 of dimension $(s_2 \times r)$ and $(n-s) \times r$ respectively.

For given β and s , the MF with s_1 SF vectors and s_2 WF vectors can be tested against the

WF by testing for the validity of the additional parameter restrictions implied by (12) using a standard LR test statistics denoted by ξ_M . No efficiency loss arises if a superconsistent estimate is substituted for the cointegrating vectors β . Under the null of the MF, ξ_M is asymptotically χ^2 -distributed with degrees of freedom being given by the number of additional parametric restrictions imposed under (12), i.e. $s_1r - s_2s_1$. This estimation procedure has been used in the empirical analysis reported in Section 5.⁶

For given r , a likelihood ratio test statistic for the null hypothesis of a SF against the alternative of a WF, for each possible common feature rank $s = \max(1, s - r + 1) \dots n - r$, is given by

$$\xi_{SW} = -T \sum_{i=1}^s \log\left\{\frac{(1 - \lambda_i)}{(1 - \tilde{\lambda}_i)}\right\}, \quad (13)$$

where the $\tilde{\lambda}_i$'s and the λ_i 's are defined as above. Conditional on known r and β , all variables involved are weakly stationary both under the null and the alternative, so that standard asymptotic theory applies. ξ_{SW} has an asymptotic χ^2 -distribution with degrees of freedom equal to the number of restrictions rs imposed under the H_0 . If the null hypothesis is rejected, one can proceed further in determining s by testing the number of zero squared canonical correlations between Z_1 and Z_2 . Note that the test statistics (9), (10) and (13) only enable to formally compare the nested models from Table 1. For "diagonal" type of model comparisons involving non-nested hypotheses, we propose to select the model which, for given p, r and β , minimizes one of the well-known model selection criteria (AIC, SBC, HQC) where, given that we have omitted deterministic terms, the number of parameters is $n(n(p-1)+r) - s(n(p-1)+r) + s(n-s)$ under the SF and $n(n(p-1)+r) - s(n(p-1)) + s(n-s)$ under the WF.⁷

4 Monte Carlo Results

In this section we present evidence on the finite sample behavior of the sequential test procedure put forward in Section 3.2. One should indeed be careful when interpreting the outcome of the three sequences of LR tests ξ_S , ξ_W and ξ_{SW} . Given that s is unknown, and given the sequential nature of the testing procedure, the significance levels of the individual tests in the

⁶Alternatively, the subsystem (8) normalized as above could be estimated by GMM, to get unrestricted estimates of the matrices $\tilde{\beta}^{*'}$ and $\tilde{\beta}'\alpha$ (for given β) and testing the rank of the matrix $\tilde{\beta}'\alpha$ (see Hecq, Palm and Urbain, 1999).

⁷These model selection criteria can be also used to select the optimal values for r and s given p (as we assumed in the preceding section) and have also recently been considered for common feature analysis by Vahid and Issler (1999) for unknown s and p .

sequence must be distinguished from the overall Type I error of the sequential testing procedure. Also, the above sequential procedures are essentially based on asymptotic properties such as the irrelevance of the reduced rank structure for the optimal estimation of β and the determination of r . A Monte Carlo experiment should shed some light on the finite sample behavior of the sequences of common features LR tests presented in the preceding section. We concentrate on two issues which we believe are particularly relevant for applications:

1. the size and power in finite samples of the common feature LR tests,
2. the possible effect of incorrectly specifying the number of cointegrating vectors and/or the lag length.

In order to address these issues we consider a simple trivariate data generating process (DGP) where we assume the existence of two common feature vectors, i.e. $s = 2$. Throughout the simulations, p is fixed either to its true value $p = 2$ or to 4. The size and power of codependence tests in the presence of either incorrectly specified lag length of the model dynamics, omission of a cointegrating vector, non-normal errors, or temporal aggregation have been extensively analyzed by Beine and Hecq (1999). Strong and weak form reduced rank structures are considered. The DGP is a Gaussian VAR of order two written in VECM form. In order to provide some motivation for the choice of the DGP, we label the three variables c_t , i_t , y_t as for consumption, investment and real output. In line with a simple form of a neo-classical model, we assume the existence of two long-run relationships: $c_t - y_t$ and $i_t - y_t$. The covariance matrix has been calibrated on quarterly real US data for consumption, investment and value added for the period 1950-1996.

$$\begin{bmatrix} \Delta c_t \\ \Delta i_t \\ \Delta y_t \end{bmatrix} = \begin{bmatrix} 0.2 & 0.1 & 0.1 \\ 0.8 & 0.4 & 0.4 \\ 0.4 & 0.2 & 0.2 \end{bmatrix} \begin{bmatrix} \Delta c_{t-1} \\ \Delta i_{t-1} \\ \Delta y_{t-1} \end{bmatrix} + \alpha \beta' \begin{bmatrix} c_{t-1} \\ i_{t-1} \\ y_{t-1} \end{bmatrix} + \begin{bmatrix} \varepsilon_{1t} \\ \varepsilon_{2t} \\ \varepsilon_{3t} \end{bmatrix} \quad (14)$$

$$\begin{bmatrix} \varepsilon_{1t} \\ \varepsilon_{2t} \\ \varepsilon_{3t} \end{bmatrix} \sim N \left(\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1.0 & 0.6 & 0.6 \\ 0.6 & 1.0 & 0.6 \\ 0.6 & 0.6 & 1.0 \end{bmatrix} \right).$$

The cofeature matrix associated with the DGP in (14) is given by

$$\tilde{\beta}' = \begin{bmatrix} 1 & -0.25 & 0 \\ 1 & 0 & -0.5 \end{bmatrix}.$$

It yields two linear combinations of the variables in the model that annihilate the short-run dynamics. In our experiments, the nature of the reduced rank structure depends on the choice of the values for α and β . Tables 2 and 3 illustrate the outcome of the simulations when the DGP has a SF reduced rank structure with $p = 2$, $r = 1$ and $s = 2$. Tables 2 and 3 present the rejection frequencies of the statistics (9), (10) and (13) for models assuming $r = 1$, $r = 2$ and r being estimated for a DGP with $s = 2$ and choosing the correct lag length $p = 2$ or setting the lag length equal to 4 respectively. Notice that for the SF and with $n = 3$, the number of cointegrating vectors is by definition bounded to be equal to one in the DGP. We therefore present simulation results for models with the correct specification of the cointegrating rank as well as over-specification of r .

In each case, we use 10,000 replications and a sample size of $T=1000$ and 100. The cointegration coefficients β are set equal to their estimated values obtained by ML estimation in a first stage (see Johansen, 1995). Conditionally on these estimates for β , α and $\tilde{\beta}$ are estimated by ML as described in Section 3.2. All simulations have been performed with GAUSS and the first 50 observations initialize the processes. The empirical (size unadjusted) power and size are given as percentage rejection frequencies. The nominal size used to obtain these rejection frequencies is fixed at 5% for each individual test.

INSERT TABLES 2-3 ABOUT HERE

Tables 2 and 3 report simulation results for a DGP with SF. Several remarks are worth to be made:

- In general, the differences between the results in Tables 2 and 3 are small. The inefficiency resulting from choosing too long lags is small.
- When the DGP has a SF and the number of cointegrating vectors is correctly specified, both ξ_S and ξ_W behave fairly well in detecting the two cofeature vectors. Note also the behavior of the sequence of the LR tests ξ_{SW} that does not show any significant size distortion.
- If we estimate the number of cointegrating vectors or fix it at a value higher than the true r , the rejection frequency of ξ_S is slightly distorted. ξ_W still behaves very well in detecting the correct number of common feature vectors. However the LR tests for SF versus WF display significant size distortions reaching 50% instead of the 5% chosen nominal level.

- Overall, the tests appear to reject too frequently the null hypothesis when the model is misspecified in some way (with the exception of lag length). The tests therefore tend to favor accepting models with fewer restrictions than the true model, implying thereby a loss of efficiency, but not a misspecification.

In the Tables 4 and 5, rejection frequencies for a DGP under WF restrictions are given.

INSERT TABLES 4-5 ABOUT HERE

We draw some conclusions from Tables 4 and 5:

- Again, the effect of overfitting the lag length is small.
- When the DGP with $r = 1$ has a WF reduced rank structure, ξ_W determines without size distortions the correct number of common feature vectors, whether r is fixed at the true value, estimated or fixed at 2. When the true value of r equals 2, ξ_W performs very well except when r is fixed at 1.
- The statistic ξ_S detects a SF reduced rank structure implied by a WF reduced rank structure ($s - r > 0$) with a rejection frequency of approximately 5% when r is correctly specified (panel one). When r is fixed at a value larger than the true one or when it is estimated, the size of ξ_S is much larger than the nominal size of 5%. For ξ_{SW} , the rejection frequencies are similar.
- It is interesting to note that the sequence of ξ_W still selects the correct number of common feature vectors without size distortions when we overspecify the number of cointegrating vectors. This is not surprising since the coefficient of a non significant $I(1)$ variable in a $I(0)$ model converges in probability to zero. ξ_S still rejects the presence of any cofeature vector since this case excludes the existence of an implied SF ($s - r = 0$).
- Overall, the likelihood ratio statistics ξ_{SW} for the null of SF against the WF has high power close to one in most cases. When $s - r > 0$ in the DGP, there are $(s - r)$ implied SF common features vectors and the rejection frequencies for $s = 1$ in Tables 4 and 5 have to be interpreted as an empirical size of the test. In these cases, the statistic ξ_{SW} rejects too frequently the (implied) null hypothesis.

Results for the statistics presented above with a small sample correction as suggested by Reinsel and Ahn (1992) for cointegration tests, where ξ_W and ξ_S are respectively premultiplied by the factors $(T - n(p - 1))/T$ and $(T - n(p - 1) - r)/T$, (for further details see Hecq, 2000),

have been obtained as well. They are available from the authors upon request. Overall, the results are similar to the corresponding results given in Tables 2-5. For $T = 100$, in some instances, the corrected version of the tests performs better than the uncorrected ones.

Table 6 contains some illustrative simulation results for a DGP with a MF reduced rank. For this purpose, the DGP is slightly modified and extended in order to account for a MF. The selected DGP is a VAR(2) with $n = 4$, $r = 2$ and $s = 3$. From Lemma 2 there is one implied cofeature vector ($s - r = 1$). The loading matrix α is chosen such that the DGP displays one additional cofeature vectors, i.e. $s_1 = 2$. The following matrices are retained:

$$\tilde{\beta} = \begin{bmatrix} 1 & 1 & 1 \\ -.25 & 0 & 0 \\ 0 & -.5 & 0 \\ 0 & 0 & -.4 \end{bmatrix}, \tilde{\beta}_\perp = \begin{bmatrix} -.1 \\ -.4 \\ -.2 \\ -.25 \end{bmatrix}, \alpha = \begin{bmatrix} -.2 & .2 \\ -.8 & 0 \\ -.4 & .8 \\ -.5 & 0 \end{bmatrix}, \beta = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 1.2 & -.8 \\ -1 & -1 \end{bmatrix}.$$

This particular choice of α implies the existence of $\tilde{\beta}'_1$ satisfying⁸ $\tilde{\beta}'_1 \alpha = 0$. As discussed in the preceding section, we report results for ξ_S and a likelihood ratio tests of the mixed form denoted by ξ_M .

In Table 6 we report rejection frequencies, based on 10,000 replications, under the correct assumption of a mixed form with $s_1 = 2$ (size of the tests) as well as those obtained when we let the parameter $\alpha_{3,1}$ successively take the values -0.45, -0.5 which implies the existence of a weak form⁹. In all the cases, the empirical power is not size adjusted.

INSERT TABLE 6 ABOUT HERE

From Table 6, we observe that ξ_S and ξ_M do not suffer from serious size distortion. With respect to the empirical powers, it appears that ξ_M perform substantially better than ξ_S . Remark that r is assumed known while β is estimated and thus the cointegrating rank is correctly specified.

The limited Monte Carlo evidence presented in this section leads us to propose the following model selection strategy.

1. Start by determining the lag length p and the number of cointegrating vectors, trying to avoid underestimation of r . In practice, Johansen's ML statistics complemented by

⁸The columns of the 4×2 matrix $\tilde{\beta}_1$ may simply be constructed by adding the first and second column of $\tilde{\beta}$ on the one hand and by adding the second and the third one on the other hand.

⁹Remark that the values of $\alpha_{3,1}$ chosen for the computation of the empirical powers only imply small deviation from the mixed form. For other values the empirical power rapidly reaches 1.

a visual inspection may prove useful to determine an upper bound for r ,

2. compute the sequences of common feature LR tests ξ_S and ξ_W and select s for the SF and WF respectively, check whether the number of WF common features exceeds r , in which case the WF implies $s - r$ SF common features,
3. for the cases where $s_1 = \max(1, s - r + 1), \dots, n - r$ compute ξ_{SW} to select the appropriate reduced rank structure,
4. for the cases where $s_1 = \max(1, s - r + 1), \dots, \min(n - r, s)$, compute a likelihood ratio MF test,
5. repeat the analysis with $r - i$ cointegrating vectors for $i = 1, \dots, r - 1$. For each case compute the various information criteria.

5 An Application

5.1 Background

A vast amount of empirical macroeconomic literature has studied the long-run implications of the real business cycle models, see e.g. Neusser (1991), King and al. (1991), Kunst and Neusser (1990). With the exception of the work of Issler and Vahid (2001), little work has however been done on short-run co-movements in neoclassical growth models. As in Issler and Vahid (2001) we analyze a simple form of the real business cycle (RBC) model which assumes common trends and common cycles between U.S. per capita real consumption, investment and output. We relax the hypotheses about the number of common feature and cointegrating vectors. More formally, consider the following trivariate system for the logarithms of income y_t , consumption c_t and investment i_t put forward by King, Plosser and Rebelo (1988) and analyzed by Issler and Vahid (2001):

$$c_t = x_t^p + \bar{c} + \pi_c \hat{k}_t \quad (15)$$

$$i_t = x_t^p + \bar{i} + \pi_i \hat{k}_t \quad (16)$$

$$y_t = x_t^p + \bar{y} + \pi_y \hat{k}_t, \quad (17)$$

where $x_t^p = x_{t-1}^p + \varepsilon_t^p$ is the common trend, that is a random walk measuring among other the impact of technology processes, \bar{y} , \bar{c} and \bar{i} are the constant steady state values, \hat{k}_t is the common cycle, that is the stationary transitory deviation of the capital stock from its steady state value and π_c, π_i and π_y are constant parameters. ε_t^p and \hat{k}_t may be correlated.

We analyze the period 1954:1 - 1996:4, that is 172 quarterly observations. Data prior to 1954:1 were used as initial observations in regressions that contain lags. Notice that we had the observations from 1948:1 onwards but we preferred, as King and al. (1991) suggested, to exclude turbulent periods during Korean War, price control and Treasury-Fed agreement. The data used are the revised (May 1997) series from the Survey of Current Business national account for the United States (source BEA). The variables are c_t : personal consumption expenditures, i_t : gross private domestic investment and the output y_t is the GDP less the government expenditures. The three variables have been divided by the size of the civilian population above sixteen years of age. Figure 1 presents the data series used in this section. The series are seasonally adjusted and transformed into natural log.

INSERT FIGURE 1 ABOUT HERE

5.2 Cointegration and Common Feature Analysis

The model that best characterizes the covariance structure of the data is a VAR of order 5 (using LR statistics) with an unrestricted intercept in the short-run. Table 7 presents the test statistics for the Johansen (1995) rank test of the number of cointegrating relationships with small sample correction and the 5% critical values.

INSERT TABLE 7 ABOUT HERE

The results in Table 7 indicate that we cannot reject the presence of two cointegrating vectors whose coefficients are not far from the theoretical ones i.e. $c_t - y_t$ and $i_t - y_t$ are both $I(0)$ ¹⁰. The estimated cointegrated relationships are respectively:

$$\begin{array}{ccc} c_t - 0.958 y_t & \text{and} & i_t - 1.103 y_t, \\ (0.016) & & (0.045) \end{array} \quad (18)$$

where asymptotic standard errors are reported in parentheses. The likelihood ratio test for unit long-run elasticities in both vectors (e.g. stationarity of the great ratios), $\chi^2_{(2)}$ -distributed under the null, yields a value of 3.07 so that these restrictions are rejected at the 5% level. For the determination of the number of common feature vectors, we use the sequential likelihood

¹⁰The traditional Augmented Dickey Fuller unit root test statistics also strongly reject the null hypothesis. For instance for and ADF(4) we get the values $\tau_\mu = -3.31$, $\tau_\tau = -3.97$ for the variable $c_t - y_t$ and $\tau_\mu = -4.14$, $\tau_\tau = -4.23$ for the variable $i_t - y_t$.

ratio tests presented in Section 3. For $p = 5$ and $r = 2$ fixed we obtain the tests statistics and the p -values reported in Table 8.

INSERT TABLE 8 ABOUT HERE

ξ_S and ξ_W respectively denote likelihood ratio tests for the number of common features for given cointegrating rank r and a given reduced rank structure. ξ_S^{cor} and ξ_W^{cor} are the small sample corrected version. $\xi_{SW} = \xi_S - \xi_W$ yields the likelihood ratio statistic for the null of SF against WF for given cointegrating and cofeature rank. For $s = 1$, the ξ_{SW} test statistic has a value of 10.06 for a $\chi^2_{(2)}$ null distribution which yields a rejection probability of .0065. We reject the SF model in favor of the WF one. The common feature relation we would retain in the SF case is $\Delta c_t + 0.106\Delta i_t - .959\Delta y_t$. In the class of WF we still have to choose between $s = 1$ and $s = 2$. The test statistic $\xi_W = 33.83$ does not reject the null hypothesis of $s \geq 2$, *i.e.* the presence of at least two common features vectors. Notice that information criteria also favor the WF assumption with $s = 2$. With $s = 2$ and $r = 2$, the WF does not imply a SF common feature since $s - r = 0$.

FIML estimates of the WF model (11), under $s = 2$ and permuting the columns of $\tilde{\beta}$ in order to find the vectors with the meaningful economic interpretation, results in the following two cofeature relationships:

$$\begin{aligned} \Delta c_t^* - 0.501 \Delta y_t^* \quad \text{and} \quad \Delta i_t^* - 4.776 \Delta y_t^*, \\ (0.087) \qquad \qquad \qquad (0.481) \end{aligned} \tag{19}$$

where asymptotic standard errors are reported in parentheses and where a * indicates that the corresponding variables are expressed in deviation from long-run effects. It is seen that these two vectors fit pretty well business cycle stylized facts, *e.g.* that consumption is smoother than output, investment is more volatile than output and there is a single synchronous cycle which can be extracted using the Stock-Watson-Beveridge-Nelson decomposition developed in Hecq, Palm and Urbain (2000). This common cyclical component is given in Figure 2 with the shaded areas indicating the NBER peak to trough periods.

INSERT FIGURE 2 ABOUT HERE

The final step is to investigate the potential presence of MF common features. We may easily obtain the matrices entering the restricted representation (12). Assuming that the first

vector is of a strong form, FIML estimation of the MF model with unknown $\tilde{\beta} = (\tilde{\beta}_1 \tilde{\beta}_2)$ leads to the following two cofeature linear combinations:

$$\begin{array}{ccc} \Delta c_t - 0.639 \Delta y_t + 0.039 \Delta i_t & \text{and} & \Delta i_t^* - 4.866 \Delta y_t^* \\ (0.191) & (0.044) & (0.475) \end{array}$$

The likelihood ratio test of the MF against the WF is distributed $\chi^2(1)$ under the null of the MF. Its value is 1.472. The likelihood ratio test for the additional restriction that the coefficient of Δi_t in the first cofeature vector is zero has a value of 0.592. It is also $\chi^2(1)$ -distributed. Consequently, we may reestimate the MF model by FIML which leads to the following two cofeature relations:

$$\begin{array}{ccc} \Delta c_t - 0.485 \Delta y_t & \text{and} & \Delta i_t^* - 4.86 \Delta y_t^* \\ (0.067) & & (0.467) \end{array}$$

Notice that a MF model that assumes that the first vector is of a weak form is rejected at any reasonable significance level (p -value less than 0.001).

At this stage it may be interesting to note the sharp reduction in the number of parameters that have to be estimated once we impose (valid) reduced rank structures. The unrestricted trivariate VAR(5) model contains 45 unknown parameters. Under cointegration with $r = 2$, the number of unknown parameters reduces to 44. With one SF cofeature vector it reduces to 32 while with two weak form cofeature vectors the number becomes 22. Note finally that the mixed form only contains 20 unknown parameters.

6 Conclusion

In this paper, we studied a linear Gaussian VAR model with nonstationary but cointegrated variables that have common cyclical features.

We introduced the concepts of strong, weak and mixed form reduced rank structures and discussed their implications for VAR modeling. SF reduced rank structures arise when the common features are such that there exists one or several linear combinations of the set of variables under investigation expressed in first differences which are white noise. The existence of a WF reduced rank structure implies that linear combinations of the first differences of the variables in the model in deviation from the long-run relationships are white noise. We showed that the constraint that the number of common features plus the number of cointegrating relationships should be less than or equal to the number of variables no longer applies under

the WF. This allows to consider more significant long-run relationships between the variables in first differences. It also yields an efficiency increase for the estimates, resulting from the reduction in the number of free parameters to be estimated.

We designed a modeling strategy and proposed likelihood ratio tests for the three types of reduced rank structures. We studied the small sample properties of the test using Monte Carlo simulations. It appeared that in particular under SF it is of great importance to correctly determine the cointegrating rank before testing SF against WF. An empirical analysis of the relationship between the consumption, investment and real GDP leads to the conclusion that the existence of a WF reduced rank structure with one common trend and one weak form common cycle is not rejected by the information in the series. The presence of two common feature vectors means that the short-run dynamics of the system is governed by a single weak form common cycle as shown in Hecq, Palm and Urbain (2000) who present the Stock-Watson-Beveridge-Nelson decompositions of y_t into permanent and transitory components for the SF and the WF reduced rank structures.

References

- [1] AHN, S. K. (1997), Inference of Vector Autoregressive Models with Cointegration and Scalar Components, *Journal of American Statistical Association*, 92, 350-356.
- [2] AHN, S. K. AND G. C. REINSEL (1988), Nested Reduced Rank Autoregressive Models for Multiple Time Series, *Journal of the American Statistical Association*, 83, 849-856.
- [3] ANDERSON, H.M. AND F. VAHID (1998), Testing Multiple Equation Systems for Common Nonlinear Components, *Journal of Econometrics*, 84, 1-36.
- [4] BEINE, M. AND A. HECQ (1999), Inference in Codependence : Some Monte Carlo Results and Applications, *Annales d'Economie et de Statistique*, 54, 69-90.
- [5] CUBADDA, G. (1999), Common Serial Correlation and Common Business Cycles: A Cautious Note, *Empirical Economics*, 24, 529-535.
- [6] ENGLE, R. F. AND S. KOZICKI (1993), Testing for Common Features (with comments), *Journal of Business and Economic Statistics*, 11, 369-395.
- [7] GOURIÉROUX, CH. AND I. PEAUCELLE (1989), Detecting a Long-Run Relationship, CEPREMAP Discussion Paper 8902.
- [8] HECQ, A. (2000), *Common Cyclical Features in Multiple Time Series and Panel Data: Methodological Aspects and Applications*, Ph. D. Thesis, University Maastricht.
- [9] HECQ, A., F.C. PALM AND J.-P. URBAIN (1999), GMM-Based Estimation Procedures and Test Criteria for Reduced Rank Models, mimeo, University Maastricht .
- [10] HECQ, A., F.C. PALM AND J.-P. URBAIN (2000), Permanent-Transitory Decomposition in VAR Models with Cointegration and Common Cycles, *Oxford Bulletin of Economics and Statistics*, 62, 511-532.
- [11] ISSLER, J. V. AND F. VAHID (2001), Common Cycles and the Importance of Transitory Shocks to Macroeconomic Aggregates, *Journal of Monetary Economics*, forthcoming.
- [12] KING, R.G., PLOSSER, C.I., STOCK, J.H. AND M.W. WATSON (1991), Stochastic Trends and Economic Fluctuations, *American Economic Review*, 81, 4, 819-840.
- [13] KING, R.G., PLOSSER, C.I. AND S. REBELO (1988), Production, Growth and Business Cycles II, *Journal of Monetary Economics*, 21, 309-341.

- [14] KUNST, R. AND K. NEUSSER (1990), Cointegration in Macroeconomic Systems, *Journal of Applied Econometrics*, 5, 351-365.
- [15] JOHANSEN, S. (1995), *Likelihood-Based Inference in Cointegrated Vector Autoregressive Models* (Oxford University Press: Oxford)
- [16] LÜTKEPOHL, H. (1991), *Introduction to Multiple Time Series Analysis*, (Springer Verlag: Berlin).
- [17] NEUSSER, K. (1991), Testing the Long-Run Implications of the Neoclassical Growth Model, *Journal of Monetary Economics*, 27, 3-37.
- [18] PHILLIPS, P.C.B. (1991), Optimal Inference in Cointegrated Systems, *Econometrica*, 59, 283-306.
- [19] REINSEL, G. C. (1993), *Elements of Multivariate Time Series Analysis*, (Springer Verlag: Berlin).
- [20] REINSEL, G. C. AND S. K. AHN (1992), Vector Autoregressive Models with Unit Roots and Reduced Rank Structure : Estimation, Likelihood Ratio Tests, and Forecasting, *Journal of Time Series Analysis*, 13, 353-375.
- [21] TIAO, G. C. AND R. S. TSAY (1989), Model Specification in Multivariate Time Series (with comments), *Journal of Royal Statistical Society, Series B*, 51, 157-213.
- [22] VAHID, F. AND R. F. ENGLE (1993), Common Trends and Common Cycles, *Journal of Applied Econometrics*, 8, 341-360.
- [23] VAHID, F. AND R. F. ENGLE (1997), Codependent Cycles, *Journal of Econometrics*, 80, 199-221.
- [24] VAHID, F. AND J.V. ISSLER (1999), The Importance of Common-Cyclical Features in VAR Analysis: A Monte-Carlo Study, paper presented at ESEM99.
- [25] VELU, R. P. , REINSEL, G. C. AND D. W. WICHERN (1986), Reduced Rank Models for Multivariate Time Series, *Biometrika*, 73, 105-118.

Table 1: Reduced Rank Hypotheses for $n = 4$

$r = 1$	$s1 = 0$	$H_{0,0}$	\supset	$H_{0,1}$	$=$	$H_{0,1}$				
		\cup		\cup		\cup				
	$s1 = 1$	$H_{1,1}$	$=$	$H_{1,1}$	\supset	$H_{1,2}$	$=$	$H_{1,2}$		
				\cup		\cup		\cup		
$r = 2$	$s1 = 2$			$H_{2,2}$	$=$	$H_{2,2}$	\supset	$H_{2,3}$	$=$	$H_{2,3}$
						\cup		\cup		\cup
	$s1 = 3$					$H_{3,3}$	$=$	$H_{3,3}$	\supset	$H_{3,4}$
$r = 3$	$s1 = 0$	$H_{0,0}$	\supset	$H_{0,1}$	\supset	$H_{0,2}$	$=$	$H_{0,2}$		
		\cup		\cup		\cup		\cup		
	$s1 = 1$	$H_{1,1}$	$=$	$H_{1,1}$	\supset	$H_{1,2}$	\supset	$H_{1,3}$	$=$	$H_{1,3}$
				\cup		\cup		\cup		\cup
$r = 3$	$s1 = 2$			$H_{2,2}$	$=$	$H_{2,2}$	\supset	$H_{2,3}$	\supset	$H_{2,4}$
	$s1 = 3$									
$r = 3$	$s1 = 0$	$H_{0,0}$	\supset	$H_{0,1}$	\supset	$H_{0,2}$	\supset	$H_{0,3}$	$=$	$H_{0,3}$
		\cup		\cup		\cup		\cup		\cup
	$s1 = 1$	$H_{1,1}$	$=$	$H_{1,1}$	\supset	$H_{1,2}$	\supset	$H_{1,3}$	\supset	$H_{1,4}$
$r = 3$	$s1 = 2$									
	$s1 = 3$									

Table 2: Empirical Rejection Frequencies¹ of the LR tests for SF, $p = 2$

DGP: SF $s = 2$	Estimated Model						
$r = 1$ $\alpha = \begin{bmatrix} -0.10 \\ -0.40 \\ -0.20 \end{bmatrix}$ $\beta' = \begin{bmatrix} 0 & 1 & -1 \end{bmatrix}$	$r = 1, p = 2$						
	$T = 1000$			$T = 100$			
		ξ_S	ξ_W	ξ_{SW}	ξ_S	ξ_W	ξ_{SW}
	$s \geq 1$	0.24	0.41	0.78	0.33	0.53	0.92
	$s \geq 2$	4.90	5.00	5.37	6.72	6.53	6.13
	$s = 3$	100.00	100.00	100.00	100.00	100.00	93.96
	$r = 2, p = 2$						
	$T = 1000$			$T = 100$			
		ξ_S	ξ_W	ξ_{SW}	ξ_S	ξ_W	ξ_{SW}
	$s \geq 1$	1.76	0.41	4.09	2.37	0.66	4.81
	$s \geq 2$	34.59	4.91	50.24	40.88	7.29	53.58
	$s = 3$	100.00	100.00	100.00	100.00	100.00	98.12
	$\hat{r} = \text{rank}(\hat{\beta}), p = 2$						
	$T = 1000$			$T = 100$			
		ξ_S	ξ_W	ξ_{SW}	ξ_S	ξ_W	ξ_{SW}
	$s \geq 1$	0.52	0.42	1.34	0.78	0.61	1.72
	$s \geq 2$	14.19	5.02	15.85	18.26	7.12	18.87
	$s = 3$	100.00	100.00	100.00	100.00	100.00	94.84

¹ The rejection frequencies are based on 10,000 replications and calculated using asymptotic critical values. The nominal level is fixed at 5%.

Table 3: Empirical Rejection Frequencies¹ of the LR tests for SF, $p = 4$

DGP: SF $s = 2$	Estimated Model
$r = 1$ $\alpha = \begin{bmatrix} -0.10 \\ -0.40 \\ -0.20 \end{bmatrix}$ $\beta' = \begin{bmatrix} 0 & 1 & -1 \end{bmatrix}$	$r = 1, p = 4$
	$T = 1000$
	$T = 100$
	$\xi_S \quad \xi_W \quad \xi_{SW}$
	$s \geq 1 \quad 0.26 \quad 0.21 \quad 2.54$
	$s \geq 2 \quad 5.52 \quad 5.49 \quad 5.36$
	$s = 3 \quad 100.00 \quad 100.00 \quad 100.00$
	$\xi_S \quad \xi_W \quad \xi_{SW}$
	$s \geq 1 \quad 0.48 \quad 0.53 \quad 3.09$
	$s \geq 2 \quad 10.38 \quad 10.17 \quad 6.57$
	$s = 3 \quad 100.00 \quad 100.00 \quad 93.45$
	$r = 2, p = 4$
	$T = 1000$
	$T = 100$
	$\xi_S \quad \xi_W \quad \xi_{SW}$
	$s \geq 1 \quad 1.10 \quad 0.23 \quad 12.94$
	$s \geq 2 \quad 21.97 \quad 5.61 \quad 49.97$
	$s = 3 \quad 100.00 \quad 100.00 \quad 100.00$
	$\xi_S \quad \xi_W \quad \xi_{SW}$
	$s \geq 1 \quad 2.20 \quad 0.77 \quad 13.99$
	$s \geq 2 \quad 32.78 \quad 11.06 \quad 51.62$
	$s = 3 \quad 100.00 \quad 100.00 \quad 97.77$
	$\hat{r} = rank(\hat{\beta}), p = 4$
	$T = 1000$
	$T = 100$
	$\xi_S \quad \xi_W \quad \xi_{SW}$
	$s \geq 1 \quad 0.63 \quad 0.22 \quad 4.51$
	$s \geq 2 \quad 11.11 \quad 5.61 \quad 15.99$
	$s = 3 \quad 100.00 \quad 100.00 \quad 100.00$
	$\xi_S \quad \xi_W \quad \xi_{SW}$
	$s \geq 1 \quad 1.14 \quad 0.67 \quad 5.43$
	$s \geq 2 \quad 18.66 \quad 11.14 \quad 19.17$
	$s = 3 \quad 100.00 \quad 100.00 \quad 94.50$

¹ The rejection frequencies are based on 10,000 replications and calculated using asymptotic critical values. The nominal level is fixed at 5%.

Table 4: Empirical Rejection Frequencies¹ of the LR tests for WF, $p = 2$

DGP: WF $s = 2$	Estimated Model
$r = 1$ $\alpha = \begin{bmatrix} -0.50 \\ 0.10 \\ 0.20 \end{bmatrix}$ $\beta' = \begin{bmatrix} 1 & 0 & -1 \end{bmatrix}$	$r = 1, p = 2$ <div style="display: flex; justify-content: space-around;"> <div> $T = 1000$ $\xi_S \quad \xi_W \quad \xi_{SW}$ </div> <div> $T = 100$ $\xi_S \quad \xi_W \quad \xi_{SW}$ </div> </div> $s \geq 1$ 5.10 0.34 10.78 6.22 0.50 12.34 $s \geq 2$ 100.00 5.03 100.00 100.00 7.59 100.00 $s = 3$ 100.00 100.00 100.00 100.00 100.00 100.00
	$r = 2, p = 2$ <div style="display: flex; justify-content: space-around;"> <div> $T = 1000$ $\xi_S \quad \xi_W \quad \xi_{SW}$ </div> <div> $T = 100$ $\xi_S \quad \xi_W \quad \xi_{SW}$ </div> </div> $s \geq 1$ 30.11 0.35 40.57 36.08 0.60 46.32 $s \geq 2$ 100.00 4.88 100.00 100.00 7.56 100.00 $s = 3$ 100.00 100.00 100.00 100.00 100.00 100.00
	$\hat{r} = rank(\hat{\beta}), p = 2$ <div style="display: flex; justify-content: space-around;"> <div> $T = 1000$ $\xi_S \quad \xi_W \quad \xi_{SW}$ </div> <div> $T = 100$ $\xi_S \quad \xi_W \quad \xi_{SW}$ </div> </div> $s \geq 1$ 11.69 0.36 17.35 16.45 0.54 22.82 $s \geq 2$ 100.00 4.98 100.00 100.00 7.80 100.00 $s = 3$ 100.00 100.00 100.00 100.00 100.00 100.00
	$r = 2, p = 2$ <div style="display: flex; justify-content: space-around;"> <div> $T = 1000$ $\xi_S \quad \xi_W \quad \xi_{SW}$ </div> <div> $T = 100$ $\xi_S \quad \xi_W \quad \xi_{SW}$ </div> </div> $s \geq 1$ 100.00 0.40 100.00 99.84 0.49 99.91 $s \geq 2$ 100.00 5.17 100.00 100.00 6.98 100.00 $s = 3$ 100.00 100.00 100.00 100.00 100.00 100.00
	$r = 1, p = 2$ <div style="display: flex; justify-content: space-around;"> <div> $T = 1000$ $\xi_S \quad \xi_W \quad \xi_{SW}$ </div> <div> $T = 100$ $\xi_S \quad \xi_W \quad \xi_{SW}$ </div> </div> $s \geq 1$ 100.00 5.18 100.00 98.00 5.24 98.61 $s \geq 2$ 100.00 100.00 100.00 100.00 97.11 99.96 $s = 3$ 100.00 100.00 100.00 100.00 100.00 100.00
	$\hat{r} = rank(\hat{\beta}), p = 2$ <div style="display: flex; justify-content: space-around;"> <div> $T = 1000$ $\xi_S \quad \xi_W \quad \xi_{SW}$ </div> <div> $T = 100$ $\xi_S \quad \xi_W \quad \xi_{SW}$ </div> </div> $s \geq 1$ 100.00 0.40 100.00 99.84 0.49 99.91 $s \geq 2$ 100.00 5.17 100.00 100.00 6.98 100.00 $s = 3$ 100.00 100.00 100.00 100.00 100.00 100.00

¹ The rejection frequencies are based on 10,000 replications and calculated using asymptotic critical values. The nominal level is fixed at 5%

Table 5: Empirical Rejection Frequencies¹ of the LR tests for WF, $p = 4$

DGP: WF $s = 2$	Estimated Model
$r = 1$ $\alpha = \begin{bmatrix} -0.50 \\ 0.10 \\ 0.20 \end{bmatrix}$ $\beta' = \begin{bmatrix} 1 & 0 & -1 \end{bmatrix}$	$r = 1, p = 4$ <div style="display: flex; justify-content: space-around;"> <div> $T = 1000$ $\xi_S \quad \xi_W \quad \xi_{SW}$ </div> <div> $T = 100$ $\xi_S \quad \xi_W \quad \xi_{SW}$ </div> </div> $s \geq 1$ 5.43 0.18 36.10 8.23 0.82 35.07 $s \geq 2$ 100.00 5.68 100.00 100.00 12.85 99.98 $s = 3$ 100.00 100.00 100.00 100.00 100.00 100.00
	$r = 2, p = 4$ <div style="display: flex; justify-content: space-around;"> <div> $T = 1000$ $\xi_S \quad \xi_W \quad \xi_{SW}$ </div> <div> $T = 100$ $\xi_S \quad \xi_W \quad \xi_{SW}$ </div> </div> $s \geq 1$ 18.56 0.19 57.14 25.35 0.62 58.96 $s \geq 2$ 100.00 5.60 100.00 100.00 11.40 100.00 $s = 3$ 100.00 100.00 100.00 100.00 100.00 100.00
	$\hat{r} = rank(\hat{\beta}), p = 4$ <div style="display: flex; justify-content: space-around;"> <div> $T = 1000$ $\xi_S \quad \xi_W \quad \xi_{SW}$ </div> <div> $T = 100$ $\xi_S \quad \xi_W \quad \xi_{SW}$ </div> </div> $s \geq 1$ 9.87 0.19 40.73 15.66 0.72 42.59 $s \geq 2$ 100.00 5.69 100.00 100.00 12.39 100.00 $s = 3$ 100.00 100.00 100.00 100.00 100.00 100.00
	$r = 2, p = 4$ <div style="display: flex; justify-content: space-around;"> <div> $T = 1000$ $\xi_S \quad \xi_W \quad \xi_{SW}$ </div> <div> $T = 100$ $\xi_S \quad \xi_W \quad \xi_{SW}$ </div> </div> $s \geq 1$ 100.00 0.18 100.00 99.42 0.59 99.95 $s \geq 2$ 100.00 5.67 100.00 100.00 11.57 100.00 $s = 3$ 100.00 100.00 100.00 100.00 100.00 100.00
	$r = 1, p = 4$ <div style="display: flex; justify-content: space-around;"> <div> $T = 1000$ $\xi_S \quad \xi_W \quad \xi_{SW}$ </div> <div> $T = 100$ $\xi_S \quad \xi_W \quad \xi_{SW}$ </div> </div> $s \geq 1$ 100.00 5.78 100.00 99.12 10.15 98.70 $s \geq 2$ 100.00 100.00 100.00 100.00 99.68 99.96 $s = 3$ 100.00 100.00 100.00 100.00 100.00 100.00
	$\hat{r} = rank(\hat{\beta}), p = 4$ <div style="display: flex; justify-content: space-around;"> <div> $T = 1000$ $\xi_S \quad \xi_W \quad \xi_{SW}$ </div> <div> $T = 100$ $\xi_S \quad \xi_W \quad \xi_{SW}$ </div> </div> $s \geq 1$ 100.00 0.18 100.00 99.43 0.65 99.95 $s \geq 2$ 100.00 5.67 100.00 100.00 12.40 100.00 $s = 3$ 100.00 100.00 100.00 100.00 100.00 100.00

¹ The rejection frequencies are based on 10,000 replications and calculated using asymptotic critical values. The nominal level is fixed at 5%.

Table 6: Empirical Rejection Frequencies¹ of the LR tests for MF, $p = 2$

T		Size $\alpha_{3,1} = -0.4$	Power $\alpha_{3,1} = -0.45$	Power $\alpha_{3,1} = -0.5$
ξ_S	100	8.22	15.04	28.29
	1000	5.15	78.20	99.35
ξ_M	100	6.87	19.28	38.93
	1000	5.03	91.42	100

¹ The nominal level is fixed at 5%. The statistics ξ_S and ξ_M use the estimated $\hat{\beta}$ under the assumption of known cointegrating rank $r = 2$.

Table 7: Cointegration Tests

	Max.Eig.Test	95% cv	Trace Test	95% cv
$r = 0$	28.08	21.0	45.96	29.7
$r \leq 1$	14.39	14.1	17.88	15.4
$r \leq 2$	3.48	3.8	3.48	3.8

Table 8: Common Feature Tests

$r = 2$	$-T \ln(1 - \lambda_i)$		df		$Pb > \chi^2_{df}$		$Pb > \chi^2_{df}$	
	ξ_S	ξ_W	ξ_S	ξ_W	ξ_S	ξ_W	ξ_S^{cor}	ξ_W^{cor}
$s \geq 1$	20.51	10.44	12	10	.058	.402	.092	.466
$s \geq 2$	(53.54)	33.83	(26)	22	(.001)	.051	(.004)	.087
$s = 3$	(150.1)	117.3	(42)	36	(< .001)	< .001	(< .001)	< .001




Figure 1: Log Levels of Macro Aggregates




Figure 2: Unique Weak Form Common Cycle and NBER Contraction Periods