

Hartung, Joachim; Argaç, Doğan

**Working Paper**

## A confidence interval to combined univariate economic forecasts

Technical Report, No. 2002,16

**Provided in Cooperation with:**

Collaborative Research Center 'Reduction of Complexity in Multivariate Data Structures' (SFB 475),  
University of Dortmund

*Suggested Citation:* Hartung, Joachim; Argaç, Doğan (2002) : A confidence interval to combined univariate economic forecasts, Technical Report, No. 2002,16, Universität Dortmund,  
Sonderforschungsbereich 475 - Komplexitätsreduktion in Multivariaten Datenstrukturen, Dortmund

This Version is available at:

<https://hdl.handle.net/10419/77163>

**Standard-Nutzungsbedingungen:**

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

**Terms of use:**

*Documents in EconStor may be saved and copied for your personal and scholarly purposes.*

*You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.*

*If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.*

# A Confidence Interval for Combined Univariate Economic Forecasts

JOACHIM HARTUNG AND DOĞAN ARGAÇ

Department of Statistics, University of Dortmund,  
D–44221 Dortmund, Germany.

## Abstract

Research in combining of economic forecasts made by several research institutes on the same economic variable has focused on estimation, hoping that the combined forecast will be improved by taking into account the expert opinions of the institutes. We provide a confidence interval on the combined forecast. A scoring of the individual institutes is proposed using their historical performance in forecasting.

*Key words:* Heterogeneous variances; Expert opinion; Confidence interval.

# 1 Introduction

Consider the situation where one has several forecasts on the same quantity, e. g. several economic research institutes forecast an important economic variable. Interest is in combining the individual forecasts to improve the accuracy of the forecast. Since by way of combining several forecasts one takes into account several expert opinions, one expects a better performance of the combined forecast. However, it is rarely observed that the simply average of the individual forecasts is beaten by more sophisticated methods. Klapper (1998, 1999, 2000) discusses rank-based techniques for combining forecasts. There is little known about confidence intervals on the combined forecast in this area.

In this paper, we give several confidence intervals on the combined forecast which are of approximate nature. The confidence intervals are derived using analysis of variance as a main tool. Some simulation results on the properties of the confidence intervals are provided. Finally, the confidence intervals are applied on German economic data.

## 2 A statistical model for combining forecasts

We consider the following model

$$y_i \sim N(\mu, \alpha_i), \quad (2.1)$$

where  $y_i$  denotes the forecast of the  $i$ -th institute, and  $\mu$  represents the true unknown quantity,  $i = 1, \dots, K$ ,  $K > 2$ . Note that we assume that the variances  $\alpha_i$  are heteroscedastic. Confidence intervals and tests are discussed in a related heteroscedastic ANOVA model by Hartung and Argaç (2002a, 2002b) and Hartung and Knapp (2000).

We estimate the mean  $\mu$  by a weighted average of the individual forecasts  $y_i$ ,

$$y = \sum_{i=1}^K b_i^2 \cdot y_i, \quad (2.2)$$

where  $b_i^2$  denotes the weight which is given to the  $i$ -th institute by some scoring process. We assume that the weights are normed

$$\sum_{i=1}^K b_i^2 = 1, \quad (2.3)$$

and we also assume that  $b_i^2 < 1/2$ , that is we exclude the possibility that one particular institute dominates the rest. Consider now the following quadratic form, see also Hartung, Böckenhoff and Knapp (2002),

$$u_{ib}^2 = b_i^2 \cdot \left( y_i - \sum_{j=1}^K b_j^2 \cdot y_j \right)^2, \quad (2.4)$$

which can be interpreted as a weighted quadratic deviation of each individual forecast from the weighted average of the individual forecasts of each institute. Let  $\beta_i = b_i^2 \cdot \alpha_i$ , and one can derive explicitly the important moments of the quadratic form  $u_{ib}^2$ ,

$$E(u_{ib}^2) = (1 - 2 \cdot b_i^2) \cdot \beta_i + b_i^2 \cdot \sum_{j=1}^K b_j^2 \cdot \beta_j =: e_i(b, \beta), \quad (2.5)$$

$$var(u_{ib}^2) = 2 \cdot e_i(b, \beta)^2, \quad (2.6)$$

$$Cov(u_{ib}^2, u_{jb}^2) = 2 \cdot b_i^2 \cdot b_j^2 \cdot \left( \sum_{k=1}^K b_k^2 \cdot \beta_k - \beta_i - \beta_j \right) =: 2 \cdot e_{ij}(b, \beta)^2. \quad (2.7)$$

We need to estimate the variance of the combined forecast and of each individual forecast, and for this purpose we use quadratic functions of the individual forecasts as estimators. The variance of the combined forecast is estimated with an unbiased positive (PSD–MINQUE) variance estimator which is given by, see also Hartung, Böckenhoff and Knapp (2002),

$$\widehat{\text{var}}(y)_{\text{psd}} = \frac{1}{1 + \sum_{k=1}^K \frac{b_k^4}{1 - 2b_k^2}} \cdot \sum_{i=1}^K \frac{b_i^2}{1 - 2b_i^2} \cdot u_{ib}^2 \quad (2.8)$$

$$= \sum_{i=1}^K d_i \cdot u_{ib}^2. \quad (2.9)$$

The estimator of the variance of each forecast is given by a positive minimum biased variance estimator (PSD–MINQMBE) given by

$$\widehat{\alpha}_i = \frac{\sum_{j=1}^K u_{jb}^2 + \sum_{j=1}^K d_j \cdot u_{jb}^2}{\sum_{j=1}^K b_j^2 \cdot \widetilde{\alpha}_j} \cdot \widetilde{\alpha}_i, \quad (2.10)$$

where

$$\widetilde{\alpha}_i = \frac{1}{b_i^2} \cdot \frac{(1 - b_i^2)^2}{(1 - b_i^2)^4 + b_i^4 \sum_{j \neq i} b_j^4} \cdot u_{ib}^2. \quad (2.11)$$

Note here also that both estimators are given explicitly, see also Hartung, Böckenhoff and Knapp (2002) for a detailed derivation. Under suitable conditions these estimators are consistent, exist always and are unique. Surprisingly, the variance of the combined forecast can be unbiasedly estimated without any information about the variance of the individual forecasts.

We will construct the confidence interval using a pivotal quantity, and for this purpose we have to determine the distribution of the variance estimators. We

will approximate the distribution of the variance estimators by a suitable  $\chi^2$ -distributions using moment matching, see Satterthwaite (1946) and Patnaik (1949). This leads to

$$\nu \cdot \sum_{i=1}^K d_i \cdot u_{ib}^2 / E\left(\sum_{i=1}^K d_i \cdot u_{ib}^2\right) \sim \chi_\nu^2, \quad (2.12)$$

where

$$\nu = 2 \frac{[E(\sum_{i=1}^K d_i \cdot u_{ib}^2)]^2}{var(\sum_{i=1}^K d_i \cdot u_{ib}^2)}. \quad (2.13)$$

Now, this gives us the following pivot (nominator and denominator are independent)

$$\frac{y - \mu}{\sqrt{\sum_{i=1}^K d_i \cdot u_{ib}^2}} \sim t_\nu, \quad (2.14)$$

where

$$\nu = \frac{\left\{ \sum_{i=1}^K d_i \cdot e_i(b, \beta) \right\}^2}{\sum_{i=1}^K d_i^2 \cdot e_i(b, \beta)^2 + \sum_{i=1}^K \sum_{j=1, j \neq i}^K d_i \cdot d_j \cdot e_{ij}(b, \beta)^2}. \quad (2.15)$$

In practice, the unknown quantities have to be replaced by estimators, and here we use  $\hat{\beta}_i = b_i^2 \cdot \hat{\alpha}_i$ .

Now, we are in the position to derive the confidence intervals which are of course of approximate nature only.

$$K_1 : \quad y \mp t_{\nu; 1-\alpha/2} \cdot \sqrt{\sum_{i=1}^K d_i \cdot u_{ib}^2} \quad (2.16)$$

$$K_2 : \quad y \mp u_{1-\alpha/2} \cdot \sqrt{\sum_{i=1}^K d_i \cdot u_{ib}^2} \quad (2.17)$$

$$K_3 : \quad y \mp t_{K-1;1-\alpha/2} \cdot \sqrt{\sum_{i=1}^K d_i \cdot u_{ib}^2} \quad (2.18)$$

The first interval is the approximate interval derived using the t-distributed pivot, the second version is obtained if one replaces the quantile of the t-distribution with the quantile of the standard normal distribution for large degrees of freedom. The last version is obtained if one ignores the heteroscedastic variances. Now, we have to specify the weights  $b_i^2$ . We suggest the following choice of the weights:

$$b_i^2 = \frac{\sum_{j=1}^L (y_{ij} - y_j^w)^{-2}}{\sum_{i=1}^K \sum_{j=1}^L (y_{ij} - y_j^w)^{-2}}, \quad (2.19)$$

where  $y_{ij}$  denotes the forecast of the  $i$ -th institute for some economic variable in the  $j$ -th time period or year and  $y_j^w$  denotes the true realized value in the  $j$ -th time period or year. Hence, each institute is scored taking into account its historical performance in forecasting in the last  $L$  time periods or years. We consider quadratic deviations of each forecast from the true value.

### 3 Monte Carlo results

Since the intervals we constructed are of approximate nature, we conducted a simulation experiment to check their validity with respect to the actual confidence coefficients and lengths of the proposed confidence intervals. We considered  $K = 7$  and  $K = 14$  institutes. The weights  $b_i^2$  are chosen to be equal, in a second scenario the weight of one institute is large compared to the other weights which are chosen to be equal, and finally, the weights are chosen to be almost equal. The variances of the individual forecasts are homoscedastic and heteroscedastic. We paired small variances with small weights and small variances with large weights. We draw samples from the normal and from a centered  $\chi^2$ -distribution with a few degrees of freedom to cover also the case of non-normal observations. The number of repetitions in the simulations is 1000. In the simulations, we give summary statistics on the degrees of freedom of the approximate t-distribution. We report the bias, the standard deviation and the root mean square error (RMSE) of the variance estimators given in (2.10). We provide the empirical confidence level of the intervals at the nominal level of 90% and 95%, the lower and the upper confidence limits and the corresponding widths of the confidence intervals.

The main result is that, in general, the intervals  $K_1$  and  $K_3$  are conservative, the confidence interval  $K_2$  also attains acceptable levels, but might become liberal under certain circumstances.

## 4 Data analysis

We applied the confidence intervals to German economic data. We used the data on GDP from seven major economic research institutes from 1984–1996. The data are described in Klapper (1998). For the first forecast, that is for 1987, we used the data from 1984–1986. The true values are given in the next table:

**Table 1:** True values of GDP.

Year	Value
1987	1.9
1988	3.7
1989	3.3
1990	4.7
1991	3.7
1992	1.6
1993	-1.7
1994	2.4
1995	1.9
1996	1.4

We give the weights  $b_i^2$ , the variance estimators given in (2.10) and (2.11), the variance of the combined forecast given in (2.8), the degrees of freedom of the approximate t-distribution, the combined forecast given in (2.2), and finally, the confidence intervals at nominal level of 90% and 95%.

## Results

```
forecast for the year: 1987
bi^2:
0.036 0.069 0.332 0.036 0.343 0.033 0.151

a_tilde:
1.119 0.085 0.405 0.240 0.507 0.112 0.293

a_hat:
0.995 0.076 0.360 0.213 0.450 0.100 0.260

variance:
0.107

df_hat:
1.447214

forecast(95%):
forecast   K1.l   K1.u   K2.l   K2.u   K3.l   K3.u
2.525   0.452 4.599 0.452 4.599 1.724 3.326

forecast(90%):
forecast   K1.l   K1.u   K2.l   K2.u   K3.l   K3.u
2.525 1.267 3.783 1.987 3.064 1.889 3.161
```

```
forecast for the year: 1988
bi^2:
0.040 0.067 0.241 0.028 0.466 0.048 0.111

a_tilde:
0.533 0.559 0.933 0.091 0.102 0.012 0.051

a_hat:
0.441 0.462 0.772 0.075 0.085 0.010 0.042

variance:
0.049

df_hat:
1.119492

forecast(95%):
forecast K1.l K1.u K2.l K2.u K3.l K3.u
1.706 -0.478 3.89 1.274 2.138 1.167 2.245

forecast(90%):
forecast K1.l K1.u K2.l K2.u K3.l K3.u
1.706 0.541 2.871 1.343 2.069 1.278 2.134

forecast for the year: 1989
bi^2:
0.040 0.067 0.241 0.028 0.464 0.048 0.111
```

```
a_tilde:  
0.005 0.036 0.007 0.194 0.012 0.579 0.006  
  
a_hat:  
0.005 0.034 0.007 0.183 0.011 0.546 0.006  
  
variance:  
0.004  
  
df_hat:  
1.270905  
  
forecast(95%):  
forecast K1.l K1.u K2.l K2.u K3.l K3.u  
2.43 1.931 2.929 2.305 2.556 2.274 2.587  
  
forecast(90%):  
forecast K1.l K1.u K2.l K2.u K3.l K3.u  
2.43 2.145 2.715 2.325 2.536 2.306 2.555  
  
forecast for the year: 1990  
bi^2:  
0.042 0.068 0.240 0.029 0.460 0.048 0.112  
  
a_tilde:  
0.234 0.001 0.002 0.001 0.003 0.059 0.033
```

```
a_hat:  
0.217 0.001 0.002 0.001 0.003 0.055 0.031  
  
variance:  
0.001  
  
df_hat:  
1.454133  
  
forecast(95%):  
forecast K1.l K1.u K2.l K2.u K3.l K3.u  
3.034 2.825 3.243 2.969 3.099 2.953 3.115  
  
forecast(90%):  
forecast K1.l K1.u K2.l K2.u K3.l K3.u  
3.034 2.907 3.161 2.979 3.089 2.969 3.099  
  
forecast for the year: 1991  
bi^2:  
0.044 0.069 0.240 0.030 0.458 0.049 0.112  
  
a_tilde:  
0.017 0.017 0.024 0.146 0.039 0.492 0.168  
  
a_hat:  
0.016 0.016 0.022 0.136 0.036 0.457 0.156
```

```
variance:  
0.013  
  
df_hat:  
1.258859  
  
forecast(95%):  
forecast K1.l K1.u K2.l K2.u K3.l K3.u  
3.373 2.484 4.262 3.153 3.593 3.099 3.647  
  
forecast(90%):  
forecast K1.l K1.u K2.l K2.u K3.l K3.u  
3.373 2.868 3.878 3.189 3.557 3.155 3.591  
  
  
  
forecast for the year: 1992  
bi^2:  
0.082 0.067 0.248 0.029 0.432 0.043 0.099  
  
a_tilde:  
1.074 0.243 0.330 0.001 0.654 0.059 0.032  
  
a_hat:  
1.002 0.227 0.308 0.001 0.611 0.055 0.030  
  
variance:  
0.17
```

```
df_hat:  
1.138552  
  
forecast(95%):  
forecast K1.l K1.u K2.l K2.u K3.l K3.u  
1.966 -1.983 5.914 1.157 2.774 0.957 2.975  
  
forecast(90%):  
forecast K1.l K1.u K2.l K2.u K3.l K3.u  
1.966 -0.161 4.092 1.287 2.644 1.164 2.767  
  
forecast for the year: 1993  
bi^2:  
0.061 0.173 0.300 0.028 0.304 0.034 0.101  
  
a_tilde:  
1.000 0.272 0.005 0.321 0.005 0.964 0.366  
  
a_hat:  
0.917 0.250 0.004 0.295 0.004 0.884 0.335  
  
variance:  
0.013  
  
df_hat:  
3.085219
```

```

forecast(95%):
forecast   K1.l   K1.u     K2.l   K2.u   K3.l   K3.u
-0.053 -0.407 0.301 -0.274 0.169 -0.329 0.224

forecast(90%):
forecast   K1.l   K1.u     K2.l     K2.u   K3.l   K3.u
-0.053 -0.316 0.21    -0.239   0.133 -0.272 0.167

forecast for the year: 1994
bi^2:
0.063 0.173 0.299 0.028 0.303 0.034 0.100

a_tilde:
1.176 0.307 0.392 0.237 0.484 0.001 0.019

a_hat:
1.029 0.269 0.343 0.207 0.423 0.001 0.016

variance:
0.085

df_hat:
1.819994

forecast(95%):
forecast   K1.l   K1.u     K2.l     K2.u   K3.l   K3.u
0.525 -0.859 1.909    -0.047   1.098 -0.19   1.24

```

```

forecast(90%):
forecast   K1.l   K1.u      K2.l      K2.u   K3.l   K3.u
    0.525 -0.388 1.439     0.045     1.006 -0.042 1.093

forecast for the year: 1995
bi^2:
0.063 0.173 0.298 0.029 0.302 0.034 0.100

a_tilde:
1.340 0.012 0.286 0.009 0.016 0.590 0.104

a_hat:
1.209 0.011 0.258 0.008 0.014 0.532 0.094

variance:
0.03

df_hat:
1.614924

forecast(95%):
forecast   K1.l   K1.u      K2.l      K2.u   K3.l   K3.u
    3.095  2.145 4.044     2.753     3.436  2.668 3.521

forecast(90%):
forecast   K1.l   K1.u      K2.l      K2.u   K3.l   K3.u
    3.095  2.493 3.696     2.808     3.381  2.756 3.433

```

```
forecast for the year: 1996
bi^2:
0.167 0.154 0.264 0.026 0.268 0.030 0.090

a_tilde:
0.826 0.001 0.011 0.544 0.081 0.548 0.453

a_hat:
0.734 0.001 0.009 0.483 0.072 0.487 0.402

variance:
0.028

df_hat:
1.904998

forecast(95%):
forecast   K1.l   K1.u      K2.l      K2.u   K3.l   K3.u
       1.78  1.029  2.531      1.454    2.106  1.373  2.187

forecast(90%):
forecast   K1.l   K1.u      K2.l      K2.u   K3.l   K3.u
       1.78  1.277  2.283      1.506    2.054  1.457  2.103
```

## Acknowledgements

The support of the German Research Community (DFG, Sonderforschungsbereich 475) is gratefully acknowledged.

## References

- Hartung, J. and Argaç, D. (2002a) Generalizing the Welch test to non-zero hypotheses on the variance component in the one-way random effects model under variance heterogeneity. *Statistics*, in press.
- Hartung, J. and Argaç, D. (2002b) Confidence intervals on the among group variance component in an unbalanced and heteroscedastic one-way random effects model. *Statistics & Decisions*, to appear.
- Hartung, J., Böckenhoff, A. and Knapp, G. (2002) Generalized Cochran-Wald statistics in combining of experiments. *Journal of Statistical Planning and Inference*, in press.
- Hartung, J. and Knapp, G. (2000) Confidence intervals for the between group variance in the unbalanced one-way random effects model of analysis of variance. *Journal of Statistical Computation and Simulation* **65**, 311–323.
- Klapper, M. (1998) Combining German macro economic forecasts using rank-based techniques. Technical Report 19/1998, SFB 475, Department of Statistics, University of Dortmund.
- Klapper, M. (1999) Combination of forecasts using rank-based techniques. *ASA Proceedings of the Business and Economic Statistics Section*, 197–202.

- Klapper, M. (2000) The combination of forecasts using rank based techniques. Josef Eul Verlag, Lohmar.
- Patnaik, P. B. (1949) The non-central  $\chi^2$ - and F-distributions and their applications. *Biometrika* **36**, 202–232.
- Satterthwaite, F. E. (1946) An approximate distribution of estimates of variance components. *Biometrics Bulletin* **2**, 110–114.
- Teilprojekt B3 (2001) *Personal communication*. SFB 475, Department of Statistics, University of Dortmund.
- Teilprojekt B4 (2001) *Personal communication*. SFB 475, Department of Statistics, University of Dortmund.

## Simulation results for K=7 (normal distribution)

```
empirical level of CI (normal distr.)  
  
bi^2 : 0.143 0.143 0.143 0.143 0.143 0.143 0.143  
variances : 2 2 2 2 2 2 2  
repetitions: 1000  
  
df:  
df_mean:  
3.275  
  
df_std:  
0.919  
  
df_quantiles:  
0%   25%   50%   75%  100%  
1.063 2.559 3.292 3.933 6.035  
  
bias(variance estimators):  
0.0849 -0.0310 -0.0120  0.0015 -0.0831 -0.1153  0.2338  
  
std(variance estimators):  
3.2120 2.7277 2.8178 3.1083 2.8052 2.6790 3.2227  
  
RMSE:  
3.2131 2.7279 2.8178 3.1083 2.8064 2.6814 3.2312
```

RMSE.total:

20.5861

conf.level(95%):

K1	K2	K3
99.8	92.5	97.7

width:

K1	K2	K3
0.452	0.27	0.337

conf.level(90%):

K1	K2	K3
96.5	84.7	92.2

width:

K1	K2	K3
0.329	0.226	0.267

empirical level of CI (normal distr.)

bi^2 : 0.143 0.143 0.143 0.143 0.143 0.143 0.143

variances : 2 4 6 8 10 12 14

repetitions: 1000

df:

df\_mean:

3.033

```
df_std:  
0.947  
  
df_quantiles:  
0%    25%    50%    75%   100%  
1.083  2.336  2.973  3.669  5.992  
  
bias(variance estimators):  
-0.0242  0.2960  0.2216  0.2688 -0.1576 -0.4110  1.0639  
  
std(variance estimators):  
2.7411  6.0115  9.0319 11.0558 13.9781 16.0869 21.2612  
  
RMSE:  
2.7412  6.0187  9.0347 11.0591 13.9790 16.0922 21.2878  
  
RMSE.total:  
80.2126  
  
conf.level(95%):  
K1      K2      K3  
99.8    92.7   98.6  
  
width:  
K1      K2      K3  
0.986  0.541  0.675
```

```
conf.level(90%):
  K1      K2      K3
97.3    84.5  92.4

width:
  K1      K2      K3
0.699   0.454  0.536

empirical level of CI (normal distr.)

bi^2 :  0.49 0.085 0.085 0.085 0.085 0.085 0.085
variances :  2 2 2 2 2 2 2
repetitions: 1000

df:
df_mean:
1.209

df_std:
0.556

df_quantiles:
  0%    25%    50%    75%   100%
  1 1.007 1.019 1.085 5.709

bias(variance estimators):
-0.0051  0.2343  0.1295  0.1702 -0.0069  0.1486  0.1629
```

```
std(variance estimators):
2.8543 3.1827 2.9185 2.9561 2.7382 3.1266 3.2277

RMSE:
2.8543 3.1913 2.9214 2.9610 2.7382 3.1301 3.2318

RMSE.total:
21.0282

conf.level(95%):
  K1      K2      K3
99.9    97.3   98.8

width:
  K1      K2      K3
5.686   0.921   1.15

conf.level(90%):
  K1      K2      K3
98.8    94.5   97.1

width:
  K1      K2      K3
2.861   0.773   0.913
```

empirical level of CI (normal distr.)

bi^2 : 0.49 0.085 0.085 0.085 0.085 0.085 0.085

variances : 2 4 6 8 10 12 14

repetitions: 1000

df:

df\_mean:

1.416

df\_std:

0.688

df\_quantiles:

0% 25% 50% 75% 100%

1.001 1.029 1.091 1.43 5.284

bias(variance estimators)(variance estimators):

-0.0938 0.2995 -0.0674 0.5797 -0.0439 -0.6084 0.0709

std(variance estimators):

2.8515 6.0550 8.7045 11.9132 13.6275 16.0967 20.2054

RMSE:

2.8530 6.0624 8.7047 11.9272 13.6275 16.1082 20.2056

RMSE.total:

79.4887

```
conf.level(95%):
```

K1	K2	K3
99.8	95	98.6

```
width:
```

K1	K2	K3
5.093	0.914	1.142

```
conf.level(90%):
```

K1	K2	K3
99.1	89.3	94.8

```
width:
```

K1	K2	K3
2.628	0.767	0.907

empirical level of CI (normal distr.)

bi^2 : 0.49 0.085 0.085 0.085 0.085 0.085 0.085 0.085

variances : 14 12 10 8 6 4 2

repetitions: 1000

```
df:
```

```
df_mean:
```

1.136

```
df_std:
```

0.411

```
df_quantiles:  
0%    25%    50%    75%   100%  
1 1.004 1.01 1.044 4.424  
  
bias(variance estimators):  
-1.2423 0.4273 0.4654 0.2545 0.1702 0.6977 0.6605  
  
std(variance estimators):  
18.3193 17.8309 15.3039 11.0667 8.5818 6.7522 3.5595  
  
RMSE:  
18.3613 17.8361 15.3110 11.0696 8.5835 6.7881 3.6202  
  
RMSE.total:  
81.5698  
  
conf.level(95%):  
K1      K2      K3  
99.9    97.9   99.5  
  
width:  
K1      K2      K3  
14.673 2.332 2.911  
  
conf.level(90%):  
K1      K2      K3  
99.6    96.4   97.9
```

width:

K1	K2	K3
7.347	1.957	2.312

empirical level of CI (normal distr.)

bi^2 : 0.163 0.163 0.163 0.128 0.128 0.128 0.128

variances : 2 2 2 2 2 2 2

repetitions: 1000

df:

df\_mean:

3.055

df\_std:

0.934

df\_quantiles:

0% 25% 50% 75% 100%

1.153 2.372 3.019 3.647 5.935

bias(variance estimators):

-0.1225 0.1601 0.1224 0.1058 0.0082 -0.1449 0.1971

std(variance estimators):

2.7621 2.9168 2.8044 2.8081 2.8088 2.5920 3.1021

RMSE:

2.7648 2.9212 2.8071 2.8101 2.8088 2.5960 3.1084

RMSE.total:

19.8164

conf.level(95%):

K1	K2	K3
99.6	91.7	97.8

width:

K1	K2	K3
0.513	0.284	0.354

conf.level(90%):

K1	K2	K3
97.1	83.4	91.3

width:

K1	K2	K3
0.365	0.238	0.281

empirical level of CI (normal distr.)

bi^2 : 0.163 0.163 0.163 0.128 0.128 0.128 0.128

variances : 2 4 6 8 10 12 14

repetitions: 1000

```
df:  
df_mean:  
3.164  
  
df_std:  
0.94  
  
df_quantiles:  
0%    25%    50%    75%   100%  
1.119  2.485  3.105  3.783  5.749  
  
bias(variance estimators):  
0.0408  0.0216  0.6190 -0.0224 -0.3049  0.3677 -0.2523  
  
std(variance estimators):  
2.9522  5.8892  9.6222 11.1032 13.5997 16.8349 20.1514  
  
RMSE:  
2.9525  5.8892  9.6421 11.1032 13.6031 16.8389 20.1530  
  
RMSE.total:  
80.1819  
  
conf.level(95%):  
K1      K2      K3  
99.7    92.1   97.9
```

```
width:  
    K1      K2      K3  
0.879  0.504  0.63  
  
conf.level(90%):  
    K1      K2      K3  
96.4   86.7  91.9  
  
width:  
    K1      K2      K3  
0.633  0.423  0.5  
  
  
empirical level of CI (normal distr.)  
  
bi^2 :  0.163 0.163 0.163 0.128 0.128 0.128 0.128  
variances :  14 12 10 8 6 4 2  
repetitions:  1000  
  
df:  
df_mean:  
2.653  
  
df_std:  
0.889  
  
df_quantiles:  
 0%    25%    50%    75%  100%  
1.108 1.993 2.579 3.255 5.986
```

```
bias(variance estimators):
0.4523 0.7456 -0.2133 0.0324 0.2825 0.0681 0.0174

std(variance estimators):
23.0188 17.4906 12.9867 11.2607 8.7093 5.8048 2.7633

RMSE:
23.0233 17.5065 12.9885 11.2607 8.7139 5.8052 2.7634

RMSE.total:
82.0614

conf.level(95%):
K1      K2      K3
99.6    93.3   98.7

width:
K1      K2      K3
1.303   0.613   0.765

conf.level(90%):
K1      K2      K3
97.5    86.5   93.1

width:
K1      K2      K3
0.881   0.514   0.608
```

## Simulation results for K=14 (normal distribution)

```
empirical level of CI (normal distr.)  
  
bi^2 :  0.071 0.071 0.071 0.071 0.071 0.071 0.071 0.071 0.071  
        0.071 0.071 0.071 0.071 0.071  
variances :  2 2 2 2 2 2 2 2 2 2 2 2 2 2  
repetitions:  1000  
  
df:  
df_mean:  
5.748  
  
df_std:  
1.424  
  
df_quantiles:  
 0%   25%   50%   75%   100%  
1.681 4.792 5.743 6.799 10.331  
  
bias(variance estimators):  
 0.0618 -0.0999  0.0779 -0.0179  0.0227  0.0256 -0.1004  0.1329  
-0.0441 -0.0642  0.0477  0.0600  0.1040  0.0923  
  
std(variance estimators):  
 2.8247 2.8607 2.9056 2.8659 2.7845 2.8692 2.6589 2.9051 2.9288  
2.7215 2.8942 2.8175 2.9424 2.9869
```

RMSE:

2.8253	2.8624	2.9067	2.8659	2.7845	2.8694	2.6608	2.9081	2.9291
2.7223	2.8946	2.8181	2.9442	2.9883				

RMSE.total:

39.9798
---------

conf.level(95%):

K1	K2	K3
98	94.7	96

width:

K1	K2	K3
0.131	0.101	0.111

conf.level(90%):

K1	K2	K3
94.7	87	91.2

width:

K1	K2	K3
0.103	0.085	0.091

empirical level of CI (normal distr.)

bi^2 : 0.071 0.071 0.071 0.071 0.071 0.071 0.071 0.071 0.071 0.071  
0.071 0.071 0.071 0.071 0.071

variances : 2 4 6 8 10 12 14 2 4 6 8 10 12 14

repetitions: 1000

df:

df\_mean:

5.097

df\_std:

1.48

df\_quantiles:

0% 25% 50% 75% 100%  
1.502 4.067 5.026 6.071 10.179

bias(variance estimators):

-0.0497 -0.3323 -0.2174 -0.2721 0.1960 0.4987 -0.0362 0.0259  
0.0075 0.1093 0.6663 0.4455 0.1394 0.2580

std(variance estimators):

2.7687 4.8413 8.4074 10.8751 14.8004 16.9099 19.2347 2.6014  
5.7870 8.5633 11.8345 14.6627 17.8034 19.9322

RMSE:

2.7692 4.8527 8.4103 10.8785 14.8017 16.9172 19.2347 2.6015  
5.7870 8.5640 11.8532 14.6695 17.8040 19.9339

```
RMSE.total:  
159.0772  
  
conf.level(95%):  
K1      K2      K3  
99.3    95.5   97.2  
  
width:  
K1      K2      K3  
0.275   0.201  0.222  
  
conf.level(90%):  
K1      K2      K3  
96.3    89.3   92.3  
  
width:  
K1      K2      K3  
0.213   0.169  0.182  
  
empirical level of CI (normal distr.)  
  
bi^2 :  0.49 0.039 0.039 0.039 0.039 0.039 0.039 0.039 0.039 0.039  
        0.039 0.039 0.039 0.039 0.039  
variances :  2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2  
repetitions: 1000
```

```
df:  
df_mean:  
1.074  
  
df_std:  
0.403  
  
df_quantiles:  
0%    25%    50%    75%   100%  
1 1.001 1.002 1.008 6.745  
  
bias(variance estimators):  
0.0039  0.2393  0.0639  0.2580  0.2457  0.1169  0.0545  0.0699  
0.0032  0.0379  0.1534  0.1163  0.2099 -0.0010  
  
std(variance estimators):  
2.7053 3.1131 2.9087 3.2083 3.2643 2.9553 2.8958 3.1724 2.9733  
2.9048 3.2233 2.9696 3.1931 2.8789  
  
RMSE:  
2.7053 3.1223 2.9094 3.2187 3.2735 2.9576 2.8964 3.1731 2.9733  
2.9051 3.2270 2.9719 3.2000 2.8789  
  
RMSE.total:  
42.4123  
  
conf.level(95%):  
K1      K2      K3  
99.9    99.2   99.5
```

```
width:  
      K1      K2      K3  
5.988    0.93  1.026
```

```
conf.level(90%):  
      K1      K2      K3  
99.7   98.8 99.1
```

```
width:  
      K1      K2      K3  
2.981  0.781 0.841
```

empirical level of CI (normal distr.)

```
bi^2 :  0.49 0.039 0.039 0.039 0.039 0.039 0.039 0.039 0.039  
        0.039 0.039 0.039 0.039 0.039  
variances :  2 4 6 8 10 12 14 2 4 6 8 10 12 14  
repetitions: 1000
```

```
df:  
df_mean:  
1.215
```

```
df_std:  
0.76
```

```
df_quantiles:  
0%   25%   50%   75% 100%  
1 1.003 1.008 1.042 7.84  
  
bias(variance estimators):  
-0.0080  0.5225  0.1500 -0.3747 -0.2556 -0.1500  0.1654  0.0679  
0.2000 -0.1521 -0.2405  0.5913  0.1039  0.1053  
  
std(variance estimators):  
2.8348  6.3810  7.9408 10.9674 13.8247 16.6863 19.8173  3.0228  
5.7593  8.1164 10.7338 14.9750 17.5106 20.4718  
  
RMSE:  
2.8349  6.4023  7.9422 10.9738 13.8271 16.6870 19.8180  3.0235  
5.7628  8.1179 10.7365 14.9867 17.5109 20.4720  
  
RMSE.total:  
159.0957  
  
conf.level(95%):  
K1      K2      K3  
99.6    98.4   98.9  
  
width:  
K1      K2      K3  
5.818   0.92   1.014
```

conf.level(90%):

K1	K2	K3
99.3	96.6	97.5

width:

K1	K2	K3
2.909	0.772	0.831

empirical level of CI (normal distr.)

bi^2 : 0.49 0.039 0.039 0.039 0.039 0.039 0.039 0.039 0.039 0.039

0.039 0.039 0.039 0.039 0.039

variances : 14 12 10 8 6 4 2 14 12 10 8 6 4 2

repetitions: 1000

df:

df\_mean:

1.066

df\_std:

0.401

df\_quantiles:

0% 25% 50% 75% 100%

1 1 1.001 1.005 7.254

```
bias(variance estimators):
-0.5070 -0.0808  2.0567  0.3658  0.5928  0.7026  0.7820  0.3977
 1.1400  0.8301  0.8617  0.8067  0.4196  0.8316

std(variance estimators):
18.8285 15.9962 16.9769 11.4816  9.6429  6.8297  3.8885 20.0774
18.9652 14.8103 12.4895  9.3268  5.7532  3.9510

RMSE:
18.8353 15.9964 17.1010 11.4874  9.6611  6.8657  3.9663 20.0813
18.9994 14.8336 12.5192  9.3616  5.7685  4.0375

RMSE.total:
169.5144

conf.level(95%):
  K1      K2      K3
 100    99.3  99.3

width:
  K1      K2      K3
 15.49    2.4  2.646

conf.level(90%):
  K1      K2      K3
 99.7   98.9  99
```

```
width:  
    K1      K2      K3  
7.706  2.014  2.169
```

```
empirical level of CI (normal distr.)
```

```
bi^2 :  0.082 0.082 0.082 0.082 0.082 0.082 0.064 0.064 0.064  
        0.064 0.064 0.064 0.064 0.064  
variances :  2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2  
repetitions: 1000
```

```
df:  
df_mean:  
5.189
```

```
df_std:  
1.534
```

```
df_quantiles:  
    0%    25%    50%    75%   100%  
1.32  4.089  5.087  6.169  9.834
```

```
bias(variance estimators):  
0.0183  0.2387 -0.0449 -0.0247  0.0637  0.0782 -0.0954  0.0967  
0.0945  0.0123  0.1122 -0.0044  0.1358 -0.0229
```

```
std(variance estimators):  
2.7978 3.2477 2.7921 2.8415 2.8609 2.8032 2.7845 2.9860 2.8381  
2.7730 2.9123 2.7994 2.9603 2.9343
```

RMSE:

```
2.7979 3.2565 2.7925 2.8416 2.8616 2.8043 2.7861 2.9876 2.8397  
2.7731 2.9145 2.7994 2.9634 2.9344
```

RMSE.total:

```
40.3525
```

conf.level(95%):

K1	K2	K3
99	93.5	96.8

width:

K1	K2	K3
0.144	0.106	0.117

conf.level(90%):

K1	K2	K3
95	85.4	89.2

width:

K1	K2	K3
0.112	0.089	0.096

empirical level of CI (normal distr.)

bi^2 : 0.082 0.082 0.082 0.082 0.082 0.082 0.064 0.064 0.064  
0.064 0.064 0.064 0.064 0.064

variances : 2 4 6 8 10 12 14 2 4 6 8 10 12 14

repetitions: 1000

df:

df\_mean:

4.913

df\_std:

1.568

df\_quantiles:

0% 25% 50% 75% 100%  
1.424 3.773 4.87 5.938 10.294

bias(variance estimators):

0.0021 0.1168 0.1399 -0.0487 0.0748 0.4408 -1.8022 -0.0412  
-0.1266 -0.2446 0.0432 -0.5136 -0.0758 -0.0226

std(variance estimators):

2.6662 5.8799 8.8537 10.5506 13.9117 18.5455 16.5929 2.8618  
5.8081 8.3145 11.2268 12.7527 17.8100 21.0710

RMSE:

2.6662 5.8811 8.8548 10.5507 13.9119 18.5507 16.6904 2.8621  
5.8094 8.3181 11.2268 12.7630 17.8102 21.0710

```
RMSE.total:  
156.9666  
  
conf.level(95%):  
K1      K2      K3  
99.4    95.5   97.6  
  
width:  
K1      K2      K3  
0.287   0.203  0.223  
  
conf.level(90%):  
K1      K2      K3  
97.1    89.9   92.7  
  
width:  
K1      K2      K3  
0.219   0.17   0.183
```

empirical level of CI (normal distr.)

```
bi^2 : 0.082 0.082 0.082 0.082 0.082 0.082 0.064 0.064 0.064  
        0.064 0.064 0.064 0.064 0.064  
variances : 14 12 10 8 6 4 2 14 12 10 8 6 4 2  
repetitions: 1000
```

```
df:  
df_mean:  
4.594  
  
df_std:  
1.457  
  
std_quantiles:  
0% 25% 50% 75% 100%  
1.317 3.526 4.569 5.552 9.163  
  
bias(variance estimators):  
1.9064 -0.5692 0.0767 0.0588 -0.2077 0.2297 0.0106 -0.5260  
-0.5235 0.9935 0.4286 -0.1073 -0.0044 0.1365  
  
std(variance estimators):  
22.1807 16.6132 13.3367 11.3082 8.4413 5.7210 2.6936 19.5792  
16.6148 15.5968 11.2331 8.4723 5.6918 2.9436  
  
RMSE:  
22.2624 16.6230 13.3369 11.3083 8.4439 5.7256 2.6937 19.5863  
16.6230 15.6284 11.2413 8.4730 5.6918 2.9467  
  
RMSE.total:  
160.5844  
  
conf.level(95%):  
K1 K2 K3  
98.9 94.1 96.9
```

```
width:  
    K1      K2      K3  
0.313  0.216  0.238
```

```
conf.level(90%):  
    K1      K2      K3  
95.8   86.8  90.7
```

```
width:  
    K1      K2      K3  
0.238  0.181  0.195
```

## Simulation results for K=7 (centered $\chi^2$ -distribution)

empirical level of CI (Chisquare)

bi^2 : 0.143 0.143 0.143 0.143 0.143 0.143 0.143

df : 1 1 1 1 1 1 1

repetitions: 1000

df:

df\_mean:

3.359

df\_std:

1.507

df\_quantiles:

0% 25% 50% 75% 100%

1.038 2.005 3.29 4.601 6.818

bias (variance estimators):

-0.4009 0.3490 -0.0869 -0.2389 0.1519 -0.1593 -0.0323

std (variance estimators):

4.2679 7.0795 6.3895 6.3215 9.8191 6.0000 6.6633

RMSE:

4.2867 7.0881 6.3900 6.3260 9.8202 6.0021 6.6633

RMSE.total:

46.5766

conf.level(95%):

K1	K2	K3
92.2	83	89.3

width:

K1	K2	K3
0.563	0.241	0.301

conf.level(90%):

K1	K2	K3
84.1	76.5	82.6

width:

K1	K2	K3
0.367	0.202	0.239

empirical level of CI (Chisquare)

bi^2 : 0.143 0.143 0.143 0.143 0.143 0.143 0.143

df : 1 2 3 4 5 6 7

repetitions: 1000

df:

df\_mean:

2.939

```
df_std:  
0.944  
  
df_quantiles:  
 0%   25%   50%   75% 100%  
1.033 2.257 2.906 3.612 5.754  
  
bias (variance estimators):  
-0.0471 -0.3609 -0.0926 -1.1107 -0.1310  1.0688 -0.7785  
  
std (variance estimators):  
8.7001  9.3991 13.6465 12.5846 17.9840 26.8254 21.7252  
  
RMSE:  
8.7003  9.4061 13.6468 12.6335 17.9845 26.8467 21.7391  
  
RMSE.total:  
110.957  
  
conf.level(95%):  
  K1      K2      K3  
99.3    89.7  96.6  
  
width:  
  K1      K2      K3  
1.028  0.515  0.643
```

```
conf.level(90%):
```

K1	K2	K3
94.8	83.2	89.3

```
width:
```

K1	K2	K3
0.707	0.433	0.511

```
empirical level of CI (Chisquare)
```

```
bi^2 : 0.49 0.085 0.085 0.085 0.085 0.085 0.085
```

```
df : 1 1 1 1 1 1 1
```

```
repetitions: 1000
```

```
df:
```

```
df_mean:
```

```
1.147
```

```
df_std:
```

```
0.447
```

```
df_quantiles:
```

0%	25%	50%	75%	100%
----	-----	-----	-----	------

1.001	1.006	1.018	1.061	5.776
-------	-------	-------	-------	-------

```
bias (variance estimators):
```

0.1116	-0.1553	0.3741	-0.0544	0.0611	0.1048	-0.0784
--------	---------	--------	---------	--------	--------	---------

```
std (variance estimators):
8.5120 5.6313 8.1074 7.3199 6.4857 8.6732 5.3924

RMSE:
8.5127 5.6334 8.1161 7.3201 6.4860 8.6738 5.3930

RMSE.total:
50.135

conf.level(95%):
  K1      K2      K3
99.7    97.5   98.7

width:
  K1      K2      K3
4.95   0.804  1.004

conf.level(90%):
  K1      K2      K3
99.1    95.2   97.3

width:
  K1      K2      K3
2.493   0.675  0.797
```

empirical level of CI (Chisquare)

bi^2 : 0.49 0.085 0.085 0.085 0.085 0.085 0.085

df : 1 2 3 4 5 6 7

repetitions: 1000

df:

df\_mean:

1.393

df\_std:

0.649

df\_quantiles:

0% 25% 50% 75% 100%

1.001 1.051 1.129 1.376 4.829

bias (variance estimators):

-0.3588 -0.3678 0.1325 1.1266 0.0367 0.6134 0.1670

std (variance estimators):

4.9605 9.5840 13.1359 19.1095 17.9362 27.2881 27.0886

RMSE:

4.9734 9.5911 13.1366 19.1426 17.9363 27.2950 27.0891

RMSE.total:

119.1641

```
conf.level(95%):
```

K1	K2	K3
99.4	92.9	97.1

```
width:
```

K1	K2	K3
4.144	0.779	0.973

```
conf.level(90%):
```

K1	K2	K3
98.3	85.4	92.9

```
width:
```

K1	K2	K3
2.165	0.654	0.772

empirical level of CI (Chisquare)

bi^2 : 0.49 0.085 0.085 0.085 0.085 0.085 0.085

df : 7 6 5 4 3 2 1

repetitions: 1000

df:

df\_mean:

1.127

df\_std:

0.374

df\_quantiles:

0%	25%	50%	75%	100%
1	1.003	1.011	1.049	4.384

bias (variance estimators):

0.0238 1.6984 1.7597 0.4762 0.9604 1.2631 1.0971

std (variance estimators):

25.0118 26.8882 22.1192 14.8699 14.1669 13.1838 11.3971

RMSE:

25.0119 26.9418 22.1891 14.8775 14.1994 13.2442 11.4498

RMSE.total:

127.9136

conf.level(95%):

K1	K2	K3
100	98.4	99.4

width:

K1	K2	K3
14.735	2.345	2.927

conf.level(90%):

K1	K2	K3
99.7	95.9	98.3

width:

K1	K2	K3
7.381	1.968	2.325

empirical level of CI (Chisquare)

bi^2 : 0.163 0.163 0.163 0.128 0.128 0.128 0.128

df : 1 1 1 1 1 1 1

repetitions: 1000

df:

df\_mean:

3.064

df\_std:

1.246

df\_quantiles:

0% 25% 50% 75% 100%

1.034 1.922 3.008 4.083 6.005

bias (variance estimators):

0.4249 0.3321 -0.2552 -0.3396 0.2818 -0.1030 -0.0455

```
std (variance estimators):
9.1910 9.8628 5.2837 4.4549 7.3616 7.5874 8.6069

RMSE:
9.2009 9.8684 5.2899 4.4678 7.3670 7.5881 8.6070

RMSE.total:
52.389

conf.level(95%):
  K1      K2      K3
95.2    82.2   90.4

width:
  K1      K2      K3
0.635   0.256   0.319

conf.level(90%):
  K1      K2      K3
85.4    75.8   81.5

width:
  K1      K2      K3
0.406   0.214   0.253
```

empirical level of CI (Chisquare)

bi^2 : 0.163 0.163 0.163 0.128 0.128 0.128 0.128

df : 1 2 3 4 5 6 7

repetitions: 1000

df:

df\_mean:

3.088

df\_std:

1.014

df\_quantiles:

0% 25% 50% 75% 100%

1.081 2.297 3.086 3.823 5.903

bias (variance estimators):

0.0201 -0.4296 0.8007 -1.4969 0.6285 0.2590 -1.0156

std (variance estimators):

6.2827 7.6358 16.2171 11.8507 21.4044 29.1257 20.8245

RMSE:

6.2828 7.6479 16.2368 11.9448 21.4137 29.1268 20.8493

RMSE.total:

113.5021

```
conf.level(95%):  
  K1      K2      K3  
 99.3    89.9  95.8
```

```
width:  
  K1      K2      K3  
 0.941   0.481  0.601
```

```
conf.level(90%):  
  K1      K2      K3  
 93.5    82.6  89.9
```

```
width:  
  K1      K2      K3  
 0.651   0.404  0.477
```

#### empirical level of CI (Chisquare)

```
bi^2 :  0.163 0.163 0.163 0.128 0.128 0.128 0.128  
df :  7 6 5 4 3 2 1  
repetitions: 1000
```

```
df:  
df_mean:  
2.536
```

```
df_std:  
0.856  
  
df_quantiles:  
 0%   25%   50%   75% 100%  
1.035 1.875 2.463 3.059 5.407  
  
bias (variance estimators):  
0.6707 0.3385 0.3536 0.2057 0.4185 -0.2960 0.3251  
  
std (variance estimators):  
28.0134 21.8782 22.0783 17.5604 14.2897 11.3791 9.6280  
  
RMSE:  
28.0214 21.8808 22.0812 17.5616 14.2959 11.3829 9.6335  
  
RMSE.total:  
124.8573  
  
conf.level(95%):  
 K1      K2      K3  
99.8    91.7    98  
  
width:  
 K1      K2      K3  
1.381  0.603  0.753
```

```
conf.level(90%):
```

K1	K2	K3
97.8	83.2	91

```
width:
```

K1	K2	K3
0.911	0.506	0.598

## Simulation results for K=14 (centered $\chi^2$ -distribution)

```
empirical level of CI (Chisquare)

bi^2 :  0.071 0.071 0.071 0.071 0.071 0.071 0.071 0.071 0.071
         0.071 0.071 0.071 0.071 0.071
df :  1 1 1 1 1 1 1 1 1 1 1 1 1 1
repetitions: 1000

df:
df_mean:
4.697

df_std:
2.708

std_quantiles:
 0%   25%   50%   75%   100%
1.082 2.549 4.02 6.323 12.344

bias (variance estimators):
0.1161 -0.1868 0.1990 0.2943 0.0806 -0.2641 -0.0478 0.0153
-0.0703 -0.2341 0.1063 0.0230 0.0951 -0.3336

std (variance estimators):
8.0426 5.0594 11.5179 11.1385 8.3347 5.6707 6.7102 6.4682
8.5697 5.4499 6.6220 7.8788 7.6240 6.9652
```

RMSE:

8.0435	5.0629	11.5196	11.1424	8.3351	5.6769	6.7104	6.4682
8.5700	5.4549	6.6228	7.8789	7.6245	6.9732		

RMSE.total:

106.0831

conf.level(95%):

K1	K2	K3
91.1	86.9	89.5

width:

K1	K2	K3
0.177	0.093	0.103

conf.level(90%):

K1	K2	K3
86.1	81.8	83.9

width:

K1	K2	K3
0.123	0.078	0.084

empirical level of CI (Chisquare)

bi^2 : 0.071 0.071 0.071 0.071 0.071 0.071 0.071 0.071 0.071 0.071  
0.071 0.071 0.071 0.071 0.071

df : 1 2 3 4 5 6 7 1 2 3 4 5 6 7

repetitions: 1000

df:

df\_mean:

4.632

df\_std:

1.675

df\_quantiles:

0% 25% 50% 75% 100%

1.256 3.363 4.602 5.915 11.366

bias (variance estimators):

0.1096 -0.2397 -0.3474 0.7535 0.3557 0.2619 -0.8744 0.2431

0.0070 0.2901 -0.4068 0.5526 0.4600 -0.7067

std (variance estimators):

8.3947 9.9169 11.7673 19.9040 19.2651 23.9713 24.3522 9.5834

10.5655 16.9995 18.5879 26.9952 26.6536 26.2663

RMSE:

8.3954 9.9198 11.7724 19.9182 19.2684 23.9727 24.3679 9.5865

10.5655 17.0020 18.5924 27.0008 26.6575 26.2758

RMSE.total:

253.2953

conf.level(95%):

K1	K2	K3
97.5	93	94.7

width:

K1	K2	K3
0.302	0.196	0.217

conf.level(90%):

K1	K2	K3
93.6	86.8	90.1

width:

K1	K2	K3
0.225	0.165	0.178

empirical level of CI (Chisquare)

bi^2 : 0.49 0.039 0.039 0.039 0.039 0.039 0.039 0.039 0.039 0.039  
0.039 0.039 0.039 0.039 0.039

df : 1 1 1 1 1 1 1 1 1 1 1 1

repetitions: 1000

```
df:  
df_mean:  
1.067  
  
df_std:  
0.401  
  
df_quantiles:  
0%   25%   50%   75%   100%  
1 1.001 1.003 1.007 7.839  
  
bias (variance estimators):  
-0.1052 0.5221 0.0120 -0.2031 0.0908 0.0649 -0.0410 -0.0910  
-0.1046 0.0694 0.2752 0.4972 0.1763 0.7091  
  
std (variance estimators):  
6.0611 12.8983 5.6311 5.5860 6.7873 6.9731 7.4593 6.7069  
6.2153 5.9869 7.1581 8.0291 7.9955 13.3629  
  
RMSE:  
6.0620 12.9088 5.6311 5.5897 6.7879 6.9734 7.4594 6.7075  
6.2162 5.9873 7.1634 8.0444 7.9975 13.3817  
  
RMSE.total:  
106.9104  
  
conf.level(95%):  
K1      K2      K3  
99.9    99.1   99.2
```

```
width:  
      K1      K2      K3  
5.048  0.785  0.866
```

```
conf.level(90%):  
      K1      K2      K3  
99.7   98.4  98.8
```

```
width:  
      K1      K2      K3  
2.514  0.659  0.71
```

#### empirical level of CI (Chisquare)

```
bi^2 :  0.49 0.039 0.039 0.039 0.039 0.039 0.039 0.039 0.039  
        0.039 0.039 0.039 0.039 0.039  
df :  1 2 3 4 5 6 7 1 2 3 4 5 6 7  
repetitions: 1000
```

```
df:  
df_mean:  
1.143
```

```
df_std:  
0.564
```

```
df_quantiles:  
0%   25%   50%   75%  100%  
1 1.006 1.011 1.029 7.869  
  
bias (variance estimators):  
-0.3721  0.0963  0.5852 -0.0775  1.1802 -0.5679 -1.1394 -0.0250  
0.6170  0.1463 -0.3930 -0.2446  1.0341  0.2649  
  
std (variance estimators):  
5.0853 10.3437 13.7329 15.2421 22.5533 21.3652 22.7735 6.1834  
11.9011 14.0667 16.4910 16.8193 28.4762 23.4579  
  
RMSE:  
5.0989 10.3442 13.7453 15.2423 22.5841 21.3728 22.8019 6.1835  
11.9171 14.0675 16.4957 16.8211 28.4949 23.4594  
  
RMSE.total:  
228.6286  
  
conf.level(95%):  
K1      K2      K3  
99.8    98.9   99.3  
  
width:  
K1      K2      K3  
4.823  0.767  0.845
```

```
conf.level(90%):
```

K1	K2	K3
99.4	98	98.7

```
width:
```

K1	K2	K3
2.416	0.644	0.693

```
empirical level of CI (Chisquare)
```

```
bi^2 : 0.49 0.039 0.039 0.039 0.039 0.039 0.039 0.039 0.039 0.039  
0.039 0.039 0.039 0.039 0.039
```

```
df : 7 6 5 4 3 2 1 7 6 5 4 3 2 1
```

```
repetitions: 1000
```

```
df:
```

```
df_mean:
```

```
1.047
```

```
df_std:
```

```
0.312
```

```
df_quantiles:
```

```
0% 25% 50% 75% 100%
```

```
1 1 1.001 1.004 6.369
```

```
bias (variance estimators):
1.5573 2.4006 0.9161 0.9757 0.3818 0.8075 0.9463 -0.2010
0.4716 0.8197 0.1363 0.9830 1.1622 0.5475

std (variance estimators):
29.1755 29.7733 21.3582 18.7988 12.6091 10.0351 6.4408 25.6372
21.5856 20.5570 15.4451 13.7668 11.4395 5.9937

RMSE:
29.2170 29.8699 21.3779 18.8241 12.6149 10.0675 6.5100 25.6380
21.5908 20.5733 15.4457 13.8018 11.4984 6.0187

RMSE.total:
243.0479

conf.level(95%):
K1      K2      K3
99.9    99.5   99.7

width:
K1      K2      K3
16.319  2.527   2.786

conf.level(90%):
K1      K2      K3
99.8    99.2   99.2
```

```
width:  
    K1      K2      K3  
8.118  2.121  2.284
```

```
empirical level of CI (Chisquare)
```

```
bi^2 :  0.082 0.082 0.082 0.082 0.082 0.082 0.064 0.064 0.064  
        0.064 0.064 0.064 0.064 0.064  
df :  1 1 1 1 1 1 1 1 1 1 1 1 1 1 1  
repetitions: 1000
```

```
df:  
df_mean:  
4.587
```

```
df_std:  
2.422
```

```
df_quantiles:  
 0%   25%   50%   75%  100%  
1.088 2.539 4.076 6.543 11.72
```

```
bias (variance estimators):  
-0.2142 -0.0946 -0.0837  0.1369 -0.2922 -0.0499 -0.2427  0.0269  
-0.3982  0.1428 -0.0146 -0.3882 -0.1507  0.0678
```

```
std (variance estimators):
5.1084 5.9582 6.3212 6.6632 5.2090 5.7063 7.2672 10.2402
4.2995 7.3587 6.6523 8.7553 6.6601 6.5954

RMSE:
5.1129 5.9590 6.3218 6.6646 5.2172 5.7066 7.2713 10.2402
4.3179 7.3601 6.6523 8.7639 6.6618 6.5957

RMSE.total:
92.8454

conf.level(95%):
K1      K2      K3
91.1    86.7   88.6

width:
K1      K2      K3
0.182   0.096  0.106

conf.level(90%):
K1      K2      K3
85.6    81.6   84.1

width:
K1      K2      K3
0.126   0.08   0.087
```

empirical level of CI (Chisquare)

bi^2 : 0.082 0.082 0.082 0.082 0.082 0.082 0.064 0.064 0.064  
0.064 0.064 0.064 0.064 0.064

df : 1 2 3 4 5 6 7 1 2 3 4 5 6 7

repetitions: 1000

df:

df\_mean:

4.515

df\_std:

1.666

df\_quantiles:

0% 25% 50% 75% 100%  
1.308 3.25 4.513 5.785 8.958

bias (variance estimators):

0.5394 -0.1500 -0.1449 -0.5557 -0.2473 0.8610 -0.6024 0.3752  
-0.4230 0.6961 0.5339 0.3338 -0.6406 1.3720

std (variance estimators):

10.3663 9.8152 13.2064 15.0877 18.9813 28.3542 22.8702 9.1346  
7.7884 16.2544 21.3924 20.6429 21.1151 35.6943

RMSE:

10.3803 9.8164 13.2072 15.0980 18.9829 28.3672 22.8782 9.1423  
7.7999 16.2693 21.3991 20.6456 21.1248 35.7206

```
RMSE.total:  
250.8317  
  
conf.level(95%):  
K1      K2      K3  
97.9    92.9  95.8  
  
width:  
K1      K2      K3  
0.315   0.2  0.221  
  
conf.level(90%):  
K1      K2      K3  
94.4    86.3  88.9  
  
width:  
K1      K2      K3  
0.233   0.168 0.181
```

```
empirical level of CI (Chisquare)  
  
bi^2 :  0.082 0.082 0.082 0.082 0.082 0.082 0.064 0.064 0.064  
        0.064 0.064 0.064 0.064 0.064  
df :  7 6 5 4 3 2 1 7 6 5 4 3 2 1  
repetitions: 1000
```

```
df:  
df_mean:  
4.336  
  
df_std:  
1.627  
  
df_quantiles:  
0%   25%   50%   75% 100%  
1.217 3.079 4.327 5.503 9.646  
  
bias (variance estimators):  
0.8380 0.4453 0.6705 -0.1161 0.7208 -0.0810 -0.2193 -0.5960  
0.5554 -1.0513 1.3427 0.5341 -0.2364 0.1986  
  
std (variance estimators):  
28.5691 25.2573 23.1433 17.6375 18.3094 12.3501 5.1858 23.4702  
23.9417 15.7911 26.2190 18.4062 10.8732 10.2991  
  
RMSE:  
28.5813 25.2612 23.1531 17.6379 18.3236 12.3504 5.1905 23.4777  
23.9481 15.8261 26.2534 18.4140 10.8758 10.3011  
  
RMSE.total:  
259.5941  
  
conf.level(95%):  
K1      K2      K3  
97.8    91.9   94.3
```

```
width:  
    K1      K2      K3  
0.346  0.213  0.234
```

```
conf.level(90%):  
    K1      K2      K3  
92.6   85.9  88.9
```

```
width:  
    K1      K2      K3  
0.253  0.179  0.192
```