

Krämer, Walter

Working Paper

The robustness of the F-test to spatial autocorrelation among regression disturbances

Technical Report, No. 2002,56

Provided in Cooperation with:

Collaborative Research Center 'Reduction of Complexity in Multivariate Data Structures' (SFB 475), University of Dortmund

Suggested Citation: Krämer, Walter (2002) : The robustness of the F-test to spatial autocorrelation among regression disturbances, Technical Report, No. 2002,56, Universität Dortmund, Sonderforschungsbereich 475 - Komplexitätsreduktion in Multivariaten Datenstrukturen, Dortmund

This Version is available at:

<http://hdl.handle.net/10419/77194>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.

The robustness of the F-test to spatial autocorrelation among regression disturbances¹

by

Walter Krämer

Fachbereich Statistik, Universität Dortmund

D-44221 Dortmund, Germany

walterk@statistik.uni-dortmund.de

Version December 2002

Abstract

It is shown that the null distribution of the F-test in a linear regression is rather non-robust to spatial autocorrelation among the regression disturbances. In particular, the true size of the test tends to either zero or unity when the spatial autocorrelation coefficient approaches the boundary of the parameter space.

Key words: F-test, size, spatial autocorrelation.

¹Research supported by Deutsche Forschungsgemeinschaft (DFG) under SFB 475. I am grateful to Anurag Banerjee for providing the GAUSS-routine for the computation of exact rejection probabilities (available in the internet under <http://www.american.deu/academics.depts/cas/econ/gaussres/Gausidx.htm>)

1 Introduction and summary

The null distribution of the F-test under nonspherical errors has concerned applied statisticians and econometricians for many decades. Let

$$y = X\beta + u = X^{(1)}\beta^{(1)} + X^{(2)}\beta^{(2)} + u \quad (1)$$

be the model under test, where y and u are $T \times 1$, X is $T \times K$ and nonstochastic of rank $K < T$, β is $K \times 1$, and the disturbance vector u is multivariate normal with mean zero and (possibly) nonscalar covariance matrix V . The design matrix is partitioned into $X^{(1)}(T \times q)$ and $X^{(2)}(T \times (K - q))$ and the null hypothesis to be tested is $H_0 : \beta^{(1)} = b^{(1)}$.

The standard F-test rejects for large values of

$$F = \frac{(\tilde{u}'\tilde{u} - \hat{u}'\hat{u})/q}{\hat{u}'\hat{u}/(T - K)}, \quad (2)$$

where $\hat{u} = y - X\hat{\beta}$, $\hat{\beta} = (X'X)^{-1}X'y$, $\tilde{u} = y - X^{(1)}b^{(1)} - X^{(2)}\tilde{\beta}^{(2)}$, $\tilde{\beta}^{(2)} = (X^{(2)'}X^{(2)})^{-1}X^{(2)'}(y - X^{(1)}b^{(1)})$. It has a central F-distribution with q and $T - K$ degrees of freedom, given H_0 and $V = \sigma^2I$, and the problem to be studied here is the robustness of this null distribution to deviations from $V = \sigma^2I$.

So far, this problem has been addressed mainly for given specific forms of V , with various bounds for the size of the test being derived as the design matrix X is allowed to vary across all $T \times K$ -matrices of rank K (Vinod 1976, Kiviet 1980). Below we take a different approach, following Krämer (1989) and Krämer et al. (1990), by fixing X and letting V vary across a certain range of disturbance covariance matrices. In particular, we allow the disturbance vector u to be generated by the spatial autoregressive scheme

$$u = \rho Wu + \varepsilon, \quad (3)$$

where ε is a $T \times 1$ normal random vector with mean zero and scalar covariance matrix σ_ε^2I , and W is some known $T \times T$ -matrix of nonnegative spatial weights with $w_{ii} = 0$ ($i = 1, \dots, T$). Such patterns of dependence are often entertained

when the objects under study are positioned in some "space", whether geographical or sociological (in some social network, say) and account for spillovers from one unit to its neighbors, whichever way "neighborhood" may be defined. They date back to Whittle (1954) and has become quite popular in econometric recently. See Anselin and Florax (1995), Kelejian and Prucha (2001) or Anselin (2001) for a convenient survey of this literature.

The coefficient ρ in (3) measures the degree of correlation, which can be both positive and negative. Below we focus on the empirically more relevant case of positive disturbance correlation, where

$$0 \leq \rho < \frac{1}{\lambda_{max}}$$

and where λ_{max} is the Frobenius-root of W (i.e. the unique positive real eigenvalue such that $\lambda_{max} \geq |\lambda_i|$ for arbitrary eigenvalues λ_i). The disturbances are then given by

$$u = (I - \rho W)^{-1} \varepsilon, \tag{4}$$

so $V := Cov(u) = \sigma_\varepsilon^2 [(I - \rho W)(I - \rho W)']^{-1}$ and $V = \sigma_\varepsilon^2 I$ whenever $\rho = 0$.

Below we consider the null distribution of the F-test for $\rho \neq 0$. This is shown to be extremely non-robust, with the size of the test tending to either zero or unity as $\rho \rightarrow 1/\lambda_{max}$. The same limits are also obtained for the power of the test. Which of these limits obtains is easily seen from X and W , which are both observed and known. Therefore, our result provides an easy guide to the interpretation of both a significant and insignificant F-test when there is possible spatial correlation among the regression disturbances: If H_0 is rejected, and X and W are such that the size of the test tends to unity, an error of the first kind has to be suspected. And if H_0 is not rejected, and X and W are such that the size and the power of the test tends to zero, one has to beware of an error of the second kind.

2 The null distribution of the F-test as auto-correlation increases

We first rewrite the test statistic (2) as

$$F = \frac{u'(M^{(2)} - M)u/q}{u'Mu/(T - K)}, \quad (5)$$

where $M = I - X(X'X)^{-1}X'$ and $M^{(2)} = I - X^{(2)}(X^{(2)'}X^{(2)})^{-1}X^{(2)'}$. Let $F_{q,T-K}^\alpha$ be the $(1 - \alpha)$ quantile of the central F-distribution with q and $T - K$ degrees of freedom, respectively, where α is the nominal size of the test. Then

$$\begin{aligned} P(F \geq F_{q,T-K}^\alpha) &= P(u'(M^{(2)} - M)u - \frac{q}{T - K}F_{q,T-K}^\alpha u'Mu \geq 0) \\ &= P(u'(M^{(2)} - dM)u \geq 0) \\ &\quad (\text{where } d = 1 + \frac{q}{T - K}F_{q,T-K}^\alpha) \\ &= P(\varepsilon'V^{1/2}(M^{(2)} - dM)V^{1/2}\varepsilon \geq 0) \\ &\quad (\text{where } \varepsilon = V^{-1/2}u \sim N(0, I)) \\ &= P\left(\sum_{i=1}^T \lambda_i \xi_i^2 \geq 0\right) \\ &= P\left((1 - \rho\lambda_{max})^2 \sum_{i=1}^T \lambda_i \xi_i^2 \geq 0\right), \end{aligned} \quad (6)$$

where the ξ_i^2 are iid $\chi_{(1)}^2$ and the λ_i are the eigenvalues of $V^{1/2}(M^{(2)} - dM)V^{1/2}$, and therefore also of $V(M^{(2)} - dM)$.

The limiting rejection probability as $\rho \rightarrow 1/\lambda_{max}$ depends upon the limiting behavior of $(1 - \rho\lambda_{max})^2 V$. Let

$$W = \sum_{i=1}^T \lambda_i \omega_i \omega_i' \quad (7)$$

be the spectral decomposition of W , with the eigenvalues λ_i in increasing order. Then

$$V = \sum_{i=1}^T \frac{1}{(1 - \rho\lambda_i)^2} \omega_i \omega_i' \quad (8)$$

is the spectral decomposition of V , and

$$\lim_{\rho \rightarrow 1/\lambda_{max}} (1 - \rho\lambda_i)^2 V = \omega'_T \omega_T, \quad (9)$$

a matrix of rank 1. Therefore, all limiting eigenvalues of $(1 - \rho\lambda_{max})^2 V (M^{(2)} - dM)$ are zero except one, which is given by

$$\text{tr}(\omega_T \omega_T' (M^{(2)} - dM)) = \omega_T' (M^{(2)} - dM) \omega_T. \quad (10)$$

If $\omega_T' (M^{(2)} - dM) \omega_T$ is positive, the rejection probability tends to one. If $\omega_T' (M^{(2)} - dM) \omega_T$ is negative, the rejection probability tends to zero.

As $\omega_T' (M^{(2)} - dM) \omega_T$ is known, it is easy to determine in practice which of these cases obtains. For illustration, figure 1 shows both an example where rejection probability tends to one, and an example where the rejection probability tends to zero. The weight matrix is 25×25 and is derived from a regular 5×5 lattice using the queen criterion, which assigns a weight of one to all cells immediately surrounding a given cell, and zero otherwise. The case where the rejection probability tends to zero corresponds to a 25×2 X -matrix with a first column of ones, and a second column given by $(1, 2, 3, \dots, 25)'$, where we test whether the coefficient of the second regressor is zero. The case where the rejection probability tends to one corresponds to an X -matrix where the second column was generated as $nid(0, 1)$ variables.

(figure 1 here)

The figure shows that convergence to zero of the rejection probability need not be monotone and that an X -matrix which induces a limiting rejection probability of zero might, for certain regions of the parameter space, engender higher rejection probabilities than X -matrices where the rejection probability eventually tends to one.

Whether a limit of the rejection probability of zero or one obtains depends on the interplay of the design matrix X , the weight matrix W , and the nominal size of the test. For $T = 25$, a nominal size of 5%, a weight matrix defined by the

queen criterion, and an X matrix given by a first column of ones, and a second column of $nid(0, 1)$ variables (whose significance is to be tested), a Monte Carlo experiment was performed with 1000 independent runs. We obtained a limiting rejection probability of one in about 10% of all cases. Ceteris paribus the incidence of this irregular behavior diminishes as sample size increases, but it is possible to find examples where the rejection probability tends to one also for larger samples and for different types of spatial weights.

3 Spatial autocorrelation and the power of the test

Let $g := X^{(1)}\beta^{(1)} - X^{(1)}b^{(1)}$. Under the alternative, $g \neq 0$, and the expression (5) becomes

$$F = \frac{[u'(M^{(2)} - M)u + 2g'M^{(2)}u + g'M^{(2)}g]/q}{u'Mu(T - K)}, \quad (11)$$

with rejection occurring if and only if

$$\begin{aligned} & (1 - \rho\lambda_{max})^2 u'[M^{(2)} - (1 + \frac{q}{T - K} F_{q, T-k}^\alpha)M]u \\ & + (1 - \rho\lambda_{max})^2 [2g'M^{(2)}u] \\ & + (1 - \rho\lambda_{max})^2 g'M^{(2)}g \geq 0. \end{aligned} \quad (12)$$

Since the last two terms in expression (12) are easily seen to tend to zero as $\rho \rightarrow 1/\lambda_{max}$, the power of the test has the same limiting behavior as the size.

References

- Anselin, L. and Florax, R. (eds.) (1995):** *New directions in spatial econometrics*. Berlin (Springer).
- Anselin, L. (2001):** "Rao's score test in spatial econometrics." *Journal of Statistical Planning and Inference* 97, 113 – 139.
- Kelejian, H. and Prucha, I. (2001):** "On the asymptotic distribution of the Moran I-test statistic with applications." *Journal of Econometrics* 104, 219 – 257.
- Kiviet, J.F. (1980):** "Effects of ARMA errors on tests for regression coefficients: comments on Vinod's article, improved and additional results." *Journal of the American Statistical Association*, 75, 333 – 358.
- Krämer, W. (1989):** "On the robustness of the F-test to autocorrelation among disturbances." *Economics Letters* 30, 37 – 40.
- Krämer, W.; Kiviet, J. and Breitung, J. (1990):** "The null distribution of the F -test in the linear regression model with autocorrelated disturbances." *Statistica* 50, 503 – 509.
- Vinod, H.D. (1976):** "Effects of ARMA errors on the significance tests for regression coefficients." *Journal of the American Statistical Association* 71, 929 – 933.
- Whittle, P. (1954):** "On stationary processes in the plane." *Biometrika* 41, 434 – 449.




Figure 1: Rejection probability of the F-Test