

Krämer, Walter; Sibbertsen, Philipp

**Working Paper**

## Testing for structural change in the presence of long memory

Technical Report, No. 2000,31

**Provided in Cooperation with:**

Collaborative Research Center 'Reduction of Complexity in Multivariate Data Structures' (SFB 475), University of Dortmund

*Suggested Citation:* Krämer, Walter; Sibbertsen, Philipp (2000) : Testing for structural change in the presence of long memory, Technical Report, No. 2000,31, Universität Dortmund, Sonderforschungsbereich 475 - Komplexitätsreduktion in Multivariaten Datenstrukturen, Dortmund

This Version is available at:

<https://hdl.handle.net/10419/77279>

**Standard-Nutzungsbedingungen:**

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

**Terms of use:**

*Documents in EconStor may be saved and copied for your personal and scholarly purposes.*

*You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.*

*If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.*

# TESTING FOR STRUCTURAL CHANGE IN THE PRESENCE OF LONG MEMORY<sup>1</sup>

by

**Walter Krämer and Philipp Sibbertsen**

Fachbereich Statistik, Universität Dortmund, D-44221 Dortmund, Germany

Version June 2000

## **Abstract**

We derive the limiting null distributions of the standard and OLS-based CUSUM-tests for structural change of the coefficients of a linear regression model in the context of long memory disturbances. We show that both tests behave fundamentally different in a long memory environment, as compared to short memory, and that long memory is easily mistaken for structural change when standard critical values are employed.

## **1 Introduction and Summary**

It is by now well known that long memory and structural change are easily confused (Lobato and Sawin 1997, Engle and Smith 1999, Granger and Hyung 1999, Diebold and Inoue 1999 and many others). Therefore it is of interest to know about both the stochastic properties of procedures for detecting and measuring long memory when there is only structural change, and of the performance of tests for structural change when there is only long memory.

---

<sup>1</sup>Research supported by Deutsche Forschungsgemeinschaft; we are grateful to A. Zeileis for expert computational assistance.

While the former problem has attracted considerable attention, there has been rather little work on the latter (Hidalgo and Robinson 1996, Wright 1998). Below we consider the behaviour of the standard and the OLS-based CUSUM-tests, whose limiting distributions are well understood in the context of various regressor-sequences and iid- or short memory disturbances (Krämer et al. 1988, Ploberger and Krämer 1992, 1996). As shown by Wright (1998) for the OLS-based CUSUM-test and the special case of polynomial regressors, these limiting distributions are not robust to departures from short memory - in fact, the OLS-based CUSUM-test has an asymptotic size of unity. The present paper allows for more general regressor sequences also covers the conventional CUSUM-test based on recursive residuals as well. We show that Wright's results concerning the behaviour under  $H_0$  essentially go through with more general regressors, and that similar results hold for the standard CUSUM-test. This is a rather negative result which confirms related theorems from the structural-change-mistaken-for-long-memory-literature: Similar to structural change being mistaken for long memory, long memory is likewise easily mistaken for structural change, and it remains an open problem to efficiently discriminate between the two<sup>2</sup>.

## 2 Two unpleasant theorems

We consider the standard linear regression model

$$y_t = \beta' x_t + \varepsilon_t, \quad (t = 1, \dots, T) \quad (1)$$

with nonstochastic, fixed regressors  $x_t$  and stationary mean zero disturbances  $\varepsilon_t$ . We assume that

$$\frac{1}{T} \sum_{t=1}^T x_t \rightarrow c < \infty \quad \text{and} \quad (2)$$

---

<sup>2</sup>There do exist solutions for some special cases, such as Künsch's (1986) procedure to discriminate between long memory and monotonic trends, but a general treatment of this problem is still missing.

$$\frac{1}{T} \sum_{t=1}^T x_t x_t' \rightarrow Q \text{ (finite, nonsingular).} \quad (3)$$

These are standard assumptions in linear regression large sample asymptotics; they exclude trending data, which require separate treatment and proofs which differ from the ones below.

We are concerned with testing the model (1) against the alternative of unspecified structural change in the regression coefficients  $\beta$ . We consider first the OLS-based CUSUM-test, as proposed by Ploberger and Krämer (1992). This test rejects the null hypothesis of no structural change for large values of

$$TS := \sup_{0 < \lambda < 1} |C_T(\lambda)|, \quad \text{where} \quad (4)$$

$$C_T(\lambda) := T^{-\frac{1}{2}} \hat{\sigma}_\varepsilon^{-1} \sum_{t=1}^{[T\lambda]} e_t, \quad (5)$$

and where  $e_t := y_t - x_t' \hat{\beta}$  are the OLS-residuals from (1).

The limiting null distribution of TS is well known for white noise and short memory disturbances. Our first theorem extends these results to stationary long memory disturbances, where the  $\varepsilon_t$  follow a stationary ARFIMA(p,d,q)-process:

$$E(\varepsilon_t \varepsilon_{t-k}) = L(k) k^{-d}, \quad (6)$$

$L(k)$  slowly varying,  $0 < d < 1/2$ .

**Theorem 1** *In the regression model (1), with disturbances as in (6) we have*

$$T^{-d} C_T(\lambda) \rightarrow B_d(\lambda) - c' Q^{-1} \xi(\lambda), \quad (7)$$

where  $B_d(\lambda)$  is fractional Brownian Motion with self-similarity parameter  $d$  and  $\xi(\lambda) \sim N(0, \lambda \sigma_\varepsilon^2 Q)$ .

PROOF: We have

$$C_T(\lambda) = T^{-\frac{1}{2}}\hat{\sigma}_\varepsilon^{-1} \left\{ \sum_{t=1}^{[T\lambda]} \varepsilon_t - \sum_{t=1}^{[T\lambda]} x'_t \left( \sum_{t=1}^T x_t x'_t \right)^{-1} \sum_{t=1}^T x_t \varepsilon_t \right\}, \text{ so} \quad (8)$$

$$\begin{aligned} T^{-d}C_T(\lambda) &= \left\{ T^{-d+\frac{1}{2}} z_{[T\lambda]} \right. \\ &\quad \left. - T^{-d+\frac{1}{2}} \sum_{t=1}^{[T\lambda]} x'_t \left( \sum_{t=1}^T x_t x'_t \right)^{-1} \sum_{t=1}^T x_t \varepsilon_t \right\} / \hat{\sigma}_\varepsilon, \end{aligned} \quad (9)$$

where  $z_t = z_{t-1} + \varepsilon_t$ ,  $z_0 = 0$ . In view of

$$T^{-d-\frac{1}{2}} z_{[T\lambda]} \rightarrow \sigma_\varepsilon B(\lambda) \quad (\text{see e.g. Marmol 1995}), \quad (10)$$

$$\frac{1}{T} \sum_{t=1}^{[T\lambda]} x_t \rightarrow \lambda c, \quad (11)$$

$$\left( \frac{1}{T} \sum_{t=1}^{[T\lambda]} x_t x'_t \right)^{-1} \rightarrow \lambda^{-1} Q^{-1}, \quad (12)$$

$$T^{-d-\frac{1}{2}} \sum_{t=1}^{[T\lambda]} x_t \varepsilon_t \rightarrow \xi(\lambda) \quad (\text{see Giraitis and Taqqu 1998}), \quad (13)$$

and

$$\hat{\sigma}_\varepsilon^2 = \sum_{t=1}^T \frac{e_t^2}{T} = \sum_{t=1}^T \frac{\varepsilon_t^2}{T} + o_P(1) \rightarrow \sigma_\varepsilon^2$$

the limiting relationship (7) follows.  $\square$

From (7), it is immediately seen that  $TS \xrightarrow{P} \infty$  under  $H_0$ , so the OLS-based CUSUM-test is extremely non-robust to long-memory disturbances, in the sense that long memory is easily mistaken for structural change when conventional critical values are employed.

Next we consider the standard CUSUM-test based on recursive residuals

$$\tilde{e}_t = \frac{y_t - x'_t \hat{\beta}^{(t-1)}}{f_t}, \quad \hat{\beta}^{(t-1)} = \left( X^{(t-1)'} X^{(t-1)} \right)^{-1} X^{(t-1)'} y^{(t-1)} \quad (14)$$

$$f_t = \left( 1 + x'_t \left( X^{(t-1)'} X^{(t-1)} \right)^{-1} x_t \right)^{\frac{1}{2}} \quad (t = K + 1, \dots, T), \quad (15)$$

where the superscript  $t - 1$  means that only observations  $1, \dots, t - 1$  are used. It rejects for large values of

$$S_T = \sup_{0 < \lambda < 1} W_T(\lambda)/(1 + 2\lambda). \quad (16)$$

where

$$W_T(\lambda) := T^{-\frac{1}{2}} \hat{\sigma}_\varepsilon^{-1} \sum_{t=K+1}^{[T\lambda]} \tilde{\varepsilon}_t. \quad (17)$$

**Theorem 2** *In the regression model (1), with disturbances as in (6) we have*

$$T^{-d} W_T(\lambda) \rightarrow B_d(\lambda), \quad (18)$$

where again  $B_d(\lambda)$  is fractional Brownian Motion with self-similarity parameter  $d$ .

PROOF: Following Krämer et al. (1988), we write  $W_T(\lambda)$  as

$$W_T(\lambda) = \frac{1}{\sqrt{T}} \sum_{t=K+1}^{[T\lambda]} \varepsilon_t - \sum_{t=K+1}^{[T\lambda]} (\hat{\beta}^{(t-1)} - \beta)' x_t. \quad (19)$$

Let  $Q_j := \frac{1}{T} \sum_{i=1}^j x_i x_i'$ . First we show that

$$\max_{K \leq t \leq T} \left\| \left( \hat{\beta}^{(t)} - \beta \right) - \sum_{j=K}^t [(y_j - x_j' \beta) x_j] Q_j^{-1} \right\| = o_p \left( T^{d+\frac{1}{2}} (\ln \ln T)^{\frac{1}{2}} \right). \quad (20)$$

Let  $S_t := \sum_{j=1}^t (y_j - x_j' \beta) x_j$ . By the law of the iterated logarithm for the sums of long memory Gaussian random variables we have for some slowly varying function  $L(T)$

$$\max_{1 \leq t \leq T} \frac{S_t}{\left[ \frac{2}{(d+\frac{1}{2})2(d+\frac{1}{2})-1} T^{2(d+\frac{1}{2})} L(T) \ln \ln(T) \right]^{\frac{1}{2}}} = O_p(1), \quad (21)$$

so (20) follows directly from Lemma 3.1 of Jureckova and Sen (1984).

Combining (19) and (20) gives

$$\begin{aligned} & \frac{1}{\sqrt{T}} \sum_{t=K+1}^{[T\lambda]} (y_t - x'_t \hat{\beta}^{(t-1)}) \\ &= \frac{1}{\sqrt{T}} \sum_{t=1}^{[T\lambda]} \sum_{j=1}^t c_{ij} (y_j - x'_j \beta) + o(T^{(d+\frac{1}{2})} \ln \ln T)^{\frac{1}{2}}, \end{aligned} \quad (22)$$

where

$$c_{ij} = \begin{cases} -x'_j Q_{(i-1)}^{-1} x_i & i > j \\ 1 & i = j \\ 0 & i < j \end{cases} \quad (23)$$

(see also Sibbertsen, 2000). In view of a result by Sen (1984) that

$$\sum_{j \leq i} c_{ij}^2 = 1 + o\left(\frac{1}{i}\right) \quad (24)$$

and theorem 5.1 of Taqqu (1975), the theorem now follows from (22).  $\square$

Theorem 2 shows that the null distribution of the standard CUSUM-test tends to infinity as well, so the standard CUSUM-test has likewise an asymptotic size of unity.

### 3 Some finite sample Monte Carlo evidence

Figure 1 below gives the empirical rejection rates, using 1000 runs and standard critical values from the iid-disturbance case, for the OLS-based CUSUM-test. When the disturbances are in fact ARFIMA(0,d,0). It confirms our theoretical results: rejection rates increase with  $d$  and sample size, and produce misleading evidence even for small  $d$  and  $T$ .




Figure 1: Empirical rejection probability of OLS-based CUSUM-test




Figure 2: Empirical rejection probability of standard CUSUM-test

Figure 2 gives the corresponding empirical rejection rates for the standard CUSUM-test. Not surprisingly, the empirical size is not as far off the mark as for the OLS-based CUSUM-test, but the test is misleading here as well.



## References

- Diebold, F.S. and Inoue, A. (1999):** "Long memory and structural change." Stern School of Business discussion paper.
- Engle, R.F. and Smith, A. (1999):** "Stochastic permanent breaks." *Review of Economics and Statistics*
- Giraitis, L. and Taqqu, M. (1998):** "Convergence of normalized quadratic forms." To appear in *Journal of Statistical Planning and Inference* 80, 15 – 35.
- Granger, C.W.J. and Hyung, N. (1999):** "Occasional structural breaks and long memory." UCSD discussion paper 99 -114.
- Hidalgo, J. and Robinson, P.M. (1996):** "Testing for structural change in a long-memory environment." *Journal of Econometrics* 70, 159 – 174.
- Jureckova, J. and Sen, P.K. (1984):** "On adaptive scale-equivariant M-estimators in linear models." *Statistics and Decision* 2, 31 – 46.
- Kunsch, H. (1986):** "Discrimination between monotonic trends and long range dependence." *Journal of Applied Probability* 23, 1025 – 1030.
- Krämer, W.; Ploberger, W. and Alt, R. (1988):** "Testing for structural change in dynamic models." *Econometrica* 56, 1355 – 1369.
- Lobato, I.N. and Savin, N.E. (1997):** "Real and spurious lon-memory properties of stock market data (with discussion and reply)." *Journal of Business and Economic Statistics* 16, 261 – 283.
- Marmol, F. (1995):** "Spurious regression between I(d)-processes." *Journal of Time Seies Analysis* 16, 313 – 321.
- Ploberger, W. and Krämer, W. (1992):** "The CUSUM-test with OLS-residuals." *Econometrica* 60, 271 – 286.
- Ploberger, W. and Krämer, W. (1996):** "A trend-resistant test for structural change based on OLS-residuals." *Journal of Econometrics* 70, 175 – 185.
- Sibbertsen, Ph. (2000):** "Robust CUSUM-M-test in the presence of long memory disturbances." *Technical Report* 19/2000, SFB 475, Universität Dortmund.
- Taqqu, M.S. (1975):** "Weak convergence of fractional Brownian Motion to the Rosenblatt process." *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete* 31, 287 – 302.

**Wright, J. (1998):** "Testing for a structural break at unknown date with long-memory disturbances." *Journal of Time Series Analysis* 19, 369 – 379.