

Wenzel, Thomas

Working Paper

Using different Pitman-closeness techniques for the linear combination of multivariate forecasts

Technical Report, No. 1999,18

Provided in Cooperation with:

Collaborative Research Center 'Reduction of Complexity in Multivariate Data Structures' (SFB 475),
University of Dortmund

Suggested Citation: Wenzel, Thomas (1999) : Using different Pitman-closeness techniques for the linear combination of multivariate forecasts, Technical Report, No. 1999,18, Universität Dortmund, Sonderforschungsbereich 475 - Komplexitätsreduktion in Multivariaten Datenstrukturen, Dortmund

This Version is available at:

<https://hdl.handle.net/10419/77298>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.

Using different Pitman-closeness techniques for the linear combination of multivariate forecasts

Thomas Wenzel

*Department of Statistics, University of Dortmund, Vogelpothsweg 87,
D-44221 Dortmund, Germany*

Abstract: We specify the Pitman-closeness criterion for the evaluation of multivariate forecasts in three categories. This is done closely to the definition of covariance adjustment techniques analysed in other articles. We also apply the Pitman-closeness techniques to an example dealing with German economic data.

Key words: Pitman-closeness, multivariate forecasting methods, covariance adjustment.

Acknowledgement: This work was supported by the Deutsche Forschungsgemeinschaft, Sonderforschungsbereich 475.

AMS 1991 Subject Classification: 62H12.

1 Introduction

We introduced the Pitman-closeness criterion for the evaluation of multivariate forecasts (see Wenzel, 1998b) and derived optimal weight matrices for the combination of forecasts. There we used real valued square matrices as weights, which is the most general case and is called ***Strong** Pitman-closeness*. Now we also analyse the situation where the weight matrices are assumed to be diagonal as well as the case where the weights for the combination of the multivariate forecasts are restricted to be scalar. For this we define the notions of ***Medium** Pitman-closeness* and the ***Weak** Pitman-closeness*. The denotations ***Strong***, ***Medium*** and ***Weak** Pitman-closeness* are close to those of the covariance adjustment techniques presented by Troschke (1999a). Finally we calculate the different combinations of forecasts for a German macro economic data set.

2 The Pitman-closeness criterion

First we give a description of the problem as in Wenzel (1998b).

Assume that

$\mathbf{Y} := (Y_1, \dots, Y_k)'$ is a random vector to be forecasted ($k \geq 2$),

$\mathbf{F}_i := (F_{i1}, \dots, F_{ik})'$ are unbiased multivariate forecasts ($i=1, \dots, n$) for \mathbf{Y} and

$\mathbf{u}_i := (Y_1 - F_{i1}, \dots, Y_k - F_{ik})'$ is the error vector of the i -th forecast method,

where $\mathbf{u} := (\mathbf{u}_1', \dots, \mathbf{u}_n')' \sim N_{n \cdot k}(\mathbf{0}, \Sigma)$, Σ p.d., and there exists a vector \mathbf{u}_i , without loss of generality $\mathbf{u}_1 = \mathbf{u}_n$, so that $\text{Cov}[(\mathbf{u}_1 - \mathbf{u}_n)', \dots, (\mathbf{u}_{n-1} - \mathbf{u}_n)']$ is p.d.

Furthermore we are again giving the definition of the component-by-component Pitman-closeness.

Definition 1. The forecast \mathbf{F}_1 is component-by-component Pitman-closer to a random vector \mathbf{Y} than the forecast \mathbf{F}_2 ($\mathbf{F}_1 \neq \mathbf{F}_2$) if and only if

$$P(|Y_j - F_{j1}| < |Y_j - F_{j2}|) > 0.5 \quad \forall j \in \{1, \dots, k\}, \text{ where } F_{j1} \neq F_{j2}.$$

The probability statement of *Definition 1* is equivalent to

$$P(|u_{j1}| < |u_{j2}|) > 0.5 \quad \forall j \in \{1, \dots, k\}, \text{ where } F_{j1} \neq F_{j2}.$$

In the following we specify the calculation procedure of optimal weights in three categories.

2.1 Strong Pitman-closeness

Here, the multivariate combination of forecasts are given as

$$\mathbf{F}_A = \sum_{i=1}^n \mathbf{A}_i \mathbf{F}_i \text{ and } \mathbf{F}_B = \sum_{i=1}^n \mathbf{B}_i \mathbf{F}_i, \text{ where}$$

$$\mathbf{A}_i := \begin{pmatrix} a_{11}^{(i)} & \cdots & a_{1k}^{(i)} \\ \vdots & \ddots & \vdots \\ a_{k1}^{(i)} & \cdots & a_{kk}^{(i)} \end{pmatrix} \in \mathbb{R}^{k \times k} \text{ and } \mathbf{B}_i := \begin{pmatrix} b_{11}^{(i)} & \cdots & b_{1k}^{(i)} \\ \vdots & \ddots & \vdots \\ b_{k1}^{(i)} & \cdots & b_{kk}^{(i)} \end{pmatrix} \in \mathbb{R}^{k \times k} \text{ are matrices of weights,}$$

$$\sum_{i=1}^n \mathbf{A}_i = \mathbf{I}_k, \sum_{i=1}^n \mathbf{B}_i = \mathbf{I}_k, \text{ and } \mathbf{I}_k \text{ denotes the identity matrix.}$$

Using the definitions of Wenzel (1998b), where

$$\Sigma := (\Sigma_{rs})_{r,s=1,\dots,n}, \quad \Sigma_{rs} := \text{Cov}(\mathbf{u}_r, \mathbf{u}_s),$$

$$\mathbf{V}_{rs} := \Sigma_{rs} + \Sigma_{nn} - \Sigma_{rn} - \Sigma_{ns}, \quad r, s = 1, \dots, n-1,$$

$$\mathbf{V} := (\mathbf{V}_{rs})_{r,s=1,\dots,n-1},$$

$$\mathbf{w}_{ji} = (\Sigma_{nn} - \Sigma_{in})\mathbf{e}_j \quad \text{and therefore} \quad \mathbf{w}_{ji}' := \mathbf{e}_j' (\Sigma_{nn} - \Sigma_{ni}), \quad j=1,\dots,k, \quad i=1,\dots,n-1,$$

$$\mathbf{w}_j := \left(\mathbf{w}_{j1}', \dots, \mathbf{w}_{j,n-1}' \right)',$$

we get the following theorem.

Theorem 1. The **Strongly Pitman-closest combination** (component-by-component Pitman-closest combination) of n multivariate forecasts for a random vector \mathbf{Y} of dimension k ($k \geq 2$) is given by the matrix of weights:

$$\mathbf{A}^{\text{opt,str}} := [\mathbf{A}_1^{\text{opt,str}}, \dots, \mathbf{A}_n^{\text{opt,str}}] = [\mathbf{W}'\mathbf{V}^{-1}, \mathbf{I}_k - \mathbf{W}'\mathbf{V}^{-1}\mathbf{I}_k^*]$$

where $\mathbf{W} := (\mathbf{w}_1, \dots, \mathbf{w}_k) \sim (n-1) \cdot k \times k$,

$$\mathbf{I}_k^* := [\mathbf{I}_k, \dots, \mathbf{I}_k]' \sim (n-1) \cdot k \times k.$$

Proof: See Wenzel (1998b).

2.2 Medium Pitman-closeness

Here we use the restriction of diagonal matrices of weights, so

$$\mathbf{A}_i := \begin{pmatrix} a_{11}^{(i)} & & 0 \\ & \ddots & \\ 0 & & a_{kk}^{(i)} \end{pmatrix} \in \mathbb{R}^{k \times k} \quad \text{and} \quad \mathbf{B}_i := \begin{pmatrix} b_{11}^{(i)} & & 0 \\ & \ddots & \\ 0 & & b_{kk}^{(i)} \end{pmatrix} \in \mathbb{R}^{k \times k} \quad \text{are the matrices of weights,}$$

$$i=1,\dots,n, \quad \text{and} \quad \sum_{i=1}^n \mathbf{A}_i = \mathbf{I}_k, \quad \sum_{i=1}^n \mathbf{B}_i = \mathbf{I}_k.$$

Using again the component-by-component Pitman-closeness definition we get the next theorem.

Theorem 2. The **Medium Pitman-closest combination** (where the matrices of weights are restricted to be diagonal matrices) of n multivariate forecasts for a random vector \mathbf{Y} of dimension k ($k \geq 2$) is given by the matrices of weights:

$$\mathbf{A}_i^{\text{opt,med}} = \begin{pmatrix} a_{11}^{(i),\text{opt,med}} & & 0 \\ & \ddots & \\ 0 & & a_{kk}^{(i),\text{opt,med}} \end{pmatrix}, \quad i=1, \dots, n,$$

where $a_{jj}^{(i),\text{opt,med}}$ is the i -th component of the vector $\frac{\Sigma_j^{-1} \mathbf{1}}{\mathbf{1}' \Sigma_j^{-1} \mathbf{1}}$ ($j=1, \dots, k$) and Σ_j denotes the covariance matrix of the individual forecast errors of the j -th component and $\mathbf{1} = (1, \dots, 1)'$ is a vector of length n .

Proof: The proof is straightforward. Calculating the optimal weights for the j -th component is equivalent to calculating the optimal univariate combination for this component, which is described in Wenzel (1998a).

2.3 Weak Pitman-closeness

Here we concentrate on the combination of multivariate forecasts using scalar weights.

Thus a forecast combination is given by $\mathbf{F}_a = \sum_{i=1}^n a_i \mathbf{F}_i$, $a_i \in \mathbf{R}$, $\sum_{i=1}^n a_i = 1$. Obvious we cannot use the definition of component-by-component Pitman-closeness to calculate optimal weights for the combination. Therefore we define the **Weak Pitman-closeness**.

Definition 2. The forecast \mathbf{F}_1 is weakly Pitman-closer to the random vector \mathbf{Y} than the forecast \mathbf{F}_2 ($\mathbf{F}_1 \neq \mathbf{F}_2$) if and only if

$$P\left(\left|\sum_{j=1}^k Y_j - F_{j1}\right| < \left|\sum_{j=1}^k Y_j - F_{j2}\right|\right) > 0.5 \quad .$$

The probability statement of *Definition 2* is equivalent to

$$P\left(\left|\sum_{j=1}^k u_{j1}\right| < \left|\sum_{j=1}^k u_{j2}\right|\right) > 0.5 \quad .$$

With *Definition 2* in mind it is possible to calculate a **Weakly Pitman-closest** forecast.

Theorem 3. The **Weakly** Pitman-closest combination (where the weights are restricted to be scalar) of n multivariate forecasts for a random vector \mathbf{Y} of dimension k ($k \geq 2$) is given by the vector of weights:

$$\mathbf{a}^{\text{opt, weak}} = (a_1^{\text{opt, weak}}, \dots, a_n^{\text{opt, weak}})' = \frac{\Sigma_{\text{sum}}^{-1} \mathbf{1}}{\mathbf{1}' \Sigma_{\text{sum}}^{-1} \mathbf{1}}, \text{ where}$$

$$\Sigma_{\text{sum}} := \left(\tilde{\mathbf{1}}' \Sigma_{r,s} \tilde{\mathbf{1}} \right)_{r,s=1,\dots,n} \quad \text{and } \tilde{\mathbf{1}} \text{ denotes the vector } (1, \dots, 1)' \text{ of length } k.$$

Proof: If we compare two forecast combinations we have to find the vector \mathbf{a} so that for any

vector \mathbf{b} ($\mathbf{a} \neq \mathbf{b}$ and $\sum_{i=1}^n b_i = 1$) the inequality

$$\begin{aligned} & P \left(\left| \sum_{j=1}^k \sum_{i=1}^n a_i u_{ji} \right| < \left| \sum_{j=1}^k \sum_{i=1}^n b_i u_{ji} \right| \right) > 0.5 \\ \Leftrightarrow & P \left(\left| \mathbf{a}' \begin{pmatrix} u_{11} + u_{21} + \dots + u_{k1} \\ \vdots \\ u_{1n} + u_{2n} + \dots + u_{kn} \end{pmatrix} \right| < \left| \mathbf{b}' \begin{pmatrix} u_{11} + u_{21} + \dots + u_{k1} \\ \vdots \\ u_{1n} + u_{2n} + \dots + u_{kn} \end{pmatrix} \right| \right) > 0.5 \quad \text{holds.} \end{aligned}$$

The covariance matrix of the vector with the sums of the u_{ji} 's is Σ_{sum} and therefore, using the conclusions as in Wenzel (1998a) we get the optimal weight vector given in the statement above.

2.4 Remarks

Consulting the paper of Troschke (1999a) we can see that the **Strongly** and **Medium** Pitman-closest combinations are equivalent to the combinations resulting by the corresponding covariance adjustment techniques. Therefore we use the same denotations, which are **Strong** Pitman-closeness and **Medium** Pitman-closeness. The **Weakly** Pitman-closest combination differs from the result of the weak covariance adjustment technique because of the problem of defining a closeness criterion for this case. We have to remark that **Weak** Pitman-closeness is a criterion using the absolute of the sum of errors, which is not really reasonable for the comparison of forecasts. Obviously, the **Strongly** Pitman-closest combination is component-by-component Pitman-closer to \mathbf{Y} than the **Medium** Pitman-closest combination and the **Weakly** Pitman-closest combination and likewise the

Medium Pitman-closest combination is component-by-component Pitman-closer to \mathbf{Y} than the **Weakly** Pitman-closest combination.

3 Analysing German economic data

Klapper (1998) investigated German macro economic data and analysed different univariate combination techniques. In another article he also used a multivariate rank approach for the combination of the individual forecasts (Klapper, 1999). We derived the Pitman-closest combinations for this problem where we assumed that the errors are normally distributed with zero mean. Using only a ten years history for estimating the covariance matrix (by the usual ML-estimator) we have the problem of a singular matrix \mathbf{V} . In the calculation of the **Strongly** Pitman-closest combination we therefore use the Moore-Penrose-Inverse \mathbf{V}^+ instead of \mathbf{V}^{-1} but then the result depends on which of the individual forecasts is defined as \mathbf{F}_n . The order of the other forecasts does not influence the result. A comprehensive discussion of this can be found in Troschke (1999b).

In the following table we present the Root Mean Square Errors (RMSE) and the Mean Absolute Deviations (MAD) of the special combinations relative to the RMSE and the MAD of the simple average of the individual forecasts.

Table 1: RMSE and *MAD* of the special combinations relative to the RMSE and *MAD* of the simple average forecast.

	GDP	Priv. Cons.	Publ. Cons.	Export	Import	Cons. Prices
SPC _{OECD}	1.37 <i>1.37</i>	1.54 <i>1.70</i>	1.11 <i>1.02</i>	1.64 <i>1.60</i>	1.13 0.93	2.50 <i>2.16</i>
SPC _{WSI}	1.37 <i>1.27</i>	1.16 <i>1.16</i>	1.43 <i>1.18</i>	1.37 <i>1.31</i>	0.85 0.83	2.72 <i>2.32</i>
SPC _{SVR}	1.41 <i>1.44</i>	1.27 <i>1.35</i>	1.14 <i>1.13</i>	1.36 <i>1.38</i>	0.97 <i>1.00</i>	2.43 <i>2.29</i>
SPC _{AWF}	1.19 <i>1.15</i>	1.36 <i>1.30</i>	1.36 <i>1.12</i>	1.38 <i>1.23</i>	0.96 0.92	2.62 <i>2.49</i>
SPC _{IFW}	0.79 0.79	1.40 <i>1.47</i>	1.30 <i>1.23</i>	1.04 <i>1.01</i>	0.72 0.70	2.52 <i>2.26</i>
SPC _{IFO}	1.26 <i>1.14</i>	1.01 0.99	1.00 0.90	1.36 <i>1.28</i>	0.98 0.91	2.86 <i>2.92</i>
SPC _{DIW}	1.61 <i>1.61</i>	1.55 <i>1.84</i>	1.34 <i>1.20</i>	1.49 <i>1.46</i>	0.99 0.96	3.24 <i>3.29</i>
MPC	1.83 <i>1.47</i>	1.75 <i>1.92</i>	2.51 <i>1.87</i>	2.32 <i>1.95</i>	1.45 <i>1.31</i>	1.80 <i>1.68</i>
WPC	2.71 <i>2.00</i>	2.69 <i>2.61</i>	4.35 <i>3.02</i>	2.19 <i>1.93</i>	1.54 <i>1.39</i>	4.42 <i>3.92</i>

SPC_{name}: **Strongly** Pitman-closest combination with \mathbf{F}_n =name
MPC: **Medium** Pitman-closest combination
WPC: **Weakly** Pitman-closest combination

At first we can see that only in 7 of the 54 cases a Pitman-closeness combination performs better than the simple average (in the sense of the RMSE; 9 of 54 in the comparison of the MAD's). Therefore we can say that for this special data the multivariate combinations analysed here do not perform very well.

Furthermore the ***Strongly Pitman-closest combinations*** always outperform the ***Weakly Pitman-closest combination***. Except for the variable Consumer Prices, where they perform worse in all cases, they always dominates the ***Medium Pitman-closest combination*** (there is only one further case where a ***Strongly Pitman-closest combination*** has a larger MAD). This indicates that one should use multivariate combination techniques instead of univariate combinations of forecasts. At last, we see that the ***Medium Pitman-closest combination*** is always better than the ***Weakly Pitman-closest combination***.

5 Conclusions

We defined the Pitman-closeness in three categories, called ***Strong***, ***Medium*** and ***Weak Pitman-closeness***. We compared the quality of these techniques by using German economic data. The theoretical conclusions that the ***Strongly Pitman-closest combination*** performs best and the ***Weakly Pitman-closest combination*** performs worst are underlined by this example. Finally, we have to remark that in some cases, especially for the variable *Import*, the ***Strongly Pitman-closest combinations*** performs very well in comparison to the simple average forecast, whereas for the variables *Consumer Prices* and *Export* the simple average outperforms all other combination techniques. Consequently further research is needed in which situations *Pitman-closeness* strategies are appropriate and when one should use simple combination techniques like e.g. the arithmetic mean.

6 References

- [1] **Klapper, M.** (1998): Combining German Macro Economic Forecasts using Rank-Based Techniques, *Technical Report 19/1998, University of Dortmund*.
- [2] **Klapper, M.** (1999): Multivariate Rank-Based Forecast Combining Techniques, *Technical Report 2/1999, University of Dortmund*.
- [3] **Troschke, S. O.** (1999a): A Simulation Study To Compare Various Covariance Adjustment Techniques, *Technical Report, University of Dortmund, to appear*.
- [4] **Troschke, S. O.** (1999b): An Extensive Analysis of German Macroeconomic Forecast Data Using Covariance Adjustment Techniques, *Technical Report, University of Dortmund, to appear*.

- [5] **Wenzel, T.** (1998a): Pitman-closeness as a measure to evaluate the quality of forecasts, *Technical Report 15/1998, University of Dortmund*.
- [6] **Wenzel, T.** (1998b): Pitman-closeness and the linear combination of multivariate forecasts, *Technical Report 34/1998, University of Dortmund*.