

Hering, Franz

Working Paper

An algorithm for constructing Hadamard matrices

Technical Report, No. 2000,61

Provided in Cooperation with:

Collaborative Research Center 'Reduction of Complexity in Multivariate Data Structures' (SFB 475), University of Dortmund

Suggested Citation: Hering, Franz (2000) : An algorithm for constructing Hadamard matrices, Technical Report, No. 2000,61, Universität Dortmund, Sonderforschungsbereich 475 - Komplexitätsreduktion in Multivariaten Datenstrukturen, Dortmund

This Version is available at:

<https://hdl.handle.net/10419/77314>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.

An Algorithm for Constructing Hadamard Matrices

Franz Hering
Department of Statistics,
University of Dortmund,
44221 Dortmund, Germany
e-mail hering@statistik.uni-dortmund.de

Abstract

We define a new class of Hadamard matrices and present some examples obtained by an algorithm using a computer program.

Introduction

A Hadamard matrix \mathbf{H}_n is an orthogonal $(n \times n)$ -matrix in which all elements are either +1 or -1. If $n > 2$ then $n \equiv 0 \pmod{4}$ is a necessary condition that a \mathbf{H}_n exists. In this paper we define a class of Hadamard matrices and give a list of examples obtained by a computer program.

Hadamard matrices play an important in various field of statistics and mathematics. E.g. in statistics: incomplete balanced block designs, orthogonal arrays, factorial designs, Taguchi methods and Hotellings weighing problem; e.g. in mathematics: geometry, combinatorics, approximation theory and coding theory. There exists a wide-spread literature on Hadamard matrices, especially on the existence problem. A survey may be found in various text books of combinatorial theory.

1 Hadamard Schemes

To $m \in \mathbb{N}$ (\mathbb{N} includes 0) we write

$$\begin{aligned}\mathbb{N}_m &= \{0, 1, \dots, m-1, m\}, \\ \mathbb{Z}_m &= \{-m, -(m-1), \dots, -1, 0, 1, \dots, m-1, m\}.\end{aligned}$$

To $u \in \mathbb{Z}$ let $\rho(u)$ denote the additive residual class of u modulo \mathbb{Z}_m . (So, to $u \in \mathbb{Z}$ there exists exactly one $z \in \mathbb{Z}$ with $\rho(u) = u + z(2m+1) \in \mathbb{Z}_m$.) Let

$$M := \{1, i, -1, -i\}.$$

Here i is the imaginary unit. M is a multiplicative cyclic group of order 4 generated by i . For the elements of M we write '+' instead of 1, '-' instead of -1 and ' j ' instead of $-i$. We use the following notation: Small boldface letters denote vectors, vectors in $M^{\mathbb{Z}_m}$ carry an asterix; matrices are denoted by boldface capital letters.

If $\hat{\mathbf{A}}$ is an $(n \times m)$ -matrix whose elements $A_{i,j}$ are $(p \times q)$ -matrices, then we call $\hat{\mathbf{A}}$ a *block-matrix*. Block matrices are marked by carrying a hat accent. k -fold iterated block matrices are also marked with a hat accent, (but with one hat only). If $\hat{\mathbf{A}}$ is an (iterated) block-matrix, then the matrix obtained from $\hat{\mathbf{A}}$ by stripping off all the block boundaries is called the *deblocked* matrix of $\hat{\mathbf{A}}$. The deblocked matrix of $\hat{\mathbf{A}}$ is denoted by deleting the hat accent. E.g. if $\hat{\mathbf{A}} = [A_{i,j}]_{i=1\dots n, j=1\dots m}$ is an $(n \times m)$ -block-matrix whose elements $\mathbf{A}_{i,j} = [a_{i,j;u,v}]_{u=1\dots p, v=1\dots q}$, $i = 1, \dots, n$, $j = 1, \dots, m$ are $(p \times q)$ -matrices, then

$$\mathbf{A} = [a_{(i-1)p+u, (j-1)q+v}]_{\substack{i=1, \dots, n, u=1, \dots, p, \\ j=1, \dots, m, v=1, \dots, q}}$$

is the deblocked matrix of $\hat{\mathbf{A}}$.

If $z = z_1 + iz_2 \in \mathbb{C}$ is a complex number then $\bar{z} = z_1 - iz_2$ is its conjugate.

The mapping $\sigma : M \times M \rightarrow \{-1, 0, 1\}$ is defined by

$$\sigma(u, v) := \begin{cases} u \cdot v & \text{if } u = v, \\ -u \cdot v & \text{if } u = -v, \\ 0 & \text{if } u \neq \pm v. \end{cases} \quad (1)$$

To $k \in \mathbb{Z}$, $\mathbf{a}^*, \mathbf{c}^* \in M^{\mathbb{Z}_m}$ we define

$$\chi_k(\mathbf{a}^*, \mathbf{c}^*) := \sum_{u \in \mathbb{Z}_m} \sigma(a_u, c_{\rho(u+k)}). \quad (2)$$

Then

$$\chi_0(\mathbf{a}^*, \mathbf{a}^*) = \sum_{u \in \mathbb{N}_m} \sigma(a_u, a_u) = 2m + 1, \quad (3)$$

$$\chi_k(\mathbf{a}^*, \mathbf{c}^*) = \chi_{\rho(k)}(\mathbf{a}^*, \mathbf{c}^*), \quad (4)$$

$$\chi_k(\mathbf{a}^*, \mathbf{c}^*) = \chi_{-k}(\mathbf{c}^*, \mathbf{a}^*). \quad (5)$$

So $\chi_k(\mathbf{a}^*, \mathbf{c}^*)$, $k \in \mathbb{Z}$ is completely determined by the indices $k = 1, \dots, m$. To $\mathbf{a}^*, \mathbf{c}^*$ we define the χ -vector of $\mathbf{a}^*, \mathbf{c}^*$:

$$\chi(\mathbf{a}^*, \mathbf{c}^*) := [\chi_k(\mathbf{a}^*, \mathbf{c}^*)]_{k \in \mathbb{N}_m}.$$

Definition 1.

- a) $\mathbf{a}^* \in M^{\mathbb{Z}_m}$ is symmetric, if $a_u = a_{-u}$ for every $u \in \mathbb{N}_m$.
- b) If $\mathbf{a} \in M^{\mathbb{N}_m}$ then

$$\epsilon(\mathbf{a}) := [a_m, a_{m-1}, \dots, a_{-1}, a_0, a_1, \dots, a_{m-1}, a_m]$$

is the symmetric extension of \mathbf{a} .

- c) If $\mathbf{a}, \mathbf{b} \in M^{\mathbb{N}_m}$ then

$$\chi(\mathbf{a}, \mathbf{b}) := \chi(\epsilon(\mathbf{a}), \epsilon(\mathbf{b})).$$

■

Definition 2. Suppose $\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d} \in M^{\mathbb{N}_m}$.

- a) $[\mathbf{a}, \mathbf{b}]$ is a Hadamard pair (of size m), if

$$\chi(\mathbf{a}, \mathbf{a}) + \chi(\mathbf{b}, \mathbf{b}) = [1, 0, \dots, 0]. \quad (6)$$

b) Suppose that $[\mathbf{a}, \mathbf{b}], [\mathbf{c}, \mathbf{d}]$ are Hadamard pairs.

$$\begin{bmatrix} \mathbf{a} & \mathbf{b} \\ \mathbf{c} & \mathbf{d} \end{bmatrix} \quad (7)$$

is a *Hadamard scheme (of size m)*, if

$$\chi(\mathbf{a}, \mathbf{c}) + \chi(\mathbf{b}, \mathbf{d}) = \mathbf{0}. \quad \blacksquare$$

HP(m) symbolizes a Hadamard pair, **HS**(m) a Hadamard scheme (of size m) and **HM**(n) an $(n \times n)$ -Hadamard matrix.

Definition 3. If $k \in \mathbb{Z}, \mathbf{x}^*, \mathbf{y}^* \in M^{\mathbb{Z}_m}, d \in M$ then

$$I(k, \mathbf{x}^*, \mathbf{y}^*; d) := \{u \in \mathbb{Z}_m : \sigma(x_u, y_{\rho(u+k)}) = d\}.$$

If $k, \mathbf{x}^*, \mathbf{y}^*$ are fixed then $I(d) := I(k, \mathbf{x}^*, \mathbf{y}^*; d)$.

To $\mathbf{x}, \mathbf{y} \in M^{\mathbb{N}_m}, d \in M$ define $I(k, \mathbf{x}, \mathbf{y}; d) := I(k, \epsilon(\mathbf{x}), \epsilon(\mathbf{y}); d)$. \blacksquare

So for every fixed $k, \mathbf{x}^*, \mathbf{y}^*$

$$\mathbb{Z}_m = I(+) + I(i) + I(-) + I(j) \quad (8)$$

is a partition of \mathbb{Z}_m .

Lemma 1. Using the notation of definition 2 we obtain

a) $[\mathbf{a}, \mathbf{b}]$ is a **HP**(m) if and only if

$$\begin{aligned} \#I(k, \mathbf{a}, \mathbf{a}; +) + \#I(k, \mathbf{b}, \mathbf{b}; +) &= \#I(k, \mathbf{a}, \mathbf{a}; -) + \#I(k, \mathbf{b}, \mathbf{b}; -), \\ \#I(k, \mathbf{a}, \mathbf{a}; i) + \#I(k, \mathbf{b}, \mathbf{b}; i) &= \#I(k, \mathbf{a}, \mathbf{a}; j) + \#I(k, \mathbf{b}, \mathbf{b}; j) \end{aligned} \quad (9)$$

for $k = 1, \dots, m$.

b) $\begin{bmatrix} \mathbf{a} & \mathbf{b} \\ \mathbf{c} & \mathbf{d} \end{bmatrix}$ is a **HS**(m) if and only if $[\mathbf{a}, \mathbf{b}]$ and $[\mathbf{c}, \mathbf{d}]$ are **HP**(m)'s and

$$\begin{aligned} \#I(k, \mathbf{a}, \mathbf{c}; +) + \#I(k, \mathbf{b}, \mathbf{d}; +) &= \#I(k, \mathbf{a}, \mathbf{c}; -) + \#I(k, \mathbf{b}, \mathbf{d}; -), \\ \#I(k, \mathbf{a}, \mathbf{c}; i) + \#I(k, \mathbf{b}, \mathbf{d}; i) &= \#I(k, \mathbf{a}, \mathbf{c}; j) + \#I(k, \mathbf{b}, \mathbf{d}; j). \end{aligned} \quad (10)$$

for $k = 0, 1, \dots, m$.

Proof.

$$\chi_k(\mathbf{a}, \mathbf{a}) + \chi_k(\mathbf{b}, \mathbf{b}) = 0$$

\iff

$$\sum_{u \in \mathbb{Z}_m} \sigma(a_u, a_{\rho(u+k)}) + \sigma(b_u, b_{\rho(u+k)}) = 0$$

\iff

$$\sum_{d \in M} \left(\sum_{u \in I(k, \mathbf{a}, \mathbf{a}; d)} \sigma(a_u, a_{\rho(u+k)}) + \sum_{u \in I(k, \mathbf{b}, \mathbf{b}; d)} \sigma(b_u, b_{\rho(u+k)}) \right) = 0$$

\iff

$$\begin{aligned} & (+) \cdot \#I(k, \mathbf{a}, \mathbf{a}; +) + (-) \cdot \#I(k, \mathbf{a}, \mathbf{a}; -) \\ & + (i) \cdot \#I(k, \mathbf{a}, \mathbf{a}; i) + (j) \cdot \#I(k, \mathbf{a}, \mathbf{a}; j) \\ & + (+) \cdot \#I(k, \mathbf{b}, \mathbf{b}; +) + (-) \cdot \#I(k, \mathbf{b}, \mathbf{b}; -) \\ & + (i) \cdot \#I(k, \mathbf{b}, \mathbf{b}; i) + (j) \cdot \#I(k, \mathbf{b}, \mathbf{b}; j) = 0 \end{aligned}$$

\iff

$$\begin{aligned} & (+) \cdot \#I(k, \mathbf{a}, \mathbf{a}; +) + (-) \cdot \#I(k, \mathbf{a}, \mathbf{a}; -) \\ & + (+) \cdot \#I(k, \mathbf{b}, \mathbf{b}; +) + (-) \cdot \#I(k, \mathbf{b}, \mathbf{b}; -) = 0 \end{aligned}$$

and

$$\begin{aligned} & + (i) \cdot \#I(k, \mathbf{a}, \mathbf{a}; i) + (j) \cdot \#I(k, \mathbf{a}, \mathbf{a}; j) \\ & + (i) \cdot \#I(k, \mathbf{b}, \mathbf{b}; i) + (j) \cdot \#I(k, \mathbf{b}, \mathbf{b}; j) = 0. \end{aligned}$$

b) Part *b* follows by a completely analogous calculation. ■

The following lemma gives a representation of $\chi(\mathbf{a})$, using indices in \mathbb{N}_m only.

Lemma 2. Suppose $\mathbf{a} \in M^{\mathbb{N}_m}$ and $k \in 1, \dots, m$. Then

$$\chi_k(\mathbf{a}) = 1 + 2 \cdot \sum_{u=1}^{\lceil \frac{k+1}{2} \rceil - 1} \sigma(a_{-(u-\lceil \frac{k+1}{2} \rceil)}, a_{u+\lceil \frac{k}{2} \rceil})$$

$$\begin{aligned}
& + 2 \cdot \sum_{u=\lceil \frac{k+1}{2} \rceil}^{m-\lceil \frac{k}{2} \rceil} \sigma(a_{u-\lceil \frac{k+1}{2} \rceil}, a_{u+\lceil \frac{k}{2} \rceil}) \\
& + 2 \cdot \sum_{u=m-\lceil \frac{k}{2} \rceil+1}^m \sigma(a_{u-\lceil \frac{k+1}{2} \rceil}, a_{-(u+\lceil \frac{k}{2} \rceil)+(2m+1)}).
\end{aligned}$$

Proof. The mapping $k \rightarrow \rho(2k)$ is bijectiv on \mathbb{Z}_m . We denote its inverse mapping by $\cdot//2$, i.e. $k \rightarrow k//2$ for every $k \in \mathbb{Z}_m$. Then

$$k//2 = \begin{cases} k/2 & : k \in \mathbb{N}_m, \quad k \text{ even}, \\ (k-1)/2 - m & : k \in \mathbb{N}_m, \quad k \text{ odd}. \end{cases} \quad (11)$$

It follows

$$\begin{aligned}
\chi_k(\mathbf{a}) &= \chi_k(\epsilon(\mathbf{a})) \\
&= \sum_{u=-m}^m \sigma(a_u, a_{\rho(u+k)}) \\
&= \sum_{u=-m}^m \sigma(a_u, a_{\rho(u+k//2+k//2)}) \\
&= \sum_{u=-m}^m \sigma(a_{\rho(u-k//2)}, a_{\rho(u+k//2)}) \\
&= \sigma(a_{-k//2}, a_{+k//2}) + 2 \cdot \sum_{u=1}^m \sigma(a_{\rho(u-|k//2|)}, a_{\rho(u+|k//2|)}) \\
&= 1 + 2 \cdot \sum_{u=1}^m \sigma(a_{|\rho(u-|k//2|)|}, a_{|\rho(u+|k//2|)|}). \quad (12)
\end{aligned}$$

If k is even, then $\rho(k//2) > 0$, if k odd, then $\rho(k//2) < 0$. Using this and (11) we determine $|\rho(u - |k//2|)|$ and $|\rho(u + |k//2|)|$.

$$|\rho(u - |k//2|)| = \begin{cases} -(u - k/2) & k \text{ even}, \quad u \leq k/2 - 1, \\ u - k/2 & k \text{ even}, \quad u \geq k/2, \\ -(u - (m - (k-1)/2)) & k \text{ odd}, \quad u \leq m - (k-1)/2, \\ u - (m - (k-1)/2) & k \text{ odd}, \quad u \geq m - (k+1)/2; \end{cases}$$

$$|\rho(u + |k//2|)| = \begin{cases} u + k/2 & k \text{ even, } u \leq m - k/2, \\ -(u + k/2) + (2m + 1) & k \text{ even, } u \geq m - k/2 + 1, \\ u + m - (k - 1)/2 & k \text{ odd, } u \leq (k - 1)/2, \\ -u + m + (k + 1)/2 & k \text{ odd, } u \geq (k + 1)/2. \end{cases}$$

Inserting these values (7) and using Gauss-brackets leads to the equation of the lemma. \blacksquare

To $\mathbf{a} \in M^{\mathbb{N}_m}$, $k \in \{1, \dots, m\}$ define

$$\psi_k(\mathbf{a}) := \frac{1}{2}(\chi_k(a) - 1). \quad (13)$$

and $\psi(\mathbf{a}) := [\psi_k(\mathbf{a})]_{k=1, \dots, m}$. Then we obtain the following version of Lemma 1:

Lemma 3. Suppose $\mathbf{a} \in M^{\mathbb{N}_m}$ and $k \in 1, \dots, m$. Then

$$\psi_k(\mathbf{a}) = \sum_{\substack{u+v=k \\ 1 \leq u < v}} \sigma(a_u, a_v) + \sum_{u=0}^{m-k} \sigma(a_u, a_{u+k}) + \sum_{\substack{u+v=2m+1-k \\ u < v \leq m}} \sigma(a_u, a_v). \quad \blacksquare$$

From (6) follows

Lemma 4. $[\mathbf{a}, \mathbf{b}]$ is a $\mathbf{HP}(m)$ if and only if

$$\psi(\mathbf{a}) + \psi(\mathbf{b}) := -\mathbf{1} \quad (14)$$

(where $-\mathbf{1} = -[1, \dots, 1]$). \blacksquare

To obtain a Hadamard scheme one needs two Hadamard pairs $[\mathbf{a}, \mathbf{b}], [\mathbf{c}, \mathbf{d}]$ satisfying $\chi(\mathbf{a}, \mathbf{c}) + \chi(\mathbf{b}, \mathbf{d}) = \mathbf{0}$. The following lemma is a crucial support for the construction of Hadamard schemes, for it delivers to a given Hadamard pair a second one so that they together form a Hadamard scheme.

Lemma 5. If $[\mathbf{a}, \mathbf{b}]$ is a $\mathbf{HP}(m)$ then

$$\left[\begin{array}{cc} \mathbf{a}, & \mathbf{b} \\ \overline{\mathbf{b}}, & -\overline{\mathbf{a}} \end{array} \right]$$

is a **HS**(m).

Proof. a) We show first, that $[\bar{\mathbf{b}}, -\bar{\mathbf{a}}]$ is a Hadamard pair:

$$\begin{aligned}
 & \chi_k(\bar{\mathbf{b}}, \bar{\mathbf{b}}) + \chi_k((-\bar{\mathbf{a}}), (-\bar{\mathbf{a}})) \\
 &= \sum_{u \in \mathbb{Z}_m} (\sigma(\bar{b}_u, \bar{b}_{\rho(u+k)}) + \sigma(-\bar{a}_u, -\bar{a}_{\rho(u+k)})) \\
 &= \overline{\sum_{u \in \mathbb{Z}_m} (\sigma(b_u, b_{\rho(u+k)} + a_u, a_{\rho(u+k)}))} \\
 &= \overline{\chi_k(\sigma(\mathbf{b}, \mathbf{b})) + \chi_k(\sigma(\mathbf{a}, \mathbf{a}))} \\
 &= \overline{0} = 0.
 \end{aligned}$$

b) Now we verify the condition b of definition 3.

$$\begin{aligned}
 & \chi_k(\mathbf{a}, \bar{\mathbf{b}}) + \chi_k(\mathbf{b}, -\bar{\mathbf{a}}) \\
 &= \sum_{u \in \mathbb{Z}_m} (\sigma(a_u, \bar{b}_{\rho(u+k)}) + \sigma(b_u, -\bar{a}_{\rho(u+k)})) \\
 &= \sum_{u \in \mathbb{Z}_m} (\sigma(a_u, b_{\rho(u+k)}) - \sigma(b_u, a_{\rho(u+k)})) \\
 &= \sum_{u \in \mathbb{Z}_m} \sigma(a_u, b_{\rho(u+k)}) - \sum_{u \in \mathbb{Z}_m} \sigma(b_{\rho(-u+k)}, a_{-u}) \\
 &= \sum_{u \in \mathbb{Z}_m} \sigma(a_u, b_{\rho(u+k)}) - \sum_{v \in \mathbb{Z}_m} \sigma(b_{\rho(v+k)}, a_v) \\
 &= 0.
 \end{aligned}$$

■

To obtain a **HS**(m) it is therefore sufficient to construct a **HP**(m).

It is sometimes desirable to confine the investigation of **HP**(m) to special normalized cases. To define them requires the following preparations: On M introduce the linear order relation $+ > i > - > j$. Then $>$ introduces canonically the lexicographical order relation $>$ on $M^{\mathbb{N}_m}$: $\mathbf{a} = [a_0, \dots, a_m] > \mathbf{b} = [b_0, \dots, b_m]$ if there exists $r \in \mathbb{N}_m$ with $a_r > b_r$ and $a_s = b_s$, $s = 0, \dots, r-1$.

$x \in \mathbb{N}_m$ is *i-leading*, if x has the following property: If $D := \{u \in \mathbb{N}_m : x_u \in \{i, j\}\} \neq \emptyset$, and \min is the now well defined mimimal index with

$x_{min} \in \{i, j\}$, then $x_{min} = i$.

Definition 4. $[\mathbf{a}, \mathbf{b}]$, $\mathbf{a}, \mathbf{b} \in M^{\mathbb{Z}_m}$ is a *normalized pair* if:

- a) $\mathbf{a} \geq \mathbf{b}$,
- b) $a_0 = b_0 = +$,
- c) \mathbf{a} and \mathbf{b} are i -leading.

■

The following lemma is trivial:

Lemma 6. To every pair $[\tilde{\mathbf{a}}, \tilde{\mathbf{b}}]$, $\tilde{\mathbf{a}} \leq \tilde{\mathbf{b}} \in \mathbb{Z}_m$ there exists a normalized pair $[\mathbf{a}, \mathbf{b}]$, $\mathbf{a}, \mathbf{b} \in \mathbb{Z}_m$ with $\chi(\mathbf{a}, \mathbf{a}) = \chi(\tilde{\mathbf{a}}, \tilde{\mathbf{a}})$. ■

2 The Construction of Hadamard Matrices

Define

$$\mathbf{H}_+ = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}, \quad \mathbf{H}_i = \begin{bmatrix} -1 & 1 \\ 1 & 1 \end{bmatrix}, \quad \mathbf{H}_- = -\mathbf{H}_+, \quad \mathbf{H}_j = -\mathbf{H}_i.$$

The matrices $\mathbf{H}_u, u \in M$ are symmetric HM(2). To

$$\mathbf{K} = [k_{p,q}]_{\substack{p=0, \dots, a-1 \\ q=0, \dots, b-1}}, \quad k_{p,q} \in M$$

define

$$\hat{\mathbf{H}}_{\mathbf{K}} := [\mathbf{H}_{k_{p,q}}]_{\substack{p=0, \dots, a-1 \\ q=0, \dots, b-1}}.$$

So the blocked matrix $\hat{\mathbf{H}}_{\mathbf{K}}$ is obtained by replacing each element $k_{p,q} \in M$ of \mathbf{K} by the corresponding (2×2) -matrix $\mathbf{H}_{k_{p,q}}$.

Definition 5. If $\mathbf{a}^* \in M^{\mathbb{Z}_m}$, then the $((2m+1) \times (2m+1))$ -matrix

$$\mathbf{Z}(\mathbf{a}^*) := [a_{\rho(j-i)}]_{i,j \in M^{\mathbb{Z}_m}}$$

is the *circulant matrix* of \mathbf{a}^* . If $\mathbf{a} \in M^{\mathbb{N}_m}$ then $\mathbf{Z}(\mathbf{a}) := \mathbf{Z}(\epsilon(\mathbf{a}))$. ■

Lemma 7. Suppose $\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d} \in M^{\mathbb{N}_m}$. If

$$\begin{bmatrix} \mathbf{a}, & \mathbf{b} \\ \mathbf{c}, & \mathbf{d} \end{bmatrix}$$

is a $\mathbf{HS}(m)$, then define the two-fold blocked (2×2) -matrix

$$\hat{\mathbf{H}} = \begin{bmatrix} \hat{\mathbf{H}}_{\mathbf{Z}(\mathbf{a})} & \hat{\mathbf{H}}_{\mathbf{Z}(\mathbf{b})} \\ \hat{\mathbf{H}}_{\mathbf{Z}(\mathbf{c})} & \hat{\mathbf{H}}_{\mathbf{Z}(\mathbf{d})} \end{bmatrix}.$$

Then the $(4(2m+1) \times 4(2m+1))$ deblocked matrix \mathbf{H} of $\hat{\mathbf{H}}$ is a $\mathbf{HM}(8m+4)$.

Proof. In a matrix $\mathbf{R} = [r_{i,j}]_{i=0, \dots, s-1, j=0, \dots, t-1}$ the submatrix consisting of the rows i_1, \dots, i_w , $0 \leq i_1 < \dots < i_w \leq s-1$ is denoted by $\mathbf{R}(i_1, \dots, i_w)$. If $0 \leq g \leq 4m$ is even, then for $\mathbf{x} \in \{\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}\}$ we obtain

$$\hat{\mathbf{H}}_{\mathbf{Z}(\mathbf{x})}(g, g+1) = [\mathbf{H}_{x_{\rho(-m-g)}}, \dots, \mathbf{H}_{x_{\rho(0-g)}}, \dots, \mathbf{H}_{x_{\rho(m-g)}}].$$

To $0 \leq p < 8m+4$ define $g(p) = 2 \cdot [p/2]$. Then $\mathbf{H}(p)$ is a row of the deblocked $(2 \times (8m+4))$ -submatrix $\mathbf{H}(g(p), g(p)+1)$ of

$$\begin{aligned} \hat{\mathbf{H}}(g(p), g(p)+1) &= \\ &[\mathbf{H}_{x_{\rho(-m-g(p))}}, \dots, \mathbf{H}_{x_{\rho(0-g(p))}}, \dots, \mathbf{H}_{x_{\rho(m-g(p))}}, \\ &\quad \mathbf{H}_{y_{\rho(-m-g(p))}}, \dots, \mathbf{H}_{y_{\rho(0-g(p))}}, \dots, \mathbf{H}_{y_{\rho(m-g(p))}}] \end{aligned}$$

where $[\mathbf{x}, \mathbf{y}] = [\mathbf{a}, \mathbf{b}]$, if $0 \leq p < 4m+2$ and $[\mathbf{x}, \mathbf{y}] = [\mathbf{c}, \mathbf{d}]$ if $4m+2 \leq p < 8m+4$.

Now suppose $0 \leq p < q < 8m+4$. We show that the rows number p and q of \mathbf{H} are orthogonal, thereby proving the theorem. If $g(p) = g(q)$, then p is even, $q = p+1$ and so $\mathbf{H}(g(p), g(p)+1)$ consists of these two rows. From

$$\mathbf{H}(g(p), g(p)+1) \cdot \mathbf{H}(g(p), g(p)+1)' = (8m+4) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

we trivially obtain their orthogonality.

Now if $g(p) \neq g(q)$, then the two $(2 \times (4m+2))$ -submatrices $\mathbf{H}(g(p), g(p)+1)$ and $\mathbf{H}(g(q), g(q)+1)$ of \mathbf{H} have no common row. Define

$$\mathbf{A}(p, q) = \mathbf{H}(g(p), g(p)+1) \cdot \mathbf{H}(g(q), g(q)+1)'$$

We show

$$\mathbf{A}(p, q) = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

thereby proving the lemma.

Define

$$[\mathbf{w}, \mathbf{x}] = \begin{cases} [\mathbf{a}, \mathbf{b}] & : 0 \leq p < 4m+2, \\ [\mathbf{c}, \mathbf{d}] & : 4m+2 \leq p < 8m+4, \end{cases}$$

and likewise

$$[\mathbf{y}, \mathbf{z}] = \begin{cases} [\mathbf{a}, \mathbf{b}] & : 0 \leq q < 4m+2, \\ [\mathbf{c}, \mathbf{d}] & : 4m+2 \leq q < 8m+4. \end{cases}$$

We use the partition (8) to obtain canonically a partition of $\mathbf{A}(p, q)$ as follows:

$$\begin{aligned} \mathbf{A}(p, q) &= \sum_{d \in M} \sum_{v \in I(k, \mathbf{w}, \mathbf{x}; d)} \mathbf{H}_{\mathbf{w}_v} \times \mathbf{H}'_{\mathbf{x}_{\rho(v+k)}} \\ &\quad + \sum_{d \in M} \sum_{v \in I(k, \mathbf{y}, \mathbf{z}; d)} \mathbf{H}_{\mathbf{y}_v} \times \mathbf{H}'_{\mathbf{z}_{\rho(v+k)}} \\ &= \sum_{v \in I(k, \mathbf{w}, \mathbf{x}; +)} \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} + \sum_{v \in I(k, \mathbf{y}, \mathbf{z}; +)} \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} \\ &\quad + \sum_{v \in I(k, \mathbf{w}, \mathbf{x}; -)} \begin{bmatrix} -2 & 0 \\ 0 & -2 \end{bmatrix} + \sum_{v \in I(k, \mathbf{y}, \mathbf{z}; -)} \begin{bmatrix} -2 & 0 \\ 0 & -2 \end{bmatrix} \\ &\quad + \sum_{v \in I(k, \mathbf{w}, \mathbf{x}; i)} \begin{bmatrix} 0 & 2 \\ -2 & 0 \end{bmatrix} + \sum_{v \in I(k, \mathbf{y}, \mathbf{z}; i)} \begin{bmatrix} 0 & 2 \\ -2 & 0 \end{bmatrix} \end{aligned}$$

$$\begin{aligned}
& + \sum_{v \in I(k, \mathbf{w}, \mathbf{x}; j)} \begin{bmatrix} 0 & -2 \\ 2 & 0 \end{bmatrix} + \sum_{v \in I(k, \mathbf{y}, \mathbf{z}; j)} \begin{bmatrix} 0 & -2 \\ 2 & 0 \end{bmatrix} \\
& = (\#I(k, \mathbf{w}, \mathbf{x}; +) + \#I(k, \mathbf{y}, \mathbf{z}; +)) \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} \\
& + (\#I(k, \mathbf{w}, \mathbf{x}; -) + \#I(k, \mathbf{y}, \mathbf{z}; -)) \begin{bmatrix} -2 & 0 \\ 0 & -2 \end{bmatrix} \\
& + (\#I(k, \mathbf{w}, \mathbf{x}; i) + \#I(k, \mathbf{y}, \mathbf{z}; i)) \begin{bmatrix} 0 & 2 \\ -2 & 0 \end{bmatrix} \\
& + (\#I(k, \mathbf{w}, \mathbf{x}; j) + \#I(k, \mathbf{y}, \mathbf{z}; j)) \begin{bmatrix} 0 & -2 \\ 2 & 0 \end{bmatrix} \\
& = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}.
\end{aligned}$$

where zeros in the last equation follow from lemma 1. \blacksquare

3 Examples

3.1 An Algorithmic Construction of Hadamard Matrices

If a $\mathbf{HP}(m) = [\mathbf{a}, \mathbf{b}]$ exists, a $\mathbf{HM}(8m + 4)$ can be obtained by the following steps:

Step 1. Build the $\mathbf{HS}(m)$

$$\mathbf{HS}(m) = \begin{bmatrix} \mathbf{a}, & \mathbf{b} \\ \overline{\mathbf{b}}, & -\overline{\mathbf{a}} \end{bmatrix}$$

Step 2. Build the cyclic extension

$$\mathbf{HS}^*(m) = \begin{bmatrix} \epsilon(\mathbf{a}), & \epsilon(\mathbf{b}) \\ \epsilon(\overline{\mathbf{b}}), & \epsilon(-\overline{\mathbf{a}}) \end{bmatrix}$$

Step 3. Build the cyclic $((2m + 1) \times (2m + 1))$ -matrices $\mathbf{Z}(\mathbf{a}), \mathbf{Z}(\mathbf{b}), \mathbf{Z}(\overline{\mathbf{b}}), \mathbf{Z}(-\overline{\mathbf{a}})$.

Step 4. Using the four cyclic $(2m + 1) \times (2m + 1)$ -block matrices $\hat{\mathbf{H}}_{\mathbf{Z}(\mathbf{a})}, \hat{\mathbf{H}}_{\mathbf{Z}(\mathbf{b})}, \hat{\mathbf{H}}_{\mathbf{Z}(\bar{\mathbf{b}})}, \hat{\mathbf{H}}_{\mathbf{Z}(-\bar{\mathbf{a}})}$, having as elements (blocks) (2×2) -matrices \mathbf{H}_u , $u \in M$ build the twofold blocked matrix

$$\hat{\mathbf{H}} = \begin{bmatrix} \hat{\mathbf{H}}_{\mathbf{Z}(\mathbf{a})}, & \hat{\mathbf{H}}_{\mathbf{Z}(\mathbf{b})} \\ \hat{\mathbf{H}}_{\mathbf{Z}(\bar{\mathbf{b}})}, & \hat{\mathbf{H}}_{\mathbf{Z}(-\bar{\mathbf{a}})} \end{bmatrix}$$

Step 5. Build the $((8m + 4) \times (8m + 4))$ -matrix deblocked $\{+1, -1\}$ matrix \mathbf{H} of $\hat{\mathbf{H}}$. Then \mathbf{H} is a $\mathbf{HM}(8m + 4)$. \blacksquare

3.2 Two Examples

We apply these steps to the sizes $m = 0$ and $m = 1$.

$$m = 0.$$

The only normalized pair is $[+]$, $[+]$, obtained from $\mathbf{a} = [a_0] = [+]$ and $\mathbf{b} = [b_0] = [+]$ is a $\mathbf{HP}(0)$.

Step 1. Obviously,

$$\begin{bmatrix} + & + \\ + & - \end{bmatrix},$$

is a $\mathbf{HS}(0)$, (which after deblocking accidentally is itself a $\mathbf{HM}(2)$.)

Step 2. Here $\epsilon(\mathbf{a}) = \mathbf{a}, \epsilon(\mathbf{b}) = \mathbf{b}$. So $\mathbf{HS}^*(0) = \mathbf{HS}(0)$.

Step 3. $\mathbf{Z}([\mathbf{a}]) = \mathbf{Z}([\mathbf{b}]) = \mathbf{Z}([\bar{\mathbf{b}}]) = \mathbf{Z}([+] = [+], \mathbf{Z}([- \bar{\mathbf{a}}]) = \mathbf{Z}([-]) = [-]$

Step 4.

$$\hat{\mathbf{H}} = \begin{bmatrix} \left[\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \right] & \left[\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \right] \\ \left[\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \right] & \left[\begin{bmatrix} -1 & -1 \\ -1 & 1 \end{bmatrix} \right] \end{bmatrix}$$

Step 5. Writing "+" instead of 1 and "-" instead of -1 we obtain the matrix

$$\mathbf{H} = \begin{bmatrix} + & + & + & + \\ + & - & + & - \\ + & + & - & - \\ + & - & - & + \end{bmatrix}$$

This is a $\mathbf{HM}(8 \cdot 0 + 4)$. ■

$$m = 1.$$

Define $\mathbf{a} = [+, -]$, $\mathbf{b} = [+, i]$. Then according to section 2 $[+, -], [+, i]$ is a normalized $\mathbf{HS}(1)$.

Following again steps 1 to 6 gives us a $\mathbf{HM}(12)$.

(Indeed, from the symmetric extensions $\epsilon(\mathbf{a}) = [-, +, -]$, $\epsilon(\mathbf{b}) = [i, +, i]$ of \mathbf{a} and \mathbf{b} we obtain $\chi_1(\mathbf{a}) = \chi_1(\epsilon(\mathbf{a})) = (-1) + 1 + (-1) = -1$, $\chi_1(\mathbf{b}) = \chi_1(\epsilon(\mathbf{b})) = 0 + 1 + 0 = 1$. This implies $\psi_1(a) + \psi_1(b) = \frac{1}{2}((\chi_k(a) - 1) + (\chi_k(b) - 1)) = -1$. So, as it be checked also easily directly $[[1, -], [1, i]]$ is a $\mathbf{HP}(1)$.)

Step 1. The correponding $\mathbf{HS}(1)$ is

$$\mathbf{HS}(1) = \left[\begin{bmatrix} + & - \\ + & j \end{bmatrix} \quad \begin{bmatrix} + & i \\ - & + \end{bmatrix} \right].$$

Step 2.

$$\mathbf{HS}^*(1) = \left[\begin{bmatrix} - & + & - \\ j & + & j \end{bmatrix} \quad \begin{bmatrix} i & + & i \\ + & - & + \end{bmatrix} \right].$$

Step 3.

$$\mathbf{Z}(\mathbf{a}) = \begin{bmatrix} - & + & - \\ - & - & + \\ + & - & - \end{bmatrix}, \quad \mathbf{Z}(\mathbf{b}) = \begin{bmatrix} i & + & i \\ i & i & + \\ + & i & i \end{bmatrix},$$

$$\mathbf{Z}(\bar{\mathbf{b}}) = \begin{bmatrix} j & + & j \\ j & j & + \\ + & j & j \end{bmatrix}, \quad \mathbf{Z}(-\bar{\mathbf{a}}) = \begin{bmatrix} + & - & + \\ + & + & - \\ - & + & + \end{bmatrix}.$$

Step 4.

$$\mathbf{H}_K = \left[\begin{array}{c|c|c|c|c|c} & \begin{bmatrix} - & - \\ - & + \\ - & - \\ - & + \\ + & + \\ + & - \\ + & - \\ - & - \\ + & - \\ + & + \\ + & - \end{bmatrix} & \begin{bmatrix} + & + \\ + & - \\ - & - \\ - & + \\ + & + \\ - & + \\ + & + \\ + & - \\ + & - \\ + & - \\ - & - \end{bmatrix} & \begin{bmatrix} - & - \\ - & + \\ + & + \\ + & - \\ - & + \\ - & - \\ + & - \\ + & + \\ + & - \\ + & + \\ - & + \end{bmatrix} & \begin{bmatrix} - & + \\ + & + \\ - & + \\ + & + \\ + & - \\ + & - \\ + & - \\ - & + \\ - & + \\ + & + \\ - & + \end{bmatrix} & \begin{bmatrix} + & + \\ + & - \\ - & + \\ + & + \\ + & - \\ - & + \\ - & - \\ + & + \\ + & - \\ + & + \\ + & - \end{bmatrix} & \begin{bmatrix} - & + \\ + & + \\ + & - \\ + & + \\ + & - \\ - & + \\ - & - \\ + & + \\ + & - \\ + & + \\ + & - \end{bmatrix} \\ \hline \end{array} \right]$$

Step 5.

$$\mathbf{H}_K = \left[\begin{array}{cccccccccccc} - & - & + & + & - & - & - & + & + & + & - & + \\ - & + & + & - & - & + & + & + & + & + & - & + \\ - & - & - & - & + & + & - & + & - & + & + & + \\ - & + & - & + & + & - & + & + & + & + & + & - \\ + & + & - & - & - & - & + & + & - & + & - & + \\ + & - & - & + & - & + & + & - & + & + & + & + \\ + & - & + & + & + & - & + & + & - & - & + & + \\ + & - & + & - & - & - & + & - & - & + & + & - \\ + & - & + & - & + & + & + & + & + & + & - & - \\ + & - & - & - & + & - & + & - & + & - & - & + \\ + & + & + & - & + & - & - & - & + & + & + & + \\ + & - & - & - & - & - & - & - & + & + & - & + \end{array} \right]$$

3.3 Tables of all normalized $\mathbf{HP}(m)$ of sizes m for $0 \leq m \leq 7$

In the following tables on the left entry the Hadamard-pairs are listed. The right entry contains the two corresponding Ψ -vectors. Every $\mathbf{HP}(m)$ generates a $\mathbf{HM}(8m + 4)$.

m=0

$$\begin{array}{|c|c|} \hline \left[\begin{array}{|c|} \hline \end{array} \right] & \left[\begin{array}{|c|} \hline \end{array} \right] \\ \hline \left[\begin{array}{|c|} \hline \end{array} \right] & \left[\begin{array}{|c|} \hline \end{array} \right] \\ \hline \end{array}$$

m=1

$$\begin{array}{|c|c|} \hline \left[\begin{array}{|c|c|} \hline i & \\ \hline - & \\ \hline \end{array} \right] & \left[\begin{array}{|c|c|} \hline 0 & \\ \hline -1 & \\ \hline \end{array} \right] \\ \hline \end{array}$$

m=2

$$\begin{array}{|c|c|} \hline \left[\begin{array}{|c|c|c|} \hline i & j & \\ \hline - & - & \\ \hline \end{array} \right] & \left[\begin{array}{|c|c|c|} \hline -1 & -1 & \\ \hline 0 & 0 & \\ \hline \end{array} \right] \\ \hline \left[\begin{array}{|c|c|c|} \hline - & i & \\ \hline i & - & \\ \hline \end{array} \right] & \left[\begin{array}{|c|c|c|} \hline -1 & 0 & \\ \hline 0 & -1 & \\ \hline \end{array} \right] \\ \hline \end{array}$$

m=3

$$\begin{array}{|c|c|c|} \hline \left[\begin{array}{|c|c|c|} \hline i & j & j \\ \hline + & - & + \\ \hline \end{array} \right] & \left[\begin{array}{|c|c|c|} \hline 0 & 0 & -2 \\ \hline -1 & -1 & 1 \\ \hline \end{array} \right] \\ \hline \left[\begin{array}{|c|c|c|} \hline i & i & j \\ \hline - & + & + \\ \hline \end{array} \right] & \left[\begin{array}{|c|c|c|} \hline 0 & -2 & 0 \\ \hline -1 & 1 & -1 \\ \hline \end{array} \right] \\ \hline \left[\begin{array}{|c|c|c|} \hline - & i & j \\ \hline i & - & - \\ \hline \end{array} \right] & \left[\begin{array}{|c|c|c|} \hline -2 & -1 & 0 \\ \hline 1 & 0 & -1 \\ \hline \end{array} \right] \\ \hline \left[\begin{array}{|c|c|c|} \hline + & i & j \\ \hline i & + & - \\ \hline \end{array} \right] & \left[\begin{array}{|c|c|c|} \hline 0 & -1 & 0 \\ \hline -1 & 0 & -1 \\ \hline \end{array} \right] \\ \hline \left[\begin{array}{|c|c|c|} \hline i & - & j \\ \hline - & i & - \\ \hline \end{array} \right] & \left[\begin{array}{|c|c|c|} \hline 0 & -2 & -1 \\ \hline -1 & 1 & 0 \\ \hline \end{array} \right] \\ \hline \left[\begin{array}{|c|c|c|} \hline i & + & j \\ \hline - & i & + \\ \hline \end{array} \right] & \left[\begin{array}{|c|c|c|} \hline 0 & 0 & -1 \\ \hline -1 & -1 & 0 \\ \hline \end{array} \right] \\ \hline \left[\begin{array}{|c|c|c|} \hline i & j & i \\ \hline + & + & - \\ \hline \end{array} \right] & \left[\begin{array}{|c|c|c|} \hline -2 & 0 & 0 \\ \hline 1 & -1 & -1 \\ \hline \end{array} \right] \\ \hline \left[\begin{array}{|c|c|c|} \hline - & - & i \\ \hline i & j & - \\ \hline \end{array} \right] & \left[\begin{array}{|c|c|c|} \hline 0 & -1 & 1 \\ \hline -1 & 0 & -2 \\ \hline \end{array} \right] \\ \hline \left[\begin{array}{|c|c|c|} \hline + & - & i \\ \hline i & j & + \\ \hline \end{array} \right] & \left[\begin{array}{|c|c|c|} \hline 0 & -1 & -1 \\ \hline -1 & 0 & 0 \\ \hline \end{array} \right] \\ \hline \end{array}$$

$$m = 4$$

$\begin{bmatrix} i & j & j & j \\ - & + & + & - \end{bmatrix}$	$\begin{bmatrix} 1 & 1 & -1 & -1 \\ -2 & -2 & 0 & 0 \end{bmatrix}$
$\begin{bmatrix} - & i & j & j \\ i & + & + & - \end{bmatrix}$	$\begin{bmatrix} -1 & 0 & -1 & -1 \\ 0 & -1 & 0 & 0 \end{bmatrix}$
$\begin{bmatrix} - & i & j & j \\ i & - & + & + \end{bmatrix}$	$\begin{bmatrix} -1 & 0 & -1 & -1 \\ 0 & -1 & 0 & 0 \end{bmatrix}$
$\begin{bmatrix} i & i & i & j \\ + & - & + & - \end{bmatrix}$	$\begin{bmatrix} 1 & -1 & -1 & 1 \\ -2 & 0 & 0 & -2 \end{bmatrix}$
$\begin{bmatrix} i & - & i & j \\ + & i & + & - \end{bmatrix}$	$\begin{bmatrix} -1 & -1 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{bmatrix}$
$\begin{bmatrix} i & - & i & j \\ - & i & + & + \end{bmatrix}$	$\begin{bmatrix} -1 & -1 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{bmatrix}$
$\begin{bmatrix} - & - & i & j \\ i & j & - & - \end{bmatrix}$	$\begin{bmatrix} -1 & -2 & 1 & 0 \\ 0 & 1 & -2 & -1 \end{bmatrix}$
$\begin{bmatrix} - & i & - & j \\ i & - & j & - \end{bmatrix}$	$\begin{bmatrix} -1 & 0 & -2 & 1 \\ 0 & -1 & 1 & -2 \end{bmatrix}$
$\begin{bmatrix} i & - & - & j \\ - & i & j & - \end{bmatrix}$	$\begin{bmatrix} 1 & -1 & -2 & 0 \\ -2 & 0 & 1 & -1 \end{bmatrix}$
$\begin{bmatrix} i & j & + & j \\ - & - & i & + \end{bmatrix}$	$\begin{bmatrix} -1 & 1 & 0 & -1 \\ 0 & -2 & -1 & 0 \end{bmatrix}$
$\begin{bmatrix} i & i & + & j \\ - & + & i & - \end{bmatrix}$	$\begin{bmatrix} 1 & -1 & 0 & -1 \\ -2 & 0 & -1 & 0 \end{bmatrix}$
$\begin{bmatrix} - & i & + & j \\ i & + & i & - \end{bmatrix}$	$\begin{bmatrix} -1 & -2 & 0 & -1 \\ 0 & 1 & -1 & 0 \end{bmatrix}$
$\begin{bmatrix} i & j & i & i \\ - & - & + & + \end{bmatrix}$	$\begin{bmatrix} -1 & 1 & -1 & 1 \\ 0 & -2 & 0 & -2 \end{bmatrix}$
$\begin{bmatrix} + & - & i & i \\ i & j & + & - \end{bmatrix}$	$\begin{bmatrix} 1 & 0 & -1 & 0 \\ -2 & -1 & 0 & -1 \end{bmatrix}$
$\begin{bmatrix} i & j & + & i \\ + & - & i & - \end{bmatrix}$	$\begin{bmatrix} -1 & -1 & 0 & 1 \\ 0 & 0 & -1 & -2 \end{bmatrix}$
$\begin{bmatrix} + & - & + & i \\ i & j & j & - \end{bmatrix}$	$\begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & -1 & -1 & -1 \end{bmatrix}$
$\begin{bmatrix} - & + & + & i \\ i & j & j & - \end{bmatrix}$	$\begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & -1 & -1 & -1 \end{bmatrix}$

m=5

$\begin{bmatrix} + & - & i & j & j \end{bmatrix}$	$\begin{bmatrix} 0 & -1 & -2 & -1 & 0 \end{bmatrix}$
$\begin{bmatrix} i & i & - & + & - \end{bmatrix}$	$\begin{bmatrix} -1 & 0 & 1 & 0 & -1 \end{bmatrix}$
$\begin{bmatrix} - & i & - & j & j \end{bmatrix}$	$\begin{bmatrix} 0 & 1 & -2 & 0 & -1 \end{bmatrix}$
$\begin{bmatrix} i & + & j & - & + \end{bmatrix}$	$\begin{bmatrix} -1 & -2 & 1 & -1 & 0 \end{bmatrix}$
$\begin{bmatrix} - & + & i & i & j \end{bmatrix}$	$\begin{bmatrix} -2 & -1 & -2 & 1 & 0 \end{bmatrix}$
$\begin{bmatrix} i & i & + & - & - \end{bmatrix}$	$\begin{bmatrix} 1 & 0 & 1 & -2 & -1 \end{bmatrix}$
$\begin{bmatrix} + & - & + & i & j \end{bmatrix}$	$\begin{bmatrix} -2 & -1 & 0 & 1 & -1 \end{bmatrix}$
$\begin{bmatrix} i & j & j & - & - \end{bmatrix}$	$\begin{bmatrix} 1 & 0 & -1 & -2 & 0 \end{bmatrix}$
$\begin{bmatrix} i & j & - & + & j \end{bmatrix}$	$\begin{bmatrix} -2 & 0 & -1 & 0 & -1 \end{bmatrix}$
$\begin{bmatrix} - & - & i & i & + \end{bmatrix}$	$\begin{bmatrix} 1 & -1 & 0 & -1 & 0 \end{bmatrix}$
$\begin{bmatrix} - & i & j & - & i \end{bmatrix}$	$\begin{bmatrix} -2 & -1 & 1 & 0 & 0 \end{bmatrix}$
$\begin{bmatrix} i & + & + & j & - \end{bmatrix}$	$\begin{bmatrix} 1 & 0 & -2 & -1 & -1 \end{bmatrix}$
$\begin{bmatrix} + & - & i & - & i \end{bmatrix}$	$\begin{bmatrix} 0 & 1 & -1 & -1 & 0 \end{bmatrix}$
$\begin{bmatrix} i & j & + & i & - \end{bmatrix}$	$\begin{bmatrix} -1 & -2 & 0 & 0 & -1 \end{bmatrix}$
$\begin{bmatrix} - & + & i & - & i \end{bmatrix}$	$\begin{bmatrix} -2 & 1 & 1 & -1 & 0 \end{bmatrix}$
$\begin{bmatrix} i & i & - & j & + \end{bmatrix}$	$\begin{bmatrix} 1 & -2 & -2 & 0 & -1 \end{bmatrix}$
$\begin{bmatrix} i & - & - & + & i \end{bmatrix}$	$\begin{bmatrix} 0 & -2 & -1 & 1 & 1 \end{bmatrix}$
$\begin{bmatrix} + & i & j & i & - \end{bmatrix}$	$\begin{bmatrix} -1 & 1 & 0 & -2 & -2 \end{bmatrix}$

m=6

$$m=7$$

m=7 (continued)

3.4 One normalized HP(8)

m=8							
$\begin{bmatrix} - & - & + & - & i & - & j & j \end{bmatrix}$				$\begin{bmatrix} -1 & 0 & 1 & -2 \\ 0 & -1 & -2 & 1 \\ -1 & 1 & -1 & -1 \\ 0 & -2 & 0 & 0 \end{bmatrix}$			
$\begin{bmatrix} i & j & i & i & + & j & - & - \end{bmatrix}$				$\begin{bmatrix} 0 & -1 & -1 & -1 \\ -1 & -1 & -2 & -2 \\ 1 & -1 & 1 & 1 \\ -2 & -2 & -2 & -2 \end{bmatrix}$			
$\begin{bmatrix} i & i & - & j & - & j & i & j \end{bmatrix}$				$\begin{bmatrix} -1 & 1 & -1 & -1 \\ 0 & 0 & 1 & 1 \\ -2 & -2 & -2 & -2 \end{bmatrix}$			
$\begin{bmatrix} - & - & i & - & i & + & + & - \end{bmatrix}$				$\begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix}$			

References

- Hadamard, J. (1893). Resolution d'une question relative aux determinants. *Bull. des Sciences Math.* (2), **17**, 240-246.
- Hering, F. {Wägeprobleme} Diplomarbeit Univ. Tübingen (1963) (unpublished).
- Heydayat, A.S., N.J.A. Sloane & J. Stufken *Orthogonal Arrays* (1999) Springer Verlag, New York.
- Sylvester, J.J.(1867). Thoughts on inverse orthogonal arrays, simultaneous sign successions, and tesselated pavements in two or more colours, with applications to Newton's rule, ornamental tile work, and the theory of numbers, *Phil. Mag.* (4), **34**, 461-475